



The Chemical Company

---

**Submission for  
Verification of AgBalance™ Analysis Under  
NSF Protocol P352, Part B**

**Corn Production in Iowa AgBalance™ Analysis  
Final Report - February 2013**



**Submitted by:**

BASF Corporation  
Product Stewardship  
100 Park Avenue, Florham Park, NJ, 07932

**Prepared by:**

Thomas Laginess, Senior Sustainability Specialist  
Martin Mascianica, Regional Project Leader  
Markus Frank, Global Sustainability Leader  
Fred Moore, Regional Marketing Manager

## Table of Contents

<b>1. Purpose and Intent of this Submission .....</b>	<b>3</b>
<b>2. Content of this Submission .....</b>	<b>3</b>
<b>3. BASF's AgBalance™ Methodology.....</b>	<b>3</b>
3.1. Overview .....	3
3.2. Preconditions .....	3
3.3. Work Flow .....	6
<b>4. Study Goals, Decision Criteria and Target Audience .....</b>	<b>6</b>
4.1. Study Goals.....	6
4.2. Design Criteria .....	7
4.3. Target Audience .....	7
<b>5. Customer Benefit, Alternatives, System Boundaries and Scenarios .....</b>	<b>8</b>
5.1. Customer Benefit.....	8
5.2. Alternatives.....	8
5.3. System Boundaries .....	9
5.4. Scenario Analyses .....	10
<b>6. Input Parameters and Assumptions .....</b>	<b>10</b>
6.1. Input Parameters.....	10
6.2. Cost Inputs.....	14
6.3. Social Inputs.....	14
<b>7. Data Sources.....</b>	<b>16</b>
7.1 Environmental.....	16
7.2 Amounts and Costs.....	16
<b>8. AgBalance™ Analysis Results and Discussion .....</b>	<b>18</b>
8.1. Environmental Impact Results .....	18
8.1.1. Primary energy consumption.....	18
8.1.2. Raw material consumption.....	18
8.1.3. Air Emissions .....	20
8.1.3.1. Global Warming Potential (GWP) .....	20
8.1.3.2. Photochemical ozone creation potential (smog).....	20
8.1.3.3. Ozone depletion potential (ODP).....	21
8.1.3.4. Acidification potential (AP) .....	21
8.1.4. Water emissions .....	23
8.1.5. Solid waste generation .....	23
8.1.6. Land Use.....	24
8.1.7. Eco-Toxicity potential .....	25
8.1.8. Risk potential.....	27
8.1.9. Consumptive Water Use .....	28
8.1.10. Biodiversity.....	29
8.1.11. Soil .....	29
8.1.12. Environmental Fingerprint.....	30
8.2. Economic Cost Fingerprint.....	31
8.3. Social Fingerprint.....	33
8.4. AgBalance™ Analysis Portfolio .....	35
8.5. Scenario Analyses .....	37
8.5.1 Scenario#1 - Reduction of yield in 2010 by 5% .....	37
8.5.2 Scenario#2 - Comparison of tillage differences in 2010 .....	39

- 8.5.3 Scenario#3 - Comparison of conservative tillage 2000 & 2010.....41
- 8.5.4 Scenario#4 – Yield for 2000 set at the same yield as 2010.....43
- 9. Data Quality Assessment.....45**
  - 9.1. Data Quality Statement.....45
- 10. Sensitivity and Uncertainty Analysis..... 46**
  - 10.1. Sensitivity and Uncertainty Considerations .....46
  - 10.2. Critical Uncertainties .....48
- 11. Limitations of AgBalance™ Study Results ..... 48**
  - 11.1. Limitations .....48
- 12. References ..... 49**
- Appendix A.....51**

## **1. Purpose and Intent of this Submission**

- 1.1. The purpose of this submission is to provide a written report of the methods and findings of BASF Corporation's "Corn Production in Iowa AgBalance™ Analysis", with the intent of having it verified under the requirements of NSF Protocol P352, Part B: Verification of AgBalance™ Analysis Studies.
- 1.2. The Corn Production AgBalance™ Analysis was performed by BASF according to the methodology validated by NSF International under the requirements of Protocol P352. More information on BASF's methodology and the NSF validation can be obtained at [http://www.nsf.org/info/eco\\_efficiency](http://www.nsf.org/info/eco_efficiency).

## **2. Content of this Submission**

- 2.1. This submission outlines the study goals, procedures, and results for the Corn Production AgBalance™ Analysis study, which was conducted in accordance with BASF Corporation's AgBalance™ methodology. This submission will provide a discussion of the basis of the eco-analysis preparation and verification work.
- 2.2. As required under NSF P352 Part B, along with this document, BASF is submitting the final computerized model programmed in Microsoft® Excel. The computerized model, together with this document, will aid in the final review and ensure that the data and critical review findings have been satisfactorily addressed.

## **3. BASF's AgBalance™ Methodology**

### *3.1. Overview:*

BASF AgBalance™ involves measuring the life cycle environmental impacts, life cycle costs and life cycle social aspects for product alternatives for a defined level of output. At a minimum, BASF AgBalance™ evaluates the environmental impact of the production, use, and disposal of a product or process in the areas of energy consumption, resource consumption, emissions, eco-toxicity, land use, water use, soil and biodiversity. The AgBalance™ analysis evaluates the life cycle costs associated with the product or process by calculating the costs related to, at a minimum, materials, labor, manufacturing, waste disposal, and energy. AgBalance™ also includes social aspects dealing with agricultural metrics, which are not part of the Protocol P352.

### *3.2. Preconditions:*

The basic preconditions of this AgBalance™ Analysis are that all alternatives that are being evaluated are being compared against a common functional unit or Customer Benefit (CB). This allows for an objective comparison between the various alternatives. The scoping and definition of the Customer Benefit are aligned with the goals and objectives of the study. Data gathering and constructing the system boundaries are consistent with the CB and consider the environmental, economic and social impacts of each alternative over their life cycle in order to achieve the specified CB. The social aspects in AgBalance™ are also based on the relevant



agricultural data for the specific CB. An overview of the scope of the environmental, economic and social assessments is defined below.

3.2.1. *Environmental Burden Metrics:*

For BASF AgBalance™ environmental burden is characterized using thirteen categories, at a minimum, including: primary energy consumption, raw material consumption, global warming potential (GWP), ozone depletion potential (ODP), acidification potential (AP), photochemical ozone creation potential (POCP), water emissions, solid waste emissions, eco-toxicity potential, land use, water use, soil and biodiversity. These are shown below in Figure 1. Metrics shown in yellow represent the eight main categories of environmental burden that are used to construct the environmental fingerprint, burdens in blue represent all elements of the emissions category, and burdens in green show the categories evaluated within air emissions.




Figure 1. Environmental Impact categories

3.2.2. *Economic Metrics:*

It is the intent of the BASF AgBalance™ methodology to assess the economics of products or processes over their life cycle and to determine an overall total cost of ownership for the defined customer benefit (\$/CB). The approaches for calculating costs vary from study to study. When chemical products of manufacturing are being compared, the sale price paid by the customer is predominately used. When different production methods are compared, the relevant costs include the purchase and installation of capital equipment, depreciation, and operating costs. The costs incurred are summed and combined in appropriate units (e.g. dollar or EURO)

without additional weighting of individual financial amounts. The BASF AgBalance™ methodology will incorporate:

- the real costs that occur in the process of creating and delivering the product to the consumer;
- the subsequent costs which may occur in the future (due to tax policy changes, for example) with appropriate consideration for the time value of money; and
- Costs having ecological aspect, such as the costs involved to treat wastewater generated during the manufacturing process.

In AgBalance™, costs are quantified for each alternative. These alternatives are then aggregated and totaled to show the total cost of each alternative as it relates to the common customer benefit (CB).

### 3.2.3. *Social Metrics:*

Social parameters are not addressed specifically in the ISO LCA standards. There are no other consensus standards that can be referenced to define the criteria for a social LCA. AgBalance™ represents BASF's best attempts to create a social LCA framework through the identification and use of relevant factors associated with life cycle principles. Even though there are no industry standards available, important developments from different groups like the UNEP/SETAC working group or existing standards in the Agro-sector like RISE were considered.

The social assessment in AgBalance™ is built on the SEEBALANCE® scheme for social LCA, which was developed in 2005 by the Universities of Karlsruhe and Jena, the Öko-Institut (Institute for Applied Ecology) Freiburg and BASF.<sup>1,2</sup>

For AgBalance™, this set of social parameters has been extended and in parts modified to address specific agricultural sustainability topics, e.g., access to land, level of organization or international trade with agricultural products. These topics were initially identified through a stakeholder process in 2009 and 2010, organized by BASF, and were subsequently discussed with leading experts. Feedback from this process was then integrated into the development of these indicators.

Social factors as part of AgBalance™ means integrating social parameters into the assessment model, taking all three pillars of sustainability into account, as originally proposed in the definition of sustainability by the UN Brundtland commission. The strength of a life cycle approach is that the social aspects are evaluated along the life cycle or a defined life cycle. The assessment of social indicators shows the sustainability risks or weaknesses, as well as strengths of any given alternative. It is worth noting that any alternatives that reflect conditions conflicting with legal rights or basic human rights will not be assessed in an AgBalance™ study.

For all social indicators, the production volumes are related quantitatively to a given industry sector (e.g., 'occupational diseases per kg product'). With this approach, it is possible to relate the inputs and outputs from the environmental life cycle assessment to the individual social indicators. For this study, the social input

data can be found in Table 5 later in this document. The additional social indicators in AgBalance™ are not part of the NSF Protocol P352.

### 3.3 Work Flow:

A representative flowchart of the overall process steps and calculations conducted for this AgBalance™ Analysis is summarized in Figure 2 below.




Figure 2. Overall process flow for BASF AgBalance™ methodology

## 4. Study Goals, Context and Target Audience

### 4.1. Study Goals:

The specific goal defined for the Corn Production AgBalance™ Analysis was to quantify the differences in life cycle environmental impacts and total life cycle costs of corn production technologies in the United States.



The study specifically compares the corn production in Iowa in 2000 vs. 2010. The study considered application of the Corn Production in the state of Iowa because roughly 20% of the United States corn production is grown in Iowa. Thus most of the data used in the study is from Iowa State University research on corn production.


The major factor influencing the environmental and cost impact between the two years is the yield increase in the production of corn. Iowa State University data shows an increase of 21.7% from 2000 to 2010 in corn production yield.<sup>3,4</sup> This University information is based on average data collected for the specific years and is published on a yearly basis.

Study results will be used as the basis to guide further product development and marketing decisions that will result in more sustainable production of corn. As well as provide the necessary information to allow a clear comparison between the environmental life cycle, total cost impacts and social aspects as measured by BASF's AgBalance™ tool. It will also facilitate the clear communications of these results to key stakeholders in the agricultural industry who are challenged with evaluating and making strategic decisions related to the sustainable development associated with production of corn.

#### 4.2 *Design Criteria:*

The context of this AgBalance™ study compared the life cycle environmental, cost impacts and social aspects for production of one metric ton (1,000 kg) of corn. The corn production study used data mainly documented by Iowa State University for the production of corn. The data in the study included general data such as yield; seeding such as seed used in corn planting; fertilizers and plant protection such as amounts of N-P-K fertilizer, herbicides, insecticides, additives and applicable emissions. Fuel use in tilling and harvesting such as diesel use for tractor, diesel use for combine and transportation were evaluated. The study relied on internal information and MSDSs were utilized for non-BASF supplier information. The study was technology driven and goals, target audience, and context for decision criteria used in this study are displayed in Figure 3.





**Figure 3.** Context of Corn Production AgBalance™ Analysis

#### 4.3. Target Audience:

The target audience for the study has been defined as agricultural consumers, such as farmers, distributors of agricultural products and trade associations within North America, focusing on the corn production in the state of Iowa. It is planned to communicate study results in marketing materials and at trade conferences.

## 5. Customer Benefit, Alternatives and System Boundaries

### 5.1. Customer Benefit:

The Customer Benefit applied to all alternatives for the base case analysis is the evaluation of the inputs required to produce one metric ton (1,000 kg) of corn in the state of Iowa, which is equivalent to 39.4 bushels of corn (56 lb. per bushel of corn) in one growing season (1 year). This study specifically evaluates all input data that affects crop yield, and is based on the yields reported in the Iowa State University studies referenced in Section 4.1. For the purposes of this study, in situations that increase yield, the amount of inputs required to achieve the CB will decrease, because the yield increase is demonstrating a more efficient use of the inputs. However, the application rate of the inputs could be higher in the alternative. The justification for selecting this CB is because the metric unit is a universally accepted or known amount and one metric ton is a large enough amount to be able to understand the concept. This amount is not small, like a bushel where the representative differences might not be expressed in the study.

### 5.2. Alternatives:

The product alternatives compared under this AgBalance™ study are (1) Corn production in Iowa in 2000 and (2) Corn production in Iowa in 2010. The study also

looks at the tillage process in both years and these are defined in the study as conventional tillage and conservative tillage. Conservative tillage is broken down into minimal tillage and no-tillage. The percentage of this are evaluated in the study alternatives and are based on data from the actual farming process reported in the specific years. These alternatives were selected as they represent technology advancement and social changes in farming.

5.3. System Boundaries.

The system boundaries define the specific elements of the production phases that were considered as part of the analysis. The elements for the use and disposal of one metric ton of corn were not evaluated in this study. The system boundaries for the two alternatives evaluated in this study are shown in Figures 4. Sections identified in gray were excluded from the analysis as they represented identical impacts for both alternatives (e.g. transportation, drying, storage, processing and secondary uses). The justification for these boundaries is that these are the major impact categories for the production of corn and the only difference between the two alternatives is the data used for the different years. The use and disposal of the corn was not evaluated because the CB of one metric ton for both alternatives was the same.

The Eco-toxicity potential of the input chemicals is defined to be evaluated in the Use phase only. This Use phase is the Use phase of the respective life cycles of the input chemicals used in the production of 1 metric ton of corn for both Year 2000 and 2010. This is not the Use phase of defined system boundaries of this study.




Figure 4. System boundaries - Corn Production 2000 & 2010

#### 5.4 Scenario Analyses:

In addition to the base case analysis, additional scenarios will be evaluated to determine the sensitivity of the studies final conclusions and results to key input parameters. Scenario#1 evaluates a 5% yield decrease for the corn production in 2010. Scenario#2 evaluates the the differences of conventional tillage in 2000, conventional tillage in 2010 and conservative tillage in 2010. Scenario #3 evalautes the conservative tillage impact in 2000 versus 2010. The results of the Scenarios will be discussed in Section 8.5:

5.4.1. *Scenario #1:* Reduction of yield in 2010 by 5%

5.4.2 *Scenario #2:* Comparison of conventional tillage in 2000 and 2010 and conservative tillage in 2010.

5.4.3 *Scenario #3:* Comparison of conservative tillage 2000 & 2010

5.4.4 *Scenario #4:* Yield for 2000 set at the same yield as 2010.

## 6. Input Parameters and Assumptions

### 6.1. Input Parameters:

A comprehensive list of input parameters were included for this study and considered all relevant material and operational characteristics. The Generic Data sources included Iowa State University, BASF's North America Agricultural Products Division, Doanes research database and United States Department of Agriculture. The input values from this data are absolute values and the data is from the defined years in the study. If data was not available for defined years, data closest to the defined years was used. For example, the USDA agricultural census data is gathered every five years so the data for 2000 was from the 2002 USDA report. The data for 2010 is from the 2007 USDS report which was the latest report published.

The corn production study evaluates the production of the Customer Benefit (CB), which is one metric ton of corn for one growing season. The production of corn is an annual process, since the seed needs to be planted, the fertilizers and herbicides need to be applied and the corn needs to be harvested to get the CB. In grain agricultural production, crops are usually rotated year after year. If corn is planted in a field one year, the next year a different crop such as soybeans or wheat is planted. The reason for this is the demand of the individual crops on the soil and the nutrients in the soil. The production of corn has a high demand for Nitrogen (N) as shown in the study. If corn is planted year after year on the same land, more N is needed the next year than in the previous year. Soybeans actually produce N during the growing process so there is no demand on N. The AgBalance™ tool does evaluate the benefits of crop rotation, however the affect of crop rotation is not evaluated in this study because the evaluation is between the two years for corn production.



### 6.1.1. Cultivation Parameters:

In this analysis, the cultivation parameters for each year are evaluated as conventional tillage and conservative tillage. Conservative tillage consists of minimal tillage and no-till. The number of passes across a field varies between the tillage processes. In conventional tillage the ground would need to be worked twice and then planted for a total of 3 passes. In conservative tillage there could be one pass for working the ground and 1 pass for planting and no-till would be 1 pass for planting. In the two alternatives, these tillage processes are broken down to the actual percentage that is reported for production of corn in Iowa. The percentage of minimal tillage and no-till in the conservative tillage is also calculated based on data from Iowa and used for the tillage fuel use.

Inputs for seeds, fertilizers and plant protection, per area unit for each of the alternatives, was used as the input amounts. The input amounts used per acre (ac) are shown in Table 1. The Base Case compares the data from Year 2000 and Year 2010.

The eco-toxicity input values were calculated based on the methodology described in Section 8.1.7. The values were then multiplied by the amounts used in the production of the CB and were only based on the active ingredients of the plant protection products. Table 2 shows the eco-toxicity values used for the active ingredients in the plant protection products.

The corn seed variety changed from 2000 to 2010 to reflect the changes made in the genetics of corn seed in the 10 years. In 2000, the Pioneer 33G26 variety was mainly planted in Iowa and this corn seed was Roundup® Ready. This meant that it had a genetic trait to be glyphosate resistance, which glyphosate is the active ingredient in Roundup®. In 2010, the Dekalb DKC52-59 variety was mainly planted in Iowa and this variety had the genetic trait of VT3, which means it had 3 genetic traits. These traits were glyphosate resistance, ear worm protection (above the ground protection) and root worm protection (below the ground protection). The main advantage of the VT3 corn seed was the elimination of insecticides in the soil during planting.



**Table 1:** Input data usage rates per acre for Base Case corn production in Iowa.

<b>All Alternatives: Corn produced in Iowa in 2000 vs 2010</b>						
% tillage method in each year		<b>42</b>	<b>58</b>	<b>29</b>	<b>71</b>	
		2000 Conv-till	2000 Cons-till	2010 Conv-till	2010 Cons-till	
<b>General Data</b>	<b>Characteristics</b>					
	Corn Variety	PIONEER 33G26	PIONEER 33G26	DKC52-59	DKC52-59	
	Yield (grain)	143	143	174	174	
	Moisture content	15	15	15	15	
	Yield (15% moisture grain)	143	143	174	174	
					bu/ac	
						%
						bu/ac
<b>Seeding</b>	<b>Seed</b>					
	Amount of corn seed	28315	28315	33620	33620	
	Seed cultivation level	0.248%	0.248%	0.242%	0.242%	
	ST Fungicide (fludioxonil)	0.00039	0.00039	0.00047	0.00047	
	ST Fungicide (mefenoxam)	0.00063	0.00063	0.00075	0.00075	
	ST Insecticide (clothianidin)	0.00000	0.00000	0.03703	0.03703	
	Same characteristics as main culture?	Y	Y	y	y	
					seeds/ac	
					seed	
					lb ai/ac	
					lb ai/ac	
					lb ai/ac	
					y / n	
<b>Tillage and</b>	<b>Tillage and planting</b>					
	Number of passes	3	1.7	3	1.54	
	Diesel use tractor	2.47	1.12	2.47	0.97	
	Lubricants for machinery	0.02	0.01	0.02	0.01	
					gal/ac	
					gal/ac	
<b>Fertilizers and plant protection</b>	<b>Fertilizer</b>					
	N-fertilizer	131	131	142	142	
	P-fertilizer	55	55	65	65	
	K-fertilizer	69	69	80	80	
	Fuel Use	0.7	0.7	0.7	0.7	
						lb N/ac
						lb P/ac
						lb K/ac
						gal/ac
	<b>Plant protection</b>	Pesticide amounts				
	Burndown					
	Roundup Weathermax	0	0.25	0	0.24	
	2-4, D	0	0.13	0	0.08	
						gal/ac
	<b>Pre-emergence</b>					
	Harness Xtra	0.42	0.45	0.34	0.35	
						gal/ac
<b>Post-emergence</b>						
Roundup Ultra	0.26	0.26	0	0		
Roundup Weathermax	0	0	0.21	0		
Roundup Powermax	0	0	0	0.21		
					gal/ac	
<b>Other additives and water</b>						
Ammonium sulfate	1.61	3.21	1.61	3.21		
Water	18.90	28.35	18.90	28.35		
					lbs/ac	
<b>Insecticide</b>						
Lorsban (chlorpyrifos)	8.17	8.17	0	0		
					lbs/ac	
Plant protection passes	2	3	2	3		
Fuel Use	0.2	0.3	0.2	0.3		
					gal/ac	
<b>Harvesting</b>	<b>Harvesting</b>					
	Diesel use combine	1.45	1.45	1.45	1.45	
	Fuel use transportation	3.05	3.05	3.05	3.05	
	Lubricants for machinery	0.03	0.03	0.03	0.03	
	Field work	2.6	2.6	2.6	2.6	
					gal/ac	
					gal/ac	
					gal/ac	
					hrs/ac	

**Table 2:** Input data values for Eco-toxicity of chemicals for corn production in Iowa.

<u>Chemicals</u>	<u>Eco-Toxicity Value</u>
Fludioxonil	2660
Mefenoxam	225
Clothianidin	97
Glyphosate-isopropylammonium	177
2,4-D, Isopropylamine Salt	18
Acetochlor	7611
Atrazine	11406
Chlorpyrifos Methyl	6605

Table 3 shows the input amounts for biodiversity and soil as part of the AgBalance™ analysis. Most of the data for soil is the same for the two alternatives since the soil has not changed in Iowa in the 10 years of the evaluation. A soil compaction score is determined based on the soil texture, number of days with field capacity, depth of impermeable layer, soil organic matter content, type of land use and stocking rate. Since the values are all the same for both alternatives, the score was the same and normalized to 1 for soil compaction. The only differences in soil between the two years are the loss from wind erosion and humus amount left on field. The humus amount is calculated based on the yield amounts from the two years.

**Table 3:** Input data for soil and biodiversity for corn production in Iowa.

<b>All Alternatives: Corn produced in Iowa in 2000 vs 2010</b>					
% tillage method in each year		42	58	29	71
		2000 Conv-till	2000 Cons-till	2010 Conv-till	2010 Cons-till
<b>Biodiversity</b>	<u>Biodiversity</u>				
	Payments received for AES	95.83	95.83	119.68	119.68
	Protected areas' share at country's area	6.39	6.39	6.60	6.60
	Maximum yield potential	195	195	195	195
	Number of endangered species	113	113	113	113
<b>Soil</b>	<u>Soil</u>				
	Soil nutrient supply class of phosphate (see table to the right)	B	B	B	B
	Nitrogen mineralized in soil	100	100	100	100
	Soil texture	Heavy clay loam and Clay soils (clay content > 27%)	Heavy clay loam and Clay soils (clay content > 27%)	Heavy clay loam and Clay soils (clay content > 27%)	Heavy clay loam and Clay soils (clay content > 27%)
	Number of days with field capacity	<125	<125	<125	<125
	Depth to impermeable layer	>80 cm	>80 cm	>80 cm	>80 cm
	Soil organic matter content	3-6%	3-6%	3-6%	3-6%
	Type of land use	Arable land	Arable land	Arable land	Arable land
	Stocking rate (if livestock is considered)	No livestock	No livestock	No livestock	No livestock
	Average precipitation (in/year)	30	30	30	30
	Typical slope of hillside in the related area	3	3	3	3
	Loss from Wind Erosion	10.374	10.374	10.127	10.127

## 6.2. Cost Inputs

### 6.2.1. User Costs

User costs were evaluated for each alternative based on the fixed costs and variable costs per acre. Table 4 lists the total cost including fixed cost and the operating costs are the variable costs. The total cost minus the variable costs was calculated as the fixed costs. The other costs in Table 3 were used for the social aspects of the corn production AgBalance™.

**Table 4:** Input data cost and revenue for corn production in Iowa.

<b>All Alternatives: Corn produced in Iowa in 2000 vs 2010</b>						
% tillage method in each year		<b>42</b>	<b>58</b>	<b>29</b>	<b>71</b>	
		2000 Conv-till	2000 Cons-till	2010 Conv-till	2010 Cons-till	
<b>Total Costs</b>	<b>Costs</b>					
	Total operating cost \$/ac	272.00	272.00	534.64	534.64	\$/ac
	Total cost including fixed \$/ac	403.85	403.85	692.75	692.75	\$/ac
	Capital investments	59.11	59.11	58.75	58.75	\$/year
	Depreciations	29.77	29.77	60.19	60.19	\$/year
	Machinery costs <i>FIXED</i>	16.51	16.51	30.11	30.11	\$/year
	Machinery costs <i>VARIABLE</i>	18.57	18.57	33.87	33.87	\$/year
	General repair costs	4.13	4.13	7.52	7.52	\$/year
	Maintainance	2.06	2.06	3.76	3.76	\$/year
	EH&S programs and regulatory costs	500.00	500.00	500.00	500.00	\$/year
	Illness & injury costs (medical, legal, lost time)	11.30	11.30	20.53	20.53	\$/year
	Property protection & warehousing costs	14.72	14.72	17.02	17.02	\$/year
	Costs for extra disposal	0	0	0	0	\$/year
	Training costs	500.00	500.00	500.00	500.00	\$/year
	Other costs	10.14	10.14	20.32	20.32	\$/year
	Direct costs	0	0	0	0	\$/t of product
	<b>Economic</b>					
	Farm profits per year	-7.45	-7.45	107.42	107.42	\$/ac
	Subsidies (not for AES)	85.58	85.58	28.50	28.50	\$/ac
	Type & conditions os subsidies	0	0	0	0	\$/ac
Gross value of production (selling prices)per unit area	264.55	264.55	642.06	642.06	\$/ac	
	0	0	0	0		

## 6.3. Social Inputs

### 6.3.1. Social metrics

Since this is an AgBalance™ study there are metrics with social factors evaluated for each alternative based on the factors defined in the AgBalance™ methodology. The input data for these metrics were based on data from farms in Iowa and if data was found for farmers raising corn, then this specific data was used. Table 5 lists the input data used in each of the alternatives based on a specific unit. In the final analysis, these values are set to the CB just as the environmental and economic values are set to the CB.

**Table 5:** Input social factors for corn production in Iowa.

<b>All Alternatives: Corn produced in Iowa in 2000 vs 2010</b>						
% tillage method in each year		42	58	29	71	
		2000 Conv-till	2000 Cons-till	2010 Conv-till	2010 Cons-till	
<b>Social</b>	<b>Social</b>					
	Working accidents	1.98227E-05	1.98227E-05	1.98227E-05	1.98227E-05	no./working hour
	Occupational diseases	8.33395E-07	8.33395E-07	8.33395E-07	8.33395E-07	no./working hour
	Workers' wages (medium qualification level)	9.3	9.3	12.96	12.96	\$/hour
	Number of employees	82991	82991	71924	71924	no. in year
	Number of full-time equivalents	30482	30482	25133	25133	no. in year
	Time spent for professional training	19193	19193	24672	24672	days/year
	Number of memberships	7000	7000	7000	7000	no. in year
	Number of different member organizations	48	48	48	48	no. in year
	% of leased land of total agricultural area	59	59	60	60	%
	Land lease	120.00	120.00	184.00	184.00	\$/ac
	Percentage of women among farm proprietors	5.85	5.85	7.96	7.96	%
	Number of people w/ disabilities employed	0	0	0	0	no./working hour
	Students /trainees in dedicated education / training for agriculture	3490	3490	4180	4180	no. in year
	Total contributions to old-age insurance	7256.73	7256.73	13243.2	13243.2	\$/year
	Total contributions to accidents insurance	5591.00	5591.00	12561.00	12561.00	\$/year
	Total contributions to health insurance	1697.14	1697.14	3730.62	3730.62	\$/year



## 7. Data Sources

### 7.1. Environmental:

The environmental impacts for the production of the two alternatives were calculated from eco-profiles (a.k.a. life cycle inventories) for the individual components and for fuel usage. Life cycle inventory data for these eco-profiles were from several data sources, including BASF specific manufacturing data and customer supplied data. For the seed treatment and chlorpyrifos, surrogate eco-profiles for the insecticides and fungicides were used from Ecoinvent since these were small amounts and data for specific chemicals were not available. Overall, the quality of the data was considered medium-high to high. None of the eco-profile data was considered to be of low data quality. A summary of the eco-profiles is provided in Table 6.

**Table 6:** Summary of eco-profiles used in the corn production AgBalance™ analysis

Eco-Profile	Source, Year	Comments
Seed Treatment	2010	Ecoinvent database
Diesel Use - US	US Avg., 1996	Boustead database <sup>5</sup>
Lubricants for Machinery	1996	Boustead database <sup>5</sup>
Urea Fertilizer	Agrium, 2005	Boustead database <sup>5</sup>
DAP Fertilizer	U of Minnesota., 2002	Boustead database <sup>5</sup>
K-Fertilizer	DE Avg., 1997	Boustead database <sup>5</sup>
Water	BASF well data, 1995	Boustead database <sup>5</sup>
Glyphosate	1997	Boustead database <sup>5</sup>
Acetochlor	DE Avg., 1997 BASF, 2006	Boustead database <sup>5</sup>
Atrazine	DE avg., 1997	Boustead database <sup>5</sup>
2,4-D herbicide	DE Avg., 1997 BASF, 2006	Boustead database <sup>5</sup>
Ammonium sulfate	DE Avg., 1996	Boustead database <sup>5</sup>
Crop oil	DE Avg., 1996	Boustead database <sup>5</sup>
Chlorpyrifos	2010	Ecoinvent database
BASF data sources are internal data, while the others are external to BASF. Internal data is confidential to BASF; however, full disclosure can be provided to NSF International for verification purposes.		

### 7.2. Amounts and Costs:

The data sources for the amounts and costs of the individual components were obtained from the BASF Agricultural Products Division. A summary of the source of this data is provided in Table 7.

**Table 7:** Summary of data sources for amounts and costs

<b>Corn Characteristics</b>	<b>Data Source:</b>
Pioneer 33G26	Pioneer Hybrids
Variety - DKC52-59	Dekalb Genetics Seed Company
Yield (grain)	Iowa State University, FM-1789 (6/01 & 12/11)
Amount of corn seed	Doane Research
<b>Plant protection</b>	
N-fertilizer	USDA & Iowa State University
P-fertilizer	USDA & Iowa State University
K-fertilizer	USDA & Iowa State University
Harness Xtra	Monsanto Company, Doanes Research
Roundup	Monsanto Company, Doanes Research
Ammonium sulfate	BASF Corp.
Lorsban	Dow AgroSciences
Water amounts	BASF Corp.
Herbicide Application	Iowa State University
Diesel use	Iowa State University, PM709
<b>Biodiversity &amp; Soil</b>	
Payments Agro-environmental schemes	Doane Research
Protected areas	Doane Research
Maximum yield potential	Iowa State University, File A1-14
Number of endangered species	www.iucnredlist.org
All soil data for Iowa	Iowa State University - ISPAID 7.3 Database
Loss from Wind Erosion	NRCS publication, "2007 National Resources Inventory, Soil erosion on cropland"
<b>Cost &amp; Revenue</b>	
Variable Cost	Iowa State University, FM-1789 (6/01 & 12/11)
Fixed Cost	Iowa State University, FM-1789 (6/01 & 12/11)
Machinery cost	Iowa State University, FM-1789 (6/01 & 12/11)
Capital cost	Iowa State University, FM-1789 (6/01 & 12/11)
Land Lease	Iowa State University, FM-1789 (6/01 & 12/11)
Other costs	Iowa State University, FM-1789 (6/01 & 12/11)
Corn prices	Iowa State University, FM-1789 (6/01 & 12/11)
<b>Social</b>	
Accidents and Diseases	U.S. Ag Census Data, AC-02-A-51 & AC-07-A-51
Workers' wages	Iowa State University, Wages and Benefits publication
Employees	U.S. Ag Census Data, AC-02-A-51 & AC-07-A-51
Professional training	Iowa State University, Extension Training
Memberships	Iowa Corn Grower Association
Land lease	U.S. Ag Census Data, AC-02-A-51 & AC-07-A-51
Women Proprietors	U.S. Ag Census Data, AC-02-A-51 & AC-07-A-51
Training for agriculture	Iowa State University
Old-age insurance	U.S. IRS
Accidents insurance	Iowa State University, FM-1789 (6/01 & 12/11)
Health insurance	U.S. IRS


## 8. AgBalance™ Results and Discussion

### 8.1. Environmental Impact Results:

The environmental impact results for the Corn Production AgBalance™ are generated as defined in Section 3.2.1., Environmental Burden Metrics. The results discussed in Section 8.1.1 through 8.3 (depicted in Figures 5 through 22) are for the Base Case only and do not represent any of the Scenarios.

#### 8.1.1. Primary energy consumption:

Energy use is predominantly affected by the fertilizer production and the amount of fertilizer that is needed per customer benefit. There was an energy reduction of 10% in 2010 operations when compared to 2000 operations. More fertilizer was used in 2010 than in 2000, but due to the increase in yield the amount of fertilizer per CB was smaller. There is also a reduction from the plant protection, harvesting and tillage and planting. The energy reduction from plant protection was in the reduction of the amounts. In the other operations, the energy reduction is mainly from less fuel use, this is due to higher yields and the amounts based on 1 ton of corn as the CB. Figure 5 shows the key drivers for the primary energy consumption. Renewable energy sources were analyzed in this study, but made up only 3% of the total energy sources.



**Figure 5.** Primary energy consumption.

#### 8.1.2. Raw material consumption:

Figure 6 shows that the key driver for the raw material or resource consumption is dominated by the production of fertilizers and the relevant energy carriers. More than 60 g of fertilizer (N, P, K) were used per kg of corn (>60 kg per CB) in 2010 and more than 65 g of fertilizer per kg of corn in 2000.

Per the BASF AgBalance™ methodology, individual raw materials are weighted according to their available reserves and current consumption profile. These weighting factors are appropriate considering the context of this study. Phosphorous is the main resource that dominates raw material consumption (apart from energy carriers like coal, lignite, oil and gas). Within the different resources assessed Phosphorous is weighted highly since it is scarce. Figure 7 shows the overall use of individual raw materials for the production of corn in 2000 and 2010.




Figure 6. Raw Material consumption by Module.




Figure 7. Raw Material consumption by Type.



8.1.3. Air Emissions:

8.1.3.1. *Global Warming Potential (GWP)*: The highest carbon footprint occurred in soil and land use. With field emissions of 8 kg N<sub>2</sub>O-N per ha a year (IPCC 2006<sup>6</sup>) from crops, this is the dominant factor. Other important sources for global warming potential emissions are N<sub>2</sub>O-emissions from N-fertilizers (1% of fertilizer N directly and 0.325% of fertilizer N indirectly through volatilization and leaching; IPCC 2006<sup>7</sup>) as well as CO<sub>2</sub>-emission from urea (worst case: 20% of urea is being emitted as CO<sub>2</sub>). Emissions in fertilizer production are mainly due to the use of fossil energy. Figure 8 shows the overall GHG emission for production of corn in 2000 and 2010.




Figure 8. Global warming potential.

8.1.3.2. *Photochemical ozone creation potential (POCP, smog)*: Emissions with Photochemical Ozone Creation Potential are dominated by fossil fuel use. The fossil fuel is used in the production of fertilizer and in diesel fuel use in harvesting, tillage and planting. There are some POCP emissions from the production of pesticides, but this is very minor in this analysis compared to the other inputs. The difference between the two years is mainly from the increase in yield in 2010, based on the defined CB. This environmental category has a very minor influence on the total study and the results are shown in Figure 9.




Figure 9. Photochemical ozone creation potential.

8.1.3.3. *Ozone depletion potential (ODP):* Overall, the ODP emissions are very small and are dominated by the production of other plant protection agents (halogenated hydrocarbons) in 2000. The plant protection agents are the insecticides and herbicides. This environmental category has a very minor influence also and the results are shown in Figure 10.




Figure 10. Ozone depletion potential.

8.1.3.4. *Acidification potential (AP):* It can be seen in Figure 11 that overall, NH<sub>3</sub>- and NO<sub>x</sub> emissions from fertilizer use are dominant. According to literature<sup>8</sup>, 2% of N-fertilizers are emitted as NH<sub>3</sub> and 2% as NO<sub>x</sub> respectively. Another

important fraction comes from fossil energy use for fertilizer production and field work (diesel and oil use / burning).




Figure 11. Acidification potential.

Figure 12 below, shows the relative impacts of the four air emissions: GHG, AP, POCP and ODP. These values are normalized and weighted based on the calculation factors (see Figure 39 for the calculation factor percentage). The calculation factor is a calculation of the relative environmental factors and the social weighting factors.




Figure 12. Overall Relative Impacts of Air Emissions



8.1.4. Water emissions:

Figure 13 displays that water emissions changes are dominated by fertilizer use and some from pesticide use. The main substances of concern emitted from fertilizers are leaching through the soil and from heavy metals. From pesticides the concern is from carbon compounds. These carbon compounds from pesticides are evaluated for the amount of water needed or the Chemical Oxygen Demand (COD) to acceptable levels. According to literature sources<sup>9</sup> mineral fertilizers contain a substantial amount of heavy metals (up to 2 g per kg). A worst case scenario was used here. Up to 10% of fertilizer N (depending on climate and region) ends up as a water emission and up to 1% of fertilizer P ends up as water emission.<sup>10</sup> Both the N-water-emissions and P-water-emissions are included as part of the AgBalance™ base case.





Figure 13. Water emissions.

8.1.5 Solid waste generation:

Solid waste emissions have minor influence on the overall result. Solid wastes (chemicals) generated in fertilizer and plant protection production are the dominating factors. These waste values include municipal, hazardous and mining waste. Hazardous waste is generated from production of pesticides, fertilizers and diesel fuel. Figure 14 displays the solid waste emissions for the two alternatives.




**Figure 14.** Solid waste generation.

8.1.6 Land use:

As displayed in Figure 15, land use is assessed for each alternative. Land use is one of the most important environmental categories for agricultural processes assessed with AgBalance™. The land use impact assessment takes into account an Ecosystem Damage Potential<sup>11,12</sup> (EDP), with damage functions and generic characterization factors for quantifying damages to ecosystems from land occupation and land transformation. EDP is based on an assessment of the impacts of land use on species diversity.

The land use in the study is quite significant to the overall study with the calculation factor being 25%. The majority of the land use changes between 2000 and 2010 are due to the increase in yield and less land is needed per CB. The land use reduction is over 35% in 2010 compared to 2000. Land use for production of fertilizers is negligible to the overall land use amounts. The units for the land use metric are square meters each year/CB (one metric ton of corn).



**Figure 15.** Land use – EDP assessment.

#### 8.1.7 *Eco-Toxicity potential:*

The Eco-toxicity potential for the corn production in Year 2000 and Year 2010 were analyzed for the Use phases only of their respective life cycles. In agricultural production, chemicals are intentionally released into the environment, i.e., fertilizers and pesticides. As a result, eco-toxicity is integrated within the AgBalance™ methodology. The method used for the determination of the eco-toxicity potential follows the basic rules of the European Union Risk Ranking System (EURAM).<sup>13</sup> This evaluation is based on water solubility, water/octanol partition coefficient, biodegradability and toxicity towards water organisms, plants, bacteria. These data are usually available in the material safety data sheets. The scoring system is based on the principles of environmental risk assessment (i.e. risk as the product of hazard and exposure) and the ultimate end-point in the environment. This includes an assessment of biodegradability, according to the OECD criteria (inherent/readily biodegradable/persistent).

For the toxicity of the chemicals used by the farmer in the Use phase, these values are assessed in the Social parameters for toxicity potential to the farmer. The toxicity scoring for the farmer uses the R-phrase for the toxicity of the final products and the relevant material quantities. This value is then assessed with other social factors for the farmer/employees and the final number is normalized to 1.0 for the worse case alternative, see Figure 22.

For the toxicity production phase of the raw materials, not only were the final toxicity of the products considered but the entire pre-chain of chemicals required to manufacture the products were considered as well. These values are assessed in AgBalance™ but are not part of the overall environmental rating, like they were in EEA.



The use of nanoparticles were not evaluated in the chemical inputs for any of the alternatives, therefore the toxicity of nanoparticles was not evaluated in the study results.

The application of the insecticide in Year 2000 to the soil has a major impact on the eco-toxicity, compared to Year 2010, when no insecticide was applied to the soil. The hybrid seed technology allows for the seed to be insect resistant and thus no insecticide is needed for the plant. The eco-toxicity of the insecticide used in 2000 was evaluated, however the eco-toxicity of the plant hybrid used in 2010 was not evaluated since the resistance is built into the plant structure.

As to be expected the application of the materials (fertilizer, herbicides, insecticides and fungicide) during the Use phase contributed the largest amount to the ecotoxicity potential for each alternative. Figure 16 shows the ecotoxicity of the two alternatives.

Figure 17 shows the overall toxicity potential score for each alternative and how the scoring is distributed across the life cycle stages. The values have been normalized and weighted. For the weighting, the human health toxicity was weighted as 50% of the total toxicity potential with the Use phase making up 70% of this total and Production phase making up 30% of this total. The eco-toxicity made up the other 50% of the total toxicity potential with all of this being the Use phase. Consistent with the discussion above, the Use phase is the most significant and disposal was not evaluated. A high safety standard was assumed for the manufacturing processes for the raw materials. For the Use phase, an allowance was made to take into consideration the open nature of the application process. For the normalization, the highest toxicity potential alternative was set to a value of 1 and the other alternative was proportioned to this value.





Figure 16. Overall Eco-toxicity potential




**Figure 17.** Overall Toxicity potential - Life Cycle Phases

#### 8.1.8 Risk potential (Occupational Illnesses and Accidents potential):

All the materials and activities accounted for in the various life cycle stages were assigned specific NACE codes. NACE (Nomenclature des Activités Economiques) is a European nomenclature which is very similar to the NAICS codes in North America. The NACE codes are utilized in classifying business establishments for the purpose of collecting, analyzing, and publishing statistical data related to the business economy and is broken down by specific industries. Specific to this impact category, the NACE codes track, among other metrics, the number of working accidents, fatalities and illnesses and diseases associated with certain industries (e.g. chemical manufacturing, petroleum refinery, inorganics etc.) per defined unit of output. By applying these incident rates to the amount of materials required for each alternative, a quantitative assessment of risk is achieved.


In Figure 18, the greatest Occupational Illnesses and Accident potential occurs in the production of fertilizers. The field work also contributes to the risk potential for occupational illnesses and accidents, but this does not include the cultivation process.



**Figure 18.** Occupational Illnesses and Accidents

8.1.9 *Consumptive Water Use:*

In AgBalance™, water use is assessed as a separate environmental impact category. The method for assessing freshwater consumption is a method described by Pfister, Köhler and Hellweg<sup>14</sup>. In this method, only consumptive water use is assessed and no green water is evaluated (precipitation and soil moisture). Consumptive water use consists of water used in production of CB and water used for irrigation. The method also includes a regionalization factor which is based on GIS data as applied at the watershed levels. Details of the corresponding regionalized damage factors are available in supplementary material provided in the Pfister et al publication. Figure 19 shows the graph of the total consumptive water used weighted with the regional factor. There was no irrigation evaluated in this study so all the water was used in application of pesticides.



**Figure 19.** Consumptive Water Use



### 8.1.10 Biodiversity:

By definition, biodiversity cannot scientifically be quantified in its totality. Therefore, any quantification of "biodiversity" is an approximation, requiring the relevant elements of biodiversity to be defined and the appropriate indicators used. In AgBalance™, the impact of agricultural activity on biodiversity is assessed as a relative function, constructed from the Biodiversity State Indicator, which is a factor based on the number of endangered species. Further indicators have the potential to increase or decrease biodiversity. The factors evaluated in this AgBalance™ study were Biodiversity State Indicator (establish baseline), Agri-Environmental Schemes, Protected Areas, Eco-toxicity, Farming intensity, Nitrogen Surplus, Potential for intermixing and Crop Rotation. Crop rotation was not evaluated in this study so the values for Potential for intermixing and Crop Rotation were the same for 2000 and 2010. The actual normalized values for the biodiversity assessments from the study are shown in Table 8.

**Table 8:** Normalized Biodiversity values for corn production in Iowa.

<b>Biodiversity</b>	<b>Year 2000</b>	<b>Year 2010</b>
Biodiversity indicator	0.94	0.94
Agri-environmental schemes	1.30	1.17
Protected areas	1.00	1.00
Crop rotation	0.60	0.60
Eco-tox potential	0.60	1.00
Farming intensity	0.67	0.63
Intermixing potential	0.83	0.83
Nitrogen Surplus	0.85	0.88
<b>Result biodiversity</b>	<b>0.21</b>	<b>0.30</b>
<b>Normalized biodiversity</b>	<b>1.00</b>	<b>0.70</b>

### 8.1.11 Soil:

The AgBalance™ methodology for the Soil impact category uses different indicators, which are designed to capture the main impacts to long-term soil quality as a result of human agricultural activity on arable land. These indicators consist of: Soil Organic Matter balance, Nutrients (N, P, K) balance, Soil Compaction Potential, and Erosion. Table 9 shows the normalized values for the soil assessment from the study.

**Table 9:** Normalized Soil values for corn production in Iowa.

<b>Soil</b>	<b>Year 2000</b>	<b>Year 2010</b>
Nutrients:N	1.00	1.00
Nutrients:P	0.83	0.81
Nutrients:K	1.00	1.00
Humus	1.00	0.59
Compaction	1.00	1.00
Erosion	1.00	0.98
<b>Result soil</b>	<b>0.97</b>	<b>0.90</b>
<b>Normalized aggregated soil</b>	<b>1.00</b>	<b>0.92</b>

#### 8.1.12 Environmental fingerprint:


Following normalization, or normalization and weighting with regards to the emissions categories, the relative impact for all eight of the environmental categories for each alternative was calculated. The actual environmental category values from the study are shown in Table 10 and the graph of these values are shown in the environmental fingerprint, Figure 20. A value of 1 represents the alternative with the highest impact in the concerning category, all other alternatives are rated in relation to 1.

The 2010 corn production is better than 2000 corn production in all the environmental categories as shown in the environmental fingerprint. As discussed previously in the individual impact categories, the higher yield in 2010 corn production clearly is the main impact on the environmental life cycle impacts due to less inputs needed to produce the CB of one ton of corn. The greatest environmental advantages in 2010 corn production over 2000 corn production can be noticed in the following categories:

- Eco-toxicity
- Land use
- Biodiversity

**Table 10:** Normalized environmental category values for corn production in Iowa.

<b>Data for environmental fingerprint</b>	<b>Year 2000</b>	<b>Year 2010</b>
<b>Energy Consumption</b>	<b>1.00</b>	<b>0.89</b>
<b>Emissions</b>	<b>1.00</b>	<b>0.88</b>
<b>Eco-Toxicity Potential</b>	<b>1.00</b>	<b>0.48</b>
<b>Resource Consumption</b>	<b>1.00</b>	<b>0.90</b>
<b>Land Use</b>	<b>1.00</b>	<b>0.65</b>
<b>Water Use (No irrigation)</b>	<b>1.00</b>	<b>0.86</b>
<b>Biodiversity</b>	<b>1.00</b>	<b>0.70</b>
<b>Soil</b>	<b>1.00</b>	<b>0.92</b>



**Figure 20.** Environmental fingerprint corn production.

8.2 *Economic Cost Fingerprint.*

The life cycle cost data for AgBalance™ are generated as defined in Section 5 of the BASF AgBalance™ methodology and described in section 6.2 above. The results of the life cycle cost analysis found that the cost in 2010 were lower due to the increase in yield. See Table 3 in section 6.2 above for production cost (fixed and variable) for 1 acre in both 2000 and 2010. Figure 21 represents the graph of the costs for each of the alternatives based on the total cost.


For each alternative, the macro-economic indicators are quantified according to the principles outlined below. The resulting values in USD/ha are then summed up, according to the formula:

$$\text{Macro-economic Indicator Result (a) [USD/ha]} = \text{Farm Profits (a) [USD/ha]} - \text{Subsidies (a) [USD/ha]} + \text{Productivity (a) [USD/ha]}.$$

Here (a) denotes the specific result for a given alternative.

The macroeconomic indicator value is aggregated with the costs to the economic score of each alternative. Figure 22 shows the graph of the fixed, variable and macro-economic “costs”. In the case of this study, the profits from 2000 were a negative number, therefore the farm net worth expressed in \$/ha was added to the equation in order to get a positive value and to show the impact of a negative profit on the net worth of the farm. Table 11 lists all the cost result values used in the economic cost fingerprint.






**Figure 21.** Corn production costs

**Table 11:** Economic cost fingerprint values for corn production in Iowa.

Fixed cost/CB	50.52	35.77	USD/CB
Variable cost/CB	104.22	120.96	USD/CB
Total cost (fix + variable)/CB	154.75	156.74	USD/CB
Profit/CB	-2.85	24.30	USD/CB
Profit/ha	-18.40	265.33	USD/ha
Iowa Farm Net Worth (Ave)	475337.00	1238211.00	USD
Iowa Farm Net Worth/ha	2123.11	5097.30	USD/ha
Subsidies (not for AES)/CB	32.79	6.45	USD/CB
Subsidies (not for AES)/ha	294.21	70.39	USD/ha
Net Worth+Profit-Subsidies/ha	2642.47	5292.24	USD/ha
Macro-Economic Costs/ha	2652.27	5303.59	USD/ha
Average Farm size in Iowa (ha)	553	600	ha



**Figure 22.** Economy Fingerprint corn production

### 8.3 Social Fingerprint.


The assessment of social impacts in the up- and downstream processes in AgBalance™ is based on the SEEBALANCE® method<sup>15,16</sup>. This approach to social assessment is based on a sectoral assessment, where key social figures from different industry segments are related to their corresponding production volumes. The resulting social profiles for processes or products then assume a format, equivalent to the eco-profiles, used in the environmental part. See Table 12 for the result values for the corn production social fingerprint data and Figure 23 for the corn production social fingerprint.

For all social indicators, a quantitative relationship is made for the production volumes of a given industry sector (e.g. 'occupational diseases per kg product'). With this approach, it is possible to relate the inputs and outputs from the environmental life cycle assessment to the social indicators. To this end, different statistical databases are combined to connect social indicators to production volumes. The link between products and corresponding social impacts is made by a sector assessment. It is based on the 'Nomenclature générale des activités économiques dans les Communautés Européennes' (NACE, general nomenclature of economical activities in the European Community), an initiative that classifies all industries into different sectors, or the ISIC, the International Standard Industrial Classification. All products can be linked to these NACE/ISIC codes, using the product classification list (CPA =

Classification of Products by Activity). The numbers for the official statistics in Europe are frequently stored in this format.

**Table 12:** Social fingerprint values for corn production in Iowa.

Ag Modul		Unit	Year 2000	Year 2010
	Working hours	h / CB	0.713	0.622
Country			USA	USA
Farmer	Working Accidents	number / CB	1.42E-05	1.17E-05
	Occupational Diseases	number / CB	5.98E-07	4.91E-07
	Toxicity Potential	points / CB	9.08E+01	8.63E+01
	Wages	PPP Dollar / CB	6.63	8.06
	Professional Training	h / CB	1.82E-03	2.83E-03
	Organization	normalized	1.00	0.87
Consumer	Residues in Food&Feed	rating	1.00	1.00
	Residues of GMO in Food	rating	0.03	0.03
Local/Nat. Commun.	Access to Land	EUR / CB	20.77	20.31
	Employment	hours / CB	1.94	1.78
	Gender Equality	%dev	44.15	42.04
	Integration	working years / CB	0.00	0.00
International Commu	Imports from Devel. Countries	EUR	-1.79E+09	-3.50E+09
	Fair Trade	EUR / CB	0.00	0.00
	Child Labor	working years / CB	0.00	0.00
Future Generations	Trainees	h / CB	5.50E-05	8.00E-05
	Social Security	EUR / CB	4.98	8.83



**Figure 23.** Social Fingerprint corn production

#### 8.4 *AgBalance™ Analysis Portfolio (Single Score):*

At the highest aggregated level, the results of the environmental, social and economic assessments are presented as Single Score diagrams. This format is also used to illustrate the total socio-eco-efficiency score of the AgBalance™ evaluation, see Figure 24. This format offers a high degree of clarity and has been introduced as a new feature within AgBalance™. Conventional portfolio-diagrams (eco-efficiency) are also created and documented in the study report, see Figure 25. The normalized values from the environmental, social, and economic fingerprint are aggregated into a single relative score through the use of relevance, societal factors and the E/C or S/C scaling factors. Given that the analysis features multiple criteria and a plethora of single results, it is vitally important to show the final conclusions in a transparent and easy-to-understand way. Otherwise, it would be impossible for the reader of an AgBalance™ study to easily aggregate.

The results in Figure 24 show the individual scores of the Ecology, Society and Economy of the AgBalance™ study. The Year 2010 has the best results compared to 2000 in all of these graphs due to the normalized value being lower. In these graphs, the better score is closer to 0.6 and a worst score is closer to 1.4. These are established based on the normalized values being centered at 1 or the individual normalized value being divided by the average score of all the alternatives. The Total Score graph shows the sum of the Ecology, Society and Economy assessments with each having equal weighting of 33.33%.

For a clearer understanding of how weighting and normalization is determined and applied please reference Section 8 of BASF's Part A submittal to P-352. Specific to this study, the worksheets "Relevance" and "Evaluation" in the Eco-efficiency model provided to NSF as part of this verification process should be consulted to see the specific values utilized and how they were applied to determine the appropriate calculation factors. Specific to the choice of environmental relevance factors and social weighting factors applied to this study, factors for the USA (national average) were utilized. The environmental relevance values utilized were last reviewed in 2007 and the social weighting factors were recently updated in 2009 by an external, qualified 3<sup>rd</sup> party.

Figure 25 displays the Base Case (BC) eco-efficiency portfolio, which shows the results when all six individual environmental categories are combined into a single relative environmental impact and combined with the life cycle cost impact. Because environmental impact and cost are equally important, the most eco-efficient alternative is the one with the largest perpendicular distance above the diagonal line and the results from this study find that corn Production in 2010 is the more eco-efficient alternative due to its combination of lower environmental burden and having the lowest life cycle cost.






Figure 24. Total socio-eco-efficiency score of the AgBalance™ – Corn Production




Figure 25. Eco-Efficiency Portfolio Base Case – Corn Production

8.5 Scenario Analysis:

In addition to the base case analysis, additional scenarios were evaluated to determine the sensitivity of the studies final conclusions and results to key input parameters. These scenarios are hypothetical technological or operational improvements.

8.5.1 Scenario #1: 5% reduction of corn production yield from the base case in 2010.

In this scenario analysis the yield for the 2010 operations is decreased by 5%, based on other data sources for Iowa corn production showing yield in 2010 at 165 bu/acre (Base case had 174 bu/acre for yield). This scenario shows that yield has a major impact on the analysis since the CB is a fixed amount and any decrease in yield would increase the environmental and cost burden. The socio-economic efficiency score can be seen in Figure 26 and the difference between the two alternatives is less than the base case. Year 2010 is still better than the Year 2000 in all the assessments. Figure 27 shows the Eco-efficiency Portfolio results of Scenario #1 and the changes from the base case. Figure 28 shows the Environmental Fingerprint of Scenario #1, with the Year 2000 being slightly better than the base case but still not better than Year 2010.




Figure 26. Total socio-economic efficiency score corn production – Scenario #1




Figure 27. Eco-Efficiency Portfolio corn production – Scenario #1




Figure 28. Environmental fingerprint corn production – Scenario #1

8.5.2 Scenario #2: Comparison of conventional tillage in 2000, conventional tillage in 2010 and conservative tillage in 2010.

In this scenario analysis, the different tillage options were analyzed individually to show the impact of the tillage processes on the study. The socio-eco-efficiency score can be seen in Figure 29. The impact of the two years tillage operations is quite interesting where Year 2000 Conv. tillage is better than Year 2010 in the Society assessment. However in Ecology and Economy, Year 2010 is still better than Year 2000 independent of the tillage operations and the Total Score is still better in Year 2010. The slight Ecology difference between Conventional and Conservative tillage in 2010 has to do with more pesticide application in Conservative tillage, since the ground is not being worked to kill weeds. There is a slight change in the environmental fingerprint compared to the base case where Year 2010 conservative tillage has a greater use and impact on water use, due to the water use on additional application of pesticides. The cost of the tillage evaluation from the base case has not changed, but the environmental impact does shift based on tillage practices. Figure 30 shows the Eco-efficiency Portfolio results of Scenario #2 and Figure 31 shows the Environmental Fingerprint of Scenario #2.






Figure 29. Total socio-eco-efficiency score corn production – Scenario #2



**Scenario 2:**  
**Production of 1 tonne of Corn in lowa comparing conventional tillage in 2000, conventional tillage in 2010, and conservative tillage in 2010.**



**Figure 30.** Eco-Efficiency Portfolio corn production – Scenario #2



**Figure 31.** Environmental fingerprint corn production – Scenario #2

8.5.3 Scenario #3: Comparison of conservative tillage operations in 2000 versus conservative tillage in 2010.

In this scenario analysis, the conservative tillage for 2000 was compared against conservative tillage in 2010. From the analysis, the conservative tillage has very minimal impact from the base case. There is a minor change in the Society assessment compared to the base case and a little shift in the portfolio where the alternatives move slightly closer. In the environmental fingerprint the scenario analysis is just the same as the base case. Figure 32 shows the socio-economic efficiency score of Scenario #3, Figure 33 shows the Eco-efficiency Portfolio results of Scenario #3 and Figure 34 shows the Environmental Fingerprint of Scenario #3.






Figure 32. Total socio-economic efficiency score corn production – Scenario #3

**Scenario 3:**  
**Production of 1 tonne of Corn in lowa comparing conservative tillage in 2000, versus conservative tillage in 2010.**



**Figure 33.** Eco-Efficiency Portfolio corn production – Scenario #3



**Figure 34.** Environmental fingerprint corn production – Scenario #3

8.5.4 Scenario #4: Yield for 2000 set at the same yield as 2010.

In this scenario analysis, the yield for both 2000 and 2010 was set to the yield data from 2010 (174 bushel/acre). From the analysis, the socio-eco-efficiency score is still better for Year 2010 over Year 2000 in all the assessments, although the Ecology and Economy are much closer and a slight change in Society as compared to Base Case. The Eco-efficiency portfolio in Scenario #4 however shows that Year 2000 was better than Year 2010 compared to the Base Case. For the environmental fingerprint Year 2000 was better for the environmental categories that affect the corn production and not the AgBalance metrics. Figure 35 shows the socio-eco-efficiency score of Scenario #4, Figure 36 shows the Eco-efficiency Portfolio results of Scenario #4 and Figure 37 shows the Environmental Fingerprint of Scenario #4.




Figure 35. Total socio-eco-efficiency score corn production – Scenario #4



**Scenario 4:**  
Production of 1 tonne of Corn in Iowa with yield for 2000 set at the same yield as 2010.




Figure 36. Eco-Efficiency Portfolio corn production – Scenario #4

### Environmental Fingerprint




Figure 37. Environmental fingerprint corn production – Scenario #4

## 9. Data Quality Assessment

### 9.1. Data Quality Statement:

The data used for parameterization of the corn production AgBalance™ was sufficient with most parameters of high data quality. Moderate data is where industry average values or assumptions pre-dominate the value. No critical uncertainties were identified within the parameters and assumptions that could have a significant effect on the results and conclusions. The data is from agricultural production of corn in the state of Iowa and most of the data is from 2000 and 2010 sources. There are a few sources with data before 2000, see Appendix A for data sources and years. Table 13 provides a summary of the data quality for the corn production study.

**Table 13:** Data quality evaluation for Corn production parameters.

<b>Corn Characteristics</b>	<b>Quality Statement</b>	<b>Data Source:</b>
Pioneer 33G26	High	Pioneer Hybrids
Variety - DKC52-59	High	Dekalb Genetics Seed Company
Yield (grain)	High	Iowa State University, FM-1789 (6/01 & 12/11)
Amount of corn seed	High	Doane Research
<b>Plant protection</b>		
N-fertilizer	High	USDA & Iowa State University
P-fertilizer	High	USDA & Iowa State University
K-fertilizer	High	USDA & Iowa State University
Harvest Xtra	High	Monsanto Company, Doanes Research
Roundup	High	Monsanto Company, Doanes Research
Ammonium sulfate	Mod-High	BASF Corp.
Lorsban	High	Dow AgroSciences
Water amounts	High	BASF Corp.
Herbicide Application	Mod-High	Iowa State University
Diesel use	Mod.-High	Iowa State University, PM709
<b>Biodiversity &amp; Soil</b>		
Payments Agro-environmental schemes	Mod.-High	Doane Research
Protected areas	Moderate	Doane Research
Maximum yield potential	Moderate	Iowa State University, File A1-14
Number of endangered species	Mod.-High	www.iucnredlist.org
All soil data for Iowa	Moderate	Iowa State University - ISPAID 7.3 Database
Loss from Wind Erosion	Moderate	NRCS publication, "2007 National Resources Inventory, Soil erosion on cropland"
<b>Cost &amp; Revenue</b>		
Variable Cost	Mod-High	Iowa State University, FM-1789 (6/01 & 12/11)
Fixed Cost	Mod-High	Iowa State University, FM-1789 (6/01 & 12/11)
Machinery cost	Moderate	Iowa State University, FM-1789 (6/01 & 12/11)
Capital cost	High	Iowa State University, FM-1789 (6/01 & 12/11)
Land Lease	High	Iowa State University, FM-1789 (6/01 & 12/11)
Other costs	Mod.-High	Iowa State University, FM-1789 (6/01 & 12/11)
Corn prices	High	Iowa State University, FM-1789 (6/01 & 12/11)
<b>Social</b>		
Accidents and Diseases	Mod.-High	U.S. Ag Census Data, AC-02-A-51 & AC-07-A-51
Workers' wages	Moderate	Iowa State University, Wages and Benefits publication
Employees	Mod.-High	U.S. Ag Census Data, AC-02-A-51 & AC-07-A-51
Professional training	Moderate	Iowa State University, Extension Training
Memberships	Mod.-High	Iowa Corn Grower Association
Land lease	High	U.S. Ag Census Data, AC-02-A-51 & AC-07-A-51
Women Proprietors	High	U.S. Ag Census Data, AC-02-A-51 & AC-07-A-51
Training for agriculture	Moderate	Iowa State University
Old-age insurance	Mod.-High	U.S. IRS
Accidents insurance	Moderate	Iowa State University, FM-1789 (6/01 & 12/11)
Health insurance	Mod.-High	U.S. IRS

## 10. Sensitivity and Uncertainty Analysis

### 10.1. Sensitivity and Uncertainty Considerations:


A sensitivity analysis of the final results indicates that the environmental impacts were more influential or relevant in determining the final relative eco-efficiency positions of the alternatives. This conclusion is supported by reviewing the BIP Relevance (or GDP-Relevance) factor calculated for the study. The BIP Relevance indicates for each individual study whether the environmental impacts or the economic impacts were more influential in determining the final results of the study. For this study, the BIP Relevance indicated that the environmental impacts were significantly more influential in impacting the results than the economic impacts (reference the "Evaluation" worksheet in the Excel model for the BIP Relevance calculation). The main assumptions and data related to environmental impacts were:

- Yield
- Emissions
- Fertilizer Application Rates


As the data quality related to these main contributors were of high to moderate high quality and scenario variations were run related to them (see section 8.4), this strengthened our confidence in the final conclusions indicated by the study. Looking at the relevance factors of the study, see Figure 38, indicates that the impact with the highest environmental relevance was land use, followed by emissions and toxicity potential. This is to be expected, as the study dealt with the production of a crop and the use of fertilizers. When the social weighting factors, Figure 39, are combined with the relevance factors, emissions had the greatest overall impact on the study at 29%, with water emissions being the greatest impact in the emissions category. In the air emissions, GWP and AP are considered the two most important air emissions. The calculation factors, Figure 40, considers both the social weighting factors and the environmental relevance factors, indicate which environmental impact categories were having the largest affect on the final outcome. Calculation factors are utilized in converting the environmental fingerprint results (Figure 20) into the final, single environmental score as reflected in the total socio-eco-efficiency score (Figure 24) as well as the portfolio (Figure 25). The impacts with the highest calculation factors were similar to the environmental relevance factors, with regards to the six main impact categories. The emissions factor was slightly higher than the land use in the calculation factors. The input parameters that were related to these impact categories have sufficient data quality to support a conclusion that this study has a low uncertainty. The social weighting factors considered for this study did influence some minor reprioritization of the impact categories represented in the emissions and air emissions sub-categories.

Most of the input parameters for this study were mainly taken from data gathered at Iowa State University, which would be considered highly credible. The production of corn is an annual process and crops are usually rotated year after

year. In this study, the evaluation was done for one growing season and the next crop was not evaluated.




**Figure 38.** Environmental Relevance factors that are used in the sensitivity and uncertainty analysis.



**Figure 39.** Social weighting factors that are used in the sensitivity and uncertainty analysis.





**Figure 40.** Calculation factors that are used in the sensitivity and uncertainty analysis.

*10.2. Critical Uncertainties:*

There were no significant critical uncertainties from this study that would limit the findings or interpretations of this study. The data quality, relevance and sensitivity of the study support the use of the input parameters and assumptions as appropriate and justified.

**11 Limitations of AgBalance™ Study Results**

*11.1. Limitations:*

These corn production AgBalance™ results and its conclusions are based on the specific comparison of the production, for the described customer benefit, alternatives and system boundaries. Transfer of these results and conclusions to other production methods or products is expressly prohibited. In particular, partial results may not be communicated so as to alter the meaning, nor may arbitrary generalizations be made regarding the results and conclusions.

## 12. References

1. Kölsch, D; Saling, P.; Kicherer A.; Grosse-Sommer, A.; Schmidt, I.; Int. J. Sustainable Development, 2008, 11, 1, 1.
2. Schmidt, I., Meurer, M., Saling, P. Reuter, W., Kicherer, A. and Gensch C-O. '*SEEBALANCE® managing sustainability of products and processes with the socio-eco-efficiency analysis by BASF*, Greener Management International; 2005.
3. Iowa State University, *2000 Iowa Cost and Returns*, FM-1789, June 2001
4. Iowa State University, *2010 Iowa Cost and Returns*, FM-1789, Revised December 2011.
5. Boustead Consulting Ltd UK, The Boustead Model 5.1.2600.2180 LCA database.
6. Intergovernmental Panel on Climate Change, *2006 IPCC Guidelines for National Greenhouse Gas Inventories, Chapter 11: N2O Emissions from Managed Soils, and CO2 Emissions from Lime and Urea Application*, Table 11.1, page 11.11, 2006.
7. Intergovernmental Panel on Climate Change, *2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 4, Chapter 11: N2O Emissions from Managed Soils, and CO2 Emissions from Lime and Urea Application*, Table 11.3, page 11.24, 2006.
8. Klöpffer, W.; Department of the Environment/Federal Environment Agency Austria: Life Cycle Assessment of Genetically Engineered Products based on Comprehensive Assessment of Possible Environmental Impacts, internal BASF Ref.-ID 3411, 1999.
9. Levels of Nonnutritive Substances in Fertilizers, Report to the Legislature, Washington State Department of Agriculture and Washington State Department of Ecology, December 2007 <http://agr.wa.gov/PestFert/Publications/docs/2007Non-NutritiveFertRptFINAL.pdf>
10. FAO Corporate Document Repository, Control of water pollution from agriculture- FAO irrigation and drainage paper 55, Chapter 3: Fertilizers as water pollutants, 1996. <http://www.fao.org/docrep/W2598e/w2598e06.htm>
11. Koellner, T., and Scholz, R., *Assessment of Land Use Impacts on the Natural Environment, Part 1: An Analytical Framework for Pure Land Occupation and Land Use Change*, International Journal LCA 12(1) 16-23, 2007.
12. Koellner, T., and Scholz, R., *Assessment of Land Use Impacts on the Natural Environment, Part 2: Generic Characterization Factors for Local Species Diversity in Central Europe*, International Journal LCA 2006.
13. Saling, P.; Maisch, R.; Silvani, M.; König, N. Int. J. LCA, 10(5), 364-371 (2005).
14. Pfister, S. Koehler, A. and Hellweg, S., *Assessing the Environmental Impacts of Freshwater Consumption in LCA*, Environmental Science & Technology Publication, American Chemical Society, April 23, 2009.

15. Kölsch, D; Saling, P.; Kicherer A.; Grosse-Sommer, A.; Schmidt, I.; Int. J. Sustainable Development, 2008, 11, 1, 1.
16. Schmidt, I., Meurer, M., Saling, P. Reuter, W., Kicherer, A. and Gensch C-O. '*SEEBALANCE®* managing sustainability of products and processes with the socio-eco-efficiency analysis by BASF, Greener Management International; 2005.

**Appendix A:****Data Sources used for input data:****General Data**

- Iowa State University, *2000 Iowa Cost and Returns*, FM-1789, June 2001.
- Iowa State University, *2010 Iowa Cost and Returns*, FM-1789, Revised December 2011.

**Seeding**

- Doane Research website database, [www.doane.com](http://www.doane.com), 2000 and 2010 data.
- Dekalb Seed Corn, DKC52-59 Brand (VT3) brochure, April 2012.
- Maxim® 4FS Fungicide Label, Syngenta Crop Protection, LLC, pages 1-20, 2011.
- Maxim® 4FS Fungicide Safety Data Sheet, Syngenta Crop Protection, LLC, pages 1-5, Nov. 2011.
- ApronXL® Fungicide Label, Syngenta Crop Protection, LLC, pages 1-15, 2012.
- ApronXL® Fungicide Safety Data Sheet, Syngenta Crop Protection, LLC, pages 1-5, Nov. 2010.
- Poncho® 600 Insecticide Label, Bayer CropScience, LP, pages 1-7, 2010.
- Poncho® 600 Insecticide, Safety Data Sheet, Bayer CropScience, LP, pages 1-7, Oct. 2006.

**Tillage and Planting**

- Iowa State University - University Extension, *Machinery Management-Fuel Required for Field Operations*, PM-709, Revised April 2001.

**Fertilizers**

- United States Department of Agriculture (USDA) Economic Research Service Website – <http://www.ers.usda.gov/data-products/fertilizer-use-and-price.aspx#26730>, *U.S. Fertilizer Use and Price*, Tables 7, 8, 10, 12, 14, (May 2011)
- USDA- National Agricultural Statistics Service, *2011 Iowa Agricultural Statistics Bulletin*, Page 71.
- Iowa State University - University Extension, *Machinery Management-Fuel Required for Field Operations*, PM-709, Revised April 2001.

**Plant protection**

- Harness® Xtra Herbicide Label, Monsanto Company, pages 1-9, 2012.
- Harness® Xtra Herbicide Safety Data Sheet, Monsanto Company, pages 1-13, May 2010.
- Roundup Ultra® Herbicide Label, Monsanto Company, pages 1-19, 2010.
- Roundup Ultra® Herbicide Safety Data Sheet, Monsanto Company, pages 1-9, Feb. 2011.
- Roundup Weathermax® Herbicide Label, Monsanto Company, pages 1-54, 2009.
- Roundup Weathermax® Herbicide Safety Data Sheet, Monsanto Company, pages 1-9, Sept. 2008.
- Roundup PowerMax® Herbicide Label, Monsanto Company, pages 1-54, 2010.
- Roundup PowerMax® Herbicide Safety Data Sheet, Monsanto Company, pages 1-9, May 2011.



- Weedone® LV4 EC Herbicide Label, Nufarm Inc., pages 1-14.
- Weedone® LV4 EC Herbicide Safety Data Sheet, Nufarm Inc, pages 1-6, Jan. 2007.
- Lorsban® 15G Insecticide Label, Dow AgroSciences LLC, pages 1-10, Sept. 2008.
- Lorsban® 15G Insecticide Safety Data Sheet, Dow AgroSciences LLC, pages 1-5, June 2006.
- Iowa State University - University Extension, *Machinery Management-Fuel Required for Field Operations*, PM-709, Revised April 2001.

## Harvesting

- Iowa State University - University Extension, *Machinery Management-Fuel Required for Field Operations*, PM-709, Revised April 2001.

## Biodiversity

- Doane Research website database, [www.doane.com](http://www.doane.com), 2000 and 2010 data.
- Iowa State University - University Extension, *Ag Decision Maker - Iowa Corn and Soybean County Yields, File A1-14*, Table 1, March 2012.
- IUCN red list website, [www.iucnredlist.org](http://www.iucnredlist.org).

## Soil

- Iowa State University website database, [www.extension.iastate.edu/soils/ispaid](http://www.extension.iastate.edu/soils/ispaid), *ISPAID 7.3 Database and ISPAID 7.3 Database Sorted by Count, 2000 and 2010*.
- National Resource Conservation Service, *2007 National Resources Inventory-Soil Erosion*, April 2010.

## Cost

- Iowa State University, *2000 Iowa Cost and Returns*, FM-1789, June 2001.
- Iowa State University, *2010 Iowa Cost and Returns*, FM-1789, Revised December 2011.
- Iowa State University - University Extension, *Estimating Farm Machinery Costs*, pages 1-8, (Nov. 2009)
- Iowa State University - University Extension, *Ag Decision Maker - Cash Corn and Soybean Prices*, page 2, Table 1, (Feb. 2010)

## Social

- U.S. Department of Agriculture, *2002 Census of Agriculture*, AC-02-A-51, June 2004.
- U.S. Department of Agriculture, *2007 Census of Agriculture*, AC-07-A-51, Dec. 2009.
- Iowa State University - University Extension, *Wages and Benefits for Farm Employees*, FM 1862, March 1999.
- Iowa State University - University Extension, *Wages and Benefits for Farm Employees*, FM 1862, June 2006.
- Iowa State University – Departmental Data, *Fact Book 2000-2001*.
- Iowa State University – Departmental Data within College, *Fact Book 2010-2011*.
- Iowa State University, *2000 Iowa Cost and Returns*, FM-1789, June 2001.
- Iowa State University, *2010 Iowa Cost and Returns*, FM-1789, Revised December 2011.