

Equivalence classes of permutations modulo excedances

JEAN-LUC BARIL, TOUFIK MANSOUR AND ARMEN PETROSSIAN

We introduce a new equivalence relation on permutations where two permutations are equivalent if and only if they coincide on their excedance sets. This paper studies equivalence classes for several subsets of permutations. Enumerating results are presented for permutations, cycles and permutations avoiding one or two patterns of length three. Also, an open question are proposed.

Keywords: permutations; equivalence class; excedance; pattern; Bell, Motzkin, Catalan, Fibonacci numbers.

1. Introduction and notations

Let S_n be the set of permutations of length n , *i.e.*, all one-to-one correspondences from $[n] = \{1, 2, \dots, n\}$ into itself. The *one-line notation* of a permutation $\pi \in S_n$ is $\pi_1\pi_2 \cdots \pi_n$ where $\pi_i = \pi(i)$ for $i \in [n]$. A *cycle* in S_n is a n -length permutation π such that there exist some indices i_1, i_2, \dots, i_n with $\pi(i_1) = i_2, \pi(i_2) = i_3, \dots, \pi(i_{n-1}) = i_n$ and $\pi(i_n) = i_1$. According to the context, a cycle will be also denoted by its *cyclic notation* $\pi = \langle i_1, i_2, \dots, i_n \rangle$. Let $C_n \subset S_n$ be the set of all cycles of length n .

Let π be a permutation in S_n . An *excedance* of π is a position i , $1 \leq i \leq n$, such that $\pi(i) > i$. The set of excedances of π will be denoted $E(\pi)$. For instance, if $\pi = 4\ 1\ 6\ 3\ 5\ 7\ 2$ then $E(\pi) = \{1, 3, 6\}$. The *graphical representation* of π is the set of points in the plane at coordinates (i, π_i) for $i \in [n]$. We define its associated rook placement $R(\pi)$ on the triangular board consisting of the cells located atop the diagonal $y = x$ in the graphical representation of the permutation. For short, $R(\pi)$ will be called *excedance board* of π . Obviously, a point (i, π_i) appears in the excedance board of π if and only if i is an excedance of π . See Figure 1 for the graphical representation of a permutation with its associated excedance board.

We consider the equivalence relation \sim on S_n in which two permutations π and σ are considered to be equivalent if they coincide on their excedance sets, *i.e.*, $E(\pi) = E(\sigma)$ and $\pi(i) = \sigma(i)$ for $i \in E(\pi)$, or again π and σ have the same excedance board. Let $\text{Eq}(\pi, S_n)$ denote the set of permutations being equivalent to π in S_n . For instance, we have $\text{Eq}(43215, S_5) = \{43215, 43125\}$. The set (resp. the cardinality of the set) of distinct equivalence classes in S_n is denoted S_n^\sim (resp. $\#S_n^\sim$). All these definitions will remain available by replacing the set S_n with a subset of S_n .




Figure 1: Graphical representation of $\sigma = 4\ 1\ 6\ 3\ 5\ 7\ 2$ with its excedance board and its associated partition.

The theory of permutation statistics has a long history and has grown at a rapid place in the last few decades. Two classical statistics have been received lot of attention, namely number of descents and number of excedances in a permutation (see [5] and references therein). Recently, Elizalde [6] studied number of permutations that avoid a fixed set T of patterns according to the number of fixed points and excedances. In this paper, we study the number of distinct equivalence classes in either S_n^{\sim} or $S_n^{\sim}(T)$ with a fixed set T of patterns.

A permutation $\pi \in S_n$ *avoids* the pattern $\tau \in S_k$ if and only if there does not exist a sequence of indices $1 \leq i_1 < i_2 < \dots < i_k \leq n$ such that $\pi(i_1)\pi(i_2)\dots\pi(i_k)$ is order-isomorphic to τ (see [13, 14]). We denote by $S_n(\tau)$ the set of permutations of S_n avoiding the pattern τ . For example, if $\tau = 123$ then $52143 \in S_5(\tau)$ while $21534 \notin S_5(\tau)$. Many classical sequences in combinatorics appear as the cardinality of pattern-avoiding permutation classes. A large number of these results were firstly obtained by West and Knuth [9, 12, 13, 14, 15, 16] (see books of Kitaev [8] and Mansour [11]).

In Section 2, we prove that equivalence classes for S_n are enumerated by the n th Bell number. We also prove that the number of classes in C_n is given by the sequence the sequence of the number of irreducible set partitions of $[n]$ (see A074664 in the on-line Encyclopedia of Integer Sequences [17]). In Section 3, we give enumerating results for permutations avoiding one pattern of length three (see Table 1). When the pattern is 132, the number of classes is given by the Motzkin numbers which provides a kind of discrete continuity between Catalan and Motzkin numbers. The case where permutations avoid the pattern 123 remains an open question. In Section 4, we investigate permutations avoiding two patterns of length three (see Table 3).

2. Equivalence classes for S_n and C_n

A *partition* of $[n]$ is any collection of non-empty pairwise disjoint subsets, called *blocks*, whose union is $[n]$. Let \mathcal{P}_n be the set of partitions of $[n]$. A partition Π is said to be in *standard form* if it is written $\Pi = B_1/B_2/\dots$, where the blocks B_i are arranged so that their smallest

Pattern	Sequence	Sloane	$a_n, n \geq 1$
{}	B_n	A000110	1, 2, 5, 15, 52, 203, 877, 4140, 21147
{132}, {213}	M_n	A001006	1, 2, 4, 9, 21, 51, 127, 323, 835
{123}	Open question	<i>New</i>	1, 2, 4, 8, 19, 41, 98, 221, 526
{312}, {321}	$\frac{1}{n+1} \binom{2n}{n}$	A000108	1, 2, 5, 14, 42, 132, 429, 1430, 4862
{231}		<i>New</i>	1, 2, 4, 9, 21, 50, 121, 296, 729

Table 1: Number of equivalence classes for permutations avoiding at most one pattern of \mathcal{S}_3 , where B_n (resp. M_n) are the Bell numbers (resp. Motzkin numbers) defined by the generating function $\sum_{k \geq 0} \frac{x^k}{\prod_{j=1}^k (1-jx)}$ (resp. $\frac{1-x-\sqrt{1-2x-3x^2}}{2x^2}$).

elements are in increasing order. For convenience, we assume also that elements in a same block are arranged in increasing order. An *atomic partition* is a partition that does not have a proper subset of blocks with a union equal to a subset $[j]$ with $1 \leq j \leq n-1$.

From a permutation $\pi \in S_n$, we associate the partition $\Pi \in \mathcal{P}_n$ defined as follows. Two elements $x < y$ belong to the same block in Π if and only if there exist $k \geq 1$ and $i_1 = x < i_2 < \dots < i_k = y$ such that $\pi_{i_j} = i_{j+1}$ for $1 \leq j \leq k-1$. For instance, the partition associated to $\pi = 4163572$ is $\Pi = 14/2/367/5$ (see Figure 1). Indeed, if $i_1 = 1$ then we have $i_1 = 1 < i_2 = \pi_{i_1} = 4$ and $\pi_{i_2} = 3 < 4$ which induces that 14 is a block; if $i_1 = 2$ then we have $\pi_{i_1} = 1 < 2$, so 2 is a block; if $i_1 = 3$ then we have $i_1 = 3 < i_2 = \pi_{i_1} = 6 < i_3 = \pi_{i_2} = 7$ and $\pi_{i_3} = 2 < 7$, so 367 forms a block, and so on. Notice that this definition appears as a counterpart for permutations of the partition associated to a rook placement on a *Ferrers board* (see Stanley [18], p. 75, or Mansour [10], p. 99).

2.1. For S_n - Bell numbers

Theorem 2.1. *The sets S_n^\sim , $n \geq 1$, are enumerated by the Bell numbers.*

Proof. Two permutations lie in a same equivalence class $\text{Eq}(\pi, S_n)$ if and only if they have the same excedance board $R(\pi)$. Then they have the same associated partition. Moreover, the associated partitions of two non-equivalent permutations are different.

Conversely, let $\Pi = B_1/B_2/\dots/B_k$ be a partition of $[n]$ in the standard form and such that each block B_i , $1 \leq i \leq k$ is arranged in increasing order. We define the permutation $\pi \in S_n$ such that $\pi_i = j$ if j is just after i into a same block (j is the smallest element greater than i in the block) and the image by π of the largest element of a block is the smallest element of this block. For instance, if $\Pi = 14/2/367/5$ we obtain the permutation $\pi = 4261573$. By construction, this process defines a permutation in S_n such that Π is its associated partition. So, there is a bijection between S_n^\sim and the set \mathcal{P}_n of partitions of $[n]$. Therefore, S_n^\sim is enumerated by the n -th Bell number. \square

2.2. For C_n - A074664

Lemma 1. *The associated partition of a cyclic permutation $\pi \in C_n$ is atomic.*

Proof. For $\pi \in C_n$, let $\Pi = B_1/B_2/\cdots/B_k$ be its associated partition in standard form. For a contradiction, we assume that Π is not atomic, *i.e.*, there exists a proper subset of blocks B_1, B_2, \dots, B_ℓ , $\ell < k$ with a union equal to a subset $[j]$ for some $j \in [n-1]$. Then for any $i \in [j]$, we have $\pi_i \in [j]$. Indeed, if there was $i \leq j$ such that $\pi_i > j$, then i would be an excedance and π_i would be in a same block. This case does not occur since the hypothesis induces that the block containing i and π_i is included in $[j]$. Therefore, the image by π of the interval $[j]$, $j < n$, still remains $[j]$, which contradicts the fact that π is a cycle in C_n . \square

Theorem 2.2. *The sets C_n^\sim , $n \geq 1$, are enumerated by the sequence A074664 in [17].*

Proof. The previous lemma proves that the associated partition of a cycle in C_n is necessarily atomic. Moreover, the associated partitions of two non-equivalent cycles are different. Now, we will prove that for any atomic partition $\Pi \in \mathcal{P}_n$, there exists a cycle $\pi \in S_n$ having Π as associated partition (see the definition on the previous page). Let $\Pi = B_1/B_2/\cdots/B_k$ be an atomic partition (not necessarily in standard form). Let us assume that $1 \in B_1$ and put $i_1 = 1$. For $1 \leq i \leq k$, we denote by $M(B_i)$ (resp. $m(B_i)$) the maximum (resp. minimum) element in the block B_i . In the case where $M(B_1) \neq n$, the atomicity of Π induces that there is a block B_{i_2} such that $1 < m(B_{i_2}) < M(B_{i_1}) < M(B_{i_2})$. We choose the block B_{i_2} having its smallest element the lowest possible. Iterating this process by replacing B_{i_1} with B_{i_2} , we exhibit a sequence of blocks $B_{i_1}, B_{i_2}, \dots, B_{i_r}$ such that $m(B_{i_j}) < m(B_{i_{j+1}}) < M(B_{i_j}) < M(B_{i_{j+1}})$ for $1 \leq j \leq r-1$ and such that B_{i_r} contains the value n . Let π be the cycle in C_n defined as follows: $\pi = \langle B_{i_1} B_{i_2} \cdots B_{i_r} A_1 A_2 \cdots A_s \rangle$ where each block B_{i_j} , $1 \leq j \leq r$ has its elements in increasing order, and the blocks A_i , $1 \leq i \leq s$, are the remaining blocks of the partition Π ordered by their decreasing minima and each of them ordered in increasing order. By construction π is a cycle in C_n , and it is straightforward to see that Π is its associated partition. Using Lemma 1, we have exhibited a constructive bijection between C_n^\sim , $n \geq 1$, and the subset of atomic partitions in \mathcal{P}_n which is enumerated by the sequence A074664 in [17] (see Bergeron and Zabrocki [4]). \square

A permutation $\pi = \pi_1 \pi_2 \cdots \pi_n \in S_n$ is called *indecomposable* if there does not exist $i < n$ such that $\pi_1 \pi_2 \cdots \pi_i$ is a permutation of $[i]$. Let Ind_n , $n \geq 1$, be the sets of indecomposable permutations of length n . Lemma 1 remains valid if we replace cyclic permutation with indecomposable permutation. Moreover, since a cyclic permutation is necessarily indecomposable, the proof of the following theorem is obtained, *mutatis mutandis*, from the proof of Theorem 2.2.

Theorem 2.3. *The sets Ind_n^\sim , $n \geq 1$, are enumerated by the sequence A074664 in [17].*

3. Equivalence classes for $S_n(\alpha)$ with $\alpha \in S_3$

In this section, we study enumerating results about the equivalence classes of sets of permutations avoiding one pattern of length three.

For the pattern 321, the excedances of $\pi \in S_n(321)$ determine π completely which means that every permutation $\pi \in S_n(321)$ is the unique element in its equivalence class. Therefore, the cardinality of the set $S_n^{\sim}(321)$ is given the n -th Catalan number (see sequence A000108 in [17]).

For the pattern 312, we use the Simion-Schmidt bijection modulo symmetry from $S_n(321)$ to $S_n(312)$ (see [13]). Since this bijection preserves the position of points (i, π_i) whenever $\pi_i > i$, we directly conclude that the cardinality of the set $S_n^{\sim}(312)$ is given by the cardinality of $S_n(312)$, that also is the n -th Catalan number. We have not succeeded to find the cardinality of $S_n^{\sim}(123)$ for any $n \geq 1$ (first values are 1, 2, 4, 8, 19, 41, 98, 221, 526). So, we leave open this problem.

In order to study other patterns, we define some other equivalence relations between two permutations by extending the definition of an excedance. Let π be a permutation in S_n . A k -*excedance* of π , $1 - n \leq k \leq n$, is a position i , $1 \leq i \leq n$, such that $\pi(i) > i - k$. The set of k -excedances of π will be denoted $E^k(\pi)$. For instance, if $\pi = 312$ then $E^{-2}(\pi) = \emptyset$, $E^{-1}(\pi) = E^0(\pi) = E^1(\pi) = \{1\}$ and $E^2(\pi) = \{3\}$. Obviously, for any $\pi \in S_n$, we have $E^0(\pi) = E(\pi)$, $E^{1-n}(\pi) = \emptyset$ and $E^n(\pi) = [n]$. For any k , $1 - n \leq k \leq n$, we define the k -*equivalence relation* \sim^k on S_n in which two permutations π and σ are equivalent if and only if they coincide on their k -excedance sets, *i.e.*, $\pi(i) = \sigma(i)$ for $i \in E^k(\pi) = E^k(\sigma)$. Clearly, when $k = 0$ we retrieve the equivalence relation defined in the introduction. Let $\text{Eq}^k(\pi, S_n)$ denote the set of permutations in S_n k -equivalent to π ; we call it the k -*class* of π . For instance, we have $\text{Eq}^0(43215, S_5) = \text{Eq}(43215, S_5) = \{43215, 43125\}$, $\text{Eq}^2(43215, S_5) = \{43215\}$ and $\text{Eq}^{-4}(43215, S_5) = S_5$. The set of distinct k -classes in S_n is denoted $S_n^{\sim^k}$.

3.1. The sets $S_n^{\sim}(132)$ and $S_n^{\sim}(213)$ - Motzkin

We focus our study on the pattern 132 and the symmetry *reverse* \rightarrow *complement* \rightarrow *inverse* will provide the result for 213.

Theorem 3.1. *The sets $S_n^{\sim}(132)$, $n \geq 1$, are enumerated by the Motzkin numbers.*

Proof. For $k \geq 0$, let M_n^k be the number of k -classes in $S_n(132)$. Let π be a permutation in $S_n(132)$. Since π avoids the pattern 132, π can be written $\pi = \sigma\gamma i$, $1 \leq i \leq n$, where γi is a subsequence of $[i]$ avoiding the pattern 132 and σ is obtained from a permutation $\sigma' \in S_{n-i}(132)$ by adding i on all its entries, *i.e.*, $\sigma(j) = i + \sigma'(j)$ for $1 \leq j \leq n - 1 - i$.

We distinguish two cases: (1) $i \geq n - k$, and (2) $i \leq n - k - 1$.

Case 1. In this case, n is either a k -excedance (*i.e.*, $\pi_n = i \geq n - k + 1$) or $\pi_n = i = n - k$. The decomposition of $\pi = \sigma\gamma i$ is illustrated by the left part of Figure 2. The blue (resp. green) square corresponds to σ (resp. γ), and the filled gray area does not contain any point (j, π_j) .

It is straightforward and crucial to see that for a given i , $i \geq n-k$, the above decomposition of π induces that:

an integer $j > n-i$ is a k -excedance of π if and only if $j-n+i$ is a $(k+i-n)$ -excedance of γi where γi is considered as a permutation in $S_i(132)$.

Using the decomposition of $\pi = \sigma\gamma i$, the number of k -classes having a representative ending by $i \geq n-k$ is equal to the product between the number of $(k+i-n)$ -classes of $S_{i-1}(132)$ and the number of subsequences σ of length $n-i$ avoiding 132, that is $M_{i-1}^{k+i-n} \cdot c_{n-i}$ where $c_{n-i} = \frac{1}{n-i+1} \binom{2(n-i)}{n-i}$ is the $(n-i)$ th Catalan number. Varying i from $n-k$ to n , the number of k -classes having a representative with a last element greater or equal than $n-k$ is exactly $M_{n-1}^k c_0 + M_{n-2}^{k-1} c_1 + \dots + M_{n-k-1}^0 c_k$.

Now, let us examine the second case.

Case 2. We assume $i \leq n-k-1$ and we will prove that each k -class contains a permutation having its last element equal to one. Let π be a permutation such that $\pi_n = i \leq n-k-1$ and let us assume that $\pi_n = i \neq 1$. Using the decomposition $\pi = \sigma\gamma i$, the value 1 appears in γ , *i.e.*, $\pi_j = 1$ implies $j \geq n-i+1$. Since π avoids 132, all values after $\pi_j = 1$ appear in increasing order in the one-line representation of π and are less or equal than i . This implies that for $\ell \geq j$, π_ℓ are less or equal than $i - (n-\ell)$.

Now, let us define the permutation $\pi' \in S_n(132)$ such that $\pi'_n = 1$, $\pi'_\ell = \pi_\ell$ if $\ell \in [j-1]$, and $\pi'_\ell = \pi_{\ell+1}$ for $\ell \in [j, n-1]$. By construction, $\pi' \in S_n(132)$ and π belongs to the same k -class. Thus, the number of k -classes (in this case) is equal to the number of $(k+1)$ -classes in $S_{n-1}(132)$ (we omit the last value 1 of π'), which is given by M_{n-1}^{k+1} (see the right part of Figure 2).

In accordance with the two previous cases, we obtain the following recursive formula for $k \geq 0$, $n \geq 1$ and $n > k$:

$$M_n^k = M_{n-1}^{k+1} + M_{n-1}^k c_0 + M_{n-2}^{k-1} c_1 + \dots + M_{n-k-1}^0 c_k$$

anchored with $M_n^n = c_n$ for $n \geq 0$, where c_n is the n th Catalan number (see A000108 in [17]).

Setting $D^n(x) = \sum_{k \geq 0} M_{k+n}^k x^k$ for $n \geq 1$, the bivariate generating function $F(x, u)$ where the coefficient of $x^k u^n$ is the number of k -classes in $S_{n+k}(132)$ satisfies $F(x, u) = \sum_{n \geq 1} D^n(x) u^n$. A simple calculation from the previous recurrence relation provides the functional equation for $n \geq 3$:

$$D^n(x) = C(x)D^{n-1}(x) + \frac{D^{n-2}(x) - D^{n-2}(0)}{x}$$

with $D^1(x) = \frac{C(x)-1}{x}$ and $D^2(x) = \frac{D^1(x)-1}{x}$ where $C(x) = \frac{1-\sqrt{1-4x}}{2x}$ is the generating function for the Catalan numbers.

With this last relation, we obtain:

$$F(x, u)(x - C(x)xu - u^2) = D^1(x)xu + D^2(x)xu^2 - C(x)x D^1(x)u^2 - u^2 F(0, u).$$

Using the kernel method (see [1, 9]), we compute

$$x = \frac{u}{2}(1 + u - 2u^2 - \sqrt{1 - 2u - 3u^2}).$$

This induces $F(0, u) = \frac{1-u-\sqrt{1-2u-3u^2}}{2u^2}$ which is the generating function for the Motzkin numbers. Therefore, M_n^0 is the n th term of the Motzkin sequence.

Finally, a simple calculation provides

$$F(x, u) = \frac{u - u\sqrt{1 - 4x} - ux + u^2 - u^2\sqrt{1 - 4x} - x + x\sqrt{1 - 2u - 3u^2}}{x(2x - u + u\sqrt{1 - 4x} - 2u^2)}.$$

□




Figure 2: Illustration of the two cases studied in the proof of Theorem 4.

The bivariate generating function obtained in the proof of Theorem 3.1 provides a kind of *discrete continuity* between the well-known Catalan and Motzkin sequences (see Table 2). See [3] for another discrete continuity between these sequences and [2] for a discrete continuity between Fibonacci and Catalan sequences. Using Maple, the first terms of the Taylor expansion of $F(x, u)$ are $u + 2u^2 + 4u^3 + 9u^4 + 21u^5 + 51u^6 + 127u^7 + 2xu + 5xu^2 + 12xu^3 + 30xu^4 + 76xu^5 + 196xu^6 + 5x^2u + 14x^2u^2 + 37x^2u^3 + \dots$

Table 2: Number M_n^k of k -classes in $S_n(132)$ for some n and k , $n > k \geq 0$.

$k \setminus n$	1	2	3	4	5	6	7
0	1	2	4	9	21	51	127
1		2	5	12	30	76	196
2			5	14	37	99	265
3				14	42	118	331
4					42	132	387
5						132	429

3.2. The set $S_n^\sim(231)$

Theorem 3.2. *The generating function for the number of classes in $S_n(231)$, $n \geq 1$, is given by the continued fraction:*

$$1 + [a_1] - \frac{1}{1 + [a_2] - \frac{x}{1 + [a_3] - \frac{x}{1 + [a_4] + \dots}}}$$

where $[a_\ell] = \frac{x^2(1-x^{\ell-1})}{1-x}$ for $\ell \geq 1$.

Proof. For $k \leq 0$, let M_n^k be the number of k -classes in $S_n(231)$. Recall that two permutations σ and π lie in a same k -class if and only if they coincide on positions i satisfying $\pi_i > i - k$ or $\sigma_i > i - k$. Let π be a permutation in $S_n(231)$ and denote by i , $1 \leq i \leq n$, the index such that $\pi_i = n$. Since π avoids the pattern 231, π can be written $\pi = \sigma n \gamma$, where σ is a permutation of $[i - 1]$ avoiding the pattern 231 and γ is obtained from a permutation $\gamma' \in S_{n-i}(231)$ by adding $i - 1$ on all its entries, i.e., $\gamma(j) = i - 1 + \gamma'(j)$ for $i + 1 \leq j \leq n$.

We distinguish two cases: (1) $i \leq n - 1 + k$, and (2) $i > n - 1 + k$.

Case 1. In this case, i is a k -excedance since $\pi_i = n > i - k + 1 > i - k$. Notice that for a given i , $i \leq n - 1 + k$, the above decomposition of π induces that:

an integer $j \neq i$ is a k -excedance of π if and only either j is a k -excedance of σ , or $j - i + 1$ is a $(k - 1)$ -excedance of γ' considered as a permutation in $S_{n-i}(231)$.

Thus, the number of k -classes having a representative with n in position i is equal to the product $M_{i-1}^k \cdot M_{n-i}^{k-1}$. Varying i from 1 to $n - 1 + k$, the number of k -classes in this case is exactly $\sum_{i=0}^{n-2+k} M_{n-1-i}^{k-1} M_i^k$.

Now, let us examine the second case.

Case 2. We have $i > n - 1 + k$ and i is not a k -excedance. We will prove that the k -class of π contains a permutation having its last element equal to n .

Let us consider the permutation π' defined by $\pi' = \sigma\gamma n$. It is straightforward to check that π' avoids 231 and that π' belongs to the same k -class of π . Thus, the number of k -classes (in this case) is equal to the number of k -classes having a representative ending with n , which is given by M_{n-1}^k . In accordance with the two previous cases, we obtain the following recursive formula for $k \leq 0$ and $n \geq 1$:

$$\begin{cases} M_n^k = M_{n-1}^k + \sum_{\ell=0}^{n-2+k} M_\ell^k M_{n-1-\ell}^{k-1} & \text{for } n-1 > -k, \\ M_n^k = 1 & \text{for } n-1 \leq -k. \end{cases}$$

Let $D^k(x) = \sum_{n \geq 0} M_n^k x^n$ be the generating function for the number of k -classes in $S_n(231)$. Some elementary calculations from the previous recurrence relation provide the following functional equation for $k \leq 0$:

$$D^k(x) \left(1 + x \left(\frac{1-x^{-k}}{1-x} - 1 \right) - x D^{k-1}(x) \right) = 1.$$

Finally, we obtain $D^k(x)$ expressed as a continued fraction

$$\frac{1}{1 + [a_{-k+1}] - \frac{x}{1 + [a_{-k+2}] - \frac{x}{1 + [a_{-k+3}] - \frac{x}{1 + [a_{-k+4}] + \dots}}}}$$

where $[a_\ell] = \frac{x^2(1-x^{\ell-1})}{1-x}$ for $\ell \geq 1$. When $k = 0$, we obtain the expected result. \square

By the equation of $D^k(x)$ (see line 4), we see that

$$\lim_{k \rightarrow -\infty} D^k(x) \left(1 - x + x \frac{1-x^{-k}}{1-x} - x D^{k-1}(x) \right) = 1.$$

Let $D(x) = \lim_{k \rightarrow -\infty} D^k(x)$ and assume that $|x| < 1$. Then $D(x) \left(1 - x + \frac{x}{1-x} - x D(x) \right) = 1$, which implies that either $D(x) = 1/(1-x)$ or $D(x) = (1-x)/x$. Since $D^k(0) = 1$ for all $k < 0$, we have that $D(0) = 1$. Hence, $D(x) = \lim_{k \rightarrow -\infty} D^k(x) = \frac{1}{1-x}$.

Finally, the first terms of the Taylor expansion of the continued fraction $D^0(x)$ can be obtained by replacing the rest of the continued fraction by its limit $\frac{1}{1-x}$. More precisely, if we

need the first five terms then we expand the rational fraction

$$\frac{1}{1 + [a_1] - \frac{x}{1 + [a_2] - \frac{x}{1 + [a_3] - \frac{x}{1 + [a_4] + \frac{x}{1-x}}}}}$$

which provides $1 + x + 2x^2 + 4x^3 + 9x^4 + \mathcal{O}(x^5)$.

4. Equivalence classes for $S_n(\alpha, \beta)$ with α and β in S_3

In this part, we give enumerating results for classes of permutations avoiding two patterns of length three. Below, we present proofs for the most interesting cases which appear in the last three rows of Table 3, *i.e.*, for the cases where the equivalence classes are not all reduced to a single permutation. All other cases can be easily proved using similar arguments or by applying classical symmetries on permutations (inverse, reverse, and complement).

4.1. The set $S_n^\sim(123, 231)$ - A096777

Theorem 4.1. *The number of classes in $S_n^\sim(123, 231)$, $n \geq 1$, is given by*

$$1 + \frac{1}{2} \cdot \lceil \frac{n-2}{3} \rceil \cdot (1 - 2n + 3 \lceil \frac{n-2}{3} \rceil) + \lfloor \frac{n}{2} \rfloor \cdot (2n - 1 - 2 \lfloor \frac{n}{2} \rfloor)$$

(see sequence A096777 in [17]).

Proof. Let π be a permutation in $S_n(123, 231)$. For $k \in [n+1]$, we define the subsequence $\sigma(k) = n(n-1) \dots (n-k+1)$ if $k \leq n$, and $\sigma(k)$ is empty if $k = n+1$. Thus, a permutation $\pi \in S_n(123, 231)$ has a unique decomposition of the form $\pi = \sigma(k)\gamma(\ell)\delta$ where $\delta = (n-k) \dots (\ell+1)$ and $\gamma(\ell) = \ell(\ell-1) \dots 21$ with $k \in [n+1]$ and $\ell \in [n-k-1]$ (see Figure 3 for an illustration of this decomposition).

Let a_n (respectively b_n^k) be the number of classes in $S_n(123, 231)$ (respectively starting with $\sigma(k)$ for $k \leq \lfloor \frac{n}{2} \rfloor$). The above decomposition induces that

$$a_n = 1 + \sum_{k=1}^{\lfloor \frac{n}{2} \rfloor - 1} b_n^k + (n-1)$$

for $n \geq 1$. Indeed, if $k \geq \lfloor \frac{n}{2} \rfloor$ then $\sigma(k)$ fills the semi-diagonal $y = n - x + 1$ in the excedance board, and π and $n(n-1) \dots 21$ belong to the same class. There is only one class whenever




Figure 3: Illustration of the decomposition of a permutation in $S_n(123, 231)$.

$k \geq \lfloor \frac{n}{2} \rfloor$. The second term $\sum_{k=1}^{\lfloor \frac{n}{2} \rfloor - 1} b_n^k$ corresponds to permutations having a non-empty $\sigma(k)$ that does not fill the semi-diagonal, *i.e.*, $1 \leq k \leq \lfloor \frac{n}{2} \rfloor - 1$; and the last term $n - 1$ enumerates classes of permutations having an empty $\sigma(k)$.

Now, we will calculate b_n^k .

For convenience, we will say that γ (resp. δ) intersects the excedance board of π if the associated block of γ (resp. δ) illustrated in red (resp. green) in Figure 3 has a non-empty intersection with the excedance board of π .

Using Figure 3, a simple observation provides that γ and δ does not intersect the excedance board if and only if $\ell \leq k + 1$ and $\ell + n - \ell - k \leq \ell + k + 1$ which is equivalent to $n - 2k - 1 \leq \ell \leq k + 1$. This case can occur whenever $k + 1 \geq n - 2k - 1$, that is $k \geq \lceil \frac{n-2}{3} \rceil$.

We distinguish two cases.

Case 1. If $k \in [1, \lceil \frac{n-2}{3} \rceil - 1]$ then there is at least one block issue from γ or δ that intersects the excedance board of π . This means that γ and δ are entirely determined by ℓ that describes the interval $[n - k - 1]$. Therefore, in this case $b_n^k = n - k - 1$ and this case contains exactly $\sum_{k=1}^{\lceil \frac{n-2}{3} \rceil - 1} (n - k - 1)$ classes.

Case 2. If $k \geq \lceil \frac{n-2}{3} \rceil$ then there exists ℓ , $n - 2k - 1 \leq \ell \leq k + 1$, such that the two blocks associated to γ and δ do not intersect the excedance board of π . Moreover, all permutations obtained by varying ℓ from $n - 2k - 1$ to $k + 1$ lie in a unique class for a given k . Now, let us assume that either $\ell \in [n - 2k - 2]$ or $\ell \geq k + 2$. If $\ell \in [n - 2k - 2]$ then the block associated to γ intersects the excedance board of π that also determines δ . The number of classes in this subcase is exactly $n - 2k - 2$ for a given k , $k \geq \lceil \frac{n-2}{3} \rceil$. If $\ell \geq k + 2$ then the block associated to δ intersects the excedance board of π that also determines γ . The number of classes in this subcase is exactly $n - k - 1 - (k + 2) + 1 = n - 2k - 2$. So, we have $b_n^k = 2(n - 2k - 2) + 1$ and the total number of classes satisfying Case 2 is exactly $\sum_{k=\lceil \frac{n-2}{3} \rceil}^{\lfloor \frac{n}{2} \rfloor - 1} (2(n - 2k - 2) + 1)$.

In accordance with these two cases, we finally obtain

$$a_n = 1 + n - 1 + \sum_{k=1}^{\lceil \frac{n-2}{3} \rceil - 1} (n - k - 1) + \sum_{k=\lceil \frac{n-2}{3} \rceil}^{\lfloor \frac{n}{2} \rfloor - 1} (2(n - 2k - 2) + 1).$$

With a simple calculation on this formula, we obtain the expected result and the sequence A096777 in [17]. \square

4.2. The set $S_n^\sim(132, 231)$ - Fibonacci

Theorem 4.2. *The sets $S_n^\sim(132, 231)$, $n \geq 1$, are enumerated by the Fibonacci sequence (the n -th Fibonacci number F_n is defined by $F_n = F_{n-1} + F_{n-2}$ with $F_0 = F_1 = 1$, see sequence A000045 in [17]).*

Proof. For $n \geq 1$, we denote by a_n the cardinality of $S_n^\sim(132, 231)$. Let π be a permutation avoiding 132 and 231. Thus, π can be written either (1) $\pi = \pi_1 \cdots \pi_{n-1}n$ or (2) $\pi = n\pi_2 \cdots \pi_n$. The number of classes in $S_n(132, 231)$ satisfying case (1) is also the number a_{n-1} (we omit the last value n). Now, let us study the second case. There exists $j \geq 2$ such that $\pi_j = 1$ and $\pi_1 = n > \pi_2 > \cdots > \pi_j = 1 < \pi_{j+1} < \cdots < \pi_n$. It is straightforward to see that $\pi_i \leq i - 1$ for $i \geq j$. So, we set $\pi' = (\pi_2 - 1) \cdots (\pi_{j-1} - 1)(\pi_{j+1} - 1) \cdots (\pi_n - 1)$. Obviously, $\pi' \in S_{n-2}(132, 231)$ and there is a one-to-one correspondence between classes of $S_n(132, 231)$ having n in first position, and classes of $S_{n-2}(132, 231)$. Finally the number of classes in $S_n(132, 231)$ (in this case) is equal to a_{n-2} (we omit n and 1 in π). According to the two previous cases, we have $a_n = a_{n-1} + a_{n-2}$ anchored with $a_1 = 1$ and $a_2 = 2$ which defines the Fibonacci sequence. \square

4.3. The set $S_n^\sim(123, 132)$ - A052955

Theorem 4.3. *The sets $S_n^\sim(123, 132)$, $n \geq 1$, are enumerated by $2^{\frac{n+1}{2}} - 1$ if n odd, and by $3 \cdot 2^{\frac{n}{2}-1} - 1$ otherwise (see sequence A052955 in [17]).*

Proof. Let π be a permutation avoiding 123 and 132. Let $i_1 < i_2 < \cdots < i_k$ be the indices of the right-to-left maxima of π , i.e., the indices i_j , $j \in [k]$, such that $\pi(i_j) > \pi(\ell)$ for $\ell > i_j$. Then π can be obtained by gluing from left to right the blocks $(\pi(i_j) - 1)(\pi(i_j) - 2) \cdots (\pi(i_{j+1}) + 1)\pi(i_j)$ for $1 \leq j \leq k$ by setting $\pi(i_1) = n$ and for convenience $i_{k+1} = n + 1$ and $\pi(i_{k+1}) = 0$.

We distinguish two cases: (1) $i_1 > \lfloor \frac{n}{2} \rfloor$, and (2) $i_1 \leq \lfloor \frac{n}{2} \rfloor$.

Case 1. We have $i_1 > \lfloor \frac{n}{2} \rfloor$. So, π can be written $\pi = (n - 1)(n - 2) \cdots (\pi(i_2) + 1)n\gamma$ where γ has all its values less or equal than $\pi(i_2)$ and so that these values do not lie in the excedance board of π . We set $\pi' = (n - 1)(n - 2) \cdots (\pi(i_2) + 1)n\pi(i_2)(\pi(i_2) - 1) \cdots 3 2 1$. Obviously, π' avoids 123 and 132, and π' and π belong to the same class. Varying i_1 from $\lfloor \frac{n}{2} \rfloor$ to n , the number of classes (in this case) is exactly $\sum_{k=\lfloor \frac{n}{2} \rfloor + 1}^n 1 = n - \lfloor \frac{n}{2} \rfloor$.

Case 2. We have $i_1 \leq \lfloor \frac{n}{2} \rfloor$. We will prove that there is a permutation π' in the class of π such that $\pi' = (n-1)(n-2) \cdots (\pi(i_2) + 1)n\delta i_1(i_1 - 1) \cdots 21$.

Let r be the smallest index j such that $\pi(i_j) \leq i_j$. Less formally, the point $(i_r, \pi(i_r))$ is the highest point that does not lie in the excedance board of π .

Since $i_1 \leq \lfloor \frac{n}{2} \rfloor$ and $\pi(i_1) = n$, we necessarily have $r \geq 2$. The permutation π can be written $\pi = \gamma (\pi(i_r) - 1) \cdots (\pi(i_{r+1}) + 1)\pi(i_r)\sigma$ where all values in γ appear in the excedance board of π , and all values in σ do not lie in the excedance board of π . So, there exists an index $t = \pi_{i_r}, i_{r-1} < t \leq i_r$, such that $\pi(t) = t = \pi_{i_r}$.

Now let us consider the permutation π' obtained from π by replacing the block $(\pi(i_r) - 1) \cdots (\pi(i_{r+1}) + 1)\pi(i_r)$ with the block

$$(\pi(i_{r-1} + 1)) \cdots \pi(t - 1)\pi(i_r)\pi(t) \cdots (\pi(i_{r+1}) + 1).$$

Moreover, we replace σ by the decreasing sequence $(\pi(i_r) - 1) \cdots 21$. It is straightforward to see that π and π' lie in the same class. Moreover, π' is in the expected form $\pi' = (n-1)(n-2) \cdots (\pi(i_2) + 1)n\delta i_1(i_1 - 1) \cdots 21$ where δ is a subsequence of length $n - 2i_1$ that avoids 123 and 132.

Thus, the number of classes in $S_n(123, 132)$ such that the point (i_1, n) lies in the excedance board is exactly the number a_{n-2i_1} of classes in $S_{n-2i_1}(123, 132)$. Varying i_1 from 1 to $\lfloor \frac{n}{2} \rfloor$, the number b_n of classes (in this case) is given by $b_n = \sum_{i_1=1}^{\lfloor \frac{n}{2} \rfloor} a_{n-2i_1}$ with $a_1 = 1$ and $a_2 = 2$.

Considering the two cases, we obtain

$$a_n = n - \lfloor \frac{n}{2} \rfloor + \sum_{i_1=1}^{\lfloor \frac{n}{2} \rfloor} a_{n-2i_1}$$

and finally a simple calculation provides $a_n = 2a_{n-2} + 1$ with $a_1 = 1$ and $a_2 = 2$ which is the sequence A052955 in [17]. □

5. Acknowledgement

I would like to thank the anonymous referees for their very careful reading of this paper and their helpful comments and suggestions.

References

- [1] C. Banderier, M. Bousquet-Mélou, A. Denise, P. Flajolet, D. Gardy, and D. Gouyou-Beauchamps. Generating functions of generating trees. *Discrete Mathematics*, 246(1-3), 29-55, 2002.

Pattern	Sequence	Sloane	$a_n, n \geq 1$
{123, 321}			1, 2, 4, 4, 0, 0, 0, ...
{132, 312}, {213, 312}, {231, 312}, {231, 321}, {321, 312}	2^{n-1}	A000079	1, 2, 4, 8, 16, 32, 64, 128, 256
{123, 312}, {132, 321}, {213, 321}	$(n-1)n/2 + 1$	A000124	1, 2, 4, 7, 11, 16, 22, 29, 37
{132, 231}, {213, 231}	F_n	A000045	1, 2, 3, 5, 8, 13, 21, 34, 55
{123, 231}	$\lfloor (n-2)^2/3 \rfloor + n$	A096777	1, 2, 3, 5, 8, 11, 15, 20, 25
{123, 132}, {123, 213}, {132, 213}	$(5 + (-1)^n)2^{\lfloor (n-1)/2 \rfloor - 1} - 1$	A052955	1, 2, 3, 5, 7, 11, 15, 23, 31

Table 3: Number of equivalence classes for permutations avoiding two patterns in \mathcal{S}_3 , where F_n is the n -th Fibonacci number defined as $F_n = F_{n-1} + F_{n-2}$ with $F_0 = F_1 = 1$.

- [2] E. Barucci, A. Bernini, M. Poneti. From Fibonacci to Catalan permutations. *Pure Mathematics And Applications*, 17, 1-17, 2006.
- [3] E. Barucci, A. Del Lungo, E. Pergola and R. Pinzani. From Motzkin to Catalan permutations. *Discrete Mathematics*, 217, 33-49, 2000.
- [4] N. Bergeron and M. Zabrocki. The Hopf algebras of symmetric functions and quasisymmetric functions in non-commutative variables are free and cofree. *J. Algebra Appl.*, 08(4), 581-600, 2009.
- [5] R. Ehrenborg and E. Steingrimsson. The Excedance set of a permutation. *Advances in Applied Mathematics*, 24, 284-299, 2000.
- [6] S. Elizalde. Statistics on Pattern-avoiding Permutations. Ph.D thesis, M.I.T. 2004, <https://www.math.dartmouth.edu/~sergi/papers/thesis.pdf>.
- [7] S. Elizalde and E. Deutsch. A simple and unusual bijection for Dyck paths and its consequences. *Ann. Comb.*, 7(3), 281-297, 2003.
- [8] S. Kitaev. Patterns in permutations and words, Springer-Verlag, 2011.
- [9] D.E. Knuth. The Art of Computer Programming. Vol. 1, Addison-Wesley, Reading MA, 1973, Third edition, 1997.
- [10] T. Mansour. Combinatorics of Set Partitions. *Discrete Mathematics and its Applications*, Chapman & Hall/CRC, 2013.

- [11] T. Mansour. Permutations avoiding a pattern from S_k and at least two patterns from S_3 *Ars. Combin.*, 62, 227-239, 2002.
- [12] T. Mansour and A. Vainshtein Restricted 132-avoiding permutations *Adv. Appl. Math.*, 26, 258-269, 2001.
- [13] R. Simion and F.W. Schmidt. Restricted permutations. *European Journal of Combinatorics*, 6, 383406, 1985.
- [14] J. West. Permutations with forbidden subsequences and stack-sortable permutations. Ph.D. Thesis, MIT, 1990.
- [15] J. West. Generating trees and forbidden subsequences. *Proceeding of sixth FPSAC*, 441-450, 1994.
- [16] J. West. Generating trees and the Catalan and Schröder numbers. *Discrete Math.*, 146, 247-262, 1995.
- [17] N.J.A. Sloane. On-Line Encyclopedia of Integer Sequences. Published electronically at <http://oeis.org/>.
- [18] R. Stanley. Enumerative Combinatorics. *Cambridge University Press*, Vol. 1, 1997.

JEAN-LUC BARIL

LE2I UMR-CNRS 6306, UNIVERSITÉ DE BOURGOGNE, B.P. 47 870, 21078 DIJON, FRANCE

E-mail address: barjl@u-bourgogne.fr

TOUFIK MANSOUR

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, 3498838 HAIFA, ISRAEL

E-mail address: tmansour@univ.haifa.ac.il

ARMEN PETROSSIAN

LE2I UMR-CNRS 6306, UNIVERSITÉ DE BOURGOGNE, B.P. 47 870, 21078 DIJON, FRANCE

E-mail address: armen.petrossian@u-bourgogne.fr

RECEIVED JANUARY 1, 0000