

TUDOR ZAMFIRESCU

On  $k$ -path hamiltonian line-graphs

---

Estratto da « Rend. Ist. di Matem. Univ. Trieste »  
Vol. IV, fasc. II (1972)

---

ISTITUTO DI MATEMATICA - UNIVERSITÀ - TRIESTE  
1972

Estratto da :  
Rend. Ist. di Matem. Univ. di Trieste  
Vol. II, fasc. IV (1972).

## ON $k$ -PATH HAMILTONIAN LINE-GRAPHS (\*)

by TUDOR ZAMFIRESCU (in Dortmund) (\*\*)

SOMMARIO. - *La nota riguarda principalmente i grafi di linee  $k$ -path hamiltoniani. Si indicano delle proprietà del grafo di linee di un grafo  $k$ -path hamiltoniano e si assegnano delle condizioni sufficienti affinché il grafo di linee di un grafo sia  $k$ -path hamiltoniano.*

SUMMARY. - *The paper mainly concerns the  $k$ -path hamiltonian line-graphs. It presents properties of the line-graph of a  $k$ -path hamiltonian graph and sufficient conditions for a graph such that its line graph is  $k$ -path hamiltonian.*

Let  $G$  be an undirected graph, without loops or multiple edges, and  $L(G)$  its line-graph. Our intentions in this note are : 1) to give necessary conditions for  $L(G)$  such that  $G$  is  $k$ -path hamiltonian [1] (Chapters 3 and 4); 2) to give sufficient conditions for  $G$  such that  $L(G)$  is  $k$ -path hamiltonian (Chapter 5). Also, sufficient conditions for a line-graph to be hamiltonian, expressed in terms of covering circuits, are presented in Chapter 2.

1.  $P(G)$  and  $E(G)$  respectively denote the point-set and the edge set of the graph  $G$ .

We say that the graph  $G'$  is included in the graph  $G$  and write  $G' \subset G$  if  $P(G') \subset P(G)$  and  $E(G') \subset E(G)$ . In particular,  $G'$  is a subgraph of  $G$  if  $G' \subset G$  and for each  $G'' \subset G$  with  $P(G'') = P(G')$ , one has  $E(G'') \subset E(G')$ .

(\*) Pervenuto in Redazione il 3 marzo 1972.

(\*\*) Indirizzo dell'Autore : Mathematisches Institut der Universität-Dortmund (Germany).

If  $A, B \subset G$ , we write  $N_A(B)$  for the set of points in  $P(A) - P(B)$  adjacent to some point in  $P(B)$ . If  $B$  has only the point  $x$ , we will write  $N_A(x)$  for  $N_A(B)$ .

In [2] we defined  $G' \subset G$  to be: 1) of type  $T_1$  in  $G$  if it has at least three lines in common with every complete subgraph of  $G$  on 4 points from  $P(G')$ ; 2) of type  $T_2$  in  $G$  if it is of type  $T_1$  in  $G$  and  $\text{card } N_{G'}(x) \leq 1$  for every point  $x \in P(G) - P(G')$ .

Now, we define  $G'$  to be of type  $T_3$  in  $G$  if  $G'$  is of type  $T_1$  in  $G$  and there exist: a) a partition  $\{P_i\}_{i=1}^n$  of all points in  $P(G) - P(G')$ , each of which is adjacent to the end-points of two non-adjacent edges of  $G'$ , both together belonging to no complete subgraph of  $G$ ; b) the distinct points  $\{p_i\}_{i=1}^{2n}$  in  $P(G) - P(G')$ , with  $p_i \notin P_j$  ( $j = 1, \dots, n$ ) and  $N_{G'}(p_{2i-1}) \neq N_{G'}(p_{2i})$  ( $i = 1, \dots, n$ ), such that each  $P_i \cup \{p_{2i-1}, p_{2i}\}$  is the point-set of a circuit in  $G$ , in which  $p_{2i-1}$  and  $p_{2i}$  are adjacent.

A path of length  $k$  is called a  $k$ -path.

A  $k$ -clique in a graph is a clique (maximal complete subgraph) with  $k$  points.

If the edges of the graph  $G$  corresponding to the vertices of the graph  $\Pi_0$  included in  $L(G)$  determine a circuit, then  $\Pi_0$  will be called a strong circuit in  $L(G)$ .

2. Let  $\Gamma$  be a line-graph,  $\Pi_0$  a strong circuit in  $\Gamma$  and  $\Pi_1, \dots, \Pi_k$  other circuits in  $\Gamma$ .

**THEOREM 1.** *If  $P(\Gamma) = \bigcup_{i=0}^k P(\Pi_i)$ ,  $P(\Pi_i) - P(\Pi_0)$  are disjoint, and  $\text{card}(P(\Pi_i) \cap P(\Pi_0)) = 1$  ( $i = 1, \dots, k$ ), then  $\Gamma$  is hamiltonian.*

**PROOF.** Let  $\{a_i\}_{i=1}^n$  be the points of  $\Pi_0$ , written consecutively. Then  $a_i, a_{i+1}$  ( $i = 1, \dots, n-1$ ) and  $a_n, a_1$  are adjacent, but  $a_i, a_{i+2}$  ( $i = 1, \dots, n-2$ ),  $a_{n-1}, a_1$  and  $a_n, a_2$  are not adjacent (if  $n > 3$ ). Consider the point  $a_j$  of  $\Pi_0$ . The set  $N_\Gamma(a_j)$  may be divided into two subsets  $A$  and  $B$  such that each of them determines a complete subgraph of  $\Gamma$ . Let  $\Pi_{\alpha_m}$  ( $m = 1, \dots, \bar{\alpha}_j$ ) be the circuits containing  $a_j$  and no point in  $B$ ,  $\Pi_{\beta_m}$  ( $m = 1, \dots, \bar{\beta}_j$ ) be those containing  $a_j$  and no point in  $A$ , and  $\Pi_{\gamma_m}$  ( $m = 1, \dots, \bar{\gamma}_j$ ) those containing  $a_j$ , and at least one point from each  $A$  and  $B$ . Replace now  $a_j$  by the path whose set of (consecutive) points is:

$$(P(\Pi_{\alpha_1}) - \{a_j\}) \cup \dots \cup (P(\Pi_{\alpha_{\bar{\alpha}_j}}) - \{a_j\}) \cup (P(\Pi_{\gamma_1}) - \{a_j\}) \cup \dots \\ \dots \cup (P(\Pi_{\gamma_{\bar{\gamma}_j}}) - \{a_j\}) \cup \{a_j\} \cup (P(\Pi_{\beta_1}) - \{a_j\}) \cup \dots \cup (P(\Pi_{\beta_{\bar{\beta}_j}}) - \{a_j\}).$$

Doing this in all the points of  $\Pi_0$ , we get the desired hamiltonian circuit of  $\Gamma$ .

Using a proof technique not different from the preceding one, we are able to establish the following stronger result too.

**THEOREM 2.** *Let  $\Gamma$  be a line-graph,  $\Pi_0$  a strong circuit in  $\Gamma$  including the path  $\Pi$  and  $\Pi_1, \dots, \Pi_k$  other circuits in  $\Gamma$  such that:  $P(\Gamma) = \bigcup_{i=0}^k P(\Pi_i)$ , the sets  $P(\Pi_i) - P(\Pi_0)$  are disjoint,  $\text{card}(P(\Pi_i) \cap P(\Pi_0)) = 1$ , and  $P(\Pi_i) \cap P(\Pi) = \emptyset$  ( $i = 1, \dots, k$ ). Then  $\Gamma$  has a hamiltonian circuit including  $\Pi$ .*

Even stronger:

**THEOREM 3.** *Suppose the hypotheses of Theorem 2 are satisfied. Moreover, consider the path  $\bar{\Pi} \subset \Pi_0$  with  $P(\bar{\Pi}) = P(\Pi) \cup N_{\Pi_0}(\Pi)$ , the end-points  $x_1, x_2$  of  $\bar{\Pi}$  and the points  $y_1, y_2 \notin P(\bar{\Pi})$  respectively adjacent in  $\Pi_0$  with  $x_1, x_2$ . Suppose there exists at most one circuit  $\Pi_{j_s}$  ( $1 \leq j_s \leq k$ ) such that  $x_s \in P(\Pi_{j_s})$  ( $s = 1, 2$ ) and, in the existence case, suppose that a point  $z_s \in P(\Pi_{j_s}) - \{x_s\}$  may be found such that the subgraph of  $\Gamma$  with point set  $\{x_s, y_s, z_s\}$  is complete. Then  $\Gamma$  has a hamiltonian circuit including  $\bar{\Pi}$ .*

**3. THEOREM 4.** *If  $G$  is  $k$ -path hamiltonian, then each  $(k+1)$ -path of type  $T_3$  in  $L(G)$  is extendable to a hamiltonian circuit of  $L(G)$ .*

**PROOF.** Let  $\Pi$  be a  $(k+1)$ -path of type  $T_3$  in  $L(G)$ . Then the set of edges  $\{e_1, \dots, e_{k+2}\} \subset E(G)$  corresponding to the vertices of  $\Pi$  is such that  $e_i$  and  $e_{i+1}$  are adjacent ( $i = 1, \dots, k+1$ ). We consider the minimal set of natural numbers  $\{n_j\}_{j=1}^l$  ( $l \leq k+2$ ) such that

$$1 = n_1 < \dots < n_l = k+2$$

and such that the edges  $e_{n_j}, e_{n_{j+1}}, \dots, e_{n_{j+1}}$  have a common end-point ( $j = 1, \dots, l-1$ ). Clearly,  $e_{n_j}$  and  $e_{n_{j+1}}$  are adjacent ( $j = 1, \dots, l-1$ ) and no three consecutive edges in  $\{e_{n_j}\}_{j=1}^l$  form a star.

To prove that  $\{e_{n_j}\}_{j=2}^{l-1}$  determines a path in  $G$ , we have only to show that no four edges  $e_{n_\alpha}, e_{n_{\alpha+1}}, e_{n_\beta}, e_{n_{\beta+1}}$  ( $1 \leq \alpha \leq \beta-3 \leq l-4$ ) form a star. Indeed, if it were not so then the corresponding points  $p_\alpha, p_{\alpha+1}, p_\beta, p_{\beta+1}$  of  $P(L(G))$ , which belong to  $P(\Pi)$ , would be such that: 1) all the six edges determined by them are in  $E(L(G))$ , 2)  $E(\Pi)$  contains from these six edges at most the edges  $(p_\alpha, p_{\alpha+1})$  and  $(p_\beta, p_{\beta+1})$  (if  $n_{\alpha+1} = n_\alpha + 1$  and  $n_{\beta+1} = n_\beta + 1$ ); this contradicts

the fact that  $\Pi$  is of type  $T_1$  in  $L(G)$ . Now, since  $\{e_{n_j}\}_{j=2}^{l-1}$  determines a path  $W$  of length at most  $k$ , we can extend it to a hamiltonian circuit  $\Theta$ , whose edges, written consecutively, are

$$e_{n_2}, \dots, e_{n_{l-1}}, f_1, \dots, f_m.$$

Since  $\Pi$  is of type  $T_3$  in  $L(G)$ , there exist: a) a partition  $\{P_i\}_{i=1}^n$  of all points in  $P(L(G)) - P(\Pi)$ , each of which is adjacent to the end-points of two non-adjacent edges of  $\Pi$ , both of them belonging to no complete subgraph of  $L(G)$ ; b) the distinct points  $\{p_i\}_{i=1}^{2n}$  in  $P(L(G)) - (P(\Pi) \cup \bigcup_{j=1}^n P_j)$  with  $N_{\Pi}(p_{2i-1}) \neq N_{\Pi}(p_{2i})$  ( $i = 1, \dots, n$ ), such that  $P_i \cup \{p_{2i-1}, p_{2i}\}$  form circuits  $\Pi_i$ , in which  $p_{2i-1}$  and  $p_{2i}$  are adjacent. Obviously, the set of points of  $L(G)$  corresponding to the edges in  $\Theta$  determines a strong circuit  $\Pi_0$  of  $L(G)$ .

We construct now a circuit  $\Pi'$  in  $L(G)$ , with  $P(\Pi') = \bigcup_{i=0}^n P(\Pi_i)$ . Consider some  $i \in \{1, \dots, n\}$ . Since for every point  $\pi \in P_i$ , there are two nonadjacent edges in  $\Pi$  belonging to no complete subgraph of  $L(G)$ , each of their 4 end-points being adjacent to  $\pi$ , it follows that the edge  $\varepsilon$  in  $G$  corresponding to  $\pi$  is adjacent to four edges  $e_j, e_{j+1}, e_k, e_{k+1}$ , where  $j+1 < k$  and not every two of these 4 edges are adjacent. It follows that  $\varepsilon$  joins two points of  $W$  (the path determined by  $\{e_{n_j}\}_{j=2}^{l-1}$ ). Because  $P_i \cup \{p_{2i-1}, p_{2i}\}$  is the point-set of a circuit, each of the points  $p_{2i-1}, p_{2i}$  is adjacent to some point in  $P_i$ , hence each of the edges  $\nu_1$  and  $\nu_2$  corresponding in  $G$  to  $p_{2i-1}$  and  $p_{2i}$  has an end-point in  $P(W)$ . Since  $p_{2i-1}, p_{2i} \notin \bigcup_{j=1}^n P_j$ , neither  $\nu_1$  nor  $\nu_2$  has both end-points in  $P(W)$ . Because  $N_{\Pi}(p_{2i-1}) \neq N_{\Pi}(p_{2i})$ , the end-points in  $P(W)$  of  $\nu_1$  and  $\nu_2$  are different. But  $p_{2i-1}$  and  $p_{2i}$  are adjacent; therefore  $\nu_1$  and  $\nu_2$  have a common end-point, not in  $P(W)$ . Since  $p_{2i-1}, p_{2i} \notin P(\Pi)$ , the edges  $\nu_1, \nu_2$  do not belong to  $E(W)$ . Suppose now  $\nu_1, \nu_2 \in E(\Theta)$ . Then  $\nu_1, \nu_2 \in \{f_1, \dots, f_m\}$ , but since  $\nu_1$  and  $\nu_2$  are adjacent, and each of them is adjacent to some edge of  $W$ , it results  $m = 2$ ; then  $\{f_1, f_2\} = \{e_1, e_{k+2}\}$ , whence  $\{\nu_1, \nu_2\} = \{e_1, e_{k+2}\}$ , which contradicts  $p_{2i} \notin P(\Pi)$ . Therefore  $\{\nu_1, \nu_2\} \not\subset E(\Theta)$ . Since  $\nu_1$  and  $\nu_2$  have a common end-point  $\omega$  not in  $P(W)$ , and since  $\{f_1, \dots, f_m\} \neq \{e_1, e_{k+2}\}$ , there exists an edge  $f_j \in E(\Theta) - \{e_1, \dots, e_{k+2}, f_1, f_m\}$ , with  $\omega$  as an end-point. Let  $\varphi$  be the point in  $L(G)$  corresponding to  $f_j$ . Since each edge corresponding in  $G$  to some point of  $P_i$  joins two points of  $W$ , but is not itself an edge of  $W$  (otherwise  $P_i$  would have a point in  $P(\Pi)$ ), it follows that  $P_i \cap P(\Pi_0) = \emptyset$ . Let us construct the circuit  $\Pi_i^{\S}$  in  $L(G)$

such that  $\text{card}(P(\Pi_i^\delta) \cap P(\Pi_0)) = 1$ ,  $P(\Pi_i^\delta) \cap P(\Pi) = \emptyset$  and also the other hypotheses of Theorem 3 are satisfied ( $i = 1, \dots, n$ ).

CASE I.  $\{v_1, v_2\} \cap E(\Theta) \neq \emptyset$ . From  $\{v_1, v_2\} \not\subset E(\Theta)$  and  $\{v_1, v_2\} \cap E(\Theta) \neq \emptyset$ , it follows that exactly one edge from  $\{v_1, v_2\}$  belongs to  $E(\Theta)$ , hence  $\text{card}(P(\Pi_i) \cap P(\Pi_0)) = 1$ . On the other hand,  $p_{2i-1}, p_{2i} \notin P(\Pi)$ , whence  $P(\Pi_i) \cap P(\Pi) = \emptyset$ . Further, if  $v_1 \in E(\Theta)$ , then  $p_{2i-1}$  is an end-point of the path  $\bar{\Pi}$  having as point-set that set of points which correspond in  $L(G)$  to the edges of  $E(W)$  plus  $f_1, f_m$ , and no circuit from  $\{P_j\}_{j=1}^n$  except  $P_i$  has  $p_{2i-1}$  as a point. Moreover,  $p_{2i-1}, p_{2i}, \varphi$  are the points of a complete subgraph of  $L(G)$ . Thus, the hypotheses of Theorem 3 are verified and we may take  $\Pi_i^\delta = \Pi_i$ .

CASE II.  $\{v_1, v_2\} \cap E(\Theta) = \emptyset$ . The points of  $P_i \cup \{p_{2i-1}, p_{2i}\}$  determine a circuit  $\Pi_i^\delta$  in  $L(G)$ , such that  $\text{card}(P(\Pi_i^\delta) \cap P(\Pi_0)) = 1$  and  $P(\Pi_i^\delta) \cap P(\bar{\Pi}) = \emptyset$ , because  $P_i \cap P(\Pi_0) = \emptyset$ ,  $\{v_1, v_2\} \cap E(\Theta) = \emptyset$ , and  $\varphi \in P(\Pi_0) - P(\bar{\Pi})$ .

Now, following Theorem 3,  $\bigcup_{j=0}^n P(\Pi_j)$  is the point-set of a circuit  $\Pi'$  in  $L(G)$ , including  $\bar{\Pi}$ .

We consider now the edges

$$e_{n_2}, \dots, e_{n_{i-1}}, f_1, g_1, \dots, g_r, f_m \in E(G)$$

corresponding to the points of  $\Pi'$ . Since  $\{f_1, \dots, f_m\} \neq \{e_1, e_{k+2}\}$ , we have  $r \neq 0$ . The circuit  $\Pi'$  of  $L(G)$  could (and is supposed to) be constructed so that for each point not in  $P(W)$  there are two consecutive edges in  $\{f_1, g_1, \dots, g_r, f_m\}$ , both of them having this point as an end-point. This fact cannot be derived from Theorem 3 and is a consequence of the fact that in the case II considered before there are two edges in  $\{f_1, \dots, f_m\}$  having  $\omega$  as an end-point and therefore the points in  $L(G)$  corresponding to them, plus  $p_{2i-1}, p_{2i}$ , form the point-set of a complete subgraph of  $L(G)$ . From the construction of  $\Pi'$ , it is also clear that

$$E(G) - \{e_1, \dots, e_{k+2}, f_1, g_1, \dots, g_r, f_m\}$$

may be partitioned into the (possibly empty) sets  $\{H_j\}_{j=1}^{r+1}$  such that all edges of  $H_j$  have a common point, which is not in  $P(W)$  and therefore is the common end point of two consecutive edges from

$\{f_1, g_1, \dots, g_r, f_m\}$ , namely of  $f_1$  and  $g_1$  if  $j = 1$ , of  $g_{j-1}$  and  $g_j$  if  $2 \leq j \leq r$ , and of  $g_r$  and  $f_m$  if  $j = r + 1$ . Consequently, if  $H_j = \{h_i\}_{i=1}^{\psi_j}$  (where  $\psi_j$  is possibly zero, i. e.  $H_j = \emptyset$ ; also possibly  $f_1 = e_{k+2}$  or  $f_m = e_1$ ), then the points in  $L(G)$  corresponding to

$$e_1, \dots, e_{k+2}, f_1, h_1, \dots, h_{\psi_1}, g_1, h_2, \dots, h_{\psi_2},$$

$$g_2, \dots, g_{r-1}, h_r, \dots, h_{\psi_r}, g_r, h_{r+1}, \dots, h_{\psi_{r+1}}, f_m$$

form the desired hamiltonian circuit of  $L(G)$ .

4. THEOREM 5. *Let  $G$  be  $k$ -path hamiltonian, and  $\Pi$  a  $(k + 1)$ -path of type  $T_1$  in  $L(G)$ , with the property that if a point in  $P(L(G)) - P(\Pi)$  is adjacent to the end-points of two nonadjacent edges of  $\Pi$ , then these edges belong to a clique of  $L(G)$ . Then  $\Pi$  is extendable to a hamiltonian circuit of  $L(G)$ .*

This result is a consequence of Theorem 4, since every path of type  $T_1$  in  $L(G)$  with the mentioned property is obviously of type  $T_3$  in  $L(G)$ . A further simplification of the statement of Theorem 4 (but with smaller degree of generality!) leads to

THEOREM 6. *If  $G$  is  $k$  path hamiltonian, then each  $(k + 1)$ -path  $\Pi$  of type  $T_1$  in  $L(G)$  such that  $\text{card } N_{\Pi}(x) \leq 3$  for every point  $x \in P(L(G)) - P(\Pi)$  is extendable to a hamiltonian circuit of  $L(G)$ .*

As an immediate consequence we obtain :

COROLLARY [2]. *If  $G$  is  $k$  path hamiltonian, then each  $(k + 1)$ -path of type  $T_1$  in  $L(G)$  whose  $(k - 1)$ -subpath obtained by removing its end-points (and adjacent edges) is of type  $T_2$  in  $L(G)$ , is extendable to a hamiltonian circuit of  $L(G)$ .*

Theorems 5 and 6 are weaker forms of Theorem 4. In order to obtain a stronger (and more natural) form of Theorem 4, we shall use instead of the types  $T_1$  and  $T_3$ , new types  $T_1'$  and  $T_3'$ , defined as follows :

A graph  $G'$  included in  $G$  is of type  $T_1'$  in  $G$  if  $E(G')$  contains at least  $k - 1$  edges of every  $k$ -clique of  $G$  with points in  $P(G')$ . Also, let  $T_3'$  be the type we obtain if, in the definition of  $T_3$ , the type  $T_1$  is replaced by  $T_1'$ .

By replacing  $T_1$  and  $T_3$  with the new types  $T_1'$  and  $T_3'$  in the statements of Theorems 4, 5, and 6, we obtain the stronger Theorems 4', 5', and 6' respectively.

5. THEOREM 7. *If  $G$  is  $k$ -path hamiltonian and has no circuit of length at most  $k + 1$ , then  $L(G)$  is  $(k + 1)$ -path hamiltonian.*

PROOF. Let  $\Pi$  be a  $(k + 1)$ -path in  $L(G)$  and use further the notations of the proof of Theorem 4. We prove that the hypotheses of Theorem 5' are satisfied. First, suppose  $\Pi$  is not of type  $T_1'$  in  $L(G)$ . Then there exists a  $k$ -clique of  $L(G)$  on  $k$  points in  $P(\Pi)$ , such that at most only  $k - 2$  of its edges are in  $E(\Pi)$ . It follows that the edges of  $E(G)$  corresponding to the points of the clique form a star, but are not consecutive in  $\{e_1, \dots, e_{k+2}\}$ . Then they may be divided into sets  $\{E_i\}_{i=1}^t$ , with  $t \geq 2$ , such that for each  $i \leq t$  there exists  $f(i) \leq l - 1$  with

$$E_i = \{e_{n_{f(i)}}, e_{n_{f(i)}+1}, \dots, e_{n_{f(i)}+1}\}.$$

But in this case the edges

$$e_{n_{f(1)}+1}, e_{n_{f(1)}+2}, \dots, e_{n_{f(2)}}$$

from  $E(\Theta)$  form a circuit of length at most  $l - 2$ , hence at most  $k$ , which contradicts the hypothesis.

Now, suppose there exists a point in  $P(L(G)) - P(\Pi)$  which is adjacent to the end-points of two nonadjacent edges of  $\Pi$ , but these edges do not belong to any clique of  $L(G)$ . Then there are four edges  $e_\zeta, e_{\zeta+1}, e_\xi, e_{\xi+1}$  which do not form a star, but are all adjacent to some other edge  $b$  of  $E(G)$ . Let  $j' < j'' \leq l$ , with the property that  $n_{j'} \leq \zeta, \zeta + 1 \leq n_{j'+1}, n_{j''} \leq \xi, \xi + 1 \leq n_{j''+1}$ . The edges

$$e_{n_{j'+1}}, e_{n_{j'+2}}, \dots, e_{n_{j''}}, b$$

determine a circuit of length at most  $k + 1$  in  $G$ , again contradicting the hypothesis.

Following Theorem 5',  $\Pi$  may be extended to a hamiltonian circuit of  $L(G)$ .

## REFERENCES

- [1] H. KRONK, *A Note on  $k$ -path Hamiltonian Graphs*, J., Combinatorial Theory 7 (1969) 104-106.
- [2] T. ZAMFIRESCU, *On  $k$ -path Hamiltonian Graphs and Line-graphs*, Rend. Sem. Mat. Univ. Padova 46 (1971) 385-389.