

## Multiple farthest points on Alexandrov surfaces

Costin Vîlcu and Tudor Zamfirescu

(Communicated by K. Strambach)

**Abstract.** The farthest point mapping on compact surfaces, associating to each point  $x$  of the surface the set of absolute maxima of the intrinsic distance from  $x$ , is for some surfaces single-valued and a homeomorphism, while for other surfaces it is not single-valued, and not surjective. These two big classes are not very well understood. For instance it is still unknown whether, say in the convex case, the second class is dense. For a  $C^2$  metric on both surfaces and the space of surfaces, the first class has, however, nonempty interior. We describe various properties of the sets of critical points, and of relative and absolute maxima of distance functions, and find several connections between them. We see for example that, on smooth surfaces homeomorphic to  $S^2$ , a point cannot be critical with respect to more than one other point. Sufficient conditions for a surface to belong to the second class will be formulated and a particular Tannery surface belonging to the boundary of both classes will be presented.

2000 Mathematics Subject Classification. 53C45

### 1 Introduction

We investigate the set of critical points with respect to distance functions on Alexandrov surfaces. Special attention will receive Alexandrov surfaces homeomorphic to the 2-dimensional sphere  $S^2$ , while we sometimes restrict ourselves to convex surfaces.

In this paper, by *surface* we always mean a compact 2-dimensional Alexandrov space with curvature bounded below (without boundary), as defined by Burago, Gromov and Perelman in [4]. Let  $\mathcal{A}$  be the space of all surfaces.

It is well-known that our surfaces are topological manifolds. Other basic facts on surfaces, such as convergence theorems on shortest paths or on angles, the generalized Toponogov theorem, and an ample description of the topological and metric structure of the cut loci, can be found in [4], [18], [11]. Some of these basic facts will be tacitly used, while in other cases we shall recall the needed facts as lemmas.

Denote by  $\mathcal{S}$  the space of all convex surfaces embedded in  $\mathbb{R}^3$ , and by  $\mathcal{A}_0, \mathcal{R}_0$  the set of all Alexandrov, respectively Riemannian, surfaces homeomorphic to  $S^2$ .

For any two points  $x, y$  on the surface  $S$ ,  $\rho(x, y)$  means the geodesic (intrinsic) distance between them (induced by the Euclidean distance for  $S \in \mathcal{S}$ ), and  $\rho_x$  the distance function from  $x$ :  $\rho_x(y) = \rho(x, y)$ . For  $x \in S$  denote by  $F_x$  the set of all farthest points from  $x$  (i.e., absolute maxima of  $\rho_x$ ) and by  $F$  the *farthest point mapping*, the multivalued mapping associating to any point  $x \in S$  the set  $F_x$ . Similarly,  $M_x$  is the set of all relative maxima of  $\rho_x$ ,  $Q_x$  the set of all critical points with respect to  $\rho_x$ , and  $M$ , respectively  $Q$ , are the corresponding mappings. Here, a point  $y \in S$  is called *critical* with respect to  $\rho_x$ , or simply to  $x$ , if for any direction  $v$  at  $y$  there exists a segment from  $y$  to  $x$  whose direction at  $y$  makes an angle not larger than  $\pi/2$  with  $v$ .

If  $f : S \rightarrow \mathcal{P}(S)$  is a multivalued mapping (with the set  $f_x$  as image of  $x$ ), we call it *injective* if  $f_x \cap f_y = \emptyset$  whenever  $x \neq y$ , and *surjective* if for every point  $y \in S$  there exists  $x \in S$  with  $y \in f_x$ . We say that  $f$  is *connected* if  $f_x$  is connected for each  $x \in S$ . When we say that  $f$  is bijective or a homeomorphism, we implicitly state that  $f$  is single-valued. Also,  $f_y^{-1} = \{x \in S : y \in f_x\}$ .

Several questions about farthest points proposed by H. Steinhaus (see Chapter A35 of the book [7] of H. T. Croft, K. J. Falconer and R. K. Guy) have been answered in the convex case by the second author ([24], [25], [28], [29], [30]). J. Rouyer [14], [13] showed that some of these results are also true in the framework of Riemannian geometry. The present paper is also contributing to the study of the farthest point mapping on surfaces, which Steinhaus had asked for.

The next section contains a description of the components of  $M_x$  and  $F_x$ , and points out that, generically in  $\mathcal{S}$ ,  $Q_x$  — and therefore  $M_x$  and  $F_x$  too — are totally disconnected. In the third section we see that, on any smooth surface in  $\mathcal{A}_0$ , no point can be critical with respect to more than one point. Section 4 establishes the especially strong relationship between the components of  $M_x$  and those of  $Q_x$ . In the following section it is proved that the family  $\mathcal{S}_2$  of those surfaces on which  $M$  is disconnected is open in  $\mathcal{A}_0$ . Two major results of the paper, both in Section 6, are Theorem 6 stating several implications between various connectedness and surjectivity properties of  $Q$ ,  $M$ , or  $F$ , and Theorem 7 providing the generic equivalence of all properties considered in Theorem 6. The paper continues with three very different sufficient conditions for a surface to belong to  $\mathcal{S}_2$ . The third brings to light an exceptional type of surface belonging to the boundary of  $\mathcal{S}_2$ . This takes a concrete form in the next section, where a special Tannery surface is investigated. The paper ends with a few open questions.

We need more definitions and additional notation.  $\text{card } A$  denotes the cardinality of the set  $A$ ,  $\text{bd } A$  denotes its boundary,  $\bar{A}$  its closure, and  $\lambda A$  its 1-dimensional Hausdorff measure (length).

For any surface  $S$  and  $A \subset S$ , put  $\rho(x, A) = \inf_{y \in A} \rho_x(y)$  and  $r_x = \sup_{y \in S} \rho_x(y)$ . Also,  $F_A = \cup_{x \in A} F_x$ . The *radius* of the surface  $S$  is defined by  $\text{rad } S = \inf_{x \in S} r_x$  and its *diameter* by  $\text{diam } S = \sup_{x \in S} r_x$ . A *domain* of  $S$  is an open connected subset of  $S$ .

Let  $S \in \mathcal{A}$ ,  $x \in S$ , and  $\varepsilon > 0$ . Any set  $B(x, \varepsilon) = \{y \in S : \rho_x(y) < \varepsilon\}$  homeomorphic to a usual open disc in the plane is called *open disc* around  $x$ . If  $\varepsilon$  is small enough,  $B(x, \varepsilon)$  is indeed an open disc (see [4]). An *arc*  $\Gamma$  is a homeomorphic image of  $[0, 1]$ ; its *interior*  $\text{int } \Gamma$  is the image of  $]0, 1[$ .

The union of two *segments* (i.e., shortest paths) from  $x \in S$  to some point  $y \in S$ , which make an angle equal to  $\pi$  at  $y$ , will be called a *loop at x*; of course, the segments

have equal lengths.

A *geodesic* is a curve which is locally a segment. More precisely, there is an interval  $I$  of  $\mathbb{R}$  and a parametrization  $i : I \rightarrow S$  such that any point  $t \in I$  has a neighbourhood  $V$  for which  $i(V)$  is a segment. A point of a surface  $S$  is called *endpoint* of  $S$  if it is not interior to any geodesic.

If, for any  $x \in S$ , each loop (if there is at least one) at  $x$  has length  $2r_x$ , we say that  $S$  is *loopy*.

If  $\sigma, \sigma'$  are two segments with precisely one common endpoint  $a$ , then  $\angle\sigma\sigma'$  denotes the angle between the tangent directions of  $\sigma$  and  $\sigma'$  at  $a$ . For  $a \neq b$ ,  $ab$  means the segment from  $a$  to  $b$  when that segment is unique or clearly identifiable from the context.  $\angle xyz$  means  $\angle xyz$ .

A *geodesic triangle* in a surface is a collection of three segments  $\gamma_1, \gamma_2, \gamma_3$  such that  $\gamma_i, \gamma_{i+1}$  have the common endpoint  $a_{i+2}$ . The indices should be taken modulo 3. We shall denote the triangle by  $(\gamma_1, \gamma_2, \gamma_3)$ .

For a point  $x \in S$ , let  $C_x$  be the set of all points joined to  $x$  by at least two segments, and  $C(x)$  the *cut locus* of  $x$ , i.e. the set of all endpoints (different from  $x$ ) of maximal (with respect to inclusion) segments starting at  $x$ . Clearly,  $C_x \subset C(x)$ , and  $C(x)$  is known to be a local tree (that is, each of its points  $y$  has a neighbourhood  $V$  in  $S$  where the component  $K_y(V)$  of  $y$  is a tree), even a tree if the surface is homeomorphic to  $S^2$  (see [10], [16], [18] for basic properties of the cut locus). Recall that a *tree* is a set  $T$  any two points of which can be joined by a unique arc included in  $T$ . The *degree* of a point  $y$  of a local tree is the number of components of  $K_y(V) \setminus \{y\}$  if  $V$  is chosen such that  $K_y(V)$  is a tree. A point  $y \in T$  is called an *extremity* of  $T$  if it has degree 1, and a *ramification point* of  $T$  if it has degree at least 3. A tree is called *nondegenerate* if it has at least one ramification point. A *Y-tree* is a tree with precisely three extremities, and a tree is *finite* if it has finitely many extremities.

We consider isometrical surfaces as not different, and equip the space  $\mathcal{A}$  with the Hausdorff–Gromov metric. In  $\mathcal{S}$ , where distinct surfaces may be isometric, the usual Pompeiu–Hausdorff distance is never less than the Hausdorff–Gromov distance between any two surfaces.

The space  $\mathcal{S}$  is Baire, and in any Baire space *most* means “all except those in a set of first category”.

**Acknowledgement.** Some theorems of this paper are refined and generalized versions of results in C. Vîlcu’s Thesis (University of Dortmund, 2003).

## 2 Arcs in $Q_x, M_x, F_x$

For any surface  $S$ , the space  $T_y$  of all unit directions at  $y \in S$  is a closed Jordan curve of length at most  $2\pi$  [4]. The point  $y$  is said to be *conical* if  $\lambda T_y < 2\pi$ , and we call  $S$  *smooth* if it has no conical points. Notice that a surface may be smooth without being differentiable. This is, for example, the case for all convex surfaces admitting 1-singular but no 0-singular points (in the terminology of R. Schneider [17]). In 1959 V. Klee proved that most convex surfaces are smooth, even of class  $C^1$  [9].

**Lemma 1.** *On  $S \in \mathcal{A}$ , let  $y \in C(x)$  and  $D$  be a component of the complement of  $C(x)$  in an open disc  $O$  around  $y$  with  $y \in \overline{D}$ . Then there exists a segment from  $x$  to  $y$  meeting  $O \setminus \{y\}$  inside  $D$ . Consequently, if the degree of  $y$  is  $d$ , then there are at least  $d$  segments from  $x$  to  $y$ .*

*Proof.* Let the sequence of points  $y_n \in D$  converge to  $y$ . Choose any segment  $\sigma_n$  from  $x$  to  $y_n$ . Since  $S$  is compact,  $\{\sigma_n\}_{n=1}^{\infty}$  has a convergent subsequence, the limit  $\sigma$  being also a segment. Clearly,  $\sigma$  joins  $x$  to  $y$  and  $\sigma \cap O \subset \overline{D}$ . By the definition of the cut locus,  $(\text{int } \sigma) \cap O \subset D$ .  $\square$

**Lemma 2.** *Let  $S \in \mathcal{A}$ ,  $x \in S$  and  $y, z \in C(x)$ . Let  $J$  be an arc joining  $y$  to  $z$  in  $C(x)$ . If  $u \in \text{int } J$  is a relative minimum of  $\rho_x|_J$  then  $u$  is the midpoint of a loop  $\Lambda$  at  $x$  and, except for the two subarcs of  $\Lambda$ , no segments connect  $x$  to  $u$ .*

This follows from the main result in [27].

For  $S \in \mathcal{S}$  and  $x \in S$ , we know that  $F_x$  is homeomorphic to a compact subset of  $\mathbb{R}$  [28], so each component of  $F_x$  is either a point or an arc. We extend this here.

**Theorem 1.** *For any surface  $S \in \mathcal{A}$  and any point  $x$  in  $S$ , each component of  $M_x$  is homeomorphic to a connected subset of the circle  $S^1$ , hence each component of  $F_x$  is a point or an arc or a closed Jordan curve. For  $S \in \mathcal{A}_0$ , each component of  $M_x$  is homeomorphic to a connected subset of  $\mathbb{R}$ , hence each component of  $F_x$  is a point or an arc.*

*Proof.* Let  $x \in S$  and take a component  $M^1$  of  $M_x$  containing more than one point. Since  $M^1$  is included in  $C(x)$  which is a local tree,  $M^1$  is itself a local tree.

Since the restriction of  $\rho_x$  to  $M^1$  is continuous, it is a constant function, and has therefore a relative minimum at each point of  $M^1$  of degree at least 2, in particular at any ramification point of  $M^1$ , if it has one. But such a point is also a ramification point of  $C(x)$  and, by Lemma 1, is joined to  $x$  by at least three segments, in contradiction to Lemma 2.

Without any ramification points,  $M^1$  must be an arc with possibly one or both endpoints removed, or a closed Jordan curve. The latter case is excluded for  $S \in \mathcal{A}_0$ , because then  $C(x)$  is a tree.  $\square$

**Lemma 3.** *Let  $S \in \mathcal{A}_0$ ,  $x \in S$  and  $y, z \in C(x)$  be distinct. Suppose  $\Gamma_y, \Gamma'_y$  are (possibly coinciding) segments from  $x$  to  $y$  and  $\Gamma_z, \Gamma'_z$  are (possibly coinciding) segments from  $x$  to  $z$ . Then there is a domain  $\Delta$  with boundary  $\Gamma_y \cup \Gamma'_y \cup \Gamma_z \cup \Gamma'_z$  and a Jordan arc  $J_{yz}$  in  $C_x \cup \{y, z\}$  joining  $y$  to  $z$ . Moreover, every point in  $\text{int } J_{yz}$  belongs to  $\Delta$  and can be joined to  $x$  by two segments the union of which separates  $y$  from  $z$ .*

Lemma 3 appears in [24], formulated for  $\mathcal{S}$  instead of  $\mathcal{A}_0$ . The extension to  $\mathcal{A}_0$  is straightforward.

**Lemma 4.** *Let  $S \in \mathcal{A}$ ,  $x \in S$ , and  $J \subset C(x)$  be an arc each point of which is joined to  $x$  by precisely two segments. Let  $y_1, y_2$  be the endpoints of  $J$ , and suppose there*

exists a domain  $\Delta$  bounded by the segments from  $x$  to  $y_1, y_2$  and containing  $\text{int } J$ . Then  $\Delta \cap C(x) \subset J$ .

In case  $S \in \mathcal{A}_0$ , the existence of  $\Delta$  is guaranteed by Lemma 3.

*Proof.* Suppose there is a point  $z$  in  $\Delta \cap C(x) \setminus J$ . Since  $C(x)$  is arcwise connected, there is a minimal subarc  $A$  of  $C(x)$  containing  $z$  and meeting  $J$ . Let  $\{u\} = A \cap J$ .

Assume first that  $u \in \Delta$ . Then  $u \in J \setminus \{y_1, y_2\}$ , so it is a ramification point of  $C(x)$ , whence it is joined to  $x$  by at least three segments by Lemma 1, and we obtained a contradiction.

Assume now that  $u \in C(x) \setminus \Delta$ . Since the interiors of the segments from  $x$  to  $y_1, y_2$  are disjoint from  $C(x)$ ,  $u$  must be one of the endpoints of  $J$ , say  $y_1$ , and  $A \setminus \{y_1\} \subset \Delta$ . By Lemma 1, there is another segment from  $x$  to  $y_1$  arriving at  $y_1$  between  $J$  and  $A$ , and a contradiction is obtained again.  $\square$

**Lemma 5.** *On most convex surfaces, most points are endpoints.*

This result was proven in [21].

**Lemma 6.** *On  $S \in \mathcal{A}$ , let  $\sigma, \sigma'$  be (possibly coinciding) segments from  $x$  to  $y$  and  $D$  a component of the complement of  $\sigma \cup \sigma'$  in an open disc around  $y$ . Further, suppose no point of  $D$  belongs to any segment from  $x$  to  $y$ . Then  $C(x) \cap \bar{D}$  admits a half-tangent at  $y$  which bisects the angle of  $\sigma, \sigma'$  at  $y$  toward  $D$ .*

This is Lemma 2.1 in [18].

The following statement is rather obvious for  $S \in \mathcal{R}_0$ .

**Lemma 7.** *For  $S \in \mathcal{A}$  and  $x \in S$ , any arc  $J \subset C(x)$  joining two points  $y, z \in S$  with  $\rho_x(y) < \rho_x(z)$  contains a point which is not critical with respect to  $\rho_x$ .*

*Proof.* Let  $j : [0, \lambda J] \rightarrow J$  be a parametrization by arc-length of  $J$ , and  $f(s) = \rho_x(j(s))$ .

By Lemma 6,  $J$  has at any point  $v \in \text{int } J$  right and left tangents with directions  $\tau^+, \tau^-$ . There are points  $\tau_1^+, \tau_2^+, \tau_1^-, \tau_2^-$  lying together with  $\tau^+, \tau^-$  in the order  $\tau_1^+, \tau^+, \tau_2^+, \tau_1^-, \tau^-, \tau_2^-$  on  $T_v$  (where possibly  $\tau_1^+ = \tau_2^-$  or  $\tau_2^+ = \tau_1^-$ ), and there are segments from  $v$  to  $x$  with directions  $\tau_1^+, \tau_2^+, \tau_1^-, \tau_2^-$  at  $v$  such that (i) the direction at  $v$  of any further segment from  $x$  to  $v$  is separated from  $\tau^+$  by  $\tau_1^+$  and  $\tau_2^+$ , and from  $\tau^-$  by  $\tau_1^-$  and  $\tau_2^-$ ; (ii)  $\angle \tau^+ \tau_1^+ = \angle \tau^+ \tau_2^+$  and  $\angle \tau^- \tau_1^- = \angle \tau^- \tau_2^-$  (by Lemma 6).

This yields the existence of right and left derivatives  $f^+$  and  $f^-$  of  $f$  everywhere in  $]0, \lambda J[$ . (More precisely,  $f^+(s) = -\cos \angle \tau^+ \tau_1^+$  and  $f^-(s) = \cos \angle \tau^- \tau_1^-$ .) Since  $\lambda T_v \leq 2\pi$ , we have  $\angle \tau^+ \tau_1^+ + \angle \tau^- \tau_1^- \leq \pi$ , which yields  $f^- \geq f^+$ .

Let  $q = \min\{x \in [0, \lambda J] : f(x) = f(\lambda J)\}$ . There exists a point  $t \in ]0, q[$  with the whole graph of  $f|_{[0, q]}$  in one of the two closed half-planes with the line through  $(t, f(t))$  of slope  $(f(q) - f(0))/q$  as boundary.

If there is such a point  $t$  with the graph above the line, then

$$f^-(t) = f^+(t) = \frac{f(q) - f(0)}{q} > 0.$$

In the contrary case, the graph is below the line, and  $f^-(t) \geq f^+(t) > 0$  or  $(t + \varepsilon, f(t + \varepsilon))$  lies below the line through  $(t, f(t))$  and  $(q, f(q))$  for  $\varepsilon > 0$  small enough. In the latter situation, repeat the argument for  $f|_{[t,q]}$  instead of  $f|_{[0,q]}$ , and we are in the previous case.

So, in any case we have found a point  $t$  satisfying  $f^-(t) \geq f^+(t) > 0$ , which means that  $\angle \tau^- \tau_1^- < \pi/2$  and  $\tau^+ \tau_1^+ > \pi/2$ . The second inequality shows that  $v$  is not a critical point.  $\square$

**Theorem 2.** *On most convex surfaces  $S$ , for any  $x \in S$  and  $a > 0$ , the set  $\rho_x^{-1}(a) \cap C(x)$  is totally disconnected; consequently,  $F_x$ , and even  $Q_x$ , are totally disconnected too.*

*Proof.* Let  $S \in \mathcal{S}$ , and suppose there exist  $x \in S$  and  $a > 0$  such that  $\rho_x^{-1}(a) \cap C(x)$  contains an arc  $J'$ .

Let  $y_1, y_2$  be two interior points of  $J'$ , and denote by  $J$  the subarc of  $J'$  joining them. Since each point  $z \in J$  is a relative minimum for  $\rho_x|_{J'}$ , we conclude, by Lemma 2, that  $z$  is the midpoint of a loop  $\Lambda_z$  at  $x$ , and no segments connect  $x$  to  $z$  excepting those in  $\Lambda_z$ . Now, denote by  $\Delta$  the domain bounded by  $\Lambda_{y_1} \cup \Lambda_{y_2}$ . By Lemma 4, we have  $\Delta \cap C(x) \subset J$ , whence no endpoint of  $S$  belongs to  $\Delta \setminus J$ , a fact which contradicts Lemma 5.

By Lemma 7,  $\rho_x$  would be constant on any arc in  $Q_x$ , and this we just showed to be impossible.  $\square$

Further, sometimes strange, properties of most convex surfaces are surveyed in [8] and [22]. (See also [32].) It is, for example, true that on most convex surfaces there are arbitrarily long geodesics without self-intersections [23] and with huge distance between consecutive conjugate points [26].

### 3 The injectivity of $Q$

The goal of this section is to show that a smooth point cannot be simultaneously critical with respect to two distinct points.

**Lemma 8.** *On  $S \in \mathcal{A}$ , let  $\sigma, \sigma'$  be (possibly coinciding) segments from  $x$  to  $y$  and  $D$  a component of the complement of  $\sigma \cup \sigma'$  in an open disc around  $y$ . Further, suppose no point of  $D$  belongs to any segment from  $x$  to  $y$ . If the angle between  $\sigma, \sigma'$  at  $y$  toward  $D$  is larger than  $\pi$ , then  $y$  is a strict relative minimum of  $\rho_x|_{C(x) \cap \bar{D}}$ .*

This is part of Proposition 2.4 in [18].

**Lemma 9.** *For any  $S \in \mathcal{A}$  and  $x \in S$  we have  $M_x \subset Q_x$ .*

For  $S \in \mathcal{R}_0$ , this is essentially Berger's Lemma (see [2]). For  $S \in \mathcal{S}$ , this is Theorem 2 in [24]. In the general case, it follows from Lemma 8.

**Lemma 10.** *If  $S \in \mathcal{A}$ ,  $x \in S$ ,  $y \in Q_x$  and  $\lambda T_y > \pi$ , then there are at least two segments from  $x$  to  $y$ .*

This useful lemma follows directly from the definition of a critical point.

**Theorem 3.** *If  $S \in \mathcal{A}_0$ ,  $x, y \in S$  are distinct, and  $z \in Q_x \cap Q_y$ , then  $z$  is a conical point. Hence, if  $S \in \mathcal{A}_0$  is smooth then  $Q$  is injective; consequently,  $M$  and  $F$  are injective too.*

*Proof.* Let  $\mathcal{X}, \mathcal{Y}$  be the family of all segments from  $x$ , respectively  $y$ , to  $z$ .

If  $y$  lies on a segment in  $\mathcal{X}$ , then  $\text{card } \mathcal{Y} = 1$ , whence  $z$  is conical by Lemma 10. If not, since any segment in  $\mathcal{X}$  meets any segment in  $\mathcal{Y}$  only at  $z$ , all segments of  $\mathcal{Y}$  lie in a single component  $D$  of  $S \setminus \bigcup_{\sigma \in \mathcal{X}} \sigma$ .

Thus, the directions of the segments in  $\mathcal{X}$  at  $z$  are separated in  $T_z$  from those of the segments in  $\mathcal{Y}$  at  $z$  by the points  $\alpha, \beta$ , say. The longer (if equally long, any) arc  $\Gamma$  from  $\alpha$  to  $\beta$  in  $T_z$  does not contain the direction  $\tau_\sigma$  at  $z$  of any segment  $\sigma$  of precisely one of the two families, say of  $\mathcal{X}$ . Then, if  $z$  were not conical, the distance in  $T_z$  from the midpoint of  $\Gamma$  to  $\tau_\sigma$  would be larger than  $\pi/2$  for any  $\sigma \in \mathcal{X}$ , contradicting  $z \in Q_x$ .

If  $S$  has no conical points, then, by Lemma 9,

$$F_x \cap F_y \subset M_x \cap M_y \subset Q_x \cap Q_y = \emptyset$$

for  $x \neq y$ . □

The preceding result is not extendable to orientable surfaces of higher genus or to nonorientable surfaces.

On any flat torus not obtainable by identification of the sides of a rectangle, every point is at diametral distance from other two points. The standard projective plane, too, has  $F$  noninjective in a strong way: for any two points  $x, y$  on it,  $F_x \cap F_y \neq \emptyset$ .

The sets  $M_y^{-1}$  containing more than a single point are investigated in [19].

#### 4 About the components of $Q_x$ and $M_x$

We start, once again, with two preparatory lemmas. The first of them complements Lemma 8.

**Lemma 11.** *On  $S \in \mathcal{A}$ , let  $\sigma, \sigma'$  be (possibly coinciding) segments from  $x$  to  $y$  and  $D$  a component of the complement of  $\sigma \cup \sigma'$  in an open disc around  $y$ . If the angle of  $\sigma, \sigma'$  at  $y$  toward  $D$  is smaller than  $\pi$ , then  $y$  is a strict relative maximum of  $\rho_x|_{\overline{D}}$ .*

This is easily proven using again Proposition 2.4 in [18].

**Lemma 12.** *On  $S \in \mathcal{A}$ , let  $\sigma, \sigma'$  be segments from  $x$  to  $y$  and  $D, D'$  the components of the complement of  $\sigma \cup \sigma'$  in an open disc around  $y$ . If  $y_n \in D, y'_n \in D', y_n \rightarrow y, y'_n \rightarrow y, \rho_x(y_n) \geq \rho_x(y)$ , and  $\rho_x(y'_n) \geq \rho_x(y)$ , then  $\sigma \cup \sigma'$  is a loop at  $x$ , and no segment joins  $x$  to  $y$  except for  $\sigma, \sigma'$ .*

This follows from Lemma 11.

**Theorem 4.** *Consider the surface  $S \in \mathcal{A}$ . For any point  $x \in S$ , every component of  $Q_x$  is a single point or an arc or a closed Jordan curve (the latter possibility being excluded if  $S \in \mathcal{A}_0$ ). If the component  $Q^1$  of  $Q_x$  is an arc then  $Q^1$ , possibly with one or both endpoints removed, is a component of  $M_x$ . Hence  $Q_x \setminus M_x$  is totally disconnected.*

*The closure of any component of  $M_x$  is a component of  $Q_x$ . This defines a natural injection from the family of components of  $M_x$  to the family of components of  $Q_x$ . Consequently, if  $Q$  is connected then  $M$  is connected too.*

*Proof.* Let the component  $Q^1$  of  $Q_x$  contain more than one point. Since  $C(x)$  is a local tree,  $Q^1$  includes an arc  $A$ . By Lemma 7,  $\rho_x$  is constant on  $A$ . So,  $\text{int } A \subset M_x$ . Since this is true for any arc in  $Q^1$ , by Theorem 1,  $Q^1$  cannot include a nondegenerate tree, so it must be an arc or a closed Jordan curve, the latter case being excluded if  $S \in \mathcal{A}_0$ .

Let now  $M^1$  be a component of  $M_x$ . By Lemma 9,  $M^1 \subset Q_x$ . If  $M^1$  is an arc with an endpoint  $e$  removed, we show that still  $e \in Q_x$ .

Let  $\sigma, \sigma'$  be the segments obtained as limits of the pairs of segments (see Lemma 2) from  $x$  to the points  $y_n \in M^1$  when  $y_n \rightarrow e$ .

Let  $O$  be a small open disc around  $e$ , with  $M^1 \setminus O \neq \emptyset$ . For  $n$  large enough,  $y_n \in O$  and the domain  $\Delta$  from Lemma 4 exists ( $J$  joining  $y_n$  to  $e$ ). Hence every point  $u \in \Delta \cap O$  lies on some segment from  $x$  to a point in  $M^1$ ; therefore  $\rho(x, u) \leq \rho(x, e)$ .

The segments  $\sigma, \sigma'$  cannot coincide, because arbitrarily close to  $e$  there are points farther than  $e$  from  $x$ . These points must be in the component of  $O \setminus (\sigma \cup \sigma')$  not meeting  $\Delta$ . By Lemma 12,  $\sigma \cup \sigma'$  is a loop at  $x$ , and  $e \in Q_x$ .

Hence  $\overline{M^1} \subset Q_x$ . To show that  $\overline{M^1}$  is a component of  $Q_x$  it suffices to exclude the existence of a second component  $M^2$  of  $M_x$  with  $\overline{M^1} \cap \overline{M^2} \neq \emptyset$ .

Suppose such a component  $M^2$  exists. Then it is an arc with the endpoint  $\{e\} = \overline{M^1} \cap \overline{M^2}$  removed, while  $M^1$  is also an arc with the endpoint  $e$  removed. As we already know, there are precisely two segments from  $x$  to  $e$ . Since this is also true for any point  $z \in M^1 \cup M^2$ , by Lemma 4,

$$C(x) \cap V \subset M^1 \cup M^2 \cup \{e\}$$

for some neighbourhood  $V$  of  $e$ . Then each point of  $V$  lies on some segment from  $x$  to  $M^1 \cup M^2 \cup \{e\}$ . Thus,  $e \in M_x$ , whence  $M^1$  is not a component of  $M_x$ , and a contradiction is obtained.  $\square$

## 5 Openness of the family $\mathcal{S}_2$ of surfaces with disconnected $M$

The set  $\mathcal{S}_2$  of all surfaces  $S \in \mathcal{A}_0$  on which there exists a point  $x$  with disconnected  $M_x$  has been introduced for  $S \in \mathcal{S}$  by the second author in [29], where it was stated that  $\mathcal{S}_2$  is open and shown that on most  $S \in \mathcal{S}_2 \cap \mathcal{S}$  there exists a point  $x$  for which  $M_x$  is infinite. We prove here that  $\mathcal{S}_2$  is open in  $\mathcal{S}$ , as well as in  $\mathcal{A}_0$ .

The set  $\mathcal{S}_1$  of all surfaces  $S \in \mathcal{A}_0$  on which  $F$  is a homeomorphism is not empty, as  $S^2 \in \mathcal{S}_1$ . In [20] we showed that, using  $\mathcal{C}^2$  metrics on both  $S$  and  $\mathcal{A}_0$ ,  $\mathcal{S}_1$  has non-empty interior. We do not know whether, equipped with the Hausdorff–Gromov distance,  $\mathcal{S}_1/\text{isometries}$  is nowhere dense or not. (See Question 1 in Section 9.) Theorem 6 will

show that  $S_1$  and  $S_2$  are disjoint. But are they complementary sets? (See Question 4 in Section 9.)

**Lemma 13.** *Let  $S \in \mathcal{A}_0$ ,  $x \in S$ , and  $P \subset C(x)$  be an arc from  $y$  to  $z$ . If  $\rho(x, P) < \rho(x, y)$  and  $\rho(x, P) < \rho(x, z)$ , then  $M_x$  has at least two closed components  $M^1, M^2$  separated by a loop  $\Lambda$  at  $x$ , such that  $\rho(x, v) < \rho(x, w)$  for any pair of points  $v \in \Lambda$ ,  $w \in M^1 \cup M^2$ .*

*Proof.* Let  $u \in P$  satisfy  $\rho(x, P) = \rho(x, u)$ . By Lemma 2, there is a loop  $\Lambda$  at  $x$  with midpoint  $u$ . Let the arc  $P^* \subset C(x)$  be a maximal extension of  $P$  with respect to inclusion. The point  $u$  divides  $P^*$  into two subarcs  $P_1, P_2$ . Let  $M^i$  be the set of all absolute maxima of  $\rho|_{P_i}$  ( $i = 1, 2$ ). Each component of  $M^i$  is closed. Moreover, since  $u \notin M^i$ , the sets  $M^1, M^2$  are separated by  $\Lambda$ . The proof ends with the remark that the components of each  $M^i$  are also components of  $M_x$ , and we find at least two such components, one in  $M^1$  and the other in  $M^2$ .  $\square$

**Lemma 14.** *Suppose  $S \in \mathcal{A}_0$  has a loop at  $x \in S$  of length less than  $2r_x$ . Then, for some point  $x' \in S$ ,  $M_{x'}$  has at least two closed components  $M^1, M^2$  separated by a loop  $\Lambda$  at  $x'$ , such that  $\rho(x', v) < \rho(x', w)$  for any pair of points  $v \in \Lambda$ ,  $w \in M^1 \cup M^2$ .*

*Proof.* Let  $\Lambda$  be the loop of length less than  $2r_x$ , with midpoint  $z$ . Let  $D$  be a component of  $S \setminus \Lambda$  containing some point  $v \in F_x$ . There are two consecutive segments  $\sigma, \sigma'$  from  $x$  to  $z$  in  $\overline{D}$  (i.e., there is no segment from  $x$  to  $z$  separating  $\text{int } \sigma$  from  $\text{int } \sigma'$  in  $\overline{D}$ ), with  $v$  between them. Possibly  $\sigma \cup \sigma' = \Lambda$ . Then  $C(z)$  contains a small arc  $A \subset \overline{D}$  starting at  $x$ , bisecting the angle of  $\sigma, \sigma'$  at  $z$ , and also bisecting the angle of suitably chosen segments  $\sigma_y, \sigma'_y$  from any  $y \in A$  to  $z$ , by Lemma 6 (see also Proposition 2.4 in [18]). Then  $\sigma_y \rightarrow \sigma$  and  $\sigma'_y \rightarrow \sigma'$  as  $y \rightarrow x$  (supposing  $\sigma, \sigma', \sigma'_y, \sigma_y$  in this order around  $z$ ). Let  $D_y$  be the component of  $S \setminus (\sigma_y \cup \sigma'_y)$  included in  $D$ .

For  $y$  close enough to  $x$ ,  $v \in D_y$  and  $\rho(y, v) > \rho(y, z)$ . Clearly, the angle of  $\sigma_y, \sigma'_y$  at  $z$  toward  $D_y$  is less than the angle of  $\sigma, \sigma'$  at  $z$  toward  $D$ , which in turn was not larger than  $\pi$ . Thus, by Lemma 11,  $z$  is a strict relative maximum of  $\rho_y|_{\overline{D}_v}$ .

Let  $P \subset C(y)$  be the arc from  $z$  to  $v$ . Clearly,  $P \subset D_v \cup \{z\}$ . Hence  $\rho(y, P)$  is smaller than both  $\rho(y, z), \rho(y, v)$ . Now, the conclusion follows from Lemma 13.  $\square$

**Theorem 5.** *The set  $S_2$  is open in  $\mathcal{A}_0$ . The set  $S_2 \cap \mathcal{S}$  is open in  $\mathcal{S}$ .*

*Proof.* We show that  $S_2 \cap \mathcal{S}$  is open in  $\mathcal{S}$ . Let  $S \in S_2 \cap \mathcal{S}$ . By Theorem 6 in the next section,  $S$  is not loopy, hence, by Lemma 14, for some  $x \in S$ ,  $M_x$  has at least two closed components, say  $M^1, M^2$ , separated by a loop  $\Lambda$  at  $x$ . Let

$$6\varepsilon \leq \min \left\{ \rho_x(M^1) - \frac{\lambda\Lambda}{2}, \rho_x(M^2) - \frac{\lambda\Lambda}{2} \right\}$$

and denote by  $\mathcal{H}$  the Pompeiu–Hausdorff distance in  $\mathcal{S}$ .

Let

$$\mathcal{O} = \{S' \in \mathcal{S} : \mathcal{H}(S, S') < \varepsilon\}.$$

Take the points

$$x = a_1, a_2, \dots, a_k, a_{k+1} = x \in \Lambda$$

with  $\rho(a_i, a_{i+1}) \leq \varepsilon$  ( $i = 1, \dots, k$ ).

Consider an arbitrary surface  $S' \in \mathcal{O}$ . We find points

$$y = b_1, b_2, \dots, b_k, b_{k+1} = y \in S'$$

with  $\|a_i - b_i\| < \varepsilon$ . This yields  $\rho(b_i, b_{i+1}) < 3\varepsilon$ . Let  $\sigma_i$  be a segment from  $b_i$  to  $b_{i+1}$  and  $\Sigma = \cup_{i=1}^k \sigma_i$ . For any point  $s \in \Sigma$ , there is some point  $a_i \in \Lambda$  such that  $\|a_i - s\| < 3\varepsilon$ . Therefore, on  $S'$ ,

$$\rho(y, s) \leq \|y - x\| + \frac{\lambda\Lambda}{2} + \|a_i - s\| < \frac{\lambda\Lambda}{2} + 4\varepsilon.$$

For  $u_i \in M^1$  there exists a point  $v_i \in S'$  satisfying  $\|u_i - v_i\| < \varepsilon$  ( $i = 1, 2$ ). For  $\varepsilon$  small enough,  $v_1$  and  $v_2$  are separated by  $\Sigma$  on  $S'$ . We have

$$\rho(y, v_i) \geq \rho(x, u_i) - \|x - y\| - \|u_i, v_i\| > \rho(M^i) - 2\varepsilon.$$

It follows that  $\rho(y, v_i) > \rho(y, s)$  for any  $s \in \Sigma$  and  $i \in \{1, 2\}$ . This proves that  $S' \in \mathcal{S}_2$ .

The same argument with the Hausdorff–Gromov metric instead of  $\mathcal{H}$  shows that  $\mathcal{S}_2$  is open in  $\mathcal{A}_0$ .  $\square$

## 6 About the connectedness of $Q$ , $M$ , $F$ , and the surjectivity of $F$

In [29] the second author introduced, for  $S \in \mathcal{S}$ , the *antipodal arc*  $J_x$ , which is the smallest arc in  $C(x)$  including  $F_x$ . Let us extend this notion for  $S \in \mathcal{A}_0$ , and call the smallest tree  $J_x$  in  $C(x)$  including  $F_x$  simply the *antipode* of  $x$ .

**Lemma 15.** *Suppose that  $S \in \mathcal{A}_0$ ,  $x \in S$  and  $\text{card } F_x \geq 2$ . Let  $z$  be an extremity of the antipode  $J_x$ , and  $\varepsilon > 0$ . Then there exist an arc  $A$  starting at  $x$  and a number  $k > 0$ , such that for any  $v \in A$  and any  $u \in B(v, k\rho(v, x))$ , we have  $F_u \subset B(z, \varepsilon)$ .*

The proof of Lemma 15 appears inside the proof of Theorem 5 in [29]. It is given for the convex case only. However, the same argument entirely applies for any  $S \in \mathcal{A}_0$ ; in particular the antipode of  $x$ , which is now a tree, should be used at the place of the antipodal arc.

**Theorem 6.** *For a surface  $S \in \mathcal{A}_0$ , the following implications hold.*

*$F$  is single-valued  $\iff F$  is continuous  $\implies F$  is surjective  $\implies M$  is surjective  $\implies S$  is loopy  $\iff Q$  is connected  $\iff M$  is connected  $\implies F$  is connected.*

*Proof.*  $F$  is single-valued  $\iff F$  is continuous.

The upper semicontinuity of arbitrary  $F$  becomes, for  $F$  single-valued, continuity. Conversely, suppose  $F$  is continuous and there exists  $x \in S$  such that  $\text{card } F_x > 1$ . By Lemma 15, there is a sequence of points  $x_n \in S \setminus \{x\}$  convergent to  $x$  such that

the sequence of sets  $F_{x_n}$  converges to a single-point set, therefore different from  $F_x$ , in contradiction with the continuity of  $F$  in  $x$ .

*F is continuous  $\implies F$  is surjective.*

Since  $F$  is continuous,  $F_S$  is closed. If  $F$  is not surjective, there is a small open disc  $D$  in  $S \setminus F_S$ . Clearly,  $F_{S \setminus D}$  is included in  $S \setminus D$ . By Brouwer's fixed point theorem,  $F|_{S \setminus D}$  has a fixed point, which is impossible.

*F is surjective  $\implies M$  is surjective.*

This follows from  $F_x \subset M_x$  for any  $x \in S$ .

*M is surjective  $\implies S$  is loopy.*

Suppose  $S$  is not loopy, i.e., there is a loop  $\Lambda$  at  $x \in S$  with midpoint  $y$  at distance less than  $r_x$  from  $x$ . By Lemma 14, for some point  $x' \in S$ ,  $M_{x'}$  has at least two closed components. Let  $A' \subset C(x')$  be the (unique) arc meeting each of the two preceding components in precisely one point. Let  $A$  be the set of all absolute minima of  $\rho|_{A'}$ . The components of  $A$  are points or arcs. Let  $u$  be a single-point component of  $A$  or, if such components are missing, an endpoint of a component of  $A$ . Then  $u \in Q_{x'} \setminus M_{x'}$  and, by Theorem 3,  $u \notin M_S$ .

*S is loopy  $\implies Q$  is connected.*

Suppose  $S$  is loopy, but  $Q_x$  has besides a component  $Q^0$  in  $F_x$  a further component  $Q^1$ . If  $y \in Q^1$  is a strict relative maximum of  $\rho_x$  or belongs to  $F_x$ , then take  $z \in Q^0$  and consider the arc  $J \subset C(x)$  joining  $y$  to  $z$ . Then  $\rho_x|_J$  has an absolute minimum at a point  $u$  different from  $y$  and  $z$ , and  $\rho_x(u) < r_x$ . By Lemma 2, there is a loop at  $x$  through  $u$  of length  $2\rho_x(u)$ , in contradiction to the assumption that  $S$  is loopy.

Suppose now that  $y \in Q^1$  is not a strict relative maximum of  $\rho_x$  and does not belong to  $F_x$ . Moreover, assume there is no loop at  $x$  with  $y$  as midpoint. Then the maximal angle at  $y$  between consecutive segments (if there are any consecutive segments) from  $x$  to  $y$  is less than  $\pi$ . By Lemma 11,  $y$  is a strict relative maximum of  $\rho_x$ , and we reached a contradiction. Hence there is a loop at  $x$  with midpoint  $y$ ; moreover its length is less than  $r_x$ . Hence  $S$  is not loopy, and a contradiction is obtained again.

*Q is connected  $\implies M$  is connected.*

This is part of Theorem 4.

*M is connected  $\implies S$  is loopy.*

This follows from Lemma 14.

*M is connected  $\implies F$  is connected.*

Indeed, for any  $x \in S$ ,  $\rho_x$  being constant on each component of  $M_x$ , it is constant on  $M_x$ ; since  $F_x \subset M_x$ , the constant is  $r_x$ , whence  $M_x = F_x$ .  $\square$

**Theorem 7.** *Let  $S$  belong to  $\mathcal{A}_0$ . If  $F_x$  is totally disconnected for each  $x \in S$ , then all implications in Theorem 6 are valid in both directions.*

*This is in particular true for polyhedral convex surfaces and for most convex surfaces.*

*Proof.* Indeed, by Theorem 1, for any surface  $S \in \mathcal{A}_0$  and any point  $x \in S$ , if  $F_x$  is connected, then it is either a point or an arc. Since the second possibility is excluded,  $F_x$  contains a single point. On most convex surfaces, there is no point  $x$  with an arc in  $F_x$ , by Theorem 2.

For a polyhedral convex surface  $P$ , one can easily see that, for any point  $x$  in  $P$ , the only points in  $F_x$  which are joined to  $x$  by precisely two segments are among the vertices of  $P$  (see also [31]). If  $F_x$  contained an arc then, by Lemma 2, all points interior to that arc would be joined to  $x$  by precisely two segments, too many points to be among the vertices of  $P$ .  $\square$

**Remark.** The set of all surfaces  $S \in \mathcal{S}$  with  $F$  surjective is closed in  $\mathcal{S}$ .

Since a geodesic arriving at an endpoint cannot go beyond it, the endpoint is a kind of “farthest point” on that geodesic. This may suggest that endpoints of  $S \in \mathcal{S}$  might always lie in  $F_x$  for some point  $x \in S$ . However, this is deeply false, as every surface  $S \in \mathcal{S}_2 \cap \mathcal{S}$  has, by Theorem 6, an open set  $O$  disjoint from  $F_S$ , while, for most surfaces in  $\mathcal{S}$ ,  $O$  contains lots of endpoints, by Lemma 5.

## 7 Sufficient conditions for a surface to belong to $\mathcal{S}_2$

We shall make use of a hinge variant of Toponogov’s well-known comparison theorem (see, for example, [6]), which will now be recalled. Let  $M_H$  be the simply connected 2-dimensional space of constant curvature  $H$ .

**Lemma 16.** *Let  $M$  be a complete manifold with sectional curvature  $K \leq H$ , and let  $(\gamma_1, \gamma_2, \gamma_3)$  be a geodesic triangle in  $M$ . If  $H > 0$ , suppose  $\lambda\gamma_i \leq \pi/\sqrt{H}$  ( $i = 1, 2$ ).*

*Then there exists in  $M_H$  a geodesic triangle  $(\bar{\gamma}_1, \bar{\gamma}_2, \bar{\gamma}_3)$  such that  $\lambda\gamma_i = \lambda\bar{\gamma}_i$  ( $i = 1, 2$ ),  $\angle\gamma_1\gamma_2 = \angle\bar{\gamma}_1\bar{\gamma}_2$ , and  $\lambda\gamma_3 \geq \lambda\bar{\gamma}_3$ .*

**Theorem 8.** *Suppose the surface  $S \in \mathcal{A}_0$  is of class  $C^2$  in a neighbourhood of a loop  $\Lambda$  of length  $l$ . If  $S$  has Gauß curvature  $K < \pi^2/l^2$  along  $\Lambda$ , then  $S \in \mathcal{S}_2$ . In particular, this is true if  $K$  is nonpositive along  $\Lambda$ .*

*Proof.* Choose  $x, y \in S$  so that  $\Lambda$  is a loop at  $x$  with midpoint  $y$ . Using the continuity of  $K$ , we find a neighbourhood  $V \subset S$  of  $\Lambda$  and a number  $k < \pi^2/l^2$  such that  $K < k$  in  $V$ .

Let  $yz \in V$  be a segment orthogonal to  $\Lambda$  at  $y$ . Let  $\bar{x}, \bar{y}, \bar{z}$  be points on the sphere  $M_k$  of curvature  $k$ , such that  $\rho_0(\bar{x}, \bar{y}) = \pi/(2\sqrt{k})$ ,  $\rho_0(\bar{y}, \bar{z}) = \rho(y, z)$  and  $\angle\bar{x}\bar{y}\bar{z} = \pi/2$ ; here,  $\rho_0$  is the standard metric of  $M_k$ . It follows that  $\bar{y}$  belongs to the equator farthest from  $\bar{x}$ , whence  $\rho_0(\bar{x}, \bar{z}) = \rho_0(\bar{x}, \bar{y})$ . By Lemma 16, we have

$$\rho(x, z) \geq \rho_0(\bar{x}, \bar{z}) = \frac{\pi}{2\sqrt{k}} > \frac{l}{2} = \rho(x, y).$$

Hence  $y \notin F_x$  and  $S$  is not loopy. By Theorem 6,  $S \in \mathcal{S}_2$ .  $\square$

**Examples.** Theorem 8 provides compact surfaces of revolution in  $\mathcal{S}_2$ , for example those with Gauß curvature  $K < 1/(4r^2)$  along an equator  $\Lambda$  of radius  $r$ . Among them we find the ellipsoids with semi-axes  $a, b, c$  verifying  $a = b$  and  $c > 2a$ .

We can also apply Theorem 8 to boundaries of rectangular boxes and to doubly covered polygons, all of which admit a loop  $\Lambda$  with a neighbourhood isometric to an open subset of a cylinder or a cone.

**Remark.** One can, of course, formulate Theorem 8 in terms of an upper bound to the curvature, in the sense of Alexandrov, in addition to the lower bound that our surfaces have by definition. But the goal being to give an easy-to-apply criterion, we chose to use Gauß curvature.

**Theorem 9.** *If on  $S \in \mathcal{A}$  there exists a point  $y \in S$  such that  $\lambda T_y < \pi$ , then  $S \in \mathcal{S}_2$ .*

*Proof.* By Lemma 11,  $y$  is a strict relative maximum of  $\rho_x$  for any point  $x \in S \setminus \{y\}$ . Choosing  $x$  close enough to  $y$  guarantees that  $y \notin F_x$ , whence  $M_x$  does not consist of the isolated point  $y$  alone, and is therefore disconnected.  $\square$

**Theorem 10.** *If  $S \in \mathcal{A}_0$  and  $\text{rad } S = \text{diam } S/2$  then  $S \in \mathcal{S}_2$ , except for the case of a surface  $S$  with precisely two conical points  $y, z$  where  $\lambda T_y = \lambda T_z = \pi$ , and with a closed geodesic  $\Lambda$  so that  $\lambda \Lambda = \text{diam } S$ ,  $F_\Lambda = S$ , and, for each point  $x \in \Lambda$ ,  $r_x = \text{rad } S$  and  $F_x$  is an arc from  $y$  to  $z$ .*

*Proof.* Suppose  $S \notin \mathcal{S}_2$ , i.e., for each point  $x \in S$ ,  $M_x$  is connected and, since  $F_x \subset M_x$  and  $\rho_x$  is constant on  $M_x$ ,  $F_x = M_x$ .

Let  $x, y, z \in S$  satisfy  $r_x = \text{rad } S$  and  $\rho(y, z) = \text{diam } S$ . Then

$$\text{diam } S = 2r_x \geq \rho(x, y) + \rho(x, z) \geq \rho(y, z) = \text{diam } S,$$

whence  $y, z \in F_x$  and  $x$  belongs to a segment  $\Gamma_{yz}$  from  $y$  to  $z$ . Thus,  $\{y, z\} \subset C(x) \setminus C_x$ , whence  $\lambda T_y$  and  $\lambda T_z$  are, by Lemma 9, not larger than  $\pi$ .

By Theorem 1,  $F_x$  is an arc. The points  $y$  and  $z$  belong to  $F_x$  and are joined to  $x$  by unique segments. Since each point  $v$  interior to  $F_x$  is a relative minimum for  $\rho_x|_{F_x}$  and hence is the mid-point of a loop  $\Lambda_v$  at  $x$  by Lemma 2,  $y$  and  $z$  must be the endpoints of the arc  $F_x$ .

Now put  $\Delta = S \setminus \Gamma_{yz}$  and apply Lemma 4. We obtain  $\Delta = \bigcup_{v \in \text{int } F_x} \Lambda_v$ , so  $S$  has no conical points, except for  $y, z$ .

Consider now the set  $E(y, z)$  of all points in  $S$  at equal distance from  $y$  and  $z$ . Each arc from  $y$  to  $z$  meets  $E(y, z)$ .

For an arbitrary point  $w \in E(y, z) \setminus \{x\}$ , denote by  $\Lambda_w$  the loop at  $x$  through  $w$ ; so  $\Lambda_w$  separates  $y$  from  $z$ . Since  $y, z$  are points of the tree  $C(w)$ , there exists a unique arc  $J_w \subset C(w)$  joining them. Consider a point  $w' \in J_w \cap \Lambda_w$ . Then

$$\rho(w, w') \leq \lambda \Lambda_w / 2 = \text{rad } S.$$

We also have

$$2 \text{rad } S = \rho(y, z) \leq \rho(w, y) + \rho(w, z) = 2\rho(w, y),$$

so we obtain  $\rho(w, w') \leq \rho(w, y)$ . If  $\rho(w, w') < \rho(w, y) = \rho(w, z)$  then, by Lemma 13,  $S \in \mathcal{S}_2$ , which, as we assumed, is not the case. Thus,

$$\rho(w, w') = \lambda \Lambda_w / 2 = \text{rad } S = \rho(w, y) = \rho(w, z).$$

From

$$\rho(y, z) = \text{diam } S = \rho(w, y) + \rho(w, z),$$

we obtain that  $w$  is the mid-point of a segment from  $y$  to  $z$ , and the segments joining  $w$  to  $y$  and  $z$  are unique. Since  $M_w$  is connected, we have  $M_w = F_w$ , and since  $y, z \in F_w$ , we get, as before,  $F_w = J_w$ .

Assume now  $w \notin F_x$ . The equality  $\rho(w, w') = \lambda\Lambda_w/2$  implies that  $\Lambda_w$  is a closed geodesic, consequently its directions at  $x$  make an angle of  $\pi$ .

Suppose there exists a point  $w^* \in E(y, z) \setminus (F_x \cup \Lambda_w)$ . The above arguments show that  $\Lambda_{w^*}$  is a closed geodesic. Then  $\Lambda_w \cap \Lambda_{w^*} = \{x\}$  implies that the directions of  $\Lambda_{w^*}$  at  $x$  do not separate in  $T_x$  those of  $\Lambda_w$  and are different from them, so their angle is less than  $\pi$ , and a contradiction is obtained. Hence

$$E(y, z) \subset F_x \cup \Lambda_w.$$

To show that  $\Lambda_w \subset E(y, z)$ , let  $x' \in \Lambda_w$  be chosen arbitrarily. Choose an arc  $A$  joining  $y$  to  $z$ , disjoint from  $(F_x \cup \Lambda_w) \setminus \{x', y, z\}$ . This arc obviously meets  $E(y, z)$  in a point different from  $y$  and  $z$ . Due to the preceding inclusion, the point must be  $x'$ . We saw that  $u \in E(y, z)$  yields

$$r_u = \text{rad } S = \rho(u, y) = \rho(u, z).$$

So, if  $u \notin \Lambda_w$ , it is separated from  $y$  or  $z$  by  $\Lambda_w$ , say from  $y$ . Then any segment from  $u$  to  $y$  meets  $\Lambda_w$  at a point at distance  $\text{rad } S$  from  $y$ , whence  $\rho(u, y) > \text{rad } S$ , and a contradiction is obtained.

Put  $\Lambda = E(y, z)$ . Finally, we have to prove that  $F_\Lambda = S$ . To see this, notice first that, by Lemma 4,  $C(w) = F_w$  for all points  $w \in \Lambda$ . Let now  $v \in S \setminus \{y, z\}$ . Since  $y, z$  are conical, they belong to  $C(v)$ . The arc  $J \subset C(v)$  joining  $y$  to  $z$  clearly meets  $\Lambda$ , say at  $v'$ . Then  $v \in C(v') = F_{v'}$ , by the preceding remark. And, obviously,  $y, z \in F_\Lambda$ , too.  $\square$

The actual existence of an exceptional surface as described in Theorem 10 is illustrated by the example considered in the next section.

## 8 Farthest points on a Tannery surface

We investigate here the special case of a Tannery surface, particularly interesting for our purposes.

A Riemannian surface  $(S, g)$  is called a  $P_l$ -surface if all of its geodesics are periodic with least common period  $l$  ([3], p. 182). A  $P_l$ -surface of revolution (i.e., having  $S^1$  as an effective isometry group) is called a *Tannery surface* (see [3], pp. 95 and 102).

On the unit sphere  $S^2$ , let  $n$  and  $s$  be the North and South poles, and consider a point  $x \in S^2 \setminus \{n, s\}$ . Denote by  $G_x$  the great circle through  $n$  and  $x$ , and let  $2r$  be the distance from  $n$  to  $x$  (realized on  $G_x$ ) and  $\theta$  the angle made by the plane of  $G_x$  with a fixed plane through  $n$  and  $s$ .

Consider, for the set  $U = S^2 \setminus \{n, s\}$ , the parametrization  $(r, \theta)$  described before, with  $r \in ]0, \pi/2[$ ,  $\theta \in [0, 2\pi[$ , and endow  $U$  with the metric  $g = 4dr^2 + \sin^2 r d\theta$ . By Proposition 4.6 in [3] p. 96, the metric  $g$  extends (only) to a  $C^0$  metric on  $S^2$ .

We obtain, from considerations in [3] and an application of Theorem 10, the following about  $(S^2, g)$ .

- Theorem 11.** a)  $(S^2, g)$  is a Tannery surface (with parameters  $p = 2, q = 1$ ).
- b)  $(S^2, g)$  may be isometrically embedded in the Euclidean space  $\mathbb{R}^3$  as a convex surface of revolution  $(S, \rho)$ , whose half-meridian (joining the images of the poles through the isometry, denoted again by  $n$  and  $s$ ) is described by  $c(R) = \pm \int_R^1 \sqrt{\frac{4}{1-u^2} - 1} du$ , where  $R = \sin r$ .
- c) Except for the subsegments of half-meridians, every geodesic extends to a closed geodesic of  $S$ . Apart from the equator  $E$ , which has length  $2\pi$ , every closed geodesic  $\Gamma$  consists of  $2q = 2$  arcs between two consecutive points of tangent contact with the parallels;  $\Gamma$  has length  $4q\pi = 4\pi$  and turns  $p = 2$  times. The length of a half-meridian is  $2\pi$ .
- d)  $\text{diam } S = 2 \text{ rad } S$ . For each point  $x \in E$ ,  $F_x$  is the half-meridian opposed to  $x$ .
- e)  $\lambda T_n = \lambda T_s = \pi$ ,  $F_E = S$ , and  $S \notin \mathcal{S}_2$ .

*Proof.* a) and c) follow from Theorems 4.11 (p. 100) and 4.13 (p. 102) in [3], applied to the particular metric  $g$ .

Concerning b), the isometric embedding of  $(S^2, g)$  in  $\mathbb{R}^3$  is a consequence of Propositions 4.18 (p. 105) and 4.20 (p. 107) (in our case  $h \equiv 0$ ), while the convexity of the surface follows from Proposition 4.23 and Remark 4.21 (pp. 108–109), all from [3].

We prove now d). The symmetry of  $S$  and  $p = 2$  imply that each closed geodesic  $\Gamma$  has precisely one self-intersection point  $x_\Gamma$ , which must lie on  $E$ , and two points  $z_\Gamma, z'_\Gamma$  on the half-meridian opposite to  $x_\Gamma$ , at which  $\Gamma$  and the half-meridian are orthogonal.

Let  $x \in E, y \in F_x$ , and  $\sigma$  be a segment from  $x$  to  $y$ . Then the closed geodesic  $\Gamma$  including  $\sigma$  has  $x_\Gamma = x$  and  $y$  on one of the four arcs joining  $x$  to  $z_\Gamma, z'_\Gamma$ , each of length  $\pi$ . Hence  $y$  equals  $z_\Gamma$  or  $z'_\Gamma$  and  $\rho(x, y) = \pi$ . By choosing  $x \in E$  and  $y$  in the half-meridian  $H_x$  opposite to  $x$ , the same argument shows that  $y \in F_x$ , so  $F_x = H_x$ .

Since  $\rho(n, s) = 2\pi$  by c), and any segment not included in a meridian extends to a closed geodesic and has therefore length at most  $2\pi$ , we have indeed  $\text{diam } S = 2 \text{ rad } S$ .

To prove e), first note that  $F_E = \bigcup_{x \in E} H_x = S$ . If  $x \in E$ , then  $M_x = H_x$ . Clearly,  $M_n = \{s\}$  and  $M_s = \{n\}$ . If  $x \notin E \cup \{n, s\}$ , then  $C(x) = H_x$ . For any point  $y \in H_x$ , there are precisely two closed geodesics passing through  $x$  and  $y$ , and they have of course different directions at  $y$ . Therefore, by Lemmas 8 and 11,  $\rho_x$  is strictly monotone on  $H_x \setminus \{n, s\}$ . It follows that  $M_x$  equals  $\{n\}$  or  $\{s\}$ . Hence  $S \notin \mathcal{S}_2$ .

Finally,  $\lambda T_n = \lambda T_s = \pi$  follows from Theorem 10, because every point but  $n$  and  $s$  is interior to a geodesic and therefore not conical.  $\square$

Our Tannery surface has  $F$  surjective and properly multivalued; thus, the converse of the second implication in Theorem 6 is not true.

## 9 Nine open questions

We restrict our open questions to the perhaps easier but certainly important convex case, but most of them make sense in  $\mathcal{A}_0$  or even  $\mathcal{A}$ . Let us start with the problem mentioned already in the Abstract.

**Question 1.** Is  $\mathcal{S}_2$  dense in  $\mathcal{S}$ ?

The case of the regular tetrahedron shows that  $F_S$  can be connected even if  $S \in \mathcal{S}_2$ , as J. Rouyer proved in [12]. In most of our examples, however, the set  $F_S$  is disconnected as soon as  $F$  is not surjective.

**Question 2.** For which convex polyhedral surfaces  $S$  is  $F_S$  connected?

**Question 3.** Consider a surface  $S \in \mathcal{S}$  such that  $F_S \neq S$ . Do there always exist points  $y \in \text{bd } F_S$  and  $x \in F_y^{-1}$  such that  $\text{card } F_x \geq 2$ ?

It is natural to ask whether  $\{\mathcal{S}_1, \mathcal{S}_2\}$  forms a partition of  $\mathcal{A}_0$ . This reduces to the following problem.

**Question 4.** Do convex surfaces with  $F$  single-valued and noninjective exist?

Several possible implications between statements in Theorem 6 are neither proved nor disproved, so far.

**Question 5.** Does connectedness of  $F$  imply connectedness of  $M$ ?

**Question 6.** Does surjectivity of  $M$  imply surjectivity of  $F$ ?

Of course, surjectivity of  $M$  implies surjectivity of  $Q$ . But what is the relationship between the surjectivity of  $Q$  and the loopiness of  $S$ ?

**Question 7.** Does surjectivity of  $Q$  imply loopiness of  $S$ ? Or vice-versa?

J. Rouyer proved that all tetrahedra in  $\mathbb{R}^3$  belong to  $\mathcal{S}_2$  (see [12], [15]).

**Question 8.** Which convex polyhedral surfaces do not belong to  $\mathcal{S}_2$ ?

We believe that there are no surfaces different from the Tannery surface described in Section 8 playing the exceptional role in Theorem 10. However this is not proven yet.

**Question 9.** Is the Tannery surface from the preceding section the unique exception in Theorem 10?

## References

- [1] A. D. Alexandrow, *Die innere Geometrie der konvexen Flächen*. Akademie-Verlag, Berlin 1955. [MR0071041 \(17,74d\) Zbl 0065.15102](#)
- [2] M. Berger, Les variétés Riemanniennes  $(1/4)$ -pincées. *Ann. Scuola Norm. Sup. Pisa* (3) **14** (1960), 161–170. [MR0140054 \(25 #3478\) Zbl 0096.15502](#)
- [3] A. L. Besse, *Manifolds all of whose geodesics are closed*, volume 93 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, Berlin 1978. [MR496885 \(80c:53044\) Zbl 0387.53010](#)

- [4] Y. Burago, M. Gromov, G. Perel'man, A. D. Aleksandrov spaces with curvatures bounded below. *Uspekhi Mat. Nauk* **47** (1992), no. 2(284), 3–51, 222.  
[MR1185284 \(93m:53035\)](#) [Zbl 0802.53018](#)
- [5] H. Busemann, *Convex surfaces*. Interscience Publ., Inc., New York 1958.  
[MR0105155 \(21 #3900\)](#) [Zbl 0196.55101](#)
- [6] J. Cheeger, D. G. Ebin, *Comparison theorems in Riemannian geometry*. North-Holland, Amsterdam 1975. [MR0458335 \(56 #16538\)](#) [Zbl 0309.53035](#)
- [7] H. T. Croft, K. J. Falconer, R. K. Guy, *Unsolved problems in geometry*. Springer-Verlag, New York 1991. [MR1107516 \(92c:52001\)](#) [Zbl 0748.52001](#)
- [8] P. M. Gruber, Baire categories in convexity. In: *Handbook of convex geometry, Vol. A, B*, 1327–1346, North-Holland, Amsterdam 1993. [MR1243011 \(94i:52003\)](#) [Zbl 0791.52002](#)
- [9] V. Klee, Some new results on smoothness and rotundity in normed linear spaces. *Math. Ann.* **139** (1959), 51–63. [MR0115076 \(22 #5879\)](#) [Zbl 0092.11602](#)
- [10] S. Kobayashi, On conjugate and cut loci. In: *Global differential geometry*, volume 27 of *MAA Stud. Math.*, 140–169, 1989 (second ed.). [MR0212737 \(35 #3603\)](#)  
[MR1013807 \(90d:53003\)](#) [Zbl 0683.53043](#)
- [11] Y. Otsu, T. Shioya, The Riemannian structure of Alexandrov spaces. *J. Differential Geom.* **39** (1994), 629–658. [MR1274133 \(95e:53062\)](#) [Zbl 0808.53061](#)
- [12] J. Rouyer, Antipodes sur un tétraèdre régulier. *J. Geom.* **77** (2003), 152–170.  
[MR2000860 \(2004k:52016\)](#) [Zbl 1032.52004](#)
- [13] J. Rouyer, On antipodes on a manifold endowed with a generic Riemannian metric. *Pacific J. Math.* **212** (2003), 187–200. [MR2016977 \(2004m:53056\)](#) [Zbl 1060.53044](#)
- [14] J. Rouyer, On farthest points on Riemannian manifolds homeomorphic to the 2-dimensional sphere. *Rev. Roumaine Math. Pures Appl.* **48** (2003), 95–103.  
[MR1998063 \(2004e:53103\)](#) [Zbl 1052.53049](#)
- [15] J. Rouyer, On antipodes on a convex polyhedron. *Adv. Geom.* **5** (2005), 497–507.  
[MR2174479](#) [Zbl 1085.52002](#)
- [16] T. Sakai, *Riemannian geometry*. American Mathematical Society, Providence, RI 1996.  
[MR1390760 \(97f:53001\)](#) [Zbl 0886.53002](#)
- [17] R. Schneider, *Convex bodies: the Brunn-Minkowski theory*. Cambridge University Press, Cambridge 1993. [MR1216521 \(94d:52007\)](#) [Zbl 0798.52001](#)
- [18] K. Shiohama, M. Tanaka, Cut loci and distance spheres on Alexandrov surfaces. In: *Actes de la Table Ronde de Géométrie Différentielle (Luminy, 1992)*, volume 1 of *Sémin. Congr.*, 531–559, Soc. Math. France, Paris 1996. [MR1427770 \(98a:53062\)](#) [Zbl 0874.53032](#)
- [19] C. Vilcu, Common maxima of distance functions on convex surfaces. To appear.
- [20] C. Vilcu, T. Zamfirescu, Symmetry and the farthest point mapping on convex surfaces. *Adv. Geometry*, **6** (2006), 379–387.
- [21] T. Zamfirescu, Many endpoints and few interior points of geodesics. *Invent. Math.* **69** (1982), 253–257. [MR674405 \(84h:53088\)](#) [Zbl 0494.52004](#)
- [22] T. Zamfirescu, Baire categories in convexity. *Atti Sem. Mat. Fis. Univ. Modena* **39** (1991), 139–164. [MR1111764 \(92c:52002\)](#) [Zbl 0780.52003](#)
- [23] T. Zamfirescu, Long geodesics on convex surfaces. *Math. Ann.* **293** (1992), 109–114.  
[MR1162676 \(93d:53082\)](#) [Zbl 0735.53032](#)

- [24] T. Zamfirescu, On some questions about convex surfaces. *Math. Nachr.* **172** (1995), 313–324. [MR1330637 \(96e:52004\)](#) [Zbl 0833.53004](#)
- [25] T. Zamfirescu, Points joined by three shortest paths on convex surfaces. *Proc. Amer. Math. Soc.* **123** (1995), 3513–3518. [MR1273530 \(96a:52001\)](#) [Zbl 0851.52003](#)
- [26] T. Zamfirescu, Conjugate points and closed geodesic arcs on convex surfaces. *Geom. Dedicata* **62** (1996), 99–105. [MR1400984 \(97g:52006\)](#) [Zbl 0866.53050](#)
- [27] T. Zamfirescu, Closed geodesic arcs in Aleksandrov spaces. *Rend. Circ. Mat. Palermo (2) Suppl.* no. **50** (1997), 425–430. [MR1603026 \(99b:53063\)](#) [Zbl 0889.53027](#)
- [28] T. Zamfirescu, Farthest points on convex surfaces. *Math. Z.* **226** (1997), 623–630. [MR1484714 \(98m:52003\)](#) [Zbl 0896.52005](#)
- [29] T. Zamfirescu, Extreme points of the distance function on convex surfaces. *Trans. Amer. Math. Soc.* **350** (1998), 1395–1406. [MR1458314 \(98i:52005\)](#) [Zbl 0896.52006](#)
- [30] T. Zamfirescu, On the critical points of a Riemannian surface. *Adv. Geom.* **6** (2006), 493–500.
- [31] T. Zamfirescu, On the number of shortest paths between points on manifolds. *Rend. Circ. Mat. Palermo, Suppl.*, to appear.
- [32] C. Zong, *Strange phenomena in convex and discrete geometry*. Springer-Verlag, New York 1996. [MR1416567 \(97m:52001\)](#) [Zbl 0865.52001](#)

Received 29 September, 2005; revised 8 February, 2006

C. Vîlcu, Institute of Mathematics of the Romanian Academy, P.O. Box 1-764, Bucharest 014700, Romania

Email: [Costin.Vilcu@imar.ro](mailto:Costin.Vilcu@imar.ro)

T. Zamfirescu, Fachbereich Mathematik, Universität Dortmund, 44221 Dortmund, Germany

Email: [tudor.zamfirescu@mathematik.uni-dortmund.de](mailto:tudor.zamfirescu@mathematik.uni-dortmund.de)