Dissipative backward stochastic differential equations with locally Lipschitz nonlinearity.

Fulvia Confortola

Dipartimento di Matematica, Politecnico di Milano piazza Leonardo da Vinci 32, 20133 Milano, Italy fulvia.confortola@polimi.it

November 4, 2018

Abstract

In this paper we study a class of backward stochastic differential equations (BSDEs) of the form

 $dY_t = -AY_t dt - f_0(t, Y_t) dt - f_1(t, Y_t, Z_t) dt + Z_t dW_t, \ 0 \le t \le T; \ Y_T = \xi$

in an infinite dimensional Hilbert space H, where the unbounded operator A is sectorial and dissipative and the nonlinearity $f_0(t, y)$ is dissipative and defined for y only taking values in a subspace of H. A typical example is provided by the so-called polynomial nonlinearities. Applications are given to stochastic partial differential equations and spin systems.

Key words Backward stochastic differential equations, stochastic evolution equations.

MSC classification. Primary: 60H15 Secondary: 35R60

1 Introduction

Let H, K be real separable Hilbert spaces with norms $|\cdot|_H$ and $|\cdot|_K$. Let W be a cylindrical Wiener process in K defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and let $\{\mathcal{F}_t\}_{t\in[0,T]}$ denote its natural augmented filtration. Let $\mathcal{L}^2(K, H)$ be the Hilbert space of Hilbert-Schmidt operators from K to H.

We are interested in solving the following backward stochastic differential equation

$$dY_t = -AY_t dt - f(t, Y_t, Z_t) dt + Z_t dW_t, \quad 0 \le t \le T, \quad Y_T = \xi \tag{1}$$

where ξ is a random variable with values in H, $f(t, Y_t, Z_t) = f_0(t, Y_t) + f_1(t, Y_t, Z_t)$ and f_0, f_1 are given functions, and the operator A is an unbounded operator with domain D(A) contained in H. The unknowns are the processes $\{Y_t\}_{t\in[0,T]}$ and $\{Z_t\}_{t\in[0,T]}$, which are required to be adapted with respect to the filtration of the Wiener process and take values in H, $\mathcal{L}^2(K, H)$ respectively.

In finite dimensional framework such type of equations has been solved by Pardoux and Peng [12] in the nonlinear case. They proved an existence and uniqueness result for the solution of the equation (1) when A = 0, the coefficient f(t, y, z) is Lipschitz continuous in both variables y and z, and the data ξ and the process $\{f(t, 0, 0)\}_{t \in [0,T]}$ are square integrable. Since this first result, many papers were devoted to existence and uniqueness results under weaker assumptions. In finite dimension, when A = 0, the Lipschitz condition on the coefficient f with respect to the variable y is replaced by a monotonicity assumption; moreover, more general growth conditions in the variable y are formulated. Let us mention the contribution of Briand and Carmona [1], for a study of polynomial growth in L^p with p > 2, and the work of Pardoux [11] for an arbitrary growth. In [13] Pardoux and Rascanu deal with a BSDE involving the subdifferential of a convex function; in particular, one coefficient is not everywhere defined for y in \mathbb{R}^k .

In other works the existence of the solution is proved when the data, ξ and the process $\{f(t,0,0)\}_{t\in[0,T]}$, are in L^p for $p \in (1,2)$. El Karoui, Peng and Quenez [4] treat the case when f is Lipschitz continuous; in [2] this result is generalized to the case of a monotone coefficient f (both for equations on a fixed and on a random time interval) and is studied even the case p = 1.

In the infinite-dimensional framework Hu and Peng [6], and Oksendal and Zhang [10] give an existence and uniqueness result for the equation with an operator A, infinitesimal generator of a strongly continous semigroup and the coefficient f Lipschitz in y and z. Pardoux and Rascanu [14] replace the operator A with the subdifferential of a convex function and assume that f is dissipative, everywhere defined and continuous with respect to y, Lipschitz with respect to z and with linear growth in y and z.

Special results deal with stochastic backward partial differential equations (BSPDEs): we recall in particular the works of Ma and Yong [8] and [9]. Earlier, Peng [16] studied a backward stochastic partial differential equation and regarded the classical Hamilton-Jacobi-Bellman equation of optimal stochastic control as special case of this problem.

Our work extends these results in a special direction. We consider an operator A which is the generator of an analytic contraction semigroup on H and a coefficient f(t, y, z) of the form $f_0(t, y) + f_1(t, y, z)$. The coefficient $f_1(t, y, z)$ is assumed to be bounded and Lipschitz with respect to y and z. The term $f_0(t, y)$ is defined for y only taking values in a suitable subspace H_{α} of H and it satisfies the following growth condition for some $1 < \gamma < 1/\alpha$, $S \geq 0$, \mathbb{P} -a.s.

$$|f_0(t,y)|_H \le S(1+||y||_{H_\alpha}^{\gamma}) \quad \forall t \in [0,T], \quad \forall y \in H_\alpha.$$

Following [6], we understand the equation (1) in the following integral form

$$Y_t - \int_t^T e^{(s-t)A} [f_0(s, Y_s) + f_1(s, Y_s, Z_s)] ds + \int_t^T e^{(s-t)A} Z_s dW_s = e^{(T-t)A} \xi,$$
(2)

requiring, in particular, that Y takes values in H_{α} . This requires generally that the final condition also takes values in the smaller space H_{α} . We take as H_{α} a real interpolation space which belongs to the class J_{α} between H and the domain of an operator A (see Section 2). Moreover $f_0(t, \cdot)$ is assumed to be locally Lipschitz from H_{α} into H and dissipative in H. We prove (Theorem 5) that if ξ takes its values in the closure of D(A) in H_{α} and is such that $||\xi||_{H_{\alpha}}$ is essentially bounded, then equation (2) has a unique mild solution, i.e. there exists a unique pair of progressively measurable processes $Y: \Omega \times [0,T] \to H_{\alpha}, Z: \Omega \times [0,T] \to \mathcal{L}^2(K;H)$, satisfying P-a.s. equality (2) for every t in [0,T] and such that $\mathbb{E} \sup_{t \in [0,T]} ||Y_t||_{H_{\alpha}}^2 + \mathbb{E} \int_0^T ||Z_t||_{\mathcal{L}^2(K,H)}^2 dt < \infty$.

This result extends former results concerning the deterministic case to the stochastic framework: see [7], where previous works of Fujita - Kato [5], Pazy [15] and others are collected. In these papers similar assumptions are made on the coefficients f_0 , f_1 and on the operator A.

The plan of the paper is as follows. In Section 2 some notations and definitions are fixed. In Section 3 existence and uniqueness of the solution of a simplified equation are proved, where f_1 is a bounded progressively measurable process which does not depend on y and z. In Section 4, applying the previous result, a fixed point argument is used in order to prove our main result on existence and uniqueness of a mild solution of (2). Section 5 is devoted to applications.

2 Notations and setting

The letters K and H will always denote two real separable Hilbert spaces. Scalar product is denoted by $\langle \cdot, \cdot \rangle$; $\mathcal{L}^2(K; H)$ is the separable Hilbert space of Hilbert-Schmidt operators from K to H endowed with the Hilbert-Schmidt norm. $W = \{W_t\}_{t \in [0,T]}$ is a cylindrical Wiener process with values in K, defined on a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$. $\{\mathcal{F}_t\}_{t \in [0,T]}$ is the natural filtration of W, augmented with the family of \mathbb{P} -null sets of \mathcal{F} .

Next we define several classes of stochastic processes with values in a Banach space X.

- $L^2(\Omega \times [0,T];X)$ denotes the space of measurable X-valued processes Y such that $\left[\mathbb{E}\int_0^T |Y_{\tau}|^2 d\tau\right]^{1/2}$ is finite, identified up to modification.
- $L^2(\Omega; C([0, T]; X))$ denotes the space of continuous X-valued processes Y such that $\left[\mathbb{E}\sup_{\tau\in[0,T]}|Y_{\tau}|^2\right]^{1/2}$ is finite, identified up to indistinguishability.
- $C^{\alpha}([0,T];X)$ denotes the space of α -Hölderian functions on [0,T] with values in X such that $[f]_{\alpha} = \sup_{0 \le x < y \le T} \frac{|f(x) f(y)|}{(y-x)^{\alpha}} < \infty.$

Now we need to recall several preliminaries on semigroup and interpolation spaces. We refer the reader to [7] for the proofs and other related results.

A linear operator A in a Banach space X, with domain $D(A) \subset X$, is called sectorial if there are constants $\omega \in \mathbb{R}$, $\theta \in (\pi/2, \pi)$, M > 0 such that

$$\begin{cases} (i) \quad \rho(A) \supseteq S_{\theta,\omega} = \{\lambda \in \mathbb{C} : \lambda \neq \omega, |arg(\lambda - \omega)| < \theta\}, \\ (ii) \quad ||(\lambda I - A)^{-1}||_{\mathcal{L}(X)} \le \frac{M}{|\lambda - \omega|} \quad \forall \lambda \in S_{\theta,\omega} \end{cases}$$
(3)

where $\rho(A)$ is the resolvent set of A. For every t > 0, (3) allows us to define a linear bounded operator e^{tA} in X, by means of the Dunford integral

$$e^{tA} = \frac{1}{2\pi i} \int_{\omega + \gamma_{r,\eta}} e^{t\lambda} (\lambda I - A)^{-1} d\lambda, \quad t > 0, \tag{4}$$

where, $r > 0, \eta \in (\pi/2, \pi)$ and $\gamma_{r,\eta}$ is the curve $\{\lambda \in \mathbb{C} : |arg\lambda| = \eta, |\lambda| \ge r\} \cup \{\lambda \in \mathbb{C} : |arg\lambda| \le \eta, |\lambda| = r\}$, oriented counterclockwise. We also set

 $e^{0A}x = x, \forall x \in X$. Since the function $\lambda \mapsto e^{t\lambda}R(\lambda, A)$ is holomorphic in $S_{\theta,\omega}$, the definition of e^{tA} is independent of the choice of r and η . If A is sectorial, the function $[0, +\infty) \to L(X), t \mapsto e^{tA}$, with e^{tA} defined by (4) is called *analytic semigroup generated by* A in X. We note that for every $x \in X$ the function $t \mapsto e^{tA}x$ is analytic (and hence continuous) for t > 0. e^{tA} is a strongly continuous semigroup if and only if D(A) is dense in X; in particular this holds if X is a reflexive space.

We need to introduce suitable classes of subspaces of X.

Definition 1. Let (α, p) be two numbers such that $0 < \alpha < 1, 1 \le p \le \infty$ or $(\alpha, p) = (1, \infty)$. Then we denote with $D_A(\alpha, p)$ the space

$$D_A(\alpha, p) = \{ x \in X : t \mapsto v(t) = ||t^{1-\alpha-1/p}Ae^{tA}x|| \in L^p(0,1) \}$$

where $||x||_{D_A(\alpha,p)} = ||x||_X + [x]_\alpha = ||x||_X + ||v||_{L^p(0,1)}.$
(We set as usual $1/\infty = 0$).

We recall here some estimates for the function $t \mapsto e^{tA}$ when $t \to 0$, which we will use in the sequel. For convenience, in the next proposition we set $D_A(0, p) = X$, $p \in [1, \infty]$.

Proposition 1. Let $(\alpha, p), (\beta, p) \in (0, 1) \times [1, +\infty] \cup \{(1, \infty)\}, \alpha \leq \beta$. Then there exists $C = C(p; \alpha, \beta)$ such that

$$||t^{-\alpha+\beta}e^{tA}||_{L(D_A(\alpha,p),D_A(\beta,p))} \le C, \quad 0 < t \le 1.$$

Definition 2. Let $0 \le \alpha \le 1$ and let D, X be Banach spaces, $D \subset X$. A Banach space Y such that $D \subset Y \subset X$ is said to belong to the class J_{α} between X and D if there is a constant C such that $||x||_Y \le C||x||_X^{1-\alpha}||x||_D^{\alpha}$, $\forall x \in D$. In this case we write $Y \in J_{\alpha}(X, D)$.

Now we give the definition of solution to the BSDE:

$$Y_t - \int_t^T e^{(s-t)A} [f_0(s, Y_s) + f_1(s, Y_s, Z_s)] ds + \int_t^T e^{(s-t)A} Z_s dW_s = e^{(T-t)A} \xi,$$
(5)

Definition 3. A pair of progressively measurable processes (Y, Z) is called mild solution of (5) if it belongs to the space $L^2(\Omega; C([0,T]; H_\alpha)) \times L^2(\Omega \times [0,T]; \mathcal{L}^2(K,H))$ and \mathbb{P} -a.s.solves the integral equation (5) on the interval [0,T].

We finally state a lemma needed in the sequel. It is a generalization of the well known Gronwall's lemma. Its proof is given in the Appendix. **Lemma 1.** Assume a, b, α, β are nonnegative constants, with $\alpha < 1, \beta > 0$ and $0 < T < \infty$. For any nonnegative process $U \in L^1(\Omega \times [0,T])$, satisfying \mathbb{P} -a.s. $U_t \leq a(T-t)^{-\alpha} + b \int_t^T (s-t)^{\beta-1} \mathbb{E}^{\mathcal{F}_t} U_s ds$ for almost every $t \in [0,T]$, it holds \mathbb{P} -a.s. $U_t \leq aM(T-t)^{-\alpha}$, for almost every $t \in [0,T]$. M is a constant depending only on b, α, β, T .

3 A simplified equation

As a preparation for the study of (2), in this section we consider the following simplified version of that equation:

$$Y_t - \int_t^T e^{(s-t)A} [f_0(s, Y_s)ds + f_1(s)]ds + \int_t^T e^{(s-t)A} Z_s dW_s = e^{(T-t)A} \xi, \quad (6)$$

for all $t \in [0, T]$.

We suppose that the following assumptions hold.

Hypothesis 2.

- **1.** $A: D(A) \subset H \to H$ is a sectorial operator. We also assume that A is dissipative, i.e. it satisfies $\langle Ay, y \rangle \leq 0, \forall y \in D(A);$
- 2. for some $0 < \alpha < 1$ there exists a Banach space H_{α} continuously embedded in H and such that
 - (i) $D_A(\alpha, 1) \subset H_\alpha \subset D_A(\alpha, \infty);$
 - (ii) the part of A in H_{α} is sectorial in H_{α} .
- **3.** the final condition ξ is an \mathcal{F}_T -measurable random variable defined on Ω with values in the closure of D(A) with respect to H_{α} -norm. We denote this set $\overline{D(A)}^{H_{\alpha}}$. Moreover ξ belongs to $L^{\infty}(\Omega, \mathcal{F}_T, \mathbb{P}; H_{\alpha})$;
- **4.** $f_0: \Omega \times [0,T] \times H_\alpha \to H$ satisfies:
 - i) $\{f_0(t,y)\}_{t\in[0,T]}$ is progressively measurable $\forall y \in H_\alpha$;
 - ii) there exist constants S > 0, $1 < \gamma < 1/\alpha$ such that \mathbb{P} -a.s. $|f_0(t, y)|_H \le S(1 + ||y||_{H_\alpha}^{\gamma}) \quad t \in [0, T], y \in H_\alpha;$
 - iii) for every R > 0 there is $L_R > 0$ such that \mathbb{P} -a.s.

$$|f_0(t,y_1) - f_0(t,y_2)|_H \le L_R ||y_1 - y_2||_{H_\alpha}$$

for $t \in [0, T]$ and $y_i \in H_\alpha$ with $||y_i||_{H_\alpha} \leq R$;

iv) there exists a number $\mu \in \mathbb{R}$ such that \mathbb{P} -a.s., $\forall t \in [0, T], y_1, y_2 \in H_{\alpha}$,

$$|\langle f_0(t,y_1) - f_0(t,y_2), y_1 - y_2 \rangle_H \le \mu |y_1 - y_2|_H^2;$$
 (7)

5. $f_1: \Omega \times [0,T] \to H$ is progressively measurable and for some constant C > 0 it satisfies \mathbb{P} -a.s. $|f_1(t)|_H \leq C$, for $t \in [0,T]$.

Remark 1. We note that the pair (Y, Z) solves the BSDE (6) with final condition ξ and drift $f = f_0 + f_1$ if and only if the pair $(\bar{Y}, \bar{Z}) := (e^{\lambda t}Y_t, e^{\lambda t}Z_t)$ is a solution of the same equation with final condition $e^{\lambda T}\xi$ and drift $f'(t, y) := f'_0(t, y) + f'_1(t)$ where $f'_0(t, y) = e^{\lambda t}(f_0(t, e^{-\lambda t}y) - \lambda y), f'_1(t) = e^{\lambda t}f_1(t)$. If we choose $\mu = \lambda$, then f'_0 satisfies the same assumption as f_0 , but with (7) replaced by $\langle f_0(t, y_1) - f_0(t, y_2), y_1 - y_2 \rangle_H \leq 0$. If this last condition holds, then f_0 is called dissipative. Hence, without loss of generality, we shall assume until the end that f_0 is dissipative, or equivalently that $\mu = 0$ in (7).

3.1 A priori estimates

We prove a basic estimate for the solution in the norm of H.

Proposition 2. Suppose that Hypothesis 2 holds; if (Y, Z) is a mild solution of (6) on the interval [a, T], $0 \le a \le T$, then there exists a constant C_1 , which depends only on $||\xi||_{L^{\infty}(\Omega;H)}$ and on the constants S of 4.ii) and C of 5. such that \mathbb{P} -a.s. $\sup_{a \le t \le T} ||Y_t||_H \le C_1$. In particular the constant C_1 is independent of a.

Proof. Let the pair $(Y,Z) \in L^2(\Omega, C([a,T]; H_\alpha) \times L^2(\Omega \times [a,T]; \mathcal{L}^2(K; H))$ satisfy (6). Let us introduce the operators $J_n = n(nI - A)^{-1}$, n > 0. We note that the operators AJ_n are the Yosida approximations of A and they are bounded. Moreover $|J_n x - x| \to 0$ as $n \to \infty$, for every $x \in H$. We set $Y_t^n = J_n Y_t$, $Z_t^n = J_n Z_t$. It is readily verified that Y^n admits the Itô differential

$$dY_t^n = -AY_t^n dt - J_n f(t, Y_t) dt - J_n f_1(t) dt + Z_t^n dW_t, \text{and} \quad Y_T^n = J_n \xi.$$

Applying the Ito formula to $|Y_t^n|_H^2$, using the dissipativity of A, we obtain

$$|Y_t^n|_H^2 + \int_t^T ||Z_s^n||_{\mathcal{L}^2(K;H)}^2 ds \le |J_n\xi|_H^2 + 2\int_t^T \langle J_n f_0(s, Y_s), Y_s^n \rangle_H ds + 2\int_t^T \langle J_n f_1(s), Y_s^n \rangle_H ds - 2\int_t^T \langle Y_s^n, Z_s^n dW_s \rangle_H.$$
(8)

We note that $\int_t^T \langle J_n f_0(s, Y_s) + J_n f_1(s), Y_s^n \rangle_H ds \to \int_t^T \langle f_0(s, Y_s) + f_1(s), Y_s \rangle_H ds$ by dominated convergence, as $n \to \infty$. Moreover by the dominated convergence theorem we have $\int_t^T ||(Z_s^n)^* Y_s^n - Z_s^* Y_s||_K^2 ds \to 0$ \mathbb{P} -a.s. and it follows that $\int_t^T \langle Y_s^n, Z_s^n dW_s \rangle_H \to \int_t^T \langle Y_s, Z_s dW_s \rangle_H$ in probability. If we let $n \to \infty$ in (8) we obtain

$$\begin{aligned} |Y_t|_H^2 + \int_t^T ||Z_s||_{\mathcal{L}^2(K;H)}^2 ds &\leq |\xi|_H^2 + 2\int_t^T \langle f_0(s,Y_s) + f_1(s), Y_s \rangle_H \, ds \\ &- 2\int_t^T \langle Y_s, Z_s dW_s \rangle_H \, . \end{aligned}$$

Recalling (7), that we assume to hold with $\mu = 0$, it follows that

$$\begin{split} Y_t|_H^2 + \int_t^T ||Z_s||_{\mathcal{L}^2(K,H)}^2 &\leq \\ &\leq |\xi|_H^2 + 2\int_t^T < f_0(s,0), Y_s >_H + 2\int_t^T < f_1(s), Y_s >_H ds + \\ &- 2\int_t^T < Y_s, Z_s dW_s >_H \\ &\leq |\xi|_H^2 + \int_t^T |f(s,0)|_H^2 ds + \int_t^T |f_1(s)|_H^2 ds + 2\int_t^T |Y_s|_H^2 ds + \\ &- 2\int_t^T < Y_s, Z_s dW_s >_H. \end{split}$$

Now, since $\sup_{0 \le t \le T} |f(t,0)|_H^2 \le S^2$ and since the stochastic integral $\int_a^t \langle Y_s, Z_s dW_s \rangle_H, t \in [a,T]$ is a martingale, if we take the conditional expectation given \mathcal{F}_t we have

$$\begin{aligned} |Y_t|_H^2 &\leq \mathbb{E}^{\mathcal{F}_t} |\xi|_H^2 + 2\mathbb{E}^{\mathcal{F}_t} \int_t^T |Y_s|_H^2 ds + \\ &+ \mathbb{E}^{\mathcal{F}_t} \int_t^T |f(s,0)|_H^2 ds + \mathbb{E}^{\mathcal{F}_t} \int_t^T |f_1(s)|_H^2 ds \\ &\leq |\xi|_{L^{\infty}(\Omega,H)}^2 + (S^2 + C^2)T + 2\int_t^T \mathbb{E}^{\mathcal{F}_t} |Y_s|_H^2 ds \end{aligned}$$

Since Y belongs to $L^2(\Omega; C([a, T]; H_\alpha))$ and, consequently, $||Y||_{H_\alpha}^2 \in L^1(\Omega \times [0, T])$, we can apply Lemma 1 to $|Y|_H^2$ and conclude that $|Y_t|_H^2 \leq (|\xi|_{L^\infty(\Omega, H)}^2 + [S^2 + C^2]T)(1 + 2Te^{2T}).$

Now we will show that the result of Proposition 2, together with the growth condition satisfied by f_0 , yields an a priori estimate on the solution in the H_{α} -norm.

Let $0 < \alpha < 1$ and let $\gamma > 1$ be given by 4.ii). We fix $\theta = \alpha \gamma$ and consider the Banach space $D_A(\theta, \infty)$ introduced in Definition 1. It is easy to check (see [7]) that, if we take $\theta \in (0, 1), \theta > \alpha$, then H_α contains $D_A(\theta, \infty)$ and belongs to the class $J_{\alpha/\theta}$ between $D_A(\theta, \infty)$ and H, hence the following inequality is satisfied:

$$|x|_{H_{\alpha}} \le c|x|_{D_{A}(\theta,\infty)}^{\frac{\alpha}{\theta}}|x|_{H}^{1-\frac{\alpha}{\theta}}, \quad x \in D_{A}(\theta,\infty).$$
(9)

Proposition 3. Suppose that Hypothesis 2 is satisfied. Let (Y, Z) be a mild solution of (6) in [a, T], $a \ge 0$ and assume that there exists two constants R > 0 and K > 0, possibly depending on a, such that, \mathbb{P} -a.s.,

$$\sup_{t \in [a,T]} ||Y_t||_{H_{\alpha}} \le R, \quad \sup_{t \in [a,T]} |Y_t|_H \le K.$$
(10)

Then the following inequality holds \mathbb{P} -a.s.:

$$|Y_t|_{L^{\infty}(\Omega, D_A(\theta, \infty))} \le C_2 \frac{1}{(T-t)^{\theta-\alpha}}, \quad a \le t < T$$
(11)

with C_2 depending on the operator A, $||\xi||_{L^{\infty}(\Omega, H_{\alpha})}$, θ , α , K, C of 5. and S of 4.ii) of Hypothesis 2.

Proof. Taking the conditional expectation given \mathcal{F}_t in equation (6) we find $Y_t = \mathbb{E}^{\mathcal{F}_t} \left(e^{(T-t)A} \xi + \int_t^T e^{(s-t)A} [f_0(s, Y_s) + f_1(s)] ds \right), \quad a \le t \le T.$

Consequently, we have

$$||Y_t||_{D_A(\theta,\infty)} \leq \mathbb{E}^{\mathcal{F}_t} ||e^{(T-t)A}\xi||_{D_A(\theta,\infty)} + \mathbb{E}^{\mathcal{F}_t} \int_t^T ||e^{(s-t)A}[f_0(s,Y_s) + f_1(s)]||_{D_A(\theta,\infty)} ds, \quad a \leq t \leq T.$$
(12)

Since $H_{\alpha} \subset D_A(\alpha, \infty)$, we have

$$\mathbb{E}^{\mathcal{F}_{t}}||e^{(T-t)A}\xi||_{D_{A}(\theta,\infty)} \leq \\
\leq \mathbb{E}^{\mathcal{F}_{t}}||e^{(T-t)A}||_{L(D_{A}(\alpha,\infty),D_{A}(\theta,\infty))}||\xi||_{L^{\infty}(\Omega,D_{A}(\alpha,\infty))} \qquad (13) \\
\leq \frac{C_{0}}{(T-t)^{\theta-\alpha}}||\xi||_{L^{\infty}(\Omega,H_{\alpha})},$$

with $C_0 = C_0(\alpha, \theta, \infty)$, where in the last inequality we use Proposition 1.

Moreover

$$\begin{split} & \mathbb{E}^{\mathcal{F}_{t}} \int_{t}^{T} ||e^{(s-t)A}[f_{0}(s,Y_{s}) + f_{1}(s)]||_{D_{A}(\theta,\infty)} ds \leq \\ & \leq \mathbb{E}^{\mathcal{F}_{t}} \int_{t}^{T} ||e^{(s-t)A}||_{L(H,D_{A}(\theta,\infty))} |f_{0}(s,Y_{s}) + f_{1}(s)|_{H} ds \leq \\ & \leq \mathbb{E}^{\mathcal{F}_{t}} \Big(\int_{t}^{T} \frac{C_{1}}{(s-t)^{\theta}} [|f_{0}(s,Y_{s})|_{H} + |f_{1}(s)|_{H}] ds \Big) \leq \\ & \leq \mathbb{E}^{\mathcal{F}_{t}} \int_{t}^{T} \frac{C_{1}}{(s-t)^{\theta}} [S(1+||Y_{s}||_{H_{\alpha}}^{\gamma}) + C] ds. \end{split}$$

In the inequality we used Hypotheses 4.ii) and 5. and Proposition 1. Recalling (9), we conclude that the last term is dominated by

$$\mathbb{E}^{\mathcal{F}_t} \int_t^T \frac{C_1}{(s-t)^{\theta}} \left[S(1+c|Y_s|_H^{\gamma(1-\alpha)/\theta} ||Y_s||_{D_A(\theta,\infty)}^{\gamma\alpha/\theta}) + C \right] ds = \\ = \mathbb{E}^{\mathcal{F}_t} \int_t^T \frac{C_1}{(s-t)^{\theta}} \left[S(1+c|Y_s|_H^{\gamma(1-\alpha)/\theta} ||Y_s||_{D_A(\theta,\infty)}) + C \right] ds,$$

by choosing $\theta = \alpha \gamma$. By the second inequality in (10) this can be estimated by

$$\int_{t}^{T} \frac{C_{1}}{(s-t)^{\theta}} S(1+cK^{\gamma(1-\alpha)/\theta} \mathbb{E}^{\mathcal{F}_{t}} ||Y_{s}||_{D_{A}(\theta,\infty)} + C) ds$$

$$\leq \int_{t}^{T} \frac{C_{1}}{(s-t)^{\theta}} (C+S) ds + \int_{t}^{T} \frac{C_{1}}{(s-t)^{\theta}} ScK^{\gamma(1-\alpha)/\theta} \mathbb{E}^{\mathcal{F}_{t}} ||Y_{s}||_{D_{A}(\theta,\infty)} ds.$$
(14)

Hence by (13) and (14) it follows

$$\begin{split} ||Y_t||_{D_A(\theta,\infty)} &\leq \frac{C_0}{(T-t)^{\theta-\alpha}} ||\xi||_{L^{\infty}(\Omega,H_{\alpha})} + \int_t^T \frac{C_1}{(s-t)^{\theta}} (C+S) ds \\ &+ \int_t^T \frac{C_1}{(s-t)^{\theta}} Sc K^{\gamma(1-\alpha)/\theta} \mathbb{E}^{\mathcal{F}_t} ||Y_s||_{D_A(\theta,\infty)} ds, \end{split}$$

and (11) follows from Lemma 1. In order to justify the application of Lemma 1, we need to prove that $||Y||_{D_A(\theta,\infty)}$ belongs to $L^1(\Omega \times [a,T])$. This also follows from(13) and (14) since, for some constant K_1 ,

$$\begin{split} ||Y_t||_{D_A(\theta,\infty)} &\leq \\ &\leq \frac{K_1}{(T-t)^{\theta-\alpha}} ||\xi||_{L^{\infty}(\Omega,H_{\alpha})} + \mathbb{E}^{\mathcal{F}_t}[\sup_{s\in[a,T]} (1+||Y_s||_{H_{\alpha}}^{\gamma}) \int_t^T \frac{ds}{(s-t)^{\theta}}] \\ &\leq \frac{K_1}{(T-t)^{\theta-\alpha}} ||\xi||_{L^{\infty}(\Omega,H_{\alpha})} + (1+R^{\gamma}) \int_t^T \frac{ds}{(s-t)^{\theta}}. \end{split}$$

3.2 Local existence and uniqueness

We prove that, under Hypothesis 2, there exists a unique solution of (6) on an interval $[T - \delta, T]$ with δ sufficiently small.

To treat the ordinary integral in the left hand side of (6), we need the following result, whose proof can be found in [7], Proposition 4.2.1 and Lemma 7.1.1.

Lemma 3. Let $\phi \in L^{\infty}((a,T);H)$, 0 < a < T and set

$$v(t) = \int_t^T e^{(s-t)A}\phi(s)ds, \quad a \le t \le T.$$

If $0 < \alpha < 1$, then $v \in C^{1-\alpha}([a,T]; D_A(\alpha,1))$ and there is $G_0 > 0$, not depending on a, such that

$$||v||_{C^{1-\alpha}([a,T];D_A(\alpha,1))} \le G_0||\phi||_{L^{\infty}((a,T);H)}.$$

Since $D_A(\alpha, 1) \subset H_\alpha$, we also have $v \in C^{1-\alpha}([a, T]; H_\alpha)$ and there is G > 0, not depending on a, such that

$$||v||_{C^{1-\alpha}([a,T];H_{\alpha})} \le G||\phi||_{L^{\infty}((a,T);H)}$$

Theorem 4. Let us assume that Hypothesis 2 holds, except possibly 4.iv). Then there exists $\delta > 0$ such that the equation (6) has a unique local mild solution $(Y, Z) \in L^2(\Omega; C([T - \delta, T]; H_\alpha)) \times L^2(\Omega \times [T - \delta, T]; \mathcal{L}^2(K; H)).$

Remark 2. The dissipativity condition 4.iv) only plays a role in obtaining the a priori estimate in H (Proposition 2) and consequently global existence, as we will see later.

Proof. Let $M_{\alpha} := \sup_{0 \le t \le T} ||e^{tA}||_{L(H_{\alpha})}$. We fix a positive number R such that $R \ge 2M_{\alpha}||\xi||_{L^{\infty}(\Omega;H_{\alpha})}$. This implies that $\sup_{0 \le t \le T} ||e^{tA}\xi||_{H_{\alpha}} \le R/2$ \mathbb{P} -a.s. Moreover, let L_R be such that

$$|f_0(t,y_1) - f_0(t,y_2)|_H \le L_R ||y_1 - y_2||_{H_\alpha} \quad 0 \le t \le T, \quad ||y_i||_{H_\alpha} \le R$$

We recall that the space $L^2(\Omega; C([T - \delta, T]; H_\alpha))$ is a Banach space endowed with the norm $Y \to (\mathbb{E} \sup_{t \in [T - \delta, T]} ||Y_t||_{H_\alpha}^2)^{1/2}$. We define

$$\mathbb{K} = \{ Y \in L^2(\Omega; C([T-\delta,T],H_\alpha)) : \sup_{t \in [T-\delta,T]} ||Y_t||_{H_\alpha} \le R, \quad a.s. \}.$$

It easy to check that \mathbb{K} is a closed subset of $L^2(\Omega; C([T - \delta, T], H_\alpha))$, hence a complete metric space (with the inherited metrics). We look for a local mild solution (Y, Z) in the space \mathbb{K} . We define a nonlinear operator $\Gamma: \mathbb{K} \to \mathbb{K}$ as follows: given $U \in \mathbb{K}, Y = \Gamma(U)$ is the first component of the mild solution (Y, Z) of the equation

$$Y_t - \int_t^T e^{(s-t)A} [f_0(s, U_s)ds + f_1(s)]ds + \int_t^T e^{(s-t)A} Z_s dW_s = e^{(T-t)A} \xi$$
(15)

for $t \in [T - \delta, T]$. Since $U \in \mathbb{K}$ we have \mathbb{P} -a.s.

$$|f_0(t, U_t) + f_1(t)|_H \le S(1 + ||U_t||_{H_\alpha}^{\gamma}) + C \le S(1 + R^{\gamma}) + C,$$
(16)

for all t in $[T - \delta, T]$. Hence $f_0(\cdot, U_{\cdot}) + f_1(\cdot)$ belongs to $L^2(\Omega \times [T - \delta, T]; H)$ and, by a result of Hu and Peng [6], there exists a unique pair $(Y, Z) \in$ $L^2(\Omega \times [T - \delta, T]; H) \times L^2(\Omega \times [T - \delta, T]; \mathcal{L}^2(K; H))$ satisfying (15). Moreover, by taking the conditional expectation given \mathcal{F}_t , Y has the following representation

$$Y_t = \mathbb{E}^{\mathcal{F}_t} \left(e^{(T-t)A} \xi + \int_t^T e^{(s-t)A} [f_0(s, U_s) + f_1(s)] ds \right).$$

We will show that Γ is a contraction for the norm of $L^2(\Omega, C([T - \delta, T]; H_\alpha))$ and maps \mathbb{K} into itself, if δ is sufficiently small; clearly, its unique fixed point is the required solution of the BSDE.

We first check the contraction property. Let $U^1, U^2 \in \mathbb{K}$. Then

$$\Gamma(U^1)_t - \Gamma(U^2)_t = Y_t^1 - Y_t^2 = \mathbb{E}^{\mathcal{F}_t} \Big[\int_t^T e^{(s-t)A} (f_0(s, U_s^1) - f_0(s, U_s^2)) ds \Big].$$

Let $v(t) = \int_t^T e^{(s-t)A} (f_0(s, U_s^1) - f_0(s, U_s^2)) ds$. Then, noting that v(T) = 0 and recalling Lemma 3, for $t \in [T - \delta, T]$

$$\begin{split} ||Y_t^1 - Y_t^2||_{H_{\alpha}} &= \\ &= ||\mathbb{E}^{\mathcal{F}_t} v(t)||_{H_{\alpha}} &\leq \mathbb{E}^{\mathcal{F}_t} ||v(t)||_{H_{\alpha}} \\ &\leq \delta^{1-\alpha} \mathbb{E}^{\mathcal{F}_t} ||v||_{C^{(1-\alpha)}([T-\delta,T],H_{\alpha})} \\ &\leq G\delta^{(1-\alpha)} \mathbb{E}^{\mathcal{F}_t} ||f_0(\cdot, U_{\cdot}^1) - f_0(\cdot, U_{\cdot}^2)||_{L^{\infty}([T-\delta,T],H)} \\ &\leq G\delta^{(1-\alpha)} L_R \mathbb{E}^{\mathcal{F}_t} \sup_{t \in [T-\delta,T]} ||U_t^1 - U_t^2||_{H_{\alpha}} =: M_t, \end{split}$$

where $\{M_t, t \in [T - \delta, T]\}$ is a martingale. Hence, by Doob's inequality

$$\mathbb{E} \sup_{t \in [T-\delta,T]} ||Y_t^1 - Y_t^2||_{H_{\alpha}}^2 \leq \mathbb{E} \sup_{t \in [T-\delta,T]} |M_t|^2 \leq 2\mathbb{E}|M_T|^2 = = 2G^2 L_R^2 \delta^{2(1-\alpha)} \mathbb{E} \sup_{t \in [T-\delta,T]} ||U_t^1 - U_t^2||_{H_{\alpha}}^2$$

If $\delta \leq \delta_0 = 2GL_R^{\frac{-1}{(1-\alpha)}}$, then Γ is a contraction with constant 1/2.

Next we check that Γ maps \mathbb{K} into itself. For each $U \in \mathbb{K}$ and $t \in [T-\delta, T]$ with $\delta \leq \delta_0$ we have

$$\sup_{t \in [T-\delta,T]} ||\Gamma(U)_t||_{H_{\alpha}} = \sup_{t \in [T-\delta,T]} ||Y_t||_{H_{\alpha}} \leq \sup_{t \in [T-\delta,T]} \mathbb{E}^{\mathcal{F}_t} ||e^{(T-t)A}\xi||_{H_{\alpha}} + \sup_{t \in [T-\delta,T]} \mathbb{E}^{\mathcal{F}_t} ||\int_t^T e^{(s-t)A} [f_0(s,U_s) + f_1(s)] ds||_{H_{\alpha}} \\ \leq R/2 + \sup_{t \in [T-\delta,T]} \mathbb{E}^{\mathcal{F}_t} \int_t^T ||e^{(s-t)A} [f_0(s,U_s) + f_1(s)]||_{H_{\alpha}} ds \\ \leq R/2 + \sup_{t \in [T-\delta,T]} \mathbb{E}^{\mathcal{F}_t} \int_t^T ||e^{(s-t)A} [f_0(s,U_s) + f_1(s)]||_{D_A(\alpha,1)} ds,$$

where in the last inequality we have used the fact that $D_A(\alpha, 1) \subset H_{\alpha}$. Now, by Proposition 1, and from 4.ii) and 5., it follows that

$$\begin{aligned} ||e^{(s-t)A}[f_0(s,U_s) + f_1(s)]||_{D_A(\alpha,1)} &\leq \\ &\leq ||e^{(s-t)A}||_{L(H,D_A(\alpha,1))}|f_0(s,U_s) + f_1(s)|_H \\ &\leq \frac{C_\alpha}{(s-t)^\alpha}[S(1+||U_s||_{H_\alpha}^{\gamma}) + C]. \end{aligned}$$

Then, since $U \in \mathbb{K}$, we arrive at

$$\sup_{t\in[T-\delta,T]} ||\Gamma(U)_t||_{H_{\alpha}} \leq \\ \leq R/2 + \sup_{t\in[T-\delta,T]} \mathbb{E}^{\mathcal{F}_t} \int_t^T \frac{C_{\alpha}}{(s-t)^{\alpha}} [S(1+||U_s||_{H_{\alpha}}^{\gamma}) + C] ds \\ \leq R/2 + \sup_{t\in[T-\delta,T]} \int_t^T \frac{C_{\alpha}}{(s-t)^{\alpha}} [S(1+R^{\gamma}) + C] ds \\ \leq R/2 + C_{\alpha} S \frac{[(1+R^{\gamma})+C]}{1-\alpha} \delta^{1-\alpha},$$

where C_{α} depends on A, α . Hence, if $\delta \leq \delta_0$ is such that $C_{\alpha}S\frac{[(1+R^{\gamma})+C]}{1-\alpha}\delta^{1-\alpha}$ is less or equal to R/2, then $\sup_{t\in[T-\delta,T]} ||\Gamma(U)_t||_{H_{\alpha}} \leq R$. Due to Lemma 3, \mathbb{P} a.s. the function $t \mapsto Y_t - \mathbb{E}^{\mathcal{F}_t} e^{(T-t)A}\xi$ belongs to $C[T-\delta,T]; H_{\alpha}$; moreover, the map $t \mapsto \mathbb{E}^{\mathcal{F}_t} e^{(T-t)A}\xi$ belongs to $C[T-\delta,T]; H_{\alpha}$, since ξ is a random variable taking values in $\overline{D(A)}^{H_{\alpha}}$. Therefore, \mathbb{P} -a.s. $Y_{\cdot} \in C([T-\delta,T]; H_{\alpha})$ and Γ maps \mathbb{K} into itself and has a unique fixed point in \mathbb{K} .

Remark 3. By Lemma 3, using properties of analytic semigroups, it can be proved that for every fixed ω the range of the map Γ is contained in $C^{1-\beta}([T-\delta, T-\epsilon]; D_A(\beta, 1))$ for every $\epsilon \in (0, \delta), \beta \in [0, 1].$

3.3 Global existence

Now we are able to prove a global existence theorem for the solution of the equation (6), using all the results presented above.

Theorem 5. If Hypothesis 2 is satisfied, the equation (6) has a unique mild solution $(Y, Z) \in L^2(\Omega; C([0, T], H_\alpha)) \times L^2(\Omega \times [0, T]); \mathcal{L}^2(K; H)).$

Proof. By Theorem 4 equation (6) has a unique mild solution $(Y^1, Z^1) \in L^2(\Omega; C([T - \delta_1, T], H_\alpha)) \times L^2(\Omega \times [T - \delta_1, T]); \mathcal{L}^2(K; H))$ on the interval $[T - \delta_1, T]$, for some $\delta_1 > 0$. By Proposition 2 we know that there exists a constant C_1 such that \mathbb{P} -a.s.

$$|Y_{T-\delta_1}|_H \le C_1. \tag{17}$$

We recall that the constant C_1 depends only on $|\xi|_{L^{\infty}(\Omega;H)}$ and on the constants S of 4.ii) and C of 5. and is independent of δ_1 . Moreover, by Proposition 3, there exists a constant C_2 such that \mathbb{P} -a.s.

$$||Y_{T-\delta_1}||_{L^{\infty}(\Omega, D_A(\theta, \infty))} \le C_2 \frac{1}{\delta_1^{\theta-\alpha}},\tag{18}$$

with C_2 depending on the operator A, $||\xi||_{L^{\infty}(\Omega,H_{\alpha})}$, θ , α , C_1 . This implies that $Y_{T-\delta_1}$ belongs to $L^{\infty}(\Omega;H_{\alpha})$ and it can be taken as final value for the problem

$$Y_t - \int_t^{T-\delta_1} e^{(s-t)A} [f_0(s, Y_s)ds + f_1(s)]ds + \int_t^{T-\delta_1} e^{(s-t)A} Z_s dW_s = e^{(T-\delta_1-t)A} Y_{T-\delta_1}$$
(19)

on an interval $[T - \delta_1 - \delta_2, T - \delta_1]$, for some $\delta_2 > 0$. As in the proof of Theorem 4, we fix a positive number R_2 such that

$$R_2 = 2M_{\alpha} \frac{C_2}{\delta_1^{\theta - \alpha}} \ge 2M_{\alpha} ||Y_{T - \delta_1}||_{L^{\infty}(\Omega, D_A(\theta, \infty))}.$$

By Theorem 4 there exists a pair of progressively measurable processes (Y^2, Z^2) in $L^2(\Omega; C([T - \delta_1 - \delta_2, T - \delta_1]; H_\alpha)) \times L^2(\Omega \times [T - \delta_1 - \delta_2, T - \delta_1]; \mathcal{L}^2(K, H))$ which solves (19) on the interval $[T - \delta_1 - \delta_2, T - \delta_1]$ where δ_2 depends on the operator A, α, R_2 . We note that the continuity in $T - \delta_1$ of Y^2 follows from the fact that $Y_{T-\delta_1}$ takes values in $D_A(\alpha, 1)$ (see Remark 3), so that $Y_{T-\delta_1}$ takes values in $\overline{D(A)}^{H_\alpha}$. Now, the process Y_t defined by Y_t^1 on the interval $[T - \delta_1, T]$ and by Y_t^2 on $[T - \delta_1 - \delta_2, T - \delta_1]$ belongs

to $L^2(\Omega; C([T - \delta_1 - \delta_2, T]; H_\alpha))$ and it easy to see that it satisfies (6) in the whole interval $[T - \delta_1 - \delta_2, T]$. Consequently, by Proposition 2, \mathbb{P} -a.s., $|Y_{T-\delta_1-\delta_2}|_H \leq C_1$ with C_1 the constant in (17), and by (18)

$$||Y_{T-\delta_1-\delta_2}||_{L^{\infty}(\Omega,D_A(\theta,\infty))} \le \frac{C_2}{(\delta_1+\delta_2)^{\theta-\alpha}} \le \frac{C_2}{\delta_1^{\theta-\alpha}},\tag{20}$$

where C_2 is the same constant as in (18). Again, $Y_{T-\delta_1-\delta_2}$ can be taken as initial value for problem

$$Y_t - \int_t^{T-\delta_1 - \delta_2} e^{(s-t)A} [f_0(s, Y_s)ds + f_1(s)]ds + \int_t^{T-\delta_1 - \delta_2} e^{(s-t)A} Z_s dW_s = e^{(T-\delta_1 - \delta_2 - t)A} Y_{T-\delta_1 - \delta_2}$$
(21)

on the interval $[T - \delta_1 - \delta_2 - \delta_3, T - \delta_1 - \delta_2]$, where δ_3 will be fixed later. In this case, by (20), we can choose

$$R_3 = R_2 = 2M_\alpha \frac{C_2}{\delta_1^{\theta - \alpha}} \ge 2M_\alpha ||Y_{T - \delta_1 - \delta_2}||_{L^\infty(\Omega, D_A(\theta, \infty))}$$

and prove that there exists a unique mild solution (Y^3, Z^3) of (21) on the interval $[T - \delta_1 - \delta_2 - \delta_3, T - \delta_1 - \delta_2]$, with $\delta_3 = \delta_2$. So we extend the solution to $[T - \delta_1 - 2\delta_2, T]$. Proceeding this way we prove the global existence to (6) on [0, T].

4 The general case

We can now study the equation:

$$Y_t - \int_t^T e^{(s-t)A} [f_0(s, Y_s) + f_1(s, Y_s, Z_s)] ds + \int_t^T e^{(s-t)A} Z_s dW_s = e^{(T-t)A} \xi$$
(22)

We require that the function f_1 satisfy the following assumptions:

Hypothesis 6.

1. there exists $K \ge 0$ such that \mathbb{P} -a.s.

$$|f_1(t, y, z) - f_1(t, y', z')|_H \le K|y - y'|_H + K||z - z'||_{\mathcal{L}^2(K;H)},$$
for every $t \in [0, T], y, y' \in H, z, z' \in \mathcal{L}^2(K; H),$

2. there exists $C \ge 0$ such that \mathbb{P} -a.s. $|f_1(t, y, z)|_H \le C$, for every $t \in [0, T], y \in H, z \in \mathcal{L}^2(K; H)$.

Theorem 7. If Hypotheses 2 and 6 hold, then equation (22) has a unique solution in $L^2(\Omega; C([0,T]; H_{\alpha})) \times L^2(\Omega \times [0,T]; \mathcal{L}^2(K; H)).$

Proof. Let \mathbb{M} be the space of progressive processes (Y, Z) in the space $L^2(\Omega \times [0, T]; H) \times L^2(\Omega \times [0, T]; \mathcal{L}^2(K; H))$ endowed with the norm

$$|||(Y,Z)|||_{\beta}^{2} = \mathbb{E} \int_{0}^{T} e^{\beta s} (|Y_{s}|_{H}^{2} + ||Z_{s}||_{\mathcal{L}^{2}(K;H)}^{2}) ds,$$

where β will be fixed later. We define $\Phi : \mathbb{M} \to \mathbb{M}$ as follows: given $(U, V) \in \mathbb{M}$, $(Y, Z) = \Phi(U, V)$ is the unique solution on the interval [0, T] of the equation

$$Y_t - \int_t^T e^{(s-t)A} [f_0(s, Y_s)ds + f_1(s, U_s, V_s)]ds + \int_t^T e^{(s-t)A} Z_s dW_s = e^{(T-t)A} \xi_s dW_$$

By Theorem 5 the above equation has a unique mild solution (Y, Z) which belongs to $L^2(\Omega; C([0, T]; H_\alpha)) \times L^2(\Omega \times [0, T]; \mathcal{L}^2(K; H))$. Therefore $\Phi(\mathbb{M}) \subset \mathbb{M}$. We will show that Φ is a contraction for a suitable choice of β ; clearly, its unique fixed point is the required solution of (22). We take another pair $(U', V') \in \mathbb{M}$ and apply Proposition 3.1 in [3] to the difference of two equations. We obtain

$$\begin{split} & \mathbb{E} \int_{0}^{T} e^{\beta s} \left[\beta |Y_{t}^{1} - Y_{t}^{2}|_{H}^{2} + \|Z_{s}^{1} - Z_{s}^{2}\|_{\mathcal{L}^{2}(K;H)}^{2} ds \right] \leq \\ & \leq 2 \mathbb{E} \int_{0}^{T} e^{\beta s} < f_{0}(s, Y_{s}^{1}) + f_{1}(s, U_{s}^{1}, V_{s}^{1}) + \\ & -f_{0}(s, Y_{s}^{2}) - f_{1}(s, U_{s}^{2}, V_{s}^{2}), Y_{s}^{1} - Y_{s}^{2} >_{H} ds \\ & \leq 2 \mathbb{E} \int_{0}^{T} e^{\beta s} K(|U_{s}^{1} - U_{s}^{2}|_{H} + ||V_{s}^{1} - V_{s}^{2}||_{\mathcal{L}^{2}(K;H)})|Y_{s}^{1} - Y_{s}^{2}|_{H} ds \\ & \leq \mathbb{E} \int_{0}^{T} e^{\beta s} (|U_{s}^{1} - U_{s}^{2}|_{H}^{2} + ||V_{s}^{1} - V_{s}^{2}||_{\mathcal{L}^{2}(K;H)})/2 + 4K^{2}|Y_{s}^{1} - Y_{s}^{2}|_{H}^{2} ds, \end{split}$$

where we have used 4.iv) of Hypothesis 2 and 1. of Hypothesis 6. Choosing $\beta = 4K^2 + 1$, we obtain the required contraction property.

5 Applications

In this section we present some backward stochastic partial differential problems which can be solved with our techniques.

5.1 The reaction-diffusion equation

Let D be an open and bounded subset of \mathbb{R}^n with a smooth boundary ∂D . We choose $K = L^2(D)$. This choice implies that dW_t/dt is the so-called "space-time white noise". Moreover, since Hilbert-Schmidt operators on $L^2(D)$ are represented by square integrable kernels, the space $\mathcal{L}^2(L^2(D), L^2(D))$ can be identified with $L^2(D \times D)$. We are given a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with a filtration $(\mathcal{F}_t)_{t \in [0,T]}$ generated by W and augmented in the usual way. Let us consider a non symmetric bilinear, coercive continuous form $a : H_0^1(D) \times H_0^1(D) \to \mathbb{R}$ defined by $a(u, v) := -\int_D \sum_{i,j}^n a_{ij}(x) D_i u(x) D_j v(x) dx$, where the coefficients a_{ij} are Lipschitz continuous and there exists $\alpha > 0$ such that $\sum_{i,j=1}^n a_{ij}(x) \xi_i \xi_j \ge \alpha |\xi|^2$ for every $x \in \overline{D}, \xi \in \mathbb{R}^n$. Let A be the operator associated with the bilinear form a such that $\langle Au, v \rangle_{L^2(D)} = a(u, v), v \in H_0^1(D)$ and $u \in D(A)$. It is known that, in this case, $D(A) = H^2(D) \cap H_0^1(D)$, where $H^2(D)$ and $H_0^1(D)$ are the usual Sobolev spaces.

We consider for $t \in [0, T]$ and $x \in D$ the backward stochastic problem written formally

$$\begin{cases} \frac{\partial Y(t,x)}{\partial t} = AY(t,x) + r(Y(t,x)) + g(t,Y(t,x),Z(t,x),x) + \\ + Z(t,x)\frac{\partial W(t,x)}{\partial t} \\ 0 & \Omega \times [0,T] \times \bar{D} \\ Y(T,x) = \xi(x) & \text{on } \Omega \times \bar{D} \\ Y(t,x) = 0 & \text{on } \Omega \times [0,T] \times \partial D \\ \end{array}$$
(23)

We suppose the following.

Hypothesis 8.

- **1.** $r : \mathbb{R} \to \mathbb{R}$ is a continuous, increasing and locally Lipschitz function;
- **2.** r satisfies the following growth condition: $|r(x)| \leq S(1+|x|^{\gamma}) \quad \forall x \in \mathbb{R}$ for some $\gamma > 1$;
- **3.** g is a measurable real function defined on $[0,T] \times \mathbb{R} \times L^2(D \times D) \times D$ and there exists a constant K > 0 such that

$$|g(t, y_1, z_1, x) - g(t, y_2, z_2, x)| \le K(|y_1 - y_2| + ||z_1 - z_2||_{L^2(D \times D)})$$

for all
$$t \in [0, T]$$
, $y_1, y_2 \in \mathbb{R}$, $z_1, z_2 \in L^2(D)$, $x \in D$;

- 4. there exists a real function h in $L^2(D \times D)$ such that \mathbb{P} -a.s. $|g(t, y, z, x)| \le K_1 h(x)$ for all $t \in [0, T], y \in \mathbb{R}, z \in L^2(D), x \in D;$
- **5.** ξ belongs to $L^{\infty}(\Omega; H^2(D) \cap H^1_0(D))$.

We define the operator A by (Ay)(x) = Ay(x) with domain $D(A) = H^2(D) \cap H^1_0(D)$. We set $f_0(t, y)(x) = -r(y(t, x))$ for $t \in [0, T]$, $x \in D$ and y in a suitable subspace of H which will be determined below. For $t \in [0, T]$, $x \in D$, $y \in L^2(D)$, $z \in L^2(D \times D)$ we define f_1 as the operator $f_1(t, y, z)(x) = -g(t, y(t, x), z(t, x), x)$. Then problem (23) can be written in abstract way as

$$dY_t = -AY_t dt - f_0(t, Y_t) dt - f_1(t, Y_t, Z_t) dt + Z_t dW_t, \quad Y_T = \xi.$$

Under the conditions in Hypothesis 8, the assumptions in Hypotheses 2, 6 are satisfied. The operator A is a closed operator in $L^2(D)$ and it is the infinitesimal generator of an analytic semigroup in $L^2(D)$ satisfying $||e^{tA}||_{\mathcal{L}(H)} \leq 1$ (see [17], Chapter 3). In particular, by Lumer-Philips theorem, A is dissipative. The non linear function $f_0(t, \cdot) : L^{2\gamma}(D) \to L^2(D)$, $y \mapsto -r(y)$ is locally Lipschitz. We look for a space of class J_{α} between H and D(A) where f_0 is well defined and locally Lipschitz. It is well known (see [18]) that the fractional order Sobolev space $W^{\beta,2}(D)$ is of class $J_{\beta/2}$ between $L^2(D)$ and $H^2(D)$ for every $\beta \in (0,2)$. Hence the space H_{α} defined by $H_{\alpha} = W^{\beta,2}(D)$ if $\beta < 1$, by $W^{\beta,2}(D) \cap H_0^1(D)$ if $\beta \ge 1$ is of class $J_{\beta/2}$ between H and D(A). Moreover the restriction of A on H_{α} is a sectorial operator ([18]). By the Sobolev embedding theorem, $W^{\beta,2}$ is contained in $L^q(D)$ for all q if $\beta \geq \frac{n}{2}$, and in $L^{2n/(n-2\beta)}(D)$ if $\beta < \frac{n}{2}$. If we choose $\beta \in (0,2)$ we have $W^{\beta,2}(D) \subset L^{2\gamma}(D)$ for $n < 4\frac{\gamma}{\gamma-1}$. It is clear that f_0 is locally Lipschitz with respect to y from H_{α} into H. It is easy to verify that f_0 satisfies 4.ii) of Hypothesis 2 with $\gamma = 2n + 1$ and that it is dissipative with constant $\mu = 0$. The function f_1 is Lipschitz uniformly with respect to y and z and it is bounded. The final condition ξ takes values in $\overline{D(A)}^{H_{\alpha}}$ and belongs to $L^{\infty}(\Omega; H_{\alpha})$. Hence we can apply the global existence theorem and state that the above problem has a unique mild solution $(Y,Z) \in L^2(\Omega; C([0,T]; H_\alpha)) \times L^2(\Omega \times [0,T]; \mathcal{L}^2(K,H)).$

5.2 A spin system

Let \mathbb{Z} be the one-dimensional lattice of integers. Its elements will be interpreted as atoms. A *configuration* is a real function y defined on \mathbb{Z} . The value y(n) of the configuration y at the point n can be viewed as the state of the atom n.

We consider an infinite system of equations

$$dY_t^n = -a_n Y_t^n dt + \sum_{|n-j| \le 1} V(Y_t^n - Y_t^j) dt + Z_t^n dW_t^n \quad n \in \mathbb{Z}, \quad 0 \le t \le T$$
(24)

 $Y_n(T) = \xi_n \quad n \in \mathbb{Z},$

where Y^n and Z^n are real processes, and $V : \mathbb{R} \to \mathbb{R}$.

Let $l^2(\mathbb{Z})$ be the usual Hilbert space of square summable sequences. To study system (24) we apply results of previous sections. To fit our assumption in Hypotheses 2 and 6, we suppose the following

Hypothesis 9.

- **1.** W^n , $n \in \mathbb{Z}$ are independent standard real Wiener processes;
- **2.** $a = \{a_n\}_{n \in \mathbb{Z}}$ is a sequence of nonnegative real numbers;
- **3.** $\xi = \{\xi_n\}_{n \in \mathbb{Z}}$ is a random variable belonging to $L^{\infty}(\Omega, l^2(\mathbb{Z}));$
- **4.** the function $V : \mathbb{R} \to \mathbb{R}$ is defined by $V(x) = x^{2k+1}$ $k \in \mathbb{N}$.

We will study system (24) regarded as a backward stochastic evolution equation for $t \in [0, T]$

$$dY_t = (AY_t + f_0(t, Y_t))dt + Z_t dW_t, \quad Y_T = \xi$$
(25)

on a properly chosen Hilbert space H of functions on \mathbb{Z} .

To reformulate problem (24) in the abstract form (25), we set $K = H = l^2(\mathbb{Z})$. We set $W_t = \{W_t^n\}_{n \in \mathbb{Z}}, t \in [0, T]$. By 1. of Hypothesis 9, W is a cylindrical Wiener process in H defined on (Ω, \mathcal{F}, P) . We define the operator A by

$$A(y) = (a_n y_n)_n, \quad D(A) = \{ y \in l^2(\mathbb{Z}) \text{ such that } \sum_{n \in \mathbb{Z}} a_n^2 y_n^2 < \infty \}.$$

It is easy to prove that A is a self-adjoint operator in $l^2(\mathbb{Z})$, hence the infinitesimal generator of a sectorial semigroup. The coefficient f_0 is given by $(f_0(t,y))_n = (V(y_{n+1}-y_n)+V(y_{n-1}-y_n)), \quad t \in [0,T], y \in D(f_0)$ where $D(f_0) = \{y \in l^2(\mathbb{Z}) \text{ such that } \sum_{n \in \mathbb{Z}} |x_{n+1} - x_n|^{2(2k+1)} < +\infty\}.$ Under Hypothesis 9, A, f_0, ξ satisfy Hypotheses 2 and 6. We observe that in this case the domain of f_0 is the whole space H: if $y \in l^2(\mathbb{Z})$ then

$$\left\{\sum_{n\in\mathbb{Z}}|y_{n+1}-y_n|^{2(2k+1)}\right\}^{\frac{1}{2(2k+1)}} \leq \left\{\sum_{n\in\mathbb{Z}}|y_{n+1}-y_n|^2\right\}^{\frac{1}{2}} \leq 2||y||_{l^2(\mathbb{Z})}.$$

Consequently, we can take H_{α} with $\alpha = 0$, i.e. $H_0 = H$. The function f_0 is dissipative. Namely

$$< f_0(t, y) - f_0(t, y'), y - y' >_{l^2(\mathbb{Z})} = = \sum_{n \in \mathbb{Z}} \{ [(y_{n+1} - y_n)^{(2k+1)} + (y_{n-1} - y_n)^{(2k+1)}] + + [(y'_{n+1} - y'_n)^{(2k+1)} + (y'_{n-1} - y'_n)^{(2k+1)}])[y_n - y'_n] = = -\sum_{n \in \mathbb{Z}} [(y_{n+1} - y_n)^{(2k+1)} - (y'_{n+1} - y'_n)^{(2k+1)}][(y_{n+1} - y_n) - (y'_{n+1} - y'_n)]$$

and the last term is negative. Moreover, f_0 satisfies 4.ii) of Hypothesis 2 with $\gamma = 2k + 1$. The map f_0 is also locally Lipschitz from H in to H. Then by Theorem 7, problem (25) has a unique mild solution (Y, Z) which belongs to $L^2(\Omega, C([0, T]; H)) \times L^2(\Omega \times [0, T]; \mathcal{L}^2(K, H)).$

6 Appendix

This section is devoted to the proof of Lemma 1. Assume first that $\beta = 1$. Using recursively the inequality $U_t \leq a(T-t)^{-\alpha} + b \int_t^T \mathbb{E}^{\mathcal{F}_t} U_s ds$ we can easily prove that

$$U_{t} \leq \leq a(T-t)^{-\alpha} + \int_{t}^{T} a \sum_{k=1}^{n-1} b^{k} \frac{(r-t)^{k-1}}{(k-1)!} \frac{1}{(T-r)^{\alpha}} dr + b \mathbb{E}^{\mathcal{F}_{t}} \int_{t}^{T} \frac{(b(r-t))^{n-1}}{(n-1)!} U_{r} dr.$$

The last term in the above inequality tends to zero as n tends to infinity for each t in the interval [0, T]. Thus

$$U_t \leq a(T-t)^{-\alpha} + a \sum_{k=1}^{\infty} \int_t^T \frac{b^k (T-t)^{k-1}}{(k-1)!} \frac{1}{(T-r)^{\alpha}} dr \leq \leq a(T-t)^{-\alpha} + abe^{b(T-t)} \int_t^T \frac{1}{(T-r)^{\alpha}} dr \leq \leq a(T-t)^{-\alpha} + abe^{b(T-t)} \frac{1}{1-\alpha} (T-t)^{1-\alpha} \leq a(T-t)^{-\alpha} M$$

where $M = 1 + be^{bT} \frac{1}{1-\alpha}T$.

In the case $\beta \neq 1$ a similar proof can be given, based on recursive use of the inequality $U_t \leq a(T-t)^{-\alpha} + b \int_t^T (s-t)^{\beta-1} \mathbb{E}^{\mathcal{F}_t} U_s ds.$

Acknowledgments:

I wish to thank Giuseppe Da Prato for hospitality at the Scuola Normale Superiore in Pisa, suggestions and helpful discussions. I would like to express my gratitude to Marco Fuhrman: I am indebted to him for his precious help and encouragement. Special thanks are due to Alessandra Lunardi, who gave me valuable advice and support.

References

- Ph. Briand and R. Carmona. BSDEs with polynomial growth generators. J. Appl. Math. Stochastic Anal., 13(3):207–238, 2000.
- [2] Ph. Briand and B. Deylon and Y. Hu and E. Pardoux and L. Stoica. L^p solutions of backward stochastic differential equations. Stochastic Process. Appl., 108(1):109–129, 2003.
- [3] F. Confortola. Dissipative backward stochastic differential equations in infinite dimensions. *Infinite Dimensional Analysis, Quantum Probability* and Related Topics, 9 (1):155–168, 2006.
- [4] N. El Karoui and S. G. Peng and M. C. Quenez. Backward Stochastic Differential equations in Finance. *Math. Finance*, 7(1):1–71, 1997.
- [5] H. Fujita and T. Kato. On the Navier-Stokes initial value problem I. Arch. Rational Mech. Anal. 16:269–315, 1964.
- [6] Y. Hu and S. G. Peng. Adapted solution of a backward semilinear stochastic evolution equation. *Stochastic Anal. Appl.*, 9(4):445–459, 1991.
- [7] A. Lunardi. Analytic semigroups and optimal regularity in parabolic problems volume 16 of Progress in Nonlinear Differential Equations and their Applications. Birkhser Verlag, Basel 1995.
- [8] J. Ma and J. Yong Adapted solution of a degenerate backward SPDE, with applications. *Stochastic Process. Appl.* 70:59–84, 1997.
- [9] J. Ma and J. Yong On linear, degenerate backward stochastic partial differential equations. *Probab. theory Related Fields* 113:135–170 1999.
- [10] B. Oksendal and T. Zhang. On backward stochastic partial differential equations, 2001. Preprint.

- [11] E. Pardoux. BSDEs, weak convergence and homogenization of semilinear PDEs. Nonlinear analysis, differential equations and control (Montreal, QC, 1998), 503–549, NATO Sci. Ser. C Math. Phys. Sci., 528, Kluwer Acad. Publ., Dordrecht, 1999.
- [12] É. Pardoux and S. Peng. Adapted solution of a backward stochastic differential equation. Systems and Control Lett. 14:55–61, 1990.
- [13] E. Pardoux and A. Răşcanu. Backward stochastic differential equations with subdifferential operator and related variational inequalities. *Stochastic Process. Appl.*, 76(2):191–215, 1998.
- [14] E. Pardoux and A. Răşcanu. Backward stochastic variational inequalities. Stochastics Stochastics Rep., 67(3-4):159–167, 1999.
- [15] A. Pazy Semigroups of linear operators and applications to partial differential equations, Springer-Verlag, (1983).
- [16] S. Peng Stochastic Hamilton-Jacobi-Bellman equations. SIAM J. Control Optim., 30:284–304, 1992.
- [17] H. Tanabe Equations of evolution. Monographs and Studies in Mathematics, 6. Pitman (Advanced Publishing Program), Boston, Mass.-London, 1979.
- [18] H. Triebel. Interpolation Theory, Function Spaces, Differential Operators vol. 18 of North-Holland Mathematical Library North-Holland Publishing Co., Amsterdam-New York, 1978