

****FULL TITLE****
*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***
****NAMES OF EDITORS****

GRO J1655-40: from ASCA and XMM-Newton Observations

Xiao-Ling Zhang¹, Shuang Nan Zhang^{2,3,4}, Gloria Sala¹,
 Jochen Greiner¹, Yuxin Feng^{3,4}, Yangsen Yao⁵

Abstract. We have analysed four ASCA observations (1994–1995, 1996–1997) and three XMM-Newton observations (2005) of this source, in all of which the source is in high/soft state. We modeled the continuum spectra with relativistic disk model *kerrbb*, estimated the spin of the central black hole, and constrained the spectral hardening factor f_{col} and the distance. If *kerrbb* model applies, for normally used value of f_{col} (1.7), the distance cannot be very small, and f_{col} changes with observations.

1. Background

GRO J1655-40, the second microquasar (after GRS 1915-105), had X-ray outbursts in 1994-1995, 1996-1997, 2005. Its geometric parameters are considered best determined: mass $M_{\text{BH}} = 7.0 \pm 0.2 M_{\odot}$, inclination angle $\theta = 69.50^{\circ} \pm 0.08$ (Orosz & Bailyn 1997), distance $D = 3.2 \pm 0.2$ kpc (Hjellming & Rupen 1995), which makes it a very good laboratory of studying black holes and environments.

The spin of the central black hole has been estimated by various authors with various methods (see, e.g., Zhang et al. 1997; Abramowicz & Kluźniak 2001; Aschenbach 2004; Shafee et al. 2006), and the reported value range from 0.2 (Abramowicz & Kluźniak 2001) to 0.996 (Aschenbach 2004).

In estimating black hole spin from continuum spectral modeling, the color correction factor $f_{\text{col}} = T_{\text{col}}/T_{\text{eff}}$, is one of the key factors. The normally used value of f_{col} is 1.7, following Shimura & Takahara (1995), while many authors believe it should not be constant (see, e.g., Merloni et al. 2000). The distance is also very important. The widely accepted value 3.2 ± 0.2 kpc was challenged by Foellmi et al. (2006), who gave an upper limit of 1.7 kpc.

2. Observations, data reduction and model fitting

We analysed three ASCA observations during the 1994–1995 and the 1996–1997 outbursts, and three XMM-Newton observations during the 2005 outburst, in all of which the source was in high/soft state. For ASCA, only GIS2 data were used, after gain correction and deadtime correction. For XMM-Newton, only Epic-pn

¹MPE, Postfach 1312, 85741 Garching, Germany, zhangx@mpe.mpg.de

²Tsinghua Univ, 100084, Beijing, China

³U. of Alabama, Huntsville, AL 35899, USA

⁴NSSTC, Sparkman Dr. 320, Huntsville AL 35805, USA

⁵MIT Kavli Inst. for Astro. and Space Research, 70 Vassar Street, Cambridge, MA 02139

data were used, after correction for rate-dependent Charge-Transfer-Efficiency (Sala et al. 2006).

The classical way of estimating black hole spin from the continuum spectral fitting is to fit the spectra with disk models, and obtain the spin directly or indirectly. All models take the source distance as parameter, and most models treat the disk as multi-temperature black-body rings and the derived spin value depends on the apparent/effective temperature ratio.

The relativistic disk model *kerrbb* in *XSPEC* was used in the fitting. We let f_{col} vary from 1.0 to 3.0, and D vary from 1.0 kpc and 3.2 kpc. For each combination of f_{col} and D , we fitted the data sets and obtained a spin value, if the fit was acceptable ($\chi^2/dof < 2$). The contour of the derived spin a over D and f_{col} are shown in the Fig. 1.

3. Conclusion

From Fig. 1 we can see, 1. for the normally used f_{col} value 1.7, *kerrbb* model does not favor small distance; 2. because the black hole spin and the source distance should be constant, f_{col} changes dramatically between these observations.




Figure 1. Contour of black hole specific angular momentum a versus distance D and f_{col} . The two dotted lines indicate $D=1.7\text{kpc}$, and $f_{\text{col}} = 1.7$.

References

- Abramowicz, M. A., & Kluźniak, W. 2001, A&A, 374, L19
- Aschenbach, B. 2004, A&A, 425, 1075
- Foellmi, C., Depagne, E., Dall, T. H., & Mirabel, I. F. 2006, A&A, 457, 249
- Hjellming, R.M. & Rupen, M.P. 1995, Nat, 375, 464
- Merloni, A., Fabian, A.C., Ross, R.R. 2000, MNRAS, 313, 193
- Orosz, J. A., & Bailyn, C. D. 1997, ApJ, 477, 876
- Sala, G., et al 2006, A&A, accepted (astro-ph/0606272)
- Shafee, R., et al. 2006, ApJ, 636, L113
- Shimura, T. & Takahara, F. 1995, ApJ, 445, 780
- Zhang, S. N., Cui, W., & Chen, W. 1997, ApJ, 482, L155