The Giant Monopole Resonance in the ^{112–124}Sn Isotopes and the Symmetry Energy Term in Nuclear Incompressibility

T. Li^a U. Garg^a Y. Liu^a R. Marks^a B. K. Nayak^a

P. V. Madhusudhana Rao^a M. Fujiwara^b H. Hashimoto^b K. Kawase^b Y. Nakanishi^b S. Okumura^b M. Yosoi^b M. Itoh^c M. Ichikawa^c R. Matsuo^c T. Terazono^c M. Uchida^d T. Kawabata^e H. Akimune^f

Y. Iwao^g T. Murakami^g H. Sakaguchi^g S. Terashima^g Y. Yasuda^g

J. Zenihiro^g M. N. Harakeh^h

^aDepartment of Physics, University of Notre Dame, Notre Dame, IN 46556, USA

^bResearch Center for Nuclear Physics, Osaka University, Osaka 567-0047, Japan

 $^{\rm c}Cyclotron$ and Radioisotope Center, Tohuku University, Sendai 980-8578, Japan

^dDepartment of Physics, Tokyo Institute of Technology, Tokyo 152-8850, Japan

^eCenter for Nuclear Study, University of Tokyo, Tokyo 113-0033, Japan

^fDepartment of Physics, Konan University, Kobe 658-8501, Japan ^gDepartment of Physics, Kyoto University, Kyoto 606-8502, Japan

^hKVI, 9747 AA Groningen, The Netherlands

Abstract

We have investigated the isoscalar giant monopole resonance (GMR) in the Sn isotopes, using inelastic scattering of 400-MeV α -particles at extremely forward angles, including 0°. A value of -550 ± 100 MeV has been obtained for the asymmetry term, K_{τ} , in the nuclear incompressibility.

Key words: PACS: 24.30.Cz; 21.65.+f; 25.55.Ci; 27.40.+z

Incompressibility of nuclear matter remains a focus of experimental and theoretical investigations because of its fundamental importance in defining the equation of state (EOS) for nuclear matter. The compressional-mode giant resonances – the Giant Monopole Resonance (GMR) and the isoscalar giant dipole resonance (ISGDR), an exotic compressional mode of nuclear oscillation – provide a direct means to experimentally determine the nuclear incompressibility. From recent measurements on the GMR and the

Preprint submitted to Elsevier

6 November 2018

ISGDR, a value of $K_{\infty} = 240 \pm 10$ MeV has been obtained, consistent with results of recent theoretical results in both relativistic and non-relativistic frameworks [1,2,3].

The asymmetry term, K_{τ} , in the expression for nuclear incompressibility is important because it is critical to determining the radii of neutron stars, and also in understanding the compressional-mode resonances in very neutron-rich nuclei, the primary focus of investigations at current and forthcoming radioactive ion beam facilities. In particular, it has been shown that the radius of a 1–1.5 M_{\odot} neutron star is mostly determined by the density dependence of the asymmetry-energy term [4,5].

In this presentation, we report on new measurements on GMR in the even-A Sn isotopes. The experiment was performed at the ring cyclotron facility of the Research Center for Nuclear Physics (RCNP), Osaka University, using inelastic scattering of 400-MeV α particles at extremely forward angles, including 0°. Details of the experimental techniques and the data analysis procedures have been provided previously [6]. Sample background-free "0°" inelastic spectra are presented in Fig. 1 (a). Giant monopole resonance strength distributions were obtained for all Sn isotopes under study using a multipole-decomposition analysis (MDA) [6]; Fig. 1 (b) shows a sampling of MDA fits to angular distribution data. Table I lists the centroid energies and widths extracted from Lorentzian-peak fits to the extracted GMR strength distributions, as well as the "standard" moment-ratios typically used in theoretical calculations.



Fig. 1. (a) 0° inelastic scattering spectrum for $^{112}\text{Sn}(\alpha,\alpha\prime)$ and $^{124}\text{Sn}(\alpha,\alpha\prime)$ at $E_{\alpha} = 400$ MeV. (b) Angular distribution of 1-MeV bins centered at $E_x=16.5$ MeV for $^{112}\text{Sn}(\alpha,\alpha\prime)$ and $^{124}\text{Sn}(\alpha,\alpha\prime)$. The solid squares are the experimental data and the solid line is the MDA fit to the data. Also shown are the contributions to the fit from L=0 (dashed line), L=1 (dotted line), L=2 (dash-dotted line) and IVGDR (dash-dot-dotted line).

Table 1									
Lorentzian-fit	parameters	and n	noment-ratios	for the	GMR strength	$\operatorname{distributions}$	in the	Sn	isotopes

Target	E_{GMR} (MeV)	Γ (MeV)	$m_1/m_0~({ m MeV})$	$\sqrt{m_3/m_1}$ (MeV)	$\sqrt{m_1/m_{-1}}$ (MeV)
$^{112}{ m Sn}_{114}{ m Sn}_{ m Sn}_{ m ^{116}Sn}_{ m ^{118}Sn}$	$16.1 \pm 0.1 \\ 15.9 \pm 0.1 \\ 15.8 \pm 0.1 \\ 15.6 \pm 0.1$	$4.0 \pm 0.4 \\ 4.1 \pm 0.4 \\ 4.1 \pm 0.3 \\ 4.3 \pm 0.4$	16.2 ± 0.1 16.1 ± 0.1 15.8 ± 0.1 15.8 ± 0.1	$16.7 \pm 0.2 \\ 16.5 \pm 0.2 \\ 16.3 \pm 0.2 \\ 16.3 \pm 0.1$	$16.1 \pm 0.1 \\ 15.9 \pm 0.1 \\ 15.7 \pm 0.1 \\ 15.6 \pm 0.1$
120 Sn 122 Sn 122 Sn 124 Sn	$\begin{array}{c} 15.0 \pm 0.1 \\ 15.4 \pm 0.2 \\ 15.0 \pm 0.2 \\ 14.8 \pm 0.2 \end{array}$	$ \begin{array}{c} 1.0 \pm 0.1 \\ 4.9 \pm 0.5 \\ 4.4 \pm 0.4 \\ 4.5 \pm 0.5 \end{array} $	15.7 ± 0.1 15.4 ± 0.1 15.3 ± 0.1	16.2 ± 0.2 15.9 ± 0.2 15.8 ± 0.1	15.5 ± 0.1 15.2 ± 0.1 15.1 ± 0.1

The incompressibility of a nucleus, K_A , may be expressed as:

$$K_A \sim K_{vol}(1 + cA^{-1/3}) + K_{\tau}((N - Z)/A)^2 + K_{Coul}Z^2A^{-4/3}$$
 (1)

For a series of isotopes, the difference $K_A - K_{Coul}Z^2A^{4/3}$ may, thence, be approximated to have a quadratic relationship with the asymmetry parameter, [(N-Z)/A]. Such a quadratic fit, using the customary moment ratio $\sqrt{m_1/m_{-1}}$ for the energy of the GMR in calculating the K_A , gives $K_{\tau} = -550 \pm 100$ MeV; the quoted uncertainty includes the uncertainties in the values of K_A and K_{Coul} , as well as the statistical uncertainties from the fitting procedure.

From the data on the compressional-mode giant resonances, we now have "experimental" values of both K_{∞} and K_{τ} which, together, can provide a means of selecting the most appropriate of the interactions used in EOS calculations. For example, this combination of values for K_{∞} and K_{τ} essentially rules out a vast majority of the Skyrme-type interactions currently in use in nuclear structure calculations [7]. This is borne out by Fig. 2 in which are plotted the values of K_{∞} and K_{τ} for a number of interactions used in both relativistic and non-relativistic calculations. It is clear that nearly all of them fall outside of the "acceptable" region defined by the values obtained in our measurements, leaving a challenge for the theorists to construct appropriate interactions that meet this criterion.



Fig. 2. Values of K_{∞} and K_{τ} calculated from the parameter sets of various interactions as labeled [7]. The vertical and horizontal lines indicate the experimental ranges of K_{∞} and K_{τ} , as determined from the GMR work.

This work has been supported in part by the U.S. National Science Foundation (Grants INT03-42942 and PHY04-57120) and by the Japan Society for Promotion of Science (JSPS).

- [2] B. G. Todd-Rutel and J. Piekarewicz, Phys. Rev. Lett. 95, 122501 (2005).
- [3] B. K. Agrawal, S. Shlomo and V. Kim Au, Phys. Rev. C 68, 031304 (2003).
- [4] J.M. Lattimer and M. Prakash, ApJ **550**, 426 (2001).
- [5] C.J. Horowitz and J. Piekarewicz, Phys. Rev. Lett. 86, 5647 (2001).

^[1] References al., Phys. Rev. C 70, 024307 (2004).

- [6] M. Uchida *et al.*, Phys. Rev. C **69**, 051301 (2004).
 [7] H. Sagawa *et al.*, arXiv:0706.0966.