Analytic derivation of the leading-order gluon distribution function $G(x, Q^2) = xg(x, Q^2)$ from the proton structure function $F_2^p(x, Q^2)$

Martin M. Block,¹ Loyal Durand,² and Douglas W. McKay³

¹Department of Physics and Astronomy, Northwestern University, Evanston, IL 60208

²Department of Physics, University of Wisconsin, Madison, WI 53706

³Department of Physics and Astronomy, University of Kansas, Lawrence, KS 66045

(Dated: November 12, 2018)

We derive a second-order linear differential equation for the leading-order gluon distribution function $G(x,Q^2) = xg(x,Q^2)$ which determines $G(x,Q^2)$ directly from the proton structure function $F_2^p(x,Q^2)$. This equation is derived from the leading order DGLAP evolution equation for $F_2^p(x,Q^2)$, and does not require knowledge of either the individual quark distributions or the gluon evolution equation. Given an analytic expression that successfully reproduces the known experimental data for $F_2^p(x,Q^2)$ in a domain $x_{\min}(Q^2) \leq x \leq x_{\max}(Q^2)$, $Q_{\min}^2 \leq Q^2 \leq Q_{\max}^2$ of the Bjorken variable x and the virtuality Q^2 in deep inelastic scattering, $G(x,Q^2)$ is uniquely determined in the same domain. We give the general solution and illustrate the method using the recently proposed Froissart bound type parametrization of $F_2^p(x,Q^2)$ of E. L. Berger, M. M. Block and C-I Tan, PRL 98, 242001, (2007). Existing leading-order gluon distributions based on power-law descriptions of individual parton distributions agree roughly with the new distributions for $x \gtrsim 10^{-3}$ as they should, but are much larger for $x \lesssim 10^{-3}$.

PACS numbers: 13.85.Hd,12.38.Bx,12.38.-t,13.60.Hb

Introduction. Parton distributions play a key role in our understanding of Standard Model processes, in our predictions for Standard Model processes at accelerators, and in our searches for new physics. In particular, accurate knowledge of gluon distribution functions at small Bjorken x will play a vital role in estimating backgrounds, and hence, our ability to search for new physics at the Large Hadron Collider.

Traditionally, gluon and quark distribution functions have been determined simultaneously by starting with a virtuality Q_0^2 , typically in the 1 to 2 GeV² range, using the two coupled integral-differential DGLAP equations [1] to evolve individual quark and gluon trial distributions to higher Q^2 . The results are adjusted to fit the overall data (mainly the experimental data for the proton structure function $F_2^p(x, Q^2)$) by adjusting the parameters in the initial parton distributions, thus determining the evolved distributions.

In this paper, we present a new and simple method for determining the gluon distribution function $G(x, Q^2) = xg(x, Q^2)$ in leading order (LO) directly from a global parametrization of the data on $F_2^p(x, Q^2)$. The method neither requires knowledge of the separate quark distributions, nor the use of the DGLAP evolution equation for $G(x, Q^2)$, in the region in which proton structure function data exist. We illustrate the method using a Froissart bound-type fit [2] to the proton structure function, and compare with other LO results.

Results on $G(x, Q^2)$ obtained this way do not obviate the need for simultaneous fits to all quark and gluon distributions which may use data other than that on $F_2^p(x, Q^2)$, but should provide a useful check on those results. Strategy for determining gluon distributions. Instead of starting with parametrizations of parton distribution functions at some Q_0^2 and then evolving from Q_0^2 to the desired Q^2 , we follow a new strategy as follows:

- We first make a global parametrization of the experimental proton structure function F₂^p simultaneously in x and Q². Berger, Block and Tan [2] give an example of such such a parametrization which gives an excellent fit to all of the available ZEUS data [3] in the domain 10⁻⁴ ≤ x ≤ .09, 0.11 ≤ Q² ≤ 1200 GeV², and also to H1 results [4] that were not used in the fit. A smooth, accurate fit allows us to calculate the derivatives of F₂^p(x, Q²) we will need below.
- 2. We next derive an inhomogeneous second-order linear differential equation for $G(x, Q^2)$ from the LO DGLAP equation that determines the evolution of the (now known) proton structure function $F_2^p(x, Q^2)$. The inhomogeneous driving term in the equation is determined entirely by $F_2^p(x, Q^2)$ and its derivatives.
- 3. Finally, we solve the differential equation explicitly for the gluon distribution function.

This approach requires no assumptions about either the shape of the gluon distribution or the shape of individual quark distributions. The method gives a global LO gluon distribution function $G(x, Q^2)$ which is unique—within experimental uncertainties—in the region in x and Q^2 space which contains the experimental data on $F_2^p(x, Q^2)$.

Differential equation for the leading order gluon distribution function. The LO DGLAP equation[1] for the evolution of the proton structure function $F_2^p(x, Q^2)$ can be written as

$$\frac{\partial F_2^p(x,Q^2)}{\partial \ln(Q^2)} = x \left[\int_x^1 \frac{F_2^p(z,Q^2) \times K_{qq}(x/z)}{z^2} dz + \sum_i \int_{x_i}^1 e_i^2(z,Q^2) \frac{G(z,Q^2) \times K_{qg}(x_i/z)}{z^2} dz \right].$$
(1)

Here $G(x, Q^2) = xg(x, Q^2)$ and $e_i^2(z, Q^2) = e_i^2\theta(z - x_i)$ where e_i is the electric charge of the quark with flavor i, $x_i = x[1 + (4M_i^2/Q^2)]$, and the step functions enforce the parton level threshold conditions $s \ge 4M_i^2$ for the production of a pair of quarks of mass M_i [5]. The sum runs over all quarks and antiquarks. K_{qq} and K_{qg} are the LO splitting functions for quarks and gluons, respectively.

To illustrate our method most simply, we will here ignore the threshold factors and suppose that we have just four active quarks, u, d, c, s, with all quarks treated as massless. Then $\sum_{i} e_i^2 = 20/9$.

We will show elsewhere that the method can be generalized within the *identical framework* to include the mass-dependent threshold effects—again without requiring knowledge of the individual quark distributions with minimal complication.

With this simplified starting point, we write Eq. (1) as

$$\begin{split} \frac{\partial F_2^p(x,Q^2)}{\partial \ln(Q^2)} &= \frac{\alpha_s}{4\pi} \left\{ 4F_2^p(x,Q^2) + \frac{16}{3} \left[F_2^p(x,Q^2) \ln \frac{1-x}{x} \right. \\ &+ x \int_x^1 \left(\frac{F_2^p(z,Q^2)}{z} - \frac{F_2^p(x,Q^2)}{x} \right) \frac{dz}{z-x} \right] \\ &- \frac{8}{3}x \int_x^1 F_2^p(z,Q^2) \left(1 + \frac{x}{z} \right) \frac{dz}{z^2} \\ &+ \frac{20}{9}x \int_x^1 G(z,Q^2) \left(\frac{x^2 + (z-x)^2}{z^2} \right) \frac{dz}{z^2} \right\}, \quad (2) \end{split}$$

following Durand and Putikka [6] in the first of their Eqs. (19).¹

Integrating the term in line 3 of Eq. (2) by parts and using the boundary condition $F_2^p(1, Q^2) = 0$, we find that

$$-\frac{8}{3}x\int_{x}^{1}F_{2}^{p}(z,Q^{2})\left(1+\frac{x}{z}\right)\frac{dz}{z^{2}} = -4F_{2}^{p}(x,Q^{2})$$
$$-\frac{4}{3}\int_{x}^{1}\frac{\partial F_{2}^{p}}{\partial z}(z,Q^{2})\left(\frac{x^{2}}{z^{2}}+\frac{2x}{z}\right)dz.$$
(3)

We rewrite the large bracket in Eq. (2) as

$$F_2^p(x, Q^2) \ln(1-x) - F_2^p(x, Q^2) \ln x$$

$$+x \int_{x}^{1} F_{2}^{p}(z,Q^{2}) \frac{dz}{z(z-x)}$$

$$-F_{2}^{p}(x,Q^{2}) \left(\int_{x}^{1} \frac{dz}{z} + \int_{x}^{1} \frac{x}{z(z-x)} dz \right) \quad (4)$$

$$= F_{2}^{p}(x,Q^{2}) \ln(1-x)$$

$$+x \int_{x}^{1} \frac{F_{2}^{p}(z,Q^{2}) - F_{2}^{p}(x,Q^{2})}{z(z-x)} dz$$

$$\int_{x}^{1} \partial F^{p} z$$

 $= \int_{x}^{1} \frac{\partial F_{2}^{2}}{\partial z}(z,Q^{2}) \ln \frac{z}{z-x} dz, \qquad (5)$ ere we used the identity $\frac{1}{z-x} = \frac{1}{z} + \frac{x}{z(z-x)}$ to get

where we used the identity $\frac{1}{z-x} = \frac{1}{z} + \frac{x}{z(z-x)}$ to get from the large bracket in Eq. (2) to Eq. (4), and partial integration, with $F_2^p(1, Q^2) = 0$, to get to Eq. (5).

Next, we combine terms and define $\mathcal{F}_2^p(x, Q^2)$ as

$$\mathcal{F}_{2}^{p}(x,Q^{2}) \equiv \frac{\partial F_{2}^{p}(x,Q^{2})}{\partial \ln(Q^{2})} \\ -\frac{\alpha_{s}}{4\pi} \left\{ \frac{16}{3} \int_{x}^{1} \frac{\partial F_{2}^{p}}{\partial z}(z,Q^{2}) \ln \frac{z}{z-x} dz \right. \\ \left. -\frac{4}{3} \int_{x}^{1} \frac{\partial F_{2}^{p}}{\partial z}(z,Q^{2}) \left(\frac{x^{2}}{z^{2}} + \frac{2x}{z} \right) dz \right\}.$$
(6)

After multiplying both sides by $4\pi/\alpha_s$, we rewrite Eq. (2) as

$$\frac{4\pi}{\alpha_s} \mathcal{F}_2^p(x, Q^2) / x = \frac{20}{9} \int_x^1 G(z, Q^2) \left(\frac{x^2 + (z - x)^2}{z^2}\right) \frac{dz}{z^2}.$$
 (7)

After successive differentiations of both sides of Eq. (7) with respect to x, multiplication by $-x^4$, and some rearranging, we find an inhomogeneous second-order differential equation which determines $G(x, Q^2)$ in terms of $\mathcal{F}_2^p(x, Q^2)$:

$$x^{2} \frac{\partial^{2}}{\partial x^{2}} G(x, Q^{2}) - 2x \frac{\partial}{\partial x} G(x, Q^{2}) + 4G(x, Q^{2})$$
$$= -\frac{4\pi}{\alpha_{s}} \frac{9}{20} x^{4} \frac{\partial^{3}}{\partial x^{3}} \left(\frac{\mathcal{F}_{2}^{p}(x, Q^{2})}{x}\right). \tag{8}$$

To simplify the notation in this and other equations, we define $\mathcal{G}_n(x, Q^2)$ for *n* active quarks by

$$\mathcal{G}_n(x,Q^2) \equiv -\frac{4\pi}{\alpha_s} x^4 \frac{\partial^3}{\partial x^3} \left(\frac{\mathcal{F}_2^p(x,Q^2)}{x}\right) \Big/ \sum_{i=1}^n e_i^2.$$
(9)

Again, the sum is over all active quarks and antiquarks. The right hand side of the Eq. (8) is then $\mathcal{G}_4(x, Q^2)$, where $\sum_{i=1}^4 e_1^2 = 20/9$.

Explicit evaluation of this term gives

$$\begin{aligned} \mathcal{G}_4(x,Q^2) &= \\ &-\frac{4\pi}{\alpha_s}\frac{9}{20}x^4\frac{\partial^4(F_2^p(x,Q^2)/x)}{\partial x^3\partial\ln Q^2} + \frac{12}{5}x\frac{\partial F_2^p}{\partial x}(x,Q^2) \end{aligned}$$

¹ The term $-\frac{1}{6}G(x,Q^2)$ in the first of Eqs. (19) in [6] should be deleted there and inserted in the second of Eqs. 19 with a coefficient $2n_f$. A term $-6x \int_x^1 dx_1 G(x_1,t)/x_1^2$ was also dropped in the transition from Eq. (18) to the second of Eqs. (19) and should be inserted. Note that in Ref. [6], the notation $G(x,Q^2)$ was used for the function $g(x,Q^2)$ of the present paper.

$$-3x^2 \frac{\partial^2 F_2^P}{\partial x^2}(x,Q^2) + \frac{9}{5}x^3 \frac{\partial^3 F_2^p}{\partial x^3}(x,Q^2)$$
$$+ \frac{12}{5}x^4 \frac{\partial^3}{\partial x^3} \frac{1}{x} \int_x^1 \frac{\partial F_2^p}{\partial z}(z,Q^2) \ln \frac{z}{z-x} dz. \quad (10)$$

The derivatives of the factor $\ln[z/(z-x)]$ in the integrand in the last term can be treated using the identity

$$\lim_{\epsilon \to 0} \frac{\partial}{\partial x} \int_{x+\epsilon}^{1} dz \ln(z-x) f(z,Q^2) = -\ln(1-x) f(1,Q^2) + \int_{x}^{1} dz \ln(z-x) \frac{\partial f}{\partial z}(z,Q^2).$$
(11)

In the present case, $F_2^p(x, Q^2)$ and its first two derivatives with respect to x are expected vanish at x = 1. The final integrand has only a logarithmic singularity.

Analytic solution for $G(x, Q^2)$. With the notation above, Eq. (8) becomes the inhomogeneous differential equation

$$x^{2} \frac{\partial^{2}}{\partial x^{2}} G(x, Q^{2}) - 2x \frac{\partial}{\partial x} G(x, Q^{2}) + 4G(x, Q^{2})$$
$$= \mathcal{G}_{4}(x, Q^{2}). \quad (12)$$

Introducing the new variable $v = \ln(1/x)$, we rewrite Eq. (12), a linear 2^{nd} order inhomogeneous equation, as

$$\left(\frac{\partial^2}{\partial v^2} + 3\frac{\partial}{\partial v} + 4\right)\hat{G}(v, Q^2) = \hat{\mathcal{G}}_4(v, Q^2), \quad (13)$$

with $\hat{G}(v,Q^2) \equiv G(e^{-v},Q^2), \ \hat{\mathcal{G}}_4(v,Q^2) \equiv \mathcal{G}_4(e^{-v},Q^2).$ Defining the operator $D \equiv \partial/\partial v$, we factor Eq. (13) as

$$(D - \lambda_{+})(D - \lambda_{-})\hat{G}(v, Q^{2}) = \hat{\mathcal{G}}_{4}(v, Q^{2})$$
 (14)

where $\lambda_{\pm} = k \pm i\omega$ with k = -3/2, $\omega = \sqrt{7}/2$.

To construct the solution of Eq. (14), we introduce the solutions $e^{\lambda \pm v}$ of the homogeneous equation and the functions

$$\mathcal{G}_{\pm}(v,Q^2) = \int_0^v dv' e^{-\lambda_{\pm}v'} \hat{\mathcal{G}}_4(v',Q^2).$$
(15)

The solution that satisfies the boundary conditions that $\hat{G}(v, Q^2)$ and $\partial \hat{G}(v, Q^2)/\partial v$ vanish at v = 0, or equivalently, that $G(x, Q^2)$ and $\partial G(x, Q^2)/\partial x$ vanish at x = 1, is then

$$\hat{G}(v,Q^2) = \frac{1}{\lambda_+ - \lambda_-} \left[e^{\lambda_+ v} \mathcal{G}_+(v,Q^2) - e^{\lambda_- v} \mathcal{G}_-(v,Q^2) \right]$$

$$= \frac{1}{\omega} \int_0^v e^{k(v-v')} \sin(\omega(v-v')) \hat{\mathcal{G}}_4(v',Q^2) \, dv'.$$
(16)

This result is completely general and gives the *exact* LO expression for $\hat{G}(v, Q^2)$ once $\hat{\mathcal{G}}_4(v, Q^2)$ is known.

We re-emphasize at this point that both $\hat{\mathcal{G}}_4(v,Q^2)$ and $\hat{G}(v,Q^2)$ are completely determined by $F_2^p(x,Q^2)$ through the expressions in Eqs. (10) and (16), and the definition $\hat{\mathcal{G}}_4(v,Q^2) = \mathcal{G}_4(e^{-v},Q^2)$. Given a smooth analytic parametrization of $F_2^p(x,Q^2)$ as a simultaneous function of x and Q^2 in a domain $x_{\min}(Q^2) \leq x \leq x_{\max}(Q^2)$, $Q_{\min}^2 \leq Q^2 \leq Q_{\max}^2$, we can calculate $\hat{\mathcal{G}}_4(v,Q^2)$ and thus determine the gluon distribution function $G(x,Q^2)$ to within the accuracy of the parametrization. Different smooth parametrizations that fit the data on $F_2^p(x,Q^2)$ equally well in the specified domain should give equivalent gluon distributions within that domain.

We stress that the exact LO equation, (16), directly builds in the Q^2 evolution for $G(x, Q^2)$ in terms of the measured $F_2^p(x, Q^2)$ in the experimental domain. Since higher order (NLO) and higher twist effects are embodied in $F_2^p(x, Q^2)$, there is no reason to to expect that LO DGLAP evolution of $G(x, Q^2)$ at fixed x should agree with our evaluation of the Q^2 dependence. In fact, the difference is an indication of corrections to the LO DGLAP evolution in this domain. Explicit NLO corrections to relation (16), and therefore to $G(x, Q^2)$, will be discussed separately. One might use the gluon evolution DGLAP equation, with its dependence on the singlet quark distribution, if one wished to extend the gluon distribution determined above beyond the experimental domain in Q^2 at fixed x.

We note in addition that the singlet quark distribution can itself be related within the experimental domain to the now-known gluon distribution, using the evolution equation for $G(x, Q^2)$ and a construction similar to that used to relate $G(x, Q^2)$ to $F_2^p(x, Q^2)$, but have so far not investigated this procedure in detail.

Example: solution for $G(x,Q^2)$ using a Froissart bounded structure function $F_2^p(x,Q^2)$. We illustrate the procedure above using the Froissart-bound type parametrization of the proton structure function given by Berger, Block, and Tan [2] for $x \leq x_P = 0.09$,

$$F_2^p(x,Q^2) = (1-x) \left[\frac{F_P}{1-x_P} + \left(a_0 + a_1 \ln(Q^2) + a_2 \ln^2(Q^2) \right) \ln \left[\frac{x_P(1-x)}{x(1-x_P)} \right] + \left(b_0 + b_1 \ln(Q^2) + b_2 \ln^2(Q^2) \right) \ln^2 \left[\frac{x_P(1-x)}{x(1-x_P)} \right], \quad (17)$$

where $F_P = 0.41$ is the value of F_2^p at the approximate fixed point in Q^2 at $x = x_P$, and

$$a_{0} = -5.381 \times 10^{-2} \pm 2.17 \times 10^{-3},$$

$$a_{1} = 2.034 \times 10^{-2} \pm 1.19 \times 10^{-3},$$

$$a_{2} = 4.999 \times 10^{-4} \pm 2.23 \times 10^{-4},$$

$$b_{0} = 9.955 \times 10^{-3} \pm 3.09 \times 10^{-4},$$

$$b_{1} = 3.810 \times 10^{-3} \pm 1.73 \times 10^{-4},$$

$$b_{2} = 9.923 \times 10^{-4} \pm 2.85 \times 10^{-5},$$
(18)

In the absence so far of a global fit to $F_2^p(x, Q^2)$ for $x_P < x \le 1$, we follow the work of Berger, Block, McKay

and Tan [7] and approximate $F_2^P(x, Q^2)$ in that region as

$$F_2^p(x,Q^2) \approx F_P\left(\frac{x}{x_P}\right)^{\rho(Q^2)} \left(\frac{1-x}{1-x_P}\right)^3,$$
 (19)

or by that form multiplied by an extra factor $[1+b(Q^2)(x-x_P)^2((1-x)/(1-x_P))^c]$. The exponent $\rho(Q^2)$ is chosen so that the values of the functions and their first derivatives match at $x = x_P$. In the extended form, the coefficient $b(Q^2)$ is used to match the second derivatives at $x = x_P$, and c is used to obtain a rough fit to the high-x data. We have also considered other parametrizations. The results for $\mathcal{G}_4(x, Q^2)$ and $G(x, Q^2)$ turn out to be insensitive to this parametrization except for x near x_P , where $G(x, Q^2)$ is already very small, and are essentially determined for $x < x_P$ (or $v > v_P$) by the experimentally determined expression for $F_2^p(x, Q^2)$ in Eq. (17).

We have found that we can parametrize the function $\hat{\mathcal{G}}_4(v,Q^2)$ calculated this way to high numerical accuracy for $v > v_P$ as a second degree polynomial in $v = \ln(1/x)$ whose coefficients are quadratic polynomials in $\ln Q^2$, i.e., as

$$\hat{\mathcal{G}}_4(v, Q^2) = \alpha(Q^2) + \beta(Q^2)v + \gamma(Q^2)v^2,$$
 (20)

with an appropriate smooth extension to $v < v_P$.

It is also possible to skip the separate evaluation of $\hat{\mathcal{G}}_4(v,Q^2)$ by integrating repeatedly by parts in the expression for $\hat{\mathcal{G}}(v,Q^2)$ in Eq. (16) to eliminate the leading derivatives in the expression for $\hat{\mathcal{G}}_4$ in Eq. (10). The resulting expression for $\hat{\mathcal{G}}(v,Q^2)$ is rather lengthy, but the leading terms depend directly on F_2^p and $\partial F_2^p/\partial \ln Q^2$ without further integration. This procedure reduces the numerical sensitivity of the results to the higher derivatives of F_2^p that appear in Eq. (10). The results for $\mathcal{G}(x,Q^2)$ obtained using these two methods agree.

In the present numerical work, we used

$$\alpha_s(Q^2) \equiv \frac{12\pi}{25\ln(Q^2/\Lambda^2)},\tag{21}$$

for four active flavors, with $\Lambda = 0.153$ GeV adjusted to give $\alpha_s(M_Z^2) = 0.118$.

Returning to the variable x, we write our final analytic answer, for $0 < x \leq x_P$, as

$$G(x,Q^2) = -\frac{1}{\omega} \int^x \frac{dz}{z} \left(\frac{z}{x}\right)^k \sin\left(\omega \ln(\frac{z}{x})\right) \mathcal{G}_4(z,Q^2)$$

= -0.459 - 0.143 ln(Q²) - 0.0155 ln²(Q²) +
[0.231+0.00971 ln(Q²) - 0.0147 ln²(Q²)] ln(1/x) +
[0.0836+0.06328 ln(Q²)+0.0112 ln²(Q²)] ln²(1/x).(22)

This is a simple quadratic polynomial in $\ln(1/x)$, with quadratic polynomial coefficients in $\ln(Q^2)$.

To make a rough estimate of the uncertainty in $G(x, Q^2)$ caused by the experimental uncertainties in the

input function $F_2^p(x, Q^2)$, we integrate Eq. (7) twice by parts with respect to $\ln(1/x)$ and drop terms that are suppressed by powers of $\ln(1/x)$ to get the zeroth order approximation

$$\tilde{G}(x,Q^2) \approx \frac{27}{40} \frac{4\pi}{\alpha_s} \frac{\partial F_2^p(x,Q^2)}{\partial \ln(Q^2)}.$$
(23)

Numerical studies show that $\mathcal{F}_2^p(x,Q^2)$ is dominated at small x by $\partial F_2^p(x,Q^2)/\partial \ln(Q^2)$. Figure 2 of Ref. [2] shows the $\pm 1\sigma$ error bands of $\partial F_2^p(x,Q^2)/\partial \ln(Q^2)$ due to the parameter errors in Eq. (18), including their correlations, at x = 0.00008, as a function of Q^2 . For Q^2 in the range 5—200 GeV², it was found [2] that $\Delta[\partial F_2^p(x,Q^2)/\partial \ln(Q^2))]/[\partial F_2^p(x,Q^2)/\partial \ln(Q^2)] \approx$ 0.015 - 0.020, leading to the rough estimate

$$\Delta G(x, Q^2) / G(x, Q^2) \big|_{x \sim 0.0001} \approx 2\%,$$
 (24)

for the statistical error associated with the uncertainties in the fit to $F_2^p(x, Q^2)$ [2] in Eq. (17).



FIG. 1: LO gluon distribution functions $G = xg(x,Q^2)$ for virtualities $Q^2 = 5$ and 20 GeV². The curves labeled $\ln^2(1/x)$ are from our LO Froissart bound fit of Eq. (22). The curves labeled CTEQ5 are LO CTEQ5 curves [8]. The vertical lines indicate the minimum x values that the ZEUS [3] collaboration achieved, for the Q^2 in the plot.

We emphasize that only experimental F_2^p data were needed to obtain the numerical results in Eq. (22) for $x \leq x_P$. The 6-parameter fit of Ref. [2] which we used was constructed using all of the available ZEUS data [3], with $x_P \geq x \gtrsim 10^{-3}$ to 10^{-4} depending in the virtuality, and $0.11 \leq Q^2 \leq 1200 \text{ GeV}^2$. The fit was excellent, with a $\chi^2/\text{d.o.f.} = 1.09$ for 169 degrees of freedom. The use of the low- Q^2 data on $F_2^p(x, Q^2)$, a virtual photon-proton scattering cross section, is important in determining the global fit. However, we use Eq. (16) or Eq. (22) to determine $G(x, Q^2)$ only in the perturbative region $Q^2 \gtrsim 2$ GeV². We show our LO results from Eq. (22) for $G(x,Q^2)$ for $Q^2 = 5$ and 20 GeV² in Fig. 1, in which we compare them to ZEUS [9] and CTEQ5 LO power law gluon distributions [8] calculated at the same virtualities. Inspection of Fig. 1 shows rough agreement for x-values $\gtrsim 10^{-3}$, where there are experimental data [3] but differ markedly for smaller x, with our extrapolated fits being much smaller than the extrapolated CTEQ5 (and other) power law fits.

We recall that the functions $\hat{\mathcal{G}}_4(v, Q^2)$ and $\mathcal{G}_4(x, Q^2)$ on the right hand sides of the analytical expressions for $G(x, Q^2)$ in Eqs. 16 and 22 depend only on numerical values of $F_2^p(x, Q^2)$ and $\alpha_s(Q^2)$, which are in turn determined by experiment. Thus, if two global fits to $F_2^p(x, Q^2)$ —using different parametrizations—both reproduce the experimental data in a given region of x and Q^2 and treat $\alpha_s(Q^2)$ and the number of active flavors equivalently, they should give the same values for the gluon distribution in that region. This is roughly true here for the CTEQ5, ZEUS and Froissart-type fits for $x \gtrsim 10^{-3}$, as seen in Fig. 1. The striking disagreements occur in the small-x region, where there are no longer any data for $F_2^p(x, Q^2)$ [3], and involve large extrapolations of the forms determined where data exist.

Conclusions. We have shown here that, for a fixed number of active quarks, all that is actually necessary to calculate LO gluon distributions in a region of x and Q^2 is knowledge of $F_2^p(x, Q^2)$ and $\alpha_s(Q^2)$ in that region. The integrals in Eq. (16) and Eq. (22) give the *exact* LO expression for $G(x, Q^2)$ in terms of a function determined by $F_2^p(x, Q^2)$. We used the complete LO splitting functions $K_{qg}^{(1)}(z)$ and $K_{qq}^{(1)}(z)$, so no approximation is made at the LO level. Further, we did not need to use the gluon evolution equation in the derivation of our differential equation for $G(x, Q^2)$ in LO, but only the evolution equation for $F_2^p(x, Q^2)$.

Using the accurate Froissart-bound type parametrization of $F_2^p(x, Q^2)$ [2] in Eq. (22), we obtained an analytic solution for $G(x, Q^2)$ in the region in which there are experimental data, with $\leq 2\%$ uncertainty due to fitting parameter errors of F_2^p . There should be *no* difference between our results and other LO solutions for $x_P \gtrsim x \gtrsim 10^{-3}$ to 10^{-4} to the extent that they all reproduce the $F_2^p(x, Q^2)$ data equally well in this x region. Presumably, the differences seen in Fig. 1 are due to CTEQ (and ZEUS) having used detailed modeldependent functional forms for the quark and gluon distributions in conjunction with the DGLAP evolution in their analysis, a procedure that involves extra assumptions, as well as possible differences in the treatment of the number of active quarks. Further, the CTEQ group [8] used additional data sets.

One of course wants information on the quark distribution functions as well as the gluon distribution for applications to high-energy particle processes, and the use of the usual DGLAP equations allows the use of data other than those on $F_2^p(x, Q^2)$, and the separate extraction of all these parton distribution functions. We believe, however, that the present technique is simple and interesting, and gives a useful check on other approaches at LO.

We will present our generalization of this technique to include the effects of quark masses elsewhere. Again, we stress that *no* knowledge of individual quark distributions is needed. We also will use this technique to explore the singlet quark distribution, using the gluon evolution equation. We also intend to explore the extent to which this technique can be expanded to obtain NLO gluon distribution functions.

Acknowledgements: M.M.B. and L.D. thank the Aspen Center for Physics for its hospitality during the time parts of this work were done. D.W.M. receives support from DOE Grant No. DE-FG02-04ER41308.

- V. N. Gribov and L. N. Lipatov, Sov. J. Nucl. Phys. 15, 438 (1972); G. Altarelli and G. Parisi, Nucl. Phys. B126, 298 (1977); Yu. L. Dokshitzer, Sov. Phys. JETP 46, 641 (1977).
- [2] M. M. Block, E. L. Berger and C-I Tan, Phys. Rev. Lett. 97, 252003 (2006); E. L. Berger, M. M. Block and C-I Tan, Phys. Rev. Lett. 98, 242001, 2007.
- [3] ZEUS Collaboration, V. Chekanov *et al.*, Eur. Phys. J. C21, 443 (2001).
- [4] H1 Collaboration, C. Adloff *et al.*, Eur. Phys. J. C21, 33 (2001)
- [5] W. K. Tung, H. L. Lai, A. Belyaev, J. Pumplin, D. Stump, and C.-P. Yuan, JHEP 0702:053,2007, hep-ph/0611254.
- [6] L. Durand and W. Putikka, Phys. Rev. D 36, 2840 (1987).
- [7] M. M. Block, E. L. Berger, D. W. McKay and C-I Tan, arXiv: 0708.1960v1 [hep-ph](2007).
- [8] CTEQ Collaboration, H. L. Lai et al., Eur. Phys. J. C 12, 375 (2000); the LO CTEQ gluon distributions were calculated using the Mathematica package CTEQ5, taken from F. Olness, http://www.phys.psu.edu/~cteq.
- [9] ZEUS Collaboration, M. Derrick *et al.*, Phys. Lett. B 345, 576 (1995).