

# Erratum: Luminosity function, sizes and FR dichotomy of radio-loud AGN

Christian R. Kaiser<sup>\*</sup> and Philip N. Best

10 November 2018

**Key words:** errata, addenda – galaxies: jets – galaxies: active – galaxies: luminosity function – radio continuum: galaxies

The paper ‘Luminosity function, sizes and FR dichotomy of radio-loud AGN’ was published in Mon. Not. R. Astron. Soc. **381**, 1548–1560, (2007). Since publication we discovered a mistakes in some of the equations in the appendix which describe the radio source evolution according to Kaiser & Alexander (1997) and Kaiser et al. (1997). The mistakes do not alter our conclusions in the original paper in any way. However, to avoid future confusion arising from their application, we correct them here.

In the expressions for the energy densities of the magnetic field,  $u_B$ , and the relativistic electrons,  $u_e$ , equations (A9) and (A10), we accidentally swapped the symbols for the energy densities. The correct expressions read

$$u_e(t_i) = \frac{p(t_i)}{(\Gamma_l - 1)(k + 1)(r + 1)}, \quad (\text{A9})$$

$$u_B(t_i) = \frac{rp(t_i)}{(\Gamma_l - 1)(r + 1)}. \quad (\text{A10})$$

Subsequently, the expression for the energy density of the magnetic field at the observing time  $t$  becomes

$$u_B(t_i) = \frac{rp(t)}{(\Gamma_l - 1)(r + 1)} \left(\frac{t}{t_i}\right)^{a_1(4/3 - \Gamma_l)/4}. \quad (\text{A11})$$

The changes also affect the expression for the Lorentz factor of the emitting electrons. We therefore need to replace equation (A15) with

$$f_\gamma = \sqrt{\frac{2\pi m_e v}{e}} \sqrt{\frac{(\Gamma_l - 1)(r + 1)}{2\mu_0 r}}. \quad (\text{A15})$$

Also, the normalisation of the electron energy spectrum changes to

$$f_n = \frac{1}{(\Gamma_l - 1)(r + 1)(k + 1)m_e c^2} \left( \frac{\gamma_{\min}^{2-m} - \gamma_{\max}^{2-m}}{m-2} - \frac{\gamma_{\min}^{1-m} - \gamma_{\max}^{1-m}}{m-1} \right)^{-1}, \quad (\text{A17})$$

for  $m \neq 2$  and

$$f_n = \frac{1}{(\Gamma_l - 1)(r + 1)(k + 1)m_e c^2}$$

$$\left( \ln \frac{\gamma_{\max}}{\gamma_{\min}} - \frac{1}{\gamma_{\min}} + \frac{1}{\gamma_{\max}} \right)^{-1}, \quad (\text{A18})$$

for  $m = 2$ .

Finally, the factor  $f_L$  in the expression for the radio luminosity should be

$$f_L = \frac{2A^{2(1-\Gamma_l)/\Gamma_l} r f_\gamma^{3-m} f_n \sigma_{TC}}{3(r+1)v}. \quad (\text{A20})$$

All other equations in the appendix and the main text remain unchanged.

## References

- Kaiser C. R., Alexander P., 1997, MNRAS, 286, 215  
 Kaiser C. R., Dennett-Thorpe J., Alexander P., 1997, MNRAS, 292, 723

<sup>\*</sup> crk@soton.ac.uk