

# A LOCAL FORM FOR THE AUTOMORPHISMS OF THE SPECTRAL UNIT BALL

PASCAL J. THOMAS

ABSTRACT. If  $F$  is an automorphism of  $\Omega_n$ , the  $n^2$ -dimensional spectral unit ball, we show that, in a neighborhood of any cyclic matrix of  $\Omega_n$ , the map  $F$  can be written as conjugation by a holomorphically varying non singular matrix. This provides a shorter proof of a theorem of J. Rostand, with a slightly stronger result.

## 1. BACKGROUND

Let  $\mathcal{M}_n$  be the set of all  $n \times n$  complex matrices. For  $A \in \mathcal{M}_n$  denote by  $sp(A)$  the spectrum of  $A$ . The spectral ball  $\Omega_n$  is the set

$$\Omega_n := \{A \in \mathcal{M}_n : \forall \lambda \in sp(A), |\lambda| < 1\}.$$

Let  $F$  be an automorphism of  $\Omega_n$ , that is to say, a biholomorphic map of the spectral ball into itself. Ransford and White [6] proved that, by composing with a natural lifting of a Möbius map of the disk, one could reduce oneself to the case where  $F(0) = 0$ , and that in that case the linear map  $F'(0)$  was a linear automorphism of  $\Omega_n$ , so that by composing with its inverse, one is reduced to the case  $F(0) = 0$ ,  $F'(0) = I$  (the identity map). We then say that the automorphism is *normalized*. Ransford and White [6] proved that such automorphisms preserve the spectrum of matrices.

We say that two matrices  $X, Y$  are *conjugate* if there exists  $Q \in \mathcal{M}_n^{-1}$  such that  $X = Q^{-1}YQ$ .

Baribeau and Ransford [1] (see also [2] for a more elementary proof) proved that every spectrum-preserving  $\mathcal{C}^1$ -diffeomorphism of an open subset of  $\mathcal{M}_n$ , and thus every normalized automorphism of the spectral ball is a pointwise conjugation:

$$(1) \quad F(X) = Q(X)^{-1}XQ(X).$$

---

2000 *Mathematics Subject Classification.* 32H02, 32A07, 15A21.

*Key words and phrases.* holomorphic automorphisms, spectral ball, symmetrized polydisc, cyclic matrices.

This paper was made possible in part by a grant from the French Ministry of Foreign Affairs in the framework of the ECO Net programme, file 10291 SL, coordinated by Ahmed Zeriahi.

Rostand's contribution [7] was to show that  $Q(X)$  could be chosen locally holomorphically in a neighborhood of every  $X$  admitting  $n$  distinct eigenvalues.

We will give a short proof of a slightly stronger result: the exceptional set of matrices where the local holomorphic choice cannot be guaranteed will be of complex codimension 2 instead of 1.

The motivation for this result was a conjecture formulated in [6] about the automorphisms of the spectral ball, which reduces to asking whether any normalized automorphisms can be written in the form (1), where  $Q$  would be globally homomorphic on  $\Omega_n$ , and depend only on the conjugacy class of  $X$ . Notice that a recent result of Zwonek [8] shows that any proper map of the spectral ball to itself is actually an automorphism of it, so that the proof of the Ransford-White conjecture would yield a description of all the proper maps of the spectral ball into itself.

I wish to thank Nikolai Nikolov, who told me about this circle of ideas. Without the fruitful discussions I had with him on this topic, this paper wouldn't have been written.

## 2. STATEMENT

**Definition 1.** *We say that a matrix  $M$  is cyclic (or non-derogatory) if there exists a cyclic vector for  $M$ , i.e.  $v \in \mathbb{C}^n$  such that  $(v, Mv, \dots, M^k v, \dots)$  spans  $\mathbb{C}^n$ , which is equivalent to the fact that  $(v, Mv, \dots, M^{n-1}v)$  is a basis of  $\mathbb{C}^n$ .*

Many equivalent definitions of this notion can be found, for instance in [3] and [4], or [5, Proposition 3]. We point one out:  $M$  is cyclic if and only if for any  $\lambda \in \mathbb{C}$ ,  $\dim \text{Ker}(M - \lambda I_n) \leq 1$ . In particular, any matrix with  $n$  distinct eigenvalues is cyclic, and for any given spectrum  $\lambda_1, \dots, \lambda_n$ , the set of non-cyclic matrices with that spectrum is the algebraic set

$$\{M : \exists j : \dim \text{Ker}(M - \lambda_j I_n) \geq 2\}.$$

Hence the set of non-cyclic matrices is of codimension 1 in the set of matrices which admit at least one multiple eigenvalue, itself of codimension 1 in  $\mathcal{M}_n$ .

**Theorem 2.** *Let  $F$  be a spectrum-preserving holomorphic map of  $\Omega_n$ . Let  $X_0 \in \Omega_n$  be a cyclic matrix. Then there exists a neighborhood  $\mathcal{V}_{X_0}$  of  $X_0$  and a map  $Q$  holomorphic from  $\mathcal{V}_{X_0}$  to  $\mathcal{M}_n^{-1}$  such that for any  $X \in \mathcal{V}_{X_0}$ ,  $F(X) = Q(X)^{-1}XQ(X)$ .*

## 3. PROOF

We fix some notation. For  $A \in \mathcal{M}_n$ , let

$$\sigma_j(A) := \sigma_j(\lambda_1, \dots, \lambda_n) := (-1)^j \sum_{1 \leq k_1 < \dots < k_j \leq n} \lambda_{k_1} \dots \lambda_{k_j}$$

and  $\lambda_1, \dots, \lambda_n$  are the (possibly equal) eigenvalues of  $A$ . Those are polynomials in the coefficients of  $A$ . Put  $\sigma := (\sigma_1, \dots, \sigma_n) : \mathcal{M}_n \rightarrow \mathbb{C}^n$ .

Conversely, given  $a := (a_1, \dots, a_n) \in \mathbb{C}^n$ , the associated *companion matrix*  $\mathcal{C}_a$  is

$$\begin{pmatrix} 0 & & & -a_n \\ 1 & 0 & & \vdots \\ & 1 & \ddots & \vdots \\ & & \ddots & 0 & -a_2 \\ & & & 1 & -a_1 \end{pmatrix}.$$

The companion matrix associated to a matrix  $A$  is  $\mathcal{C}_{\sigma(A)}$ . They have the same characteristic polynomial, or equivalently  $\sigma(\mathcal{C}_{\sigma(A)}) = \sigma(A)$ .

Now given a matrix  $X_0$  as in the Theorem, and a vector  $v_0$  cyclic for  $X_0$ , let

$$\mathcal{U}_{X_0} := \{M \in \mathcal{M}_n : \det(v_0, Mv_0, \dots, M^{n-1}v_0) \neq 0\}.$$

This is a neighborhood of  $X_0$ . Let  $P_{v_0}(M)$  be the matrix with columns  $(v_0, Mv_0, \dots, M^{n-1}v_0)$ ; this depends polynomially on the entries of  $M$ , and is invertible. One can see that for  $X \in \mathcal{U}_{X_0}$ ,

$$P_{v_0}(X)^{-1}XP_{v_0}(X) = \mathcal{C}_{\sigma(X)}$$

(the  $n - 1$ st columns coincide, and they have the same characteristic polynomial).

By the Baribeau-Ransford theorem [1]  $F(X_0)$  is conjugate to  $X_0$ , therefore cyclic. So there is a neighborhood  $\mathcal{U}_{F(X_0)}$  where the relation  $P_{w_0}(Y)^{-1}YP_{w_0}(Y) = \mathcal{C}_{\sigma(Y)}$  holds. Take  $\mathcal{V}_{X_0} \subset \mathcal{U}_{X_0}$  small enough so that  $F(\mathcal{V}_{X_0}) \subset \mathcal{U}_{F(X_0)}$ . For any  $X \in \mathcal{V}_{X_0}$ , using the fact that  $F$  is spectrum-preserving,

$$\begin{aligned} F(X) &= P_{w_0}(F(X))\mathcal{C}_{\sigma(F(X))}P_{w_0}(F(X))^{-1} \\ &= P_{w_0}(F(X))\mathcal{C}_{\sigma(X)}P_{w_0}(F(X))^{-1} \\ &= P_{w_0}(F(X))P_{v_0}(X)^{-1}XP_{v_0}(X)P_{w_0}(F(X))^{-1}, \end{aligned}$$

so that we have the theorem with  $Q(X) = P_{v_0}(X)P_{w_0}(F(X))^{-1}$ .  $\square$

Notice that there is no hope to make a global holomorphic choice of  $v_0$  on the whole open of cyclic matrices. Indeed, since the complement of this is of codimension 2, we could then extend it to the whole of

$\Omega_n$  by Hartog's phenomenon, but it would mean that all matrices are cyclic, which is obviously false.

## REFERENCES

- [1] L. Baribeau and T. J. Ransford, *Non-linear spectrum-preserving maps*, Bull. London Math. Soc., 32 (2000), 8–14.
- [2] L. Baribeau and S. Roy, *Caractérisation spectrale de la forme de Jordan*, Linear ALgebra Appl. 320 (2000), 183–191.
- [3] R. A. Horn, C. R. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, New York, Melbourne, 1985.
- [4] R. A. Horn, C. R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, New York, Melbourne, 1991.
- [5] N. Nikolov, P. J. Thomas, W. Zwonek, *Discontinuity of the Lempert function and the Kobayashi–Royden metric of the spectral ball*, Preprint, 2007 (arXiv:math.CV/0704.2470).
- [6] T. J. Ransford and M. C. White, *Holomorphic self-maps of the spectral unit ball*, Bull. London Math. Soc., 23 (1991), 256–262.
- [7] J. Rostand, *On the automorphisms of the spectral unit ball*, Studia Math. 155 (3) (2003), 207–230.
- [8] W. Zwonek, *Proper holomorphic mappings of the spectral unit ball*, preprint, 2007 (arXiv:math.CV/0704.0614).

INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UMR CNRS 5219, UNIVERSITÉ PAUL SABATIER, 118 ROUTE DE NARBONNE, F-31062 TOULOUSE CEDEX, FRANCE

*E-mail address:* pthomas@cict.fr