## arXiv:0802.3776v2 [nlin.CD] 11 Jun 2008 [arXiv:0802.3776v2 \[nlin.CD\] 11 Jun 2008](http://arxiv.org/abs/0802.3776v2)

## Universal intermittent properties of particle trajectories in highly turbulent flows

International Collaboration for Turbulence Research, A. Arnèodo,<sup>1</sup> R. Benzi,<sup>2</sup> J. Berg,<sup>3</sup> L. Biferale<sup>\*</sup>,<sup>4</sup> E.

Bodenschatz,<sup>5</sup> A. Busse,<sup>6</sup> E. Calzavarini,<sup>7</sup> B. Castaing,<sup>1</sup> M. Cencini<sup>\*</sup>,<sup>8</sup> L. Chevillard,<sup>1</sup> R. T. Fisher,<sup>9</sup> R. Grauer,<sup>10</sup>

H. Homann,<sup>10</sup> D. Lamb,<sup>9</sup> A. S. Lanotte<sup>\*</sup>,<sup>11</sup> E. Lévèque,<sup>1</sup> B. Lüthi,<sup>12</sup> J. Mann,<sup>3</sup> N. Mordant,<sup>13</sup> W.-C. Müller,<sup>6</sup> S.

Ott,<sup>3</sup> N. T. Ouellette,<sup>14</sup> J.-F. Pinton,<sup>1</sup> S. B. Pope,<sup>15</sup> S. G. Roux,<sup>1</sup> F. Toschi<sup>\*</sup>,<sup>16,17</sup> H. Xu,<sup>5</sup> and P. K. Yeung<sup>18</sup>

<sup>1</sup>Laboratoire de Physique de l'École Normale Supérieure de Lyon, 46 allée d'Italie F-69007 Lyon, France

<sup>2</sup>*University of Tor Vergata and INFN, via della Ricerca Scientifica 1, 00133 Rome, Italy*

<sup>3</sup>*Wind Energy Department Risø National Laboratory - DTU Denmark*

<sup>4</sup>*University of Tor Vergata and INFN, Via della Ricerca Scientifica 1, 00133 Rome, Italy*

<sup>5</sup>*Max Planck Institute for Dynamics and Self-Organization, Am Fassberg 17, D-37077 Goettingen, Germany*

<sup>6</sup>*Max Planck Institute for Plasma Physics, Garching, Germany*

<sup>7</sup>*Department of Applied Physics, University of Twente, 7500 AE Enschede, The Netherlands*

<sup>8</sup> INFM-CNR, SMC Dipartimento di Fisica, Università di Roma "La Sapienza", Piazzale A. Moro, 2 I-00185 Rome, Italy

<sup>9</sup>*Department of Astronomy and Astrophysics, The University of Chicago, Chicago, IL 60637, USA*

<sup>10</sup>*Institute for Theoretical Physics I, Ruhr-University Bochum, Germany*

<sup>11</sup>*CNR-ISAC, Via Fosso del Cavaliere 100, 00133 Rome and INFN, Sezione di Lecce, 73100 Lecce, Italy*

<sup>12</sup>*IFU, ETH Z¨urich, Switzerland*

<sup>13</sup>Laboratoire de Physique Statistique de l'Ecole Normal Supérieure CNRS 24 Rue Lhomond, 75231 Paris Cedex 05, France

<sup>14</sup>*Haverford College, 370 Lancaster Avenue Haverford, PA 19041, USA*

<sup>15</sup>*Sibley School of Mechanical and Aerospace Engineering 254 Upson Hall,*

*Cornell University, Ithaca, NY 14853-7501 USA*

<sup>16</sup>*Istituto per le Applicazioni del Calcolo CNR, Viale del Policlinico 137, 00161 Roma, Italy*

<sup>17</sup>*INFN, Sezione di Ferrara, Via G. Saragat 1, 44100 Ferrara, Italy*

<sup>18</sup>*School of Aerospace Engineering, Georgia Institute of Technology, 279 Ferst Drive Atlanta, GA 30332-0150, USA*

(Dated:)

We present a collection of eight data sets, from state-of-the-art experiments and numerical simulations on turbulent velocity statistics along particle trajectories obtained in different flows with Reynolds numbers in the range  $R_{\lambda} \in [120:740]$ . Lagrangian structure functions from all data sets are found to collapse onto each other on a wide range of time lags, pointing towards the existence of a universal behaviour, within present statistical convergence, and calling for a unified theoretical description. Parisi-Frisch Multifractal theory, suitable extended to the dissipative scales and to the Lagrangian domain, is found to capture intermittency of velocity statistics over the whole three decades of temporal scales here investigated.

PACS numbers:

Understanding the statistical properties of particle tracers advected by turbulent flows is a challenging theoretical and experimental problem [\[1](#page-3-0), [2\]](#page-3-1). It is a key ingredient for the development of stochastic models [\[3](#page-3-2), [4\]](#page-3-3), in such diverse contexts as turbulent combustion, industrial mixing, pollutant dispersion and cloud formation [\[5\]](#page-3-4). The main difficulty of Lagrangian investigations, following particle trajectories, stems from the necessity to resolve the wide range of time scales driving different particle behaviours: from the longest,  $T_L$ , given by the stirring mechanism, to the shortest  $\tau_n$ , typical of viscous dissipation. Indeed the ratio,  $T_L/\tau_\eta \sim R_\lambda$ , grows with the Taylor Reynolds number,  $R_{\lambda}$ , that varies up to few thousands in laboratory flows. Some aspects of Lagrangian statistics have been experimentally measured: particle accelerations [\[2\]](#page-3-1), velocity fluctuations in the inertial range [\[6,](#page-3-5) [7](#page-3-6)] and two-particle dispersion [\[8](#page-3-7), [9\]](#page-3-8). Others, connected to the entire range of motions, have long been restricted to numerical simulations [\[10](#page-3-9), [11,](#page-3-10) [12](#page-3-11), [13,](#page-3-12) [14\]](#page-3-13). A fundamental open question is connected to intermittency, i.e. the observed strong deviations from Gaussian statistics, becoming larger and larger when considering fluctuations at smaller and smaller scales. Besides, the dependency of velocity statistics at various temporal scales on large scale forcing and boundary conditions is the so-called problem of universality. Thus, universality features are linked to the degree of anisotropy and non-homogeneities of turbulent statistics [\[15](#page-3-14)]. Similar problems have already been explored measuring the velocity fluctuations in the laboratory frame (Eulerian statistics), where clear evidence of universality have been obtained [\[16](#page-3-15)].

To build a general theory of turbulent statistics, universality is the first requirement and, if proved, may open the possibility for effective stochastic modeling [\[17](#page-3-16)] in many applied situations.

This Letter aims at investigating intermittency and universality properties of velocity temporal fluctuations by quantitatively comparing data obtained from the most advanced laboratory [\[6,](#page-3-5) [7](#page-3-6), [8\]](#page-3-7) and numerical [\[10](#page-3-9), [11,](#page-3-10) [12](#page-3-11), [13](#page-3-12), [18\]](#page-3-17) experiments. Main outcomes of our analysis are twofold. First, we show that data collapse on a common functional form, providing evidence for universality of ve-

$EXP R_\lambda $	$\tau_{\eta}$ (s)	meas. vol. $(\eta^3)$	$N_{tr}$	Tech. Ref.	
	$124 8.5\times10^{-2} $	$340^{3}$	$1.6 \times 10^6$ PTV [8]		
	690 $9 \times 10^{-4}$	$1700^3$	$6.0 \times 10^6$ PTV [7]		
	740 $2 \times 10^{-4}$	$6600^3$	$9.5 \times 10^3$ AD		

TABLE I: Experiments. By columns: 1- experiment label; 2- Taylor Reynolds number; 3- Kolmogorov time scale  $\tau_n$ ; 4measurement volume in unit of the Kolmogorov length scale  $\eta$ ; 5-  $N_{tr}$  total number of Lagrangian trajectories measured; 6- measurement technique: Particle Tracking Velocimetry (PTV) and Acoustic Doppler (AD); 7- Reference where information on the way the corresponding dataset was obtained can be found.

locity fluctuations –up to moments currently achievable with high statistical accuracy. At intermediate and inertial scales, data show an intermittent behaviour. Second, we propose a stochastic phenomenological modelisation in the entire range of scales, using a Multifractal description linking Eulerian and Lagrangian statistics.

We analyse the probability distribution of velocity fluctuations at all scales, focusing on moments of these distributions, namely the Lagrangian Velocity Structure Functions (LVSF) of positive integer order  $p$ :

$$
S_i^{(p)}(\tau) = \langle [v_i(t+\tau) - v_i(t)]^p \rangle = \langle (\delta_\tau v_i)^p \rangle, \qquad (1)
$$

where  $i = x, y, z$  are the velocity components along a single particle path, and the average is defined over the ensemble of trajectories. As stationarity and homogeneity is assumed, moments of velocity increments only depend on the time lag  $\tau$ . In the inertial range, for  $\tau_{\eta} \ll \tau \ll T_L$ , non-linear energy transfer governs the dynamics. Thus, from a dimensional viewpoint, only the scale  $\tau$  and the average energy dissipation rate for unit mass  $\epsilon$  should matter for the structure function behaviour. The only admissible choice is  $S_i^{(p)}$  $\epsilon_i^{(p)}(\tau) \sim (\epsilon \tau)^{p/2}$ , but it does not take into account the fluctuating nature of energy dissipation. Empirical studies have indeed shown that the tails of the probability density functions of  $\delta_{\tau} v$  become increasingly non-Gaussian at decreasing  $\tau/T_L$ . In terms of moments of the velocity fluctuations, intermittency reveals itself in the anomalous scaling exponents, i.e. a breakdown of the dimensional law for which we have that

<span id="page-1-0"></span>
$$
S_i^{(p)}(\tau) \sim \tau^{\xi(p)}\,,\tag{2}
$$

with  $\xi(p) \neq p/2$ . Notice that when dissipative effects dominate, typically for scales  $\tau \sim \tau_{\eta}$  and smaller, the power-law behaviour [\(2\)](#page-1-0) is no longer valid, and refined arguments have to be employed, as we will see in the following.

The statistics of velocity fluctuations at varying time lag  $\tau$  can be quantitatively captured by the logarithmic derivatives of  $S_i^{(p)}(\tau)$  versus  $S_i^{(2)}(\tau)$  [\[19](#page-3-18), [20,](#page-3-19) [21](#page-3-20)]. This

$DNS R_\lambda $	$N^3$	$N_{tr}$		Diss. Tech. Ref.	
	$140 \mid 256^3$	$5 \times 10^5$			[11]
	$320 1024^3 $	$5 \times 10^6$	N		$\left\lceil 13 \right\rceil$
3		$400\,2048^3\, \,3\times10^5$			[10]
		$600 1856^3 1.6\times10^7 $	C		(18)
$\mathbf{5}$		650 $ 2048^3 $ 4 × 10 <sup>5</sup>			

TABLE II: Direct numerical simulations. By columns: 1 numerical simulation label; 2- Taylor Reynolds number  $R_{\lambda}$ ; 3- number of collocation points  $N^3$ ; 4- total number of Lagrangian tracers  $N_p$ ; 5- characteristic of dissipation: normal viscous terms (N), weakly compressible code (C); 6- interpolation technique for Lagrangian integration: linear interpolation (L), tricubic interpolation (T); cubic-splines (CS); 7- Reference where information on the way the corresponding dataset was obtained can be found.

defines the local scaling exponents

$$
\zeta_i(p,\tau) = \frac{\mathrm{d}\log S_i^{(p)}(\tau)}{\mathrm{d}\log S_i^{(2)}(\tau)}.
$$
\n(3)

For statistically isotropic turbulence, all components are equivalent, so that their spread quantifies the degree of anisotropy present in teh flow. The  $\tau$ -dependence of  $\zeta_i(p,\tau)$  allows for a scale-by-scale characterisation of intermittency.

Figure 1 shows the local exponents of order  $p = 4$  from a collection of eight data sets, see Table I and II, for different Reynolds numbers. Most of these data sets are new, as well as completely new is the performed analysis, here presented. We focused on the fourth order moment, since it is the highest order achievable with statistical convergence for all data sets. Two observations can be done. First, all data sets show a similar strong variation around the dissipative time  $\tau/\tau_n \sim O(1)$  that depends on the Reynolds number, and then a clear tendency toward a plateau for larger lags  $\tau > 10\tau_{\eta}$ . Second, all data sets, with comparable Reynolds numbers, well agree in the whole range of time lags. The relative scatter increases only for large  $\tau$ , due to the combined effects of the lack of statistics, the anisotropy of the flows and the different valus of  $R_{\lambda}$ . In particular, finite volume effects in experimental particle tracking can produce a small – but systematic – downward shift of the points at long-lag times [\[21](#page-3-20), [22\]](#page-3-21). It is worth noticing that error bars estimated from anisotropic contributions decrease by going to small  $\tau$ , indicating that isotropy tends to be recovered at sufficiently small scales, i.e. large scale anisotropic contributions becomes less and less important. In addition, the fact that, at comparable Reynolds numbers, all data sets recover the same behaviour by going to smaller and smaller time lags provides a clear indication of Lagrangian universality of the energy cascade. Such an agreement has not been observed before and is comparable with that found for the corresponding Eulerian quantities [\[16\]](#page-3-15).



FIG. 1: (Color online) Log-Lin plot of the fourth order local exponent,  $\zeta(4,\tau)$ , averaged over velocity components, as a function of the normalised time lag  $\tau/\tau_n$ . Data sets come from three experiments (EXP) (see Table 1) and five direct numerical simulations (DNS) (see Table 2). Error bars are estimated from the spread between the three components, except in EXP3 where only two components were measured. Each data set is plotted only in the time range where recognised experimental/numerical limitations are not affecting the results. In particular, for each data set, the largest time lag always satisfies  $\tau < T_L$ . The minimal time lag is set by the highest fully resolved frequency. The shaded area displays the prediction obtained by the MF model by using  $D_{Lo}(h)$  or  $D_{Tr}(h)$ , with  $\beta = 4$ , for a range of  $R_{\lambda} \in [150 : 800]$ , comparable with the range of  $R_{\lambda}$  in the data. Notice that the MF predictions have been obtained by fixing equal to 7, the multiplicative constant in the definition of  $\tau_n$ . The straight dashed line corresponds to the dimensional non-intermittent value  $\zeta(4, \tau) = 2$ , achieved at small time lags where structure functions do become differentiable. Notice that two among the DNS are sufficiently resolved to get the mentioned dimensional scaling in the high frequency limit.

The quality of data shown in (Fig. 1) opens the possibility to quantitatively test phenomenological models for LVSF, scale-by-scale. Parisi-Frisch Multifractal (MF) model of the inertial range [\[23\]](#page-3-22), and its generalization to the dissipative range [\[24](#page-4-0), [25](#page-4-1), [26,](#page-4-2) [27\]](#page-4-3), has proved to give a satisfactory description of Eulerian and Lagrangian fluctuations [\[14,](#page-3-13) [28,](#page-4-4) [29](#page-4-5), [30\]](#page-4-6). It is thus appealing to search for a link between Eulerian and Lagrangian statistics [\[14,](#page-3-13) [28](#page-4-4), [29](#page-4-5), [30\]](#page-4-6), since this points to a unique interpretation of turbulent fluctuations. Moreover, it would reduce the number of free parameters. According to the MF model, Eulerian velocity increments at inertial scales are characterised by a local Hölder exponent  $h$ , i.e.  $\delta_r u \sim r^h$ , whose probability  $\mathcal{P}(h) \sim r^{3-D(h)}$  is weighted by the Eulerian fractal dimension  $D(h)$  of the set where h is observed [\[23\]](#page-3-22). The dimensional relation  $\tau \sim r/\delta_r u$ bridges Lagrangian fluctuations over a time lag  $\tau$  to the Eulerian ones at scale  $r$ . Following Refs. [\[27,](#page-4-3) [28\]](#page-4-4), it is shown in Ref. [\[30\]](#page-4-6) how to extend the MF framework to get a unified description at all time scales for Lagrangian turbulence. Accordingly, Lagrangian increments display a continuous and differentiable behaviour at the transition from the dissipative to the inertial range,

<span id="page-2-0"></span>
$$
\delta_{\tau}v(h) = V_0 \frac{\tau}{T_L} \left[ \left(\frac{\tau}{T_L}\right)^{\beta} + \left(\frac{\tau_{\eta}}{T_L}\right)^{\beta} \right]^{\frac{2h-1}{\beta(1-h)}}, \quad (4)
$$

 $\beta$  being a free parameter controlling the crossover around  $\tau \sim \tau_n$ , and  $V_0$  the root mean square velocity. In order to get a prediction for the behaviour of the LVSF, given by

<span id="page-2-1"></span>
$$
\langle (\delta_\tau v)^p \rangle \sim \int dh P_h(\tau, \tau_\eta) [\delta_\tau v(h)]^p , \qquad (5)
$$

we have to consider, in [\(4\)](#page-2-0), the intermittent fluctua-tions of the dissipative scale [\[14,](#page-3-13) [28,](#page-4-4) [30](#page-4-6)],  $\tau_{\eta}(h)/T_L \sim$  $R_{\lambda}^{2(h-1)/(1+h)}$  $\lambda^{2(n-1)/(1+n)}$ . The last necessary ingredient is to specify the probability of observing fluctuations of  $h$ . This is done in analogy to Eq. [\(4\)](#page-2-0):

$$
P_h(\tau, \tau_\eta) = \mathcal{Z}^{-1}(\tau) \left[ \left( \frac{\tau}{T_L} \right)^\beta + \left( \frac{\tau_\eta}{T_L} \right)^\beta \right]^{\frac{3 - D(h)}{\beta (1 - h)}}, \quad (6)
$$

where  $\mathcal Z$  is a normalizing function [\[30](#page-4-6)] and  $D(h)$  the fractal dimension of the support of the exponents  $h$ . Once specified the Reynolds number, we are left with two parameters - the expoent  $\beta$  and a multiplicative constant in the definition of  $\tau_n$  -, while the function  $D(h)$  comes from the knowledge of the Eulerian statistics.

Eulerian Velocity Structure Functions (EVSF) have been measured in the last two decades (see Ref.[\[16\]](#page-3-15) for a data collection) providing a way to estimate the function  $D(h)$ based on empirical data. Many functional forms have been proposed in the literature [\[23](#page-3-22)] that are consistent with data, up to statistical uncertainties. Eulerian velocity statistics can be measured in terms of longitudinal or transverse fluctuations. Fluid velocity along particle paths is naturally sensitive to both kinds of fluctuations. We thus evaluated the LVSF in [\(5\)](#page-2-1) using the fractal dimensions  $D_{Lo}(h)$  and  $D_{Tr}(h)$  obtained by longitudinal [\[16](#page-3-15)] and transverse [\[31\]](#page-4-7) moments of Eulerian luctuations, respectively.

The shaded area in (Fig. 1) represents the range of variation of the MF prediction computed from  $D_{Lo}(h)$ or  $D_{Tr}(h)$ , measured in the Eulerian statistics (see below), and at changing Reynolds numbers. This must be interpreted as our uncertainty. The prediction works very well: all data fall within the shaded area. The role of the parameters is clear. Changing  $\beta$  modifies the sharpness and shape of the dip region at  $\tau_{\eta}$  – the larger  $\beta$  the more pronounced the dip; while, changing the multiplicative constant in the definition of  $\tau_{\eta}$  has no effect on the curve shape, but it rigidly shifts the whole curve along the time axis.

Increasing the Reynolds number  $R_{\lambda}$ , the flat region at large lags develops a longer plateau. In the limit  $R_{\lambda} \to \infty$ the MF model predicts  $\zeta(4) \simeq 1.71$  from  $D_{Lo}(h)$  and  $\zeta(4) \simeq 1.59$  from  $D_{Tr}(h)$  statistics.

For the Eulerian  $D(h)$ , we used the following log-Poisson [\[23\]](#page-3-22) functional form,

$$
D(h) = \frac{3(h - h^{*})}{\log(\gamma)} \left[ \log \left( \frac{3(h^{*} - h)}{d^{*} \log(\gamma)} \right) - 1 \right] + 3 - d^{*}.
$$
 (7)

Different couples of parameters,  $(h^*, \gamma)$ , have been chosen to fit longitudinal and transverse Eulerian fluctuations. The parameter  $d^* = (1-3h^*)/(1-\gamma)$  is fixed by imposing the exact relation for third order EVSF. For the longitu-dinal exponents [\[16\]](#page-3-15), we used  $(h_{Lo}^* = 1/9, \gamma_{Lo} = 2/3)$ [\[23\]](#page-3-22). For the transverse exponents, we used  $(h_{Tr}^*)$  $1/9, \gamma_{Tr} = 1/2$ ) which fits the data in Ref. [\[31](#page-4-7)] (see [\[32](#page-4-8)] for details.)

This comprehensive comparison of the best available experiments and direct numerical simulations provides strong evidence of the universality of Lagrangian statistics. One important open question is the effect of a mean flow, as in turbulent jets [\[33\]](#page-4-9) and wall bounded turbulence, where strong persistence of anisotropy may break the recovery of small-scale universality. We showed that a Multifractal description is in good agreement with data, even in the dissipative range where intermittency is significantly increased. The Multifractal description captures the intermittency at all scales with only a few parameters, independent of the Reynolds number. This is the universal feature of Lagrangian turbulence revealed by this study. There exists a long debate on the statistical importance of vortex filaments around dissipative time and length scales [\[23,](#page-3-22) [34](#page-4-10)]. Simulations [\[10,](#page-3-9) [20,](#page-3-19) [35\]](#page-4-11) show that the dip region for  $\tau \sim \tau_{\eta}$  can be depleted/enhanced by decreasing/increasing the probability of particles being trapped in vortex filaments. The Multifractal model is able to capture the intermittency around  $\tau_n$  with the help of the free parameter  $\beta$ . Different values of  $\beta$  should then correspond to different statistical weights of vortex filaments along particle trajectories.

Only further advances in both experimental techniques and numerical power will allow us to test the same questions here addressed also for the higher order statistics.

Helpful discussions with U. Frisch and L. P. Kadanoff are gratefully acknowledged. F.T. thanks the DEISA Consortium (co-funded by the EU), for support within the DEISA Extreme Computing Initiative; L.B., M.C., A.S.L. and F.T. thank CINECA (Bologna, Italy) for technical support.

∗ ictr.rome@roma2.infn.it

- <span id="page-3-0"></span>[1] G. Falkovich and K. R. Sreenivasan, Phys. Today 59, 43 (2006).
- <span id="page-3-1"></span>[2] A. La Porta et al., Nature 409, 1017 (2001).
- <span id="page-3-2"></span>[3] S. B. Pope, *Turbulent Flows* (Cambridge Univ. Press, Cambridge, UK, 2000).
- <span id="page-3-3"></span>[4] A. G. Lamorgese, S. B. Pope, P.K. Yeung and B. L. Sawford, J. Fluid Mech. 582, 423 (2007).
- <span id="page-3-4"></span>[5] R. A. Shaw, Ann. Rev. Fluid Mech. 35, 183 (2003).
- <span id="page-3-5"></span>[6] N. Mordant, P. Metz, O. Michel and J.-F. Pinton, Phys. Rev. Lett. 87, 214501 (2001).
- <span id="page-3-6"></span>[7] H. Xu et al., Phys. Rev. Lett. 96, 024503 (2006).
- <span id="page-3-7"></span>[8] J. Berg et al., Phys. Rev. E **74**, 016304 (2006).
- <span id="page-3-8"></span>[9] M. Bourgoin et al., Science 311, 835-838 (2006).
- <span id="page-3-9"></span>[10] L. Biferale et al., Phys. Fluids **17**, 021701 (2005).
- <span id="page-3-10"></span>[11] N. Mordant, E. Lévèque and J.-F. Pinton, New J. Phys. 6, 34 (2004).
- <span id="page-3-11"></span>[12] P. K. Yeung, S.B. Pope and B. L. Sawford, J. Turb. 7 (58), 1 (2006).
- <span id="page-3-12"></span>[13] H. Homann et al., J. Plasma Physics 73, 821 (2007).
- <span id="page-3-13"></span>[14] L. Biferale et al., Phys. Rev. Lett. **93**, 064502 (2004).
- <span id="page-3-14"></span>[15] L. Biferale and I. Procaccia. Phys Rep. 414, 43 (2005).
- <span id="page-3-15"></span>[16] A. Arnèodo et al., Europhys. Lett. **34**, 411 (1996).
- <span id="page-3-16"></span>[17] B. L. Sawford, Phys. Fluids A 3, 1577 (1991).
- <span id="page-3-17"></span>[18] R. T. Fisher et al., IBM Journ. Res. Devel. 52 Num. (1/2), 127 (2008).
- <span id="page-3-18"></span>[19] R. Benzi et al., Phys. Rev. E 48, R29 (1993).
- <span id="page-3-19"></span>[20] I. Mazzitelli and D. Lohse, New J. Phys. 6, 203 (2004).
- <span id="page-3-20"></span>[21] L. Biferale et al., Phys. Fluids 20, 065103 (2008).
- <span id="page-3-21"></span>[22] S. Ott and J. Mann, New J. Phys. **7**, 142 (2005).
- <span id="page-3-22"></span>[23] U. Frisch, *Turbulence: the legacy of A.N. Kolmogorov*

(Cambridge University Press, Cambridge UK, 1995).

- <span id="page-4-0"></span>[24] G. Paladin and A. Vulpiani, Phys. Rev. A 35, R1971 (1987).
- <span id="page-4-1"></span>[25] M. Nelkin, Phys. Rev. A **42**, 7226 (1990).
- <span id="page-4-2"></span>[26] U. Frisch and M. Vergassola, Europhys. Lett. 14, 439 (1991).
- <span id="page-4-3"></span>[27] C. Meneveau, Phys. Rev. E **54**, 3657 (1996).
- <span id="page-4-4"></span>[28] M. S. Borgas, Phil. Trans. R. Soc. London A 342, 379 (1993).
- <span id="page-4-5"></span>[29] G. Boffetta, F. De Lillo and S. Musacchio, Phys. Rev. E 66, 066307 (2002).
- <span id="page-4-6"></span>[30] L. Chevillard et al., Phys. Rev. Lett. 91, 214502 (2003).
- <span id="page-4-7"></span>[31] T. Gotoh, D. Fukayama and T. Nakano, Phys. Fluids 14, 1065 (2002).
- <span id="page-4-8"></span>[32] The functions  $D_{Lo}(h)$  and  $D_{Tr}(h)$  are convex; the strongest fluctuations, corresponding to the smallest

Hölder exponent h, are realised for  $h = h_{Lo}^*$  and  $h_{Tr}^*$ , respectively. The maxima of fractal dimensions,  $D_{Lo}(h_{Lo}^{max}) = D_{Tr}(h_{Tr}^{max}) = 3$ , are attained for  $h_{Lo}^{max} =$ 0.382, and  $h_{Tr}^{max} = 0.57$ . Since only the range  $h^* \leq h \leq$  $h^{max}$  is relevant to the fit of EVSF of positive order [\[23\]](#page-3-22), the extremes of integration in Eq. [\(5\)](#page-2-1) have been set in the interval,  $[h_{Lo}^*, h_{Lo}^{max}]$  and  $[h_{Tr}^*, h_{Tr}^{max}]$ , respectively. Modifying  $h^{max}$  alters the MF prediction for  $\tau \sim T_L$ , but does not change the behaviour of the curves in the inertial range.

- <span id="page-4-9"></span>[33] P. Gervais et al., Exp. Fluids 42, 371 (2007).
- <span id="page-4-10"></span>[34] S. Douady, Y. Couder, and M.E. Brachet, Phys. Rev. Lett. 67, 983 (1991).
- <span id="page-4-11"></span>[35] J. Bec et al., Phys. Fluids **18**, 081702 (2006).