Estimation of Gaussian mixtures in small sample studies using l_1 penalization

Stéphane Chrétien *

October 9, 2014

Abstract

Many experiments in medicine and ecology can be conveniently modeled by finite Gaussian mixtures but face the problem of dealing with small data sets. We propose a robust version of the estimator based on self-regression and sparsity promoting penalization in order to estimate the components of Gaussian mixtures in such contexts. A space alternating version of the penalized EM algorithm is obtained and we prove that its cluster points satisfy the Karush-Kuhn-Tucker conditions. Monte Carlo experiments are presented in order to compare the results obtained by our method and by standard maximum likelihood estimation. In particular, our estimator is seen to perform better than the maximum likelihood estimator.

Keywords: finite Gaussian mixtures, maximum likelihood estimation, Kullback Proximal Point algorithms, EM algorithm, l_1 penalization, LASSO, sparsity, regression mixtures, model based clustering

1 Introduction

Finite Gaussian mixture models are widely used in a great number of application fields as a means to perform model based classification. From pattern recognition to biology, from quality control to finance, many examples have shown the pertinence of the Gaussian mixture model approach. The book [13] is the most comprehensive reference for finite non necessarily Gaussian mixture models with many application examples. In Gaussian mixture models, the data Y_1, \ldots, Y_n are assumed i.i.d. and to be drawn from the density

$$\sum_{k=1}^{K} p_k^* f^{(d)}(y; \mu_k, \Sigma_k)$$
(1)

where

$$f^{(d)}(y;\mu,\Sigma) = \frac{1}{\sqrt{(2\pi)^d \det(\Sigma^*)}} \exp\left(-\frac{1}{2}(y-\mu^*)^t \Sigma^{*-1}(y-\mu^*)\right)$$
(2)

*Mathematics Department, Université de Franche Comté and UMR CNRS-6623, 16 route de Gray, 25030 Besançon, France.

and where the vector $\theta^* = (p_1^*, \ldots, p_K^*, \mu_1^*, \ldots, \mu_K^*, \Sigma_1^*, \ldots, \Sigma_k^*)$ is an unknown multidimensional parameter. To this model, we traditionally associate an extended model using the notion of complete data. In mixture models, the complete data are independent and identically distributed couples of the form (Y_i, Z_i) where Z_i is a multinomial random variable taking values in $\{1, \ldots, K\}$ with $P(Z_i = k) = p_k^*$ and which represents the index of the mixture component from which observation *i* was drawn. We assume that conditionally on the event $Z_i = k, Y_i$ has density $\frac{1}{\sqrt{(2\pi)^d \det(\Sigma_k^*)}} \exp\left(-\frac{1}{2}(y-\mu_k^*)^t \Sigma_k^{*-1}(y-\mu_k^*)\right)$. The variables Z_1, \ldots, Z_n being unobserved, they are usually called latent variables.

The standard approach for estimating θ^* is the maximum likelihood methodology which consists of finding $\hat{\theta}$ which maximizes the log-likelihood function

$$l(\theta) = \sum_{i=1}^{n} \log\left(\sum_{k=1}^{K} p_k f^{(d)}(y; \mu_k, \Sigma_k)\right)$$
(3)

over the set

$$\Theta = \left\{ (p_1, \dots, p_K, \mu_1, \dots, \mu_K, \Sigma_1, \dots, \Sigma_K) \mid p_k \in \mathbb{R}_+, \ \mu_k \in \mathbb{R}^d, \ \Sigma_k \in \mathbb{S}_d^+, \right.$$

and
$$\sum_{k=1}^K p_k = 1 \right\}$$

where \mathbb{S}_d^+ denotes the set of all symmetric positive semidefinite matrices and \mathbb{R}_+ is the set of nonnegative real numbers.

Interestingly enough, the supremum of the log-likelihood function over Θ is equal to $+\infty$ and is obtained for singular covariance matrices. A study of the one dimensional case was made in [2]. However, many researchers and practitioners have noticed that some local maximizer of the log-likelihood function is in fact consistent in practice. From the numerical viewpoint, local maximizers of the log-likelihood function are usually obtained using the Expectation-Maximization (EM) algorithm of Dempster Laird and Rudin [8]. This algorithm is a nice procedure with closed form expression of each iteration in the Gaussian mixture case. The EM algorithm for mixture models is available in the MIXMOD package [1] for instance.

Beside the question of finding the right local optimizer of the likelihood function, one of the main problems about estimating θ^* is the one of having a sufficiently large sample size. Instances where the sample size is large enough can be found in a number of applications such as pattern recognition or financial time series analysis. On the other hand, in many other fields, e.g. ecology, the sample size may be very small in situations where finite mixture models are suspected to be very pertinent due to the biological context. The goal of this paper is to remedy this problem by proposing a new methodology for Gaussian mixture model estimation in the case where the sample size is extremely small. Our approach aims at providing a certain amount of robustness. In the same spirit as for the median in the one dimensional case, the main idea is to express the estimators of the μ_k 's as a combination of a small number of data in the middle of each cluster. This is simply done by restricting the search to the data's span, i.e. to obtain the μ_k 's as a regression with covariates the data themselves and to impose an additional sparsity constraint on the regression vectors. In order to simplify the analysis, we will assume the covariance matrices to be of the form $\sigma_k^2 I$, where I denotes the identity matrix. The σ_k 's and the p_k 's can also be estimated using for instance a maximum likelihood approach conditioned on the estimated value of the μ_k 's.

The whole procedure is formally equivalent to joint variable selection and estimation in a mixture of regression model. Variable selection and estimation are performed using l_1 -penalized EM steps which reduce the complexity of the regression model just as for the LASSO [16]. Encouraging simulations results show that the proposed approach correctly estimates the class of 8 over 10 points on average for a mixture of 3 Gaussians in dimension two. Monte Carlo experiments are performed for samples sizes of 10 points and dimension growing up to to 50 showing a good behavior of the method which outperforms the standard maximum likelihood estimator.

2 Presentation of the method

2.1 Recalls on regression mixtures

The Gaussian regression mixture assumes that the observations are couples of the form (Y, X)where Y takes values in \mathbb{R}^d , X takes values in \mathbb{R}^p , and conditionally on X, the random variable Y follows the mixture density

$$f_{Y|X}(y) = \sum_{k=1}^{K} p_k f^{(d)}(y; M_k X, \Sigma_k),$$
(4)

where M is a matrix in $\mathbb{R}^{d \times p}$.

Such mixture models are frequent in econometrics and chemometrics as described in the introduction of [11]. Estimation in these models can be performed using likelihood maximization as in [13] using the EM algorithm or a Bayesian methodology as studied in [11] using Monte Carlo Markov Chain techniques.

One way to perform model selection in such a model is to use a non-differentiable penalty such as the ℓ_1 -norm, i.e. the sum of the absolute values of the regression coefficients.

2.2 Our proposal: The mixture of self-regression with sparsity constraint

In the present paper, we only intend to perform unsupervised clustering and thus, our setting seems far from the mixture of regression framework. The originality of the proposed approach is to introduce an artificial mixture of regressions for the simpler problem of clustering. The main idea is as follows: instead of estimating the $\mu'_k s$ for $k = 1, \ldots, K$, it should be easier to estimate only the coefficients of a sparse linear combinations of the $X'_i s$ for all the datas belonging to the same cluster. This strategy should give even better results as the dimension d of the problem increases if the sparsity of the involved linear combinations stays constant and small. Let us formalize our method in the next subsection.

2.2.1 The estimator

In all what follows, we will assume that the data have been centered. Our proposal relies on the following simple idea: if the cluster proportions p_k^* 's, the class indices Z_i , i = 1, ..., n and

the variances Σ_k^* , k = 1, ..., K where known ahead of time, the estimators of the μ_k 's could be chosen, in the small sample setting, as linear combinations of the datas themselves.

A simple example of such an idea is based on the notion of medoid. In clustering, the medoids play the role of the centers for each cluster, but are selected among the data themself. In what follows, instead of choosing only one medoid, we propose to select a linear combination of the data for each cluster. In order to stay robust as the dimension the space grows, we may impose that the linear combination be sparse, e.g. only onvolves 3 or 4 datas or less for each cluster.

The main difficulty with such an approach is that choosing the right sample vectors to represent each cluster seems a priori a very hard task. Fortunately, one might rely on the recent discoveries concerning variable selection in order to overcome this problem: just as in the LASSO, a simple idea may be to use a regression formalism for estimating each μ_k , $k = 1, \ldots, K$, using a sparsity enforcing penalty like e.g. the ℓ_1 norm of the coefficients.

In the more general case where the indices Z_i , i = 1, ..., n are unobserved, and the cluster proportions p_k^* and the covariance matrices Σ_k^* , k = 1, ..., K are unknown, one can consider maximizing the l_1 -penalized log-likelihood like function given by

$$\sum_{i=1}^{n} \log \left(\sum_{k=1}^{K} p_k f^{(d)}(Y_i; \mu_k, \Sigma_k) \right) - \lambda \sum_{k=1}^{K} \|\beta_k\|_1$$
(5)

under the data-driven constraints $\mu_k = Y\beta_k$ for k = 1, ..., n where the matrix Y is given by $Y = [Y_1, ..., Y_n]$. The parameter λ is called the relaxation parameter. In other words, we would like to maximize the l_1 -penalized likelihood function

$$\tilde{l}_{pen}(\theta) = \tilde{l}(\theta) - \lambda \sum_{k=1}^{K} \|\beta_k\|_1,$$
(6)

where

$$\tilde{l}(\theta) = \sum_{i=1}^{n} \log \left(\sum_{k=1}^{K} p_k f^{(d)}(Y_i; Y\beta_k, \Sigma_k) \right).$$
(7)

2.2.2 The Space-Alternating l_1 -EM algorithm

Optimizing the l_1 -penalized function (5) can be performed using an EM-type algorithm. The Expectation Step consists of computing the conditional expectation of the complete l_1 -penalized likelihood like function given the observations Y_1, \ldots, Y_n where the distribution of the latent variables is taken to be their marginal density parametrized by the approximation $\bar{\theta}$ of the true parameter θ^* . The resulting quantity is traditionally denoted by $Q(\theta, \bar{\theta})$ and we will use the same notation in our l_1 -penalized context.

More precisely, the complete l_1 -penalized log-likelihood like function $\tilde{l}_{pen}^c(\theta)$, i.e. the penalized log-likelihood like function of the complete data $(Y_1, Z_1), \ldots, (Y_n, Z_n)$ is given by

$$\tilde{l}_{pen}^c(\theta) = \sum_{i=1}^n \log\left(p_{Z_i} f^{(d)}(Y_i; Y\beta_k, \Sigma_k)\right) - \lambda \sum_{k=1}^K \|\beta_k\|_1.$$
(8)

Thus, we obtain

$$Q(\theta,\bar{\theta}) = \sum_{i=1}^{n} \sum_{k=1}^{K} \log\left(p_k f^{(d)}(Y_i; Y\beta_k, \Sigma_k)\right) \tau_{i,k} - \lambda \sum_{k=1}^{K} \|\beta_k\|_1$$
(9)

where we used the standard notation $\tau_{i,k} = P_{\bar{\theta}}(Z_i = k \mid Y_1, \dots, Y_n).$

The Maximization Step consists of maximizing $Q(\theta, \theta)$. In order to simplify the practical implementation, the p_k 's, β_k 's and Σ_k 's can be optimized alternatively in the manner of the Gauss-Seidel approach. In fact, the separability of the problem into two subproblems, the first being optimization over the p_k 's and the second being optimization over the β_k 's and Σ_k 's is already well known and the solution to the first of these subproblems is of the form

$$p_k = \frac{\sum_{i=1}^n \tau_{i,k}}{\sum_{i=1}^n \sum_{k=1}^K \tau_{i,k}}.$$
(10)

On the other hand, joint optimization in β_k 's and the Σ_k 's is not separable and space alternating option can be helpful in order to keep the computational complexity of each step at a low level. In order to address this problem, we need a generalization of the EM algorithm allowing for componentwise optimization at each step. Such penalized EM algorithms have been recently studied in the broader framework of Space Alternating Kullback Proximal Point Algorithms in [7]. Optimizing successively over the β_k 's at one iteration and over the Σ_k 's at the next iteration should be reasonably efficient in most applications. Here, we will also optimize one cluster at a time in order to obtain the injectivity conditions which are needed in the theoretical analysis of the algorithm. A simple way to accelerate the proposed version of the Gauss-Seidel methodology could be to average the new iterates $\beta^{(l)}$ and $\Sigma^{(l)}$ with the previous respective iterates so that to smooth the algorithm's trajectory.

In what follows, we will restrict the analysis to the case where the covariance matrices are multiple of the identity but the method can easily be implemented and studied with general covariance matrices. The details of the method are summarized in Algorithm 1 below. The convergence analysis is provided in the Appendix (Section 5).

3 Simulation results

In this section, we address the question of testing the algorithm on simulated datasets. The Space Alternating l_1 -EM was first tested on simulated data sets. The experiments were built as follows: 10 samples in \mathbb{R}^2 were generated from three different Gaussian distributions with the objective to recover the index of the distribution they were drawn from up to some index permutation. The class probabilities were taken as $p_1 = .3$, $p_2 = .2$ and $p_3 = .5$ and the variances as $\sigma_1^2 = 5$, $\sigma_2^2 = 7$ and $\sigma_3^2 = 10$ without change through all the simulation experiments. Various experiments were performed using different values for the expectation vectors μ_1 , μ_2 and μ_3 since it could be easily suspected that the distance between them would play a major role in the class index recovery problem. The results presented below were obtained using the following Monte Carlo scheme: the expectations were isotropic dilations of three points in \mathbb{R}^2 drawn uniformly at random in the cube $[-\frac{1}{2}, \frac{1}{2}]^3$. We ran the code for dilation factors d going from 10 to 100 by steps of 10.

Algorithm 1 Space-Alternating l_1 -EM algorithm

Input $L \in \mathbb{N}_*$ Choose initial iterate $\theta^{(0)} = (p_1^{(0)}, \dots, p_k^{(0)}, \beta_1^{(0)}, \dots, \beta_K^{(0)}, \sigma_1^{(0)}, \dots, \sigma_K^{(0)})$ l = 1while $l \leq L$ do (E-Step) Compute the conditional probabilities $P_{\theta^{(l-1)}}(Z_i = k \mid Y)$ given the observations

(E-Step) Compute the conditional probabilities $P_{\theta^{(l-1)}}(Z_i = k \mid Y)$ given the observations Y_1, \ldots, Y_n for $i = 1, \ldots, n$ and $k = 1, \ldots, K$ using the following formula

$$\tau_{i,k}^{(l)} = \frac{p_k^{(l-1)} f^{(d)} \left(Y_i; \mu_k^{(l-1)}, \sigma_k^{(l-1)} I\right)}{\sum_{k=1}^K p_k^{(l-1)} f^{(d)} \left(Y_i; \mu_k^{(l-1)}, \sigma_k^{(l-1)} I\right)}$$
(11)

compute

-either the $p_k^{(l)}$'s by the formula

$$p_k^{(l)} = \frac{\sum_{i=1}^n \tau_{i,k}^{(l)}}{\sum_{i=1}^n \sum_{k=1}^K \tau_{i,k}^{(l)}}$$
(12)

 $-\mathbf{or}~\beta_k^{(l)}$ as the solution of the LASSO-like optimization problem

$$\beta_{k}^{(l)} \in \operatorname{argmin}_{b \in \mathbb{R}^{n}} \| \left(\sum_{i=1}^{n} Y_{i} \tau_{i,k}^{(l)} \right) - Yb \|_{2}^{2} - \lambda \| b \|_{1}.$$
(13)

for the index k updated in cyclic order along iterations. –or $\sigma_k^{(l)}$ using the formula

$$\sigma_k^{(l)} = \frac{1}{\sum_{i=1}^k \tau_{i,k}^{(l)}} \sum_{i=1}^n \|Y_i - Y\beta_k^{(l-1)}\|_2^2 \tau_{i,k}^{(l)}.$$
 (14)

for one index k updated in cyclic order along iterations. cyclically end while Output $p_k^{(L)}$, $\beta_k^{(L)}$ and $\sigma_k^{(L)}$ for k = 1, ..., K.

3.1 Two dimensional data

An example of the type of result we obtained is given in Figure 1 below where the 10 points were correctly classified.



Figure 1: A result obtained with the LASSO (or space alternating l_1)-EM for centers drawn uniformly inside the cube $[-30, 30]^3$.

Here is another example when the expectation vectors are chosen closer to each other and 8 points over 10 were correctly classified.

The average number of correctly recovered class indices (ANCRCI) for 1000 Monte Carlo experiments in the case of 10 points as a function of the box into which the expectation vectors have been uniformly drawn are given in Table 2 below.

Initial cube	$[-5,5]^2$	$[-10, 10]^2$	$[-15, 15]^2$	$[-20, 20]^2$	$[-25, 25]^2$
ANCRCI	6.11	6.98	7.49	7.89	8.23
Initial cube	$[-30, 30]^2$	$[-35, 35]^2$	$[-40, 40]^2$	$[-45, 45]^2$	$[-50, 50]^2$
ANCRCI	8.41	8.38	8.61	8.71	8.74

Table 1: The average number of correctly recovered class indices (ANCRCI) over the 1000 Monte Carlo experiments similar to the one shown in Figure 1 is given for increasing sizes of the initial cubes where the expectation vectors are chosen uniformy at random.



Figure 2: A result obtained with the space alternating l_1 -EM for centers drawn uniformly in the cube $[-15, 15]^3$.

3.2 Higher dimensional data

We performed Monte Carlo experiments in dimensions 5, 10 and 15. The results are presented in Table 2. In order to compare with the standard likelihood approach for finite Gaussian mixtures, we gathered the results obtained for the same experiments in Table 3⁻¹. Table 4 shows the results obtained using the Classification EM (CEM) algorithm.

	$[-20, 20]^d$	$[-25, 25]^d$	$[-30, 30]^d$	$[-35, 35]^d$	$[-40, 40]^d$	$[-45, 45]^d$	$[-50, 50]^d$
d = 5	8.18	8.37	8.54	8.6	8.69	8.73	8.63
d = 10	7.83	8.17	8.42	8.43	8.6	8.47	8.6
d = 15	7.58	8.07	8.15	8.25	8.35	8.34	8.42

Table 2: The average number of correctly recovered class indices (ANCRCI) using our robust estimator over the 1000 Monte Carlo experiments shown in Figuremc is given for increasing sizes of the initial cubes where the expectation vectors are chosen uniformy at random and for increasing dimension of the sample space.

	$[-20, 20]^d$	$[-25, 25]^d$	$[-30, 30]^d$	$[-35, 35]^d$	$[-40, 40]^d$	$[-45, 45]^d$	$[-50, 50]^d$
d = 5	6.75	6.86	6.9	7.06	7.11	7.04	7.07
d = 10	6.64	6.82	6.88	6.81	6.9	6.98	7.01
d = 15	6.56	6.65	6.93	6.85	6.8	6.9	7.01

Table 3: The average number of correctly recovered class indices (ANCRCI) using the standard maximum likelihood estimator (provided by the standard EM algorithm) over the 1000 Monte Carlo experiments is given for increasing sizes of the initial cubes where the expectation vectors are chosen uniformy at random and for increasing dimension of the sample space.

	$[-20, 20]^d$	$[-25, 25]^d$	$[-30, 30]^d$	$[-35, 35]^d$	$[-40, 40]^d$	$[-45, 45]^d$	$[-50, 50]^d$
d = 5	8.055	8.27	8.36	8.37	8.33	8.42	8.41
d = 10	8.14	8.15	8.22	8.30	8.17	8.29	8.22
d = 15	8.12	8.14	8.32	8.22	8.21	8.29	8.33

Table 4: The average number of correctly recovered class indices (ANCRCI) for the output of the Classification EM (CEM) algorithm, over the 1000 Monte Carlo experiments is given for increasing sizes of the initial cubes where the expectation vectors are chosen uniformy at random and for increasing dimension of the sample space.

 $^{^1\}mathrm{We}$ used the EM algorithm for Gaussian mixtures with covariance matrices equal to multiples of the identity matrix

As Table 2 shows, the class index recovery rate is still quite good in dimension 15 for well separed mixtures. A look at Table 3 shows that our method compares quite well with the standard likelihood approach for Gaussian mixtures estimation, especially in the higher dimensions where the average number of well classified data is better by often more than one unit. The proposed l_1 -penalized approach also compares favorably with the estimator given by the CEM algorithm as shown in Table 4. Giving a rigourous argument justifying these observations is currently under investigation but more experiments should be performed in order to explore in finer details the behavior of the method in more realistic context.

4 Conclusion

The goal of this paper was to propose a robust version of the maximum likelihood strategy for the estimation of finite Gaussian mixtures. Our approach is based on self-regression and sparse variable selection. Sparsity was promoted by using an l_1 penalty as in the LASSO. We developed a space alternating version of the penalized EM algorithm and proved that the interesting cluster points satisfy the Karush-Kuhn-Tucker optimality conditions. Our method was then tested on simulated datasets. In particular, the Monte Carlo experiments showed that cluster identification was more robust with our approach than by using the standard maximum likelihood estimator. Theoretical justifications of these observations ought to be investigated in a near future in order to increase our understanding of the strengths and weaknesses of this approach.

Acknowledgement. The author would like to thank Christophe Biernacki and Amelie Vaniscotte for very helpful discussions on the results of this paper.

5 Appendix: Convergence analysis of the Space-Alternating EM algorithm

In what follows, we will restrict our attention to the case where $\Sigma_k^2 = \sigma^2 I$, k = 1, ..., K since more general forms of the covariance matrix will often be intractable in the small sample setting.

When using a maximum likelihood approach, incorporation of a nondifferentiable penalty in the EM algorithm may cause some technical difficulties. A rigorous analysis has been proposed in [7] in the case of general nondifferentiable penalties and space alternating optimization versions of the EM algorithm. The convergence analysis is made easier after interpreting the EM algorithm as a Proximal Point Algorithm which was first done in [5] (see also [6] for more precise results).

In our special case, we only need to show that our Space-Alternating l_1 -EM is a Space-Alternating Kullback Proximal Point Algorithm of the form studied in [7]².

² for a definition of the Clarke subdifferential, see the Appendix of [7]

5.1 Recalls on Kullback-Proximal methods

5.1.1 Background

Proximal point methods have been introduced by Martinet [14] and Rockafellar [15] in the seventies. A relationship between Proximal Point algorithms and EM algorithms was discovered in Chrétien and Hero (2000) (see also Chrétien and Hero (2008) for details). We review the EM analogy to KPP methods to motivate the space alternating generalization. Assume that a family of conditional densities $\{k(x|y;\theta)\}_{\theta\in\mathbb{R}^p}$ is such that the Radon-Nikodym derivative $\frac{k(x|y;\theta)}{k(x|y;\theta)}$ exists for all $\theta, \bar{\theta}$. We can define the following Kullback Leibler divergence:

$$I_{y}(\theta,\bar{\theta}) = \mathsf{E}\left[\log\frac{k(x|y,\bar{\theta})}{k(x|y;\theta)}|y;\bar{\theta}\right].$$
(15)

Let Φ be a function to be maximized. Let us define D_{ϕ} as the domain of Φ , $D_{I,\theta}$ the domain of $I_y(\cdot, \theta)$ and D_I the domain of $I_y(\cdot, \cdot)$. Using the distance-like function I_y , the Kullback Proximal Point algorithm is defined by

$$\theta^{k+1} = \operatorname{argmax}_{\theta \in D_{\Phi} \cap D_{I,\theta}} \left\{ \Phi(\theta) - \beta_k I_y(\theta, \bar{\theta}) \right\}.$$
(16)

The following was proved in Chrétien and Hero (2000).

Proposition 5.1 [Chrétien and Hero (2000) Proposition 1]. In the case where Φ is the loglikelihood, the EM algorithm is a special instance of the Kullback-proximal algorithm with $\beta_k =$ 1, for all $k \in \mathbb{N}$.

5.1.2 The Space Alternating Penalized Kullback-Proximal method

In what follows, and in anticipation of component-wise implementations of penalized Kullback Proximal Point algorithm, we will use the notation $\Theta_{\alpha}(\theta)$ for the local decomposition at θ defined by $\Theta_{\alpha}(\theta) = \Theta \cap \theta + S_{\alpha}$, $\alpha = 1, \ldots, R$ where S_1, \ldots, S_R are subspaces of \mathbb{R}^p and $\mathbb{R}^p = \bigoplus_{\alpha=1}^R S_\alpha$. Let \mathcal{R} be a list with finite cardinality R and let ρ denote any bijection from \mathcal{R} to $\{1, \ldots, R\}$.

Then, the Space Alternating Penalized Proximal Point Algorithm is defined as follows.

Definition 5.1 Let l_y denote a function to be maximized (e.g. the log-likelihood or an arbitrary proxy). Let $\psi \colon \mathbb{R}^p \mapsto S_1 \times \cdots \times S_R$ be a continuously differentiable mapping and let ψ_{α} denote its α^{th} coordinate. Let $(\nu_k)_{k\in\mathbb{N}}$ be a sequence of positive real numbers and ζ_r , $r \in \mathcal{R}$ be non-negative real numbers. Let p_n be a nonnegative possibly nonsmooth locally Lipschitz penalty function with bounded Clarke-subdifferential on compact sets. Then, the Space Alternating Penalized Kullback Proximal Algorithm is defined by

$$\theta^{k+1} = \operatorname{argmax}_{\theta \in \Theta_{k-1}(\text{mod } R)+1}(\theta^k) \cap D_l \cap D_{I,\theta^k} \left\{ l_y(\theta) - \sum_{r \in \mathcal{R}} \zeta_r p_n(\psi_{\rho(r)}(\theta)) - \nu_k I_y(\theta, \theta^k) \right\}, \quad (17)$$

where D_l is the domain of l_y and $D_{I,\theta}$ is the domain of $I_y(\cdot, \theta)$.

In most practical situations, the mappings $\psi_{\rho(r)}$ will simply be the projection onto the subspace Θ_r , $r \in \mathcal{R}$.

Proposition 5.2 The Space Alternating l_1 -EM algorithm (defined in Section 2.2.2) is a particular instance of the Space Alternating Kullback Proximal Point Algorithm as defined by (17).

Proof. First, we adopt the decomposition of the parameter space into the cartesian product of the p_k 's space, the β_k 's space and the σ_k 's space. More precisely Θ_1 is the simplex in \mathbb{R}^K and $\mathcal{S}_1 = \mathbb{R}^K$, $\Theta_{2,k} = \mathbb{R}^n = \mathcal{S}_{2,k}$, and $\Theta_{3,k} = \mathbb{R}_+$ and $\mathcal{S}_{3,k} = \mathbb{R}$ for $k = 1, \ldots, K$. Thus r takes its values in the list $\mathcal{R} = \{1, (2, 1), \ldots, (2, K), (3, 1), \ldots, (3, K)\}$.

Then the mappings Ψ_r are just the orthogonal projections onto S_r for $r \in \mathcal{R}$. Moreover $\zeta_1 = 0$ and $\zeta_{(3,k)} = 0$ for $k = 1, \ldots, K$ because the class probabilities and the variances are not penalized. Moreover we set $\zeta_{(2,k)} = \lambda$ for $k = 1, \ldots, K$.

Next, the Q-function can be written ³

$$Q(\theta,\bar{\theta}) = \hat{I}(\theta) - I_y(\theta,\bar{\theta})$$
(18)

with

$$I_y(\theta,\bar{\theta}) = \sum_{i=1}^n \sum_{k=1}^K t_{ik}(\bar{\theta}) \log\left(\frac{t_{ik}(\bar{\theta})}{t_{ik}(\theta)}\right).$$
(19)

where

$$t_{ik}(\theta) = \frac{p_k f^{(d)}(Y_i; Y\beta_k, \sigma_k I)}{\sum_{l=1}^{K} p_l f^{(d)}(Y_i; Y\beta_l, \sigma_l I)}.$$
(20)

Thus, the space alternating LASSO-EM algorithm is a special case of the Space Alternating Kullback Proximal Point Algorithm for which the sequence $(\nu_k)_{k\in\mathbb{N}}$ is constant and the terms are all equal to one.

We then have the following theorem.

Theorem 5.1 Let θ^* be a cluster point of the Space Alternating Penalized Kullback Proximal sequence. If θ^* lies in the interior of $D_{\tilde{l}}$, then θ^* satisfies the following property: there exists a set of subsets $I_r^{**} \subset I^*$ where I^* denotes the index of the active constraints at θ^* , i.e. $I^* = \{(i, j) \text{ s.t. } t_{i,j}(\theta^*) = 0, \text{ and there is a family of real numbers } \gamma_{ij}, (i, j) \in \mathcal{I}_r^{**}, r \in \mathcal{R} \text{ such that the following Karush-Kuhn-Tucker condition for optimality holds at cluster point <math>\theta^*$:

$$0 \in \nabla \tilde{l}(\theta^*) - \sum_{r \in \mathcal{R}} \zeta_r \partial p_n(\psi_{\rho(r)}(\theta^*)) + \sum_{r \in \mathcal{R}} \sum_{(i,j) \in \mathcal{I}_r^{**}} \gamma_{ij}^* \nabla t_{ij}(\theta^*).$$

Proof. We start by verifying that Assumptions 2.2.1, 2.2.3 and Assumptions 2.2.4 of [7] hold in our case. The differentiability requirement in Assumptions 2.2.1.(i). is obvious. However, if one β_k belongs to the kernel of Y, it may be of any arbitrary large norm without leading the log-likelihood towards $-\infty$. However, note that, as is well known in Gaussian mixture models, \tilde{l} tends to $+\infty$ only at finite number of degenerate points. Thus, since, the penalization terms p_n tend to $+\infty$ as the norm of any β_k tends to $+\infty$, the function $Q(\theta, \theta^{(k)})$ tends to $-\infty$ if the norm of any β_k goes to $+\infty$. Moreover, as easily checked on the expression of the likelihood, the function $Q(\theta, \theta^{(k)})$ also goes to $-\infty$ if any variance σ_k^2 goes to $+\infty$.

The domain $D_{\tilde{l}}$ is defined by the fact that the term inside the log in (3) must be positive. On the other hand, for any $\bar{\theta}$ in $\Theta = \Theta_1 \times \Theta_{2,1} \times \cdots \times \Theta_{2,K} \times \Theta_{3,1} \times \cdots \times \Theta_{3,K}$, the domain

³see Section 4.1 of [7] for more details.

 $D_{I_y,\bar{\theta}}$ is the set of the θ 's for which the $t_{ik}(\theta)$ are positive, and therefore, does not depend on $\bar{\theta}$. Moreover, the set of θ 's for which the $t_{ik}(\theta)$ are positive is $D_{\tilde{l}}$. Thus, the projection of D_I onto the first coordinate is $D_{\tilde{l}}$ and Assumptions 2.2.1.(ii). are satisfied.

Assumptions 2.2.1.(iii). is immediate since here the relaxation sequence (denoted here by $(\nu_k)_{k\in\mathbb{N}}$) is constant. Assumptions 2.2.1.(iv). is also straightforward since the mappings Ψ_r are orthogonal projections onto S_r , $r \in \mathcal{R}$.

In our context, based on (20), we have $\phi = t \log(t) - 1$ and Assumptions 2.2.3.(i)-(iii). are easily verified. Injectivity of the mapping t when restricted to $\bigcup_{j=1}^{3} \Theta_{j,k}$ is proved in [4] and thus, injectivity holds on each $\Theta_{1,k}, \ldots, \Theta_{3,k}$ and Assumption 2.2.3.(iv) holds.

Moreover, since $t_{ik}(\theta) = 0$ implies that $p_k = 0$ and $p_k = 0$ implies

$$\frac{\partial t_{ik}}{\partial \beta_{jl}}(\theta) = 0 \tag{21}$$

for all $j = 1, \ldots, p$ and $l = 1, \ldots, K$ and

$$\frac{\partial t_{ik}}{\partial \sigma^2}(\theta) = 0, \tag{22}$$

it follows that $P_{\mathcal{S}_r}(\nabla t_{ik}(\theta^*)) = \nabla t_{ik}(\theta^*)$ if \mathcal{S}_r is the vector space generated by the probability vectors p and $P_{\mathcal{S}_r}(\nabla t_{ik}(\theta^*)) = 0$ otherwise.

Let θ^* be a cluster point in the interior of $D_{\bar{l}}$. Since the t_{ik} are clearly continuously differentiable around such a θ^* , Corollary 1 in [7] gives that θ^* satisfies the following property: there exists a set of subsets $I_r^* \subset I^*$ and a family of real numbers γ_{ij} , $(i, j) \in \mathcal{I}_r^*$, $r \in \mathcal{R}$ such that the following Karush-Kuhn-Tucker condition for optimality holds at cluster point θ^* :

$$0 \in \nabla \tilde{l}(\theta^*) - \sum_{r \in \mathcal{R}} \zeta_r \partial p_n(\psi_r(\theta^*)) + \sum_{r \in \mathcal{R}} \sum_{(i,j) \in \mathcal{I}_r^{**}} \gamma_{ij}^* \nabla t_{ij}(\theta^*),$$

which is the desired result.

The meaning of this theorem is simply that a Karush-Kuhn-Tucker condition is satisfied at any cluster point in the domain of definition of the log-likelihood.

References

- C. Biernacki, G. Celeux, G. Govaert and F. Langrognet, (2006) "Model-based cluster and discriminant analysis with the MIXMOD software", Computational Statistics and Data Analysis, Vol. 51, 2, 587–600.
- [2] C. Biernacki and S. Chrétien, (2003) "Degeneracy in the maximum likelihood estimation of univariate Gaussian mixtures with EM", Statist. Probab. Lett. 61, no. 4, 373–382.
- [3] E. Candès and Y. Plan, (2009) "Near ideal model selection by l_1 penalization", The Annals of Statistics, to appear.
- [4] G. Celeux, S. Chrétien, F. Forbes and A. Mkhadri (2001) "A Component-Wise EM Algorithm for Mixtures" Journal of Computational and Graphical Statistics, vol. 10, no. 4, 697-712.

- S. Chrétien and A. Hero, (2000) "Kullback proximal algorithms for maximum-likelihood estimation". Information-theoretic imaging. IEEE Trans. Inform. Theory 46, no. 5, 1800– 1810
- S. Chrétien and A. Hero, (2008) " On EM algorithms and their proximal generalizations". ESAIM P&S 12, 308–326
- [7] S. Chrétien, A. Hero and H. Perdry (2008) "Space Alternating Penalized Kullback Proximal Point Algorithms for Maximing Likelihood with Nondifferentiable Penalty", Annals Inst. Stat. Math., to appear. Available at http: //arxiv.org/abs/0901.0017
- [8] A. P. Dempster, N. M. Laird, and D. B. Rubin, (1977) "Maximum likelihood from incomplete data via the EM algorithm," J. Royal Statistical Society, Ser. B, vol. 39, no. 1, pp. 1–38.
- [9] J. Fan and R. Li (2001) "Variable selection via non-concave penalized likelihood and its oracle properties", Journal of the American Statistical Association, 96, 1348–1360.
- [10] J. A. Fessler, and A. O. Hero, (1994)"Space-alternating generalized expectationmaximization algorithm", IEEE Trans. Signal Processing, vol. 42, no. 10, pp. 2664–2677.
- [11] M. Hurn M, A. Justel and C.P. Robert, (2003) "Estimating Mixtures of Regressions", Journal of Computational and Graphical Statistics, 12, 55–79.
- [12] A. Khalili and J. Chen, (2007) "Variable Selection in Finite Mixture of Regression Models", Journal of the American Statistical Association, Volume 102, Number 479, pp. 1025-1038.
- [13] G.J. McLachlan and D. Peel. (2000) Finite Mixture Models. Wiley
- [14] B. Martinet (1970). Régularisation d'inéquation variationnelles par approximations successives. Revue Francaise d'Informatique et de Recherche Operationnelle, vol. 3, pp. 154–179.
- [15] R. T. Rockafellar (1976). Monotone operators and the proximal point algorithm. SIAM Journal on Control and Optimization, vol. 14, pp. 877–898.
- [16] R. Tibshirani, (1996) "Regression shrinkage and selection via the LASSO", Journal of the Royal Statistical Society, Series B, vol. 58, no. 1, pp. 267–288.