

The triple system HIP 96515: a low-mass eclipsing binary with a DB white dwarf companion[★]

N. Huélamo¹, L.P.R. Vaz², C. A. O. Torres³, P. Bergeron⁴, C. H. F. Melo⁵, G. R. Quast³, D. Barrado y Navascués¹, M. F. Sterzik⁶, G. Chauvin⁷, H. Bouy^{8**}, and N. R. Landin²

¹ LAEX-CAB (INTA-CSIC); Postal address: LAEFF, P.O. Box 78, E-28691 Villanueva de la Cañada, Madrid, Spain
e-mail: nhuelamo@laeff.inta.es

² Depto. de Física, Universidade Federal de Minas Gerais, C.P.702, 30161-970 – Belo Horizonte, MG, Brazil

³ Laboratório Nacional de Astrofísica/MCT, Rua Estados Unidos 154, 37504-364 Itajubá, Brazil

⁴ Département de Physique, Université de Montréal, C.P. 6128, Succ. Centre-Ville, Montréal, Québec H3C 3J7, Canada

⁵ European Southern Observatory, Karl-Schwarzschild-Strasse 2, D-85748 Garching bei Muenchen, Germany

⁶ European Southern Observatory, Alonso de Cordova 3107, Casilla 19, Santiago, Chile

⁷ Laboratoire d'Astrophysique, Observatoire de Grenoble, BP 53, 38041 Grenoble, Cedex 9, France

⁸ Instituto de Astrofísica de Canarias, C Vía Láctea, s/n, E38205 - La Laguna (Tenerife), Spain

Received; accepted

ABSTRACT

Context. HIP 96515 A is a double-lined spectroscopic binary included in the SACY catalog as a potential young star, and classified as an eclipsing binary in the ASAS catalog. It has a visual companion (CCDM 19371-5134 B, HIP 96515 B) at 8''.6. If bound to the primary, the optical and infrared colors of this wide companion are consistent with those of a white dwarf.

Aims. The aim of this work is to characterize the system HIP 96515 A&B by studying each of its components.

Methods. We have analyzed spectroscopic and photometric observations of HIP 96515 A and its visual companion, HIP 96515 B. To confirm the system as a common proper motion pair, we have analyzed the astrometry of the components using high-angular resolution infrared observations obtained within a time span of two years and archival astrometry.

Results. The high-resolution optical spectrum of HIP 96515 A has been used to derive a mass ratio, M_2/M_1 , close to 0.9, with the components showing spectral types of M1 and M2. The optical light-curve of HIP 96515 A shows periodic variations with $P_{\text{orb}}=2.3456$ days, confirming that HIP 96515 A is an eclipsing binary, with preliminary parameters of $i=89^\circ.0\pm 0^\circ.2$, and $M_1=0.59\pm 0.03 M_\odot$ and $M_2=0.54\pm 0.03 M_\odot$, for the primary and secondary, respectively, at an estimated distance of 42 ± 3 pc. This is a new eclipsing binary with component masses below $0.6 M_\odot$.

Multi-epoch observations of HIP 96515 A&B show that the system is a common proper motion pair. The optical spectrum of HIP 96515 B is consistent with a pure helium atmosphere (DB) white dwarf. The comparison with evolutionary cooling sequence models provides $T_{\text{eff,WD}}=19,126\pm 195$ K, $\log g_{\text{WD}}=8.08$, $M_{\text{WD}}/M_\odot=0.6$, and a distance of ~ 46 pc. The estimated WD cooling age is ~ 100 Myr and the total age of the object (including the main-sequence phase) is ~ 400 Myr. Finally, if HIP 96515 A&B are coeval, and assuming a common age of ~ 400 Myr, the comparison of the masses of the eclipsing binary members with evolutionary tracks shows that they are underestimated by $\sim 15\%$ and 10% , for the primary and secondary, respectively.

Key words. Stars: white dwarfs – Stars: binaries: eclipsing – Stars: individual (HIP 96515)

1. Introduction

In recent years, several infrared surveys have been focused on the detection of substellar companions around young and nearby stars using high-contrast imaging techniques (e.g. Chauvin et al. 2003; Lowrance et al. 2005; Masciadri et al. 2005; Lafreniere et al. 2007). Young nearby stars are privileged targets for these kind of studies due to their proximity (below 100 pc) which allows to inspect very small separations, and their youth, which enhances the contrast between the central star and the potential companion.

In 2006 we started an infrared survey to detect substellar objects around young stars from the SACY catalog (Torres et al. 2003, 2006). Our sample included HIP 96515, a nearby

($d=43.9\pm 9$ pc, van Leeuwen 2007) M1-type star classified as a double-line spectroscopic binary (SB2) by Torres et al. (2006). The object has a visual companion at an angular separation of $\rho\sim 8''.6$ (CR23, CCDM 19371-5134 B, Worley & Douglass 1997; Dommanget & Nys 2000, HIP 96515 B hereafter). This visual binary has been detected in different infrared and optical surveys, although its properties have never been studied in detail.

As a first step to unveil the nature of the wide visual companion to HIP 96515 A, we compared our K_s -band photometry with evolutionary tracks (Baraffe et al. 2003; Chabrier et al. 2000). The comparison suggested that HIP 96515 B may be of substellar nature if the age of the primary is between 0.07-1 Gyr and the two objects are co-moving. However, the optical and infrared colors retrieved from archived observations are not consistent with a substellar object but with a white dwarf companion.

Motivated by this preliminary result, we decided to perform a detailed study of HIP 96515. In this work, we have analyzed new and archived observations of the triple system and we have de-

Send offprint requests to: N. Huélamo

[★] Based on observations collected at the Paranal Observatory under programs 77.C-0483(A) and 81.C-0826(A).

^{**} Marie Curie Outgoing International Fellow

Table 1. Main properties of HIP 96515 A.

α (J2000)	δ (J2000)	Spectral Type	Parallax ¹ [mas]	μ_α^1 [mas yr ⁻¹]	μ_δ^1 [mas yr ⁻¹]
19 ^h 37 ^m 08 ^s .7	-51°34'00".9	M1 ²	22.79 ±4.87	91.07 ±4.93	-21.94 ±4.62

Notes: ¹ van Leeuwen (2007); ² Torres et al. (2006)

Table 2. Photometric data of the visual binary HIP 96515 A&B

Target	V [mag]	R [mag]	J [mag]	H [mag]	K _s [mag]
HIP 96515 A	11.78 ¹ ±0.04	10.90 ¹ ±0.03	8.82 ² ±0.03	8.15 ² ±0.04	7.96 ² ±0.03
HIP 96515 B	13.0 ³	14.3 ⁴ ±0.3	14.54 ² ±0.18	14.44 ² ±0.51	14.52 ² , 14.94 ⁵ ±0.31 ±0.08

Notes: ¹ Torres et al. (2006); ² 2MASS catalog (Cutri et al. 2003); ³ Visual Double stars in Hipparcos catalog (Dommangeat & Nys 2000); ⁴ UCAC2 catalog; ⁵ NACO/VLT data (this work).

rived the main properties of each source. The paper is organized as follows: Section 2 describes the main observational properties of HIP 96515 A&B. In section 3, we analyze spectroscopic and photometric data of the primary star HIP 96515 A, while the true nature of the visual component, HIP 96515 B is unveiled in section 4. A discussion of the system is provided in Section 5, and the main results are summarized in Section 6.


2. Observational properties of HIP 96515 A&B

The main stellar properties of HIP 96515 A are displayed in Table 1, while its optical and infrared photometry are shown in Table 2.

The analysis of the optical spectrum of HIP 96515 A obtained during the SACY survey has allowed Torres et al. (2006) to classify it as an SB2, with almost equal-mass components (see Table 3). From the cross-correlation function (CFF), we have estimated a magnitude difference of $\Delta V = 0.55$ mag between the binary components.

HIP 96515 is included in the *ROSAT* All-Sky Bright Star catalog (Voges et al. 1999). The source was detected by *ROSAT* with a count rate of 0.172 ± 0.036 counts s⁻¹. We have derived the X-ray flux using the conversion factor definition by Schmitt et al. (1995). The X-ray luminosity, assuming a distance of 44 pc (see Table 1), is $L_x = (3.61 \pm 0.75) 10^{29}$ erg s⁻¹. Using the bolometric correction (*BC*) by Kenyon & Hartmann (1995) for an M1-type star, and the *V*-mag listed in Table 2, we derive a $\log(L_x/L_{bol})$ ratio of -3.07 . If we assume that the two components of the spectroscopic binary (HIP 96515 Aa & Ab) are emitting in X-rays, so that $L_{x,Aa} = L_{x,Ab} = L_x/2$, and using the *V*-mag of each of the binary members (see Table 3) and the *BC* for a M1 and M2 star, respectively, we derive $\log(L_x/L_{bol})$ ratios of -3.17 and -3.04 for Aa and Ab. These values are comparable to those found in very active pre-main sequence (PMS) late-type stars (e.g. Barrado y Navascués et al. 1999; Stelzer & Neuhäuser 2001) and in main-sequence (MS) late-type stars in very close binaries (see e.g. López-Morales 2007).

The age of HIP 96515 A is not known. The lithium content is normally used as a youth indicator in the case of late-type stars. However, it is not always present in young M-type stars since it is depleted too fast (e.g. Basri et al. 1996; Neuhäuser 1997; Torres et al. 2008a). In fact, none of the binary

**Fig. 1.** Hipparcos and ASAS V-band light-curves of HIP 96515 A

members of HIP 96515 A show the lithium absorption doublet ($\lambda = 6707.8$ Å) in the optical spectrum. Torres et al. (2008a) have studied the lithium content as a function of the *V-I* color in different young associations with ages between 5-70 Myr. The comparison of HIP 96515 A with similar M-type stars (with *V-I* ~ 1.8, Torres et al. (2006)) in those associations allows to compute a lower limit to the age of ≥ 70 Myr. By following the temporal evolution of lithium in stellar models (eg. those by Landin et al. 2006) for stars with the masses of HIP 96515 A (Sect. 3), Li should be exhausted by an age between 30 Myr (for the MLT parameter $\alpha=2$) and 50 Myr ($\alpha=1$).

Finally, HIP 96515 B, is included in the AC catalog (Urban et al. 1998) as a visual companion to HIP 96515 A. The object has been detected in different infrared and optical surveys and multi-wavelength photometry is available in different public catalogs. We have included all the available data in Table 2. Note that, in most cases, the photometric measurements show large uncertainties.

3. Characterization of HIP 96515 A: a M1+M2 eclipsing binary

In order to characterize the SB2 in more detail, we retrieved the photometric data of HIP 96515 A from both the Hipparcos archive (Perryman et al. 1997, 96 points) and from the "All Sky Automated Survey", ASAS¹ (Pojmanski & Maciejewski 2005, 818 points). The Hipparcos observations, in its own photometric system (*Hip*), were corrected with the relation $V_j = Hip - 0.08$, suggested for the *V-I* of this object ($V-I=1.85$; Torres et al. 2006).

The observations of the source are displayed in Figure 1, showing significant variability. The object has been classified as an eclipsing binary by Paczyński et al. (2006), using the ASAS data alone. We have applied the method by Lafler & Kinman (1965) to search for any periodicity in the data. The best period, 2.345 days, have been used to phase-fold both the ASAS and Hipparcos data, confirming the SB2 as an eclipsing system. In the next subsections, we describe our full analysis of the photometric data and derive the main properties of the eclipsing components, Aa and Ab.

3.1. Photometric data: Period search

The method by Lafler & Kinman (1965) was used to search for periods in the Hipparcos V-band observations of HIP 96515 A. We searched for periods from 0.5 day to 63 days, by using a relative period step of $\delta P/P = 4 \times 10^{-7}$. Since M stars may flare, and

¹ <http://www.astro.uw.edu.pl/~gp/asas/asas.html>

Table 3. Spectroscopic data of the eclipsing binary HIP 96515 A

Star	V [mag]	Spt. Type	v_{rad} [km s ⁻¹]	H α EW [Å]	$v \cdot \sin i$ [km s ⁻¹]
96515Aa	12.29	M1Ve	84.1	0.9	14.8 \pm 1.5
96515Ab	12.84	M2:Ve	-81.3	2.0	15.8 \pm 1.6

HIP 96515 is a strong X-ray emitter, we excluded the 4 brightest measurements (from Hipparcos measurements) in this period search, in order to avoid these clearly non-periodic events, if they are present. Around 60 significant periods were found, but the most significant was $P_{\text{orb}} = (2.344451 \pm 0.000002)$ d. In the analysis of the photometry with a version of the WD model (see Sect. 3.2), we adjusted a phase shift to the folded observations, and obtained the ephemeris (HJD; the digits in parentheses affect the last digits)

$$\text{Min I} = 2,448,493.80515(10) + 2.3444507(15)E, \quad (1)$$

which we use in our analysis. According to Eq. (1) the secondary minimum occurs very close to phase 0.5, indicating an orbit with a small (if any) eccentricity, and in the lack of further evidence, we assume in the preliminary analysis, a circular orbit. We note that circular orbits are consistent with short orbital (and rotational) periods, like the one found in HIP 96515 A (e.g. Zahn 1989). The V light-curves (Hipparcos and ASAS), folded with these ephemeris, are shown in Fig. 2. The stellar eclipses are clearly seen in both datasets.

3.2. Preliminary analysis of the HIP 96515 A light-curve

We have analyzed the Hipparcos optical light-curve with a version of the WD model (Wilson & Devinney 1971; Wilson 1979, 1993) extensively improved as described in Vaz et al. (2007) and references therein. Our version uses stellar atmospheres and models the radiated flux of both components using the PHOENIX atmosphere models (Allard & Hauschildt 1995; Allard et al. 1997; Hauschildt et al. 1997b,a). The (linear-law) limb-darkening coefficients for both components have been taken from Claret (2000), and interpolated using a bi-linear scheme for the current values of $\log g$ and T_{eff} at each iteration. The reflection albedos for both components have been fixed to a value of 0.5, which is appropriate for stars with convective envelopes. The gravity-brightening exponent, β , has been computed using the local value of T_{eff} for each point on the stellar surface and taking into account mutual illumination following Alencar & Vaz (1997) and Alencar et al. (1999). Our procedure combines the WD modified code with several UNIX scripts and auxiliary programs to guarantee the consistency of the solutions in all steps.

We have a single measurement of the radial velocity for each component at the phase 0.7715, according to Eq. (1), by chance, very close to the quadrature (0.75 in a circular orbit), when the line separation is maximum. Since we do not have any estimation of the center of mass velocity of the system, the mass ratio is undefined. Therefore, to investigate the solution space, we have decided to make a grid of solutions for different values of the mass ratio, $q (= M_2/M_1)$, varying from 1.000 to 0.850 in steps of 0.025. By making this grid we can determine the velocity of the center of mass of the system for each q , and adjust the rest of parameters to reproduce the individual heliocentric velocities measured at phase 0.7715.

We have assigned an arbitrary initial value of 89° to the system inclination, and we have used the WD model to find initial




Fig. 2. The phase-folded light-curves of HIP 96515 A from ASAS (+ symbols) and Hipparcos (× symbols), consisting of 818 and 96 observations, respectively, with our theoretical solution for $q=0.9$ obtained in Sect. 3.2. The Hipparcos observations were shifted by 0.6 mag, in order to separate the light curves. Note that ASAS observations were not corrected from the long term variation of the maxima, shown in Fig. 1.

Table 4. Grid of solutions for 7 values of the mass ratio, $q (= M_2/M_1)$, for HIP 96515 A. Note that V_1 and V_2 , below are the theoretical velocities for the primary and the secondary at the orbital phase $\phi=0.7715$. All the solutions have the reduced χ^2 the closest to 1 possible, for both data sets.

$q (= M_2/M_1)$	1.000	0.975	0.950	0.925	0.900	0.875	0.850
V_1 (km s ⁻¹)	82.70	81.65	80.60	79.48	78.35	77.19	75.99
V_2 (km s ⁻¹)	-82.70	-83.75	-84.82	-85.92	-87.05	-88.21	-89.40
V_γ (km s ⁻¹)	1.40	2.45	3.52	4.62	5.75	6.91	8.10
K_1 (km s ⁻¹)	83.46	82.41	81.32	80.21	79.07	77.90	76.69
K_2 (km s ⁻¹)	83.46	84.51	85.60	86.71	87.85	89.02	90.23
M_1 (M_\odot)	0.56	0.57	0.58	0.59	0.59	0.60	0.61
R_1 (R_\odot)	0.64	0.65	0.65	0.64	0.64	0.64	0.64
$\log g_1$ (cgs)	4.57	4.57	4.58	4.59	4.60	4.60	4.61
M_2 (M_\odot)	0.56	0.56	0.55	0.54	0.53	0.53	0.52
R_2 (R_\odot)	0.53	0.53	0.53	0.54	0.55	0.52	0.52
$\log g_2$ (cgs)	4.74	4.74	4.73	4.71	4.69	4.72	4.72
a (R_\odot)	7.73	7.73	7.73	7.73	7.73	7.73	7.73
Δm	0.875	0.893	0.872	0.832	0.794	0.891	0.888
i ($^\circ$)	89.27	89.45	89.28	89.08	89.00	89.51	89.62
d (pc)	42	42	42	42	42	42	41

values to the gravitational pseudo-potentials (which define the size and form of the components) in order to reproduce the optical light-curve. Finally, the initial values of the effective temperatures have been adopted from the tabulated values listed in Popper (1980), and $T_{\text{eff,pri}}$ was kept fixed at the adopted value of $3,714 \pm 150$ K. In the lack of further information, the eclipsing




Fig. 3. The residuals O–C between the V light curves of HIP 96515 A from Hipparcos (\times symbols) and ASAS (+ symbols) shown in Fig. 2, and our theoretical solution for $q=0.9$ obtained in Sect. 3.2. The standard error (1 observation) is shown in the upper left part of each panel.

components were considered to be synchronized with the orbital motion, an assumption that may not be true (see below).

Despite the presence of a visual companion to HIP 96515 A at $8''.6$, we do not expect any contamination in the Hipparcos light-curve of the primary, since HIP 96515 B is outside the Hipparcos field of view. On the other hand, the resolution of the ASAS setup is $14''/2/\text{pixel}$ and the light from HD 96515 B is definitely included in their measurements. Therefore, we will assume a small amount of third light contribution in the analysis of ASAS light curve. From the appearance of the folded light curve (using our ephemeris) and, in the lack of a well determined radial velocity curve, we have assumed a circular orbit for our analysis.

Taking into account the above mentioned initial values and assumptions, we have modeled the orbital inclination, the gravitational pseudo-potentials, the secondary effective temperature, a phase shift of the primary minimum and the primary internal luminosity for each point of the grid. The most interesting results from our modeling are displayed in Table 4.

After convergence of the solutions of Table 4, we adjusted the amount of third light of the $q=0.900$ solution, due to the evidence that the visual companion, HIP 96515 B, contaminates the ASAS measurements. The third light parameter, in units of the eclipsing system light at quadrature, determined was $\mathcal{L}_3=0.005\pm 0.038$ and the fit was essentially indistinguishable from the solution with $\mathcal{L}_3=0$.

In order to reproduce the observed radial velocities, we have adjusted the semi-major orbital axis which, together with the orbital inclination and period, fixed the theoretical radial velocity amplitudes, K_1 and K_2 (in km s^{-1}). The apparent V -magnitude retrieved from SIMBAD is 11.10 mag. However, the Hipparcos measurement converted to Johnson’s V gives 11.73 mag and our measurement is 11.78 mag (Torres et al. 2006), value adopted for the distance estimation. By using bolometric corrections for both components from Popper (1980), we can derive the absolute dimensions and, finally, estimate the distance to be $(42\pm 3)\text{pc}$. The interstellar absorption in such a short distance should be negligible. The parallax for the system is $(22.79\pm 4.87)\text{mas}$ measured by Hipparcos, which gives $(44\pm 9)\text{pc}$, in very good agreement with our determination.

All solutions of Table 4 are essentially indistinguishable from the curve fitting point of view, as can be seen by the rms of the residuals (σ , 1 obs) in magnitudes. The probable intrinsic variability of one or both components makes it necessary to follow the system photometrically closely in time, preferably

in different passbands and simultaneously with the spectroscopy for radial velocity measurements. Only with more precise light-curves in different photometric bands and with a precise radial velocity curve it will be possible to eliminate this degeneracy of the solutions in the mass ratio. For the purpose of this paper, we will assume that the components are in the main-sequence and make use of the mass-luminosity relation to fix q .

Figure 2 shows our solution for $q=0.9$ and Fig. 3 shows the corresponding residuals O–C. The steady and slow decreasing of the maxima present in ASAS observations was not corrected for in this analysis. It is common that M dwarfs present variations (e.g. Rockenfeller et al. 2006) mainly due to spots on their surfaces. However, we cannot say, with the data we analyzed, if the variability is due to one or to both eclipsing components. We searched for periodicities in the O–C residuals from 0.08 day to 50 days, using the phase dispersion method by Lafler & Kinman (1965) and relative period steps of $\delta P/P=3.d-3$, and found a significant period of (0.24965 ± 0.00001) day (about 6 hrs), as can be seen in Fig. 4. It is interesting to note that the variation pattern is present in both the Hipparcos and the ASAS data sets. The cadence of these observations (one or two observations a day, being common large intervals without observations) is not adequate for detecting these systematic variations in the O–C, with the annoying consequence of introducing a multitude of aliases in the analysis. It is, however, clear that the period of 0.24965 day, if real, is not the only one: Fig. 1 gives indication for longer period variations and we, as a matter of fact, estimate at least another possible period in the O–C of $(4,100\pm 200)$ days, but with little significance level, due to the relatively short interval of the observations ($\sim 6,630$ days).

The period of 0.24965 day detected may be due either to pulsation or to spots on the surface of one or both of the eclipsing components. Rockenfeller et al. (2006) list other M dwarfs with 6 hrs period. However, if the variation is due to spots on HIP 96515 A, it would indicate that the spotted component(s) spins significantly (more than 9 times) faster than the orbital motion (see e.g. CV Boo Torres et al. 2008b). This is probably not the case in HIP 96515 A: using an orbital period of 2.345 days, and the radius of the binary components from Table 5, we estimate rotational velocities of $\sim 14\text{ km.s}^{-1}$ and $\sim 12\text{ km.s}^{-1}$ for Aa and Ab, respectively. These values are comparable with the ones derived from the optical spectrum (Table 3, and assuming an inclination of 89 degrees), indicating that the system is synchronized. Therefore, the origin of the 6h-period remains unclear with the current data-set.

By combining the results of Table 4, and taking into account the maximum variations of the variables along the grid, we can derive the preliminary absolute dimensions for HIP 96515 A, which are listed in Table 5. The masses of the binary components, HIP 96515 Aa and Ab, are below $0.6 M_{\odot}$. Hence, HIP 96515 A is the fourteenth confirmed EB with component masses below $0.7 M_{\odot}$ (see e.g. Shkolnik et al. 2008), which are important to calibrate theoretical evolutionary models.

Finally, we note that the eclipsing system definitely deserves more observations, both multi-band photometry with a cadence that allows determination of precise light curves, and spectroscopy, in order to obtain reliable radial velocity curves and spectral types of both components. If possible, HIP 96515 B, should also be observed simultaneously, in order to give consistency to the analysis.




Fig. 4. The residuals $O-C$ between the V light curves of HIP 96515 A from ASAS (+ symbols) and Hipparcos (grey circles) shown in Fig. 3, folded with the period of (0.24965 ± 0.00001) day. Note that the pattern is present in both Hipparcos and ASAS data sets (two cycles are plotted).

Table 5. Preliminary physical parameters for HIP 96515 Aa&Ab

	Primary	Secondary
Absolute dimension:		
Mass (M_{\odot})	0.59 ± 0.03	0.54 ± 0.03
Radius (R_{\odot})	0.64 ± 0.01	0.55 ± 0.03
$\log g$ (c.g.s.)	4.59 ± 0.03	4.69 ± 0.05
Photometric data:		
$\log T_{\text{eff}}$ (K)	3.571 ± 0.018	3.555 ± 0.019
$\log L/L_{\odot}$	-1.15 ± 0.07	-1.35 ± 0.08
M_{bol}	7.63 ± 0.18	8.13 ± 0.20
M_V	9.10 ± 0.18	9.90 ± 0.20
$L_{\text{sec}}/L_{\text{pri}}$	0.48 ± 0.24	
P_{orb} (days)	2.3456	
i (deg)	89.0 ± 0.2	
Distance (pc)	42 ± 3	

4. Characterization of HIP 96515 B

In the next subsections, we will focus on the characterization of HIP 96515 B. First, we have investigated if the object forms a common proper motion pair with HIP 96515 A. To do this, we have analyzed diffraction-limited infrared observations of the pair obtained in the course of our program to detect substellar object around SACY targets (Huelamo et al. 2008, in prep.). As a second step, we have obtained an optical spectrum to derive the main properties of the object.

4.1. NACO/VLT observations of HIP 96515

We observed HIP 96515 twice with NAOS-CONICA (NACO), the Adaptive Optics facility at the Very Large Telescope (VLT), the nights of the 25th May 2006 and 16th June 2008. The target was observed with the visible wavefront sensor and the Ks-band filter. We used the S27 objective (nominal plate scale of $\sim 0.027''/\text{pixel}$) which provides a total field-of-view of $\sim 27'' \times 27''$. The total exposure time was ~ 12 minutes on source.

The data were obtained using a random jitter between the exposures to compute the sky emission. The images have been reduced with the *Eclipse* reduction package (Devillard 1997)




Fig. 5. VLT/NACO K_s -band image of HIP 96515 A and its visual companion, HIP 96515 B, obtained in June 2008. The projected separation is of 378 AU at a distance of 44 pc. Note that the eclipsing binary (Aa & Ab) is not spatially resolved by NACO.

following the standard steps: dark-subtraction, flat-field division and image shifting and stacking. The final image of HIP 96515 is displayed in Figure 5. Apart from HIP 96515 A, we only detected the already known visual companion, HIP 96515 B, at a projected separation of $8''.6$ (~ 378 AU at 44 pc) from the primary. Note that the eclipsing binary is not spatially resolved in the NACO image.

The difference in magnitude between HIP 96515 A&B is 6.98 ± 0.07 mag. Taking into account the brightness of the primary ($K_s = 7.9$ mag, see Table 2), we can estimate a K_s -mag of 14.94 ± 0.08 for the secondary.

The NACO images have been used to derive the separation and position angle of the two objects in two different epochs. To derive accurate astrometry of HIP 96515 B in the two images, we have calibrated the plate scale and orientation of the infrared detector, CONICA, with the astrometric calibrator θ^1 Ori C (McCaughrean & Stauffer 1994). The results are displayed in Table 6.

4.2. HIP96515 A & B: A common proper motion pair

The astrometric accuracy of the NACO/VLT data is high enough to investigate if HIP 96515 A&B are co-moving. For completeness, we have compared the NACO/VLT data with older (and less accurate) astrometric measurements from public surveys (e.g. Dommanget & Nys 2000; Cutri et al. 2003). The visual binary is included in the Washington Double Star Catalog (WDSC, Worley & Douglass 1997) with three observations between 1910 and 2000. Table 6 includes the angular separation and position angle of the two components at five different epochs. As seen, both remain almost constant within a time span of 98 years.

Figure 6 (top panel) shows the difference in right ascension (RA) and declination (DEC) of the binary members measured in five epochs between 1910 and 2008. We have also overplotted the expected difference in RA and DEC of a background object taking into account the proper motion of the primary. If we focus on the NACO/VLT data only (Fig. 6, bottom panel), which provides the best astrometric accuracy, we can confirm




Fig. 6. Differences in right ascension and declination between HIP 96515 A & B at different epochs (black circles). The expected position of a background object (grey circles and labels) at each epoch is also displayed. **Top:** All the astrometric data from 1910 to 2008. **Bottom:** Only the NACO data, which provides the best astrometric accuracy. The figure shows that the two objects are co-moving, that is, they are a common proper motion pair.

that HIP 96515 A&B are comoving, that is, they form a common proper motion pair.

4.3. The Optical Spectrum of HIP 96515 B: observations and modeling

To unveil the nature of HIP 96515 B, we have obtained an optical spectrum of the target with the ESO Multi-Mode Instrument (EMMI) on the New Technology Telescope (NTT) on October 2007. The wavelength range goes from 380 nm to 700 nm, with a spectral resolution of $R \sim 1100$. The spectrum has been reduced with standard routines within IRAF, including bias subtraction and flat-fielding, and has been flux-calibrated with a spectrophotometric standard observed during the night.

Table 6. Relative astrometry of HIP 96515 A & B

Obs. Date	Separation [arcsec]	PA [deg]	Reference
2008-06-16	8.586 ± 0.008	224.3 ± 0.2	NACO/VLT, this paper
2006-05-23	8.586 ± 0.010	224.2 ± 0.2	NACO/VLT, this paper
2000-10-10	8.3 ± 0.2	224.6	2MASS; Cutri et al. (2003)
1974.620	8.4	223	Dommanget & Nys (2000)
1910.705	8.9 ± 0.3	222	Worley & Douglass (1997)




Fig. 7. Top: NTT/EMMI Optical spectrum of HIP 96515 B. We have marked all the identified He I lines. **Bottom:** Fit of the optical spectrum using pure helium model atmospheres. The best fit parameters are displayed at the top right corner.

Table 7. Best fit parameters for HIP 96515 B

T_{eff} (K)	19126 ± 195
$\log g$ (cgs)	8.08 ± 0.07
Mass (M_{\odot})	0.648
Radius (R_{\odot})	0.01212
M_V (mag)	10.89
M_J (mag)	11.33
distance (pc)	46
Progenitor Mass (M_{\odot})	3.1 ± 0.8
WD Cooling age (Myr)	106 ± 13
MS lifetime (Myr)	~ 300 ¹
Total age (Myr)	~ 400

¹ see text

The optical spectrum of HIP 96515 B is displayed in Figure 7. The continuum increases towards bluer wavelengths and the main detected spectral features are the He I absorption lines, typical of helium (DB) white dwarfs.

We have compared the optical spectrum of HIP 96515 B with atmospheric models of DB white dwarfs, in order to derive the physical parameters of the object. Our grid of model at-

mospheres and synthetic spectra for DB stars is described in Beauchamp et al. (1996). These include the improved Stark profiles of neutral helium of Beauchamp et al. (1997). The model atmospheres used here assume a pure helium composition and cover a range of $T_{\text{eff}} = 10,000$ (1000) 16,000 (2000) 30,000 K and $\log g = 7.0$ (0.5) 9.0. Our fitting technique relies on the nonlinear least-squares method of Levenberg-Marquardt (Press et al. 1986), which is based on a steepest descent method. The model spectra (convolved with a Gaussian instrumental profile) and the optical spectrum are first normalized to a continuum set to unity. The calculation of χ^2 is then carried out in terms of these normalized line profiles only. We adopt a pure helium composition based on the lack of any $H\alpha$ absorption feature; additional models with mixed helium and hydrogen compositions allow us to set a limit of $N(\text{H})/N(\text{He}) < 10^{-5}$.

As usual, there is a cool and a hot solution on each side of the temperature at which the line strengths reach their maximum, but the hot solution near $T_{\text{eff}}=25,000$ K can easily be ruled out based on the slope of the observed energy distribution. Our final solution, $T_{\text{eff}}=19,130\pm 190$ K and $\log g=8.08\pm 0.07$, is displayed in Figure 7. These atmospheric parameters can be converted into a mass of $M=0.65\pm 0.04 M_{\odot}$ using evolutionary models with C/O cores, $q(\text{He}) \equiv \log M_{\text{He}}/M_{\star}=10^{-2}$, and $q(\text{H})=10^{-10}$, which are representative of helium-atmosphere white dwarfs².

By using the best fit parameters of the model we can derive the absolute magnitudes in different photometric bands (see Table 7). Using M_J and the 2MASS J -band brightness of HIP 96515 B, which is the measurement with the smallest photometric error, we estimate the distance to the source of $d=45.8$ pc, which is in good agreement with both the Hipparcos parallax and the distance derived from the analysis of HIP 96515 A.

The evolutionary sequences described above, as well as empirical relations (see e.g. Fontaine et al. 2001), allow to estimate the mass of the white dwarf progenitor and its cooling age. These data are provided in Table 7. We obtain two different estimates of the mass of the white dwarf progenitor. First, we use the empirical initial-final mass relationship (IFMR) from Wood (1992), which is based on the study of the white dwarf luminosity function and mass distribution (see also Leggett et al. 1998; Fontaine et al. 2001). This yields a value of $M_{\text{MS}} = 3.8\pm 0.5 M_{\odot}$. Alternatively, we use the IFMR determined by Ferrario et al. (2005) based on white dwarfs observed in open clusters (see in particular their Fig. 1). This second relation yields a significantly smaller final mass of $M_{\text{MS}} = 2.5 \pm 0.5 M_{\odot}$. However, as discussed by Ferrario et al., the IFMR of most clusters appears to have an intrinsic spread (i.e. there is a range of initial masses that gives rise to a given final mass). Hence a value of M_{MS} as high as $3.8 M_{\odot}$ cannot be necessarily ruled out. In the following, we thus adopt a mean value of $M_{\text{MS}} = 3.1 \pm 0.8 M_{\odot}$ to allow for the possible mass range of the progenitor.

We have derived the main-sequence lifetime of HIP 96515 B using two sets of evolutionary tracks for a $3.1\pm 0.8 M_{\odot}$ star. First, we have used the ATON 2.4 models, which are a modified version of the models by D'Antona & Mazzitelli (D'Antona & Mazzitelli 1994, 1997), and are fully described in Landin et al. (2006). According to these evolutionary models, a $3.1\pm 0.8 M_{\odot}$ star spends 280^{+346}_{-125} Myr on the main-sequence, i.e., from the zero-age main-sequence (ZAMS) to the terminal age main-sequence (TAMS). On the other hand, Claret (2004) models ($Z=0.02$) predict a span of ~ 317 Myr for the MS time of a $3.1 M_{\odot}$. Assuming a MS lifetime of ~ 300 Myr and a cooling

time of ~ 100 Myr (see Table 7), the total age of the system would be ~ 400 Myr, being the errors dominated by the uncertainty in the mass.

5. Discussion

The comparison of HIP 96515 Aa & Ab with evolutionary tracks by Baraffe et al. (1998) provides an age of ~ 50 Myr for masses of $0.5 M_{\odot}$ (with $\alpha=1$), and places the binary on the PMS (see Fig. 8). The PMS nature of HIP 96515 A is also suggested by tracks generated with the ATON 2.4 stellar evolutionary code, where we adopted the solar chemical composition ($Z=0.0175$ and $Y=0.27$) and α , α , equal to 1. The tracks in Fig. 8 generated with ATON2.4 code cover the PMS, MS and earlier stages of post main-sequence phases. The best age, according to ATON 2.4, is ~ 60 Myr (Fig. 8), although with large uncertainties.

On the other hand, our astrometric study shows that the white dwarf and the EB are co-moving. If we assume that HIP 96515 A&B are coeval, the age of the triple system should be ~ 400 Myr, which is significantly older than the age provided by the PMS tracks for HIP 96515 A, although within the observational uncertainties of the primary star (see Fig. 8).

To shed light on the evolutionary status of HIP 96515 A, we have studied the kinematical properties of the object. Using an average distance of 45 pc (coming from the EB and WD independent studies presented here), the RV derived by Torres et al. (2006), and the proper motions measured by Hipparcos (see Table 1), we can derive the UVW components of the Galactic space velocity vector: $UVW=-6.0, -0.5, -21.6 \text{ km s}^{-1}$. We have compared them with the UVW vectors of young moving groups in the solar neighborhood included in Zuckerman & Song (2004) and Torres et al. (2008a). We conclude that HIP 96515 A does not share the kinematical properties of any of these associations.

Assuming an age of 400 Myr for the triple system, a possible explanation to the age of 60 Myr derived with PMS evolutionary tracks is the different evolution of EBs and very magnetically active stars in comparison with non-active objects. EBs are normally fast rotators ($P_{\text{rot}} < 3$ days) and they exhibit strong X-ray emission related with the presence of magnetic fields (López-Morales 2007). As explained by e.g. Chabrier et al. (2007), rapid rotation and/or magnetic fields reduce the efficiency of large-scale thermal convection in their interior leading to less efficient heat transport. The reduction of the stars radiating surface due to the presence of cold spots can yield to smaller T_{eff} and larger radii, thus departing significantly from the models predictions (see e.g. Torres & Ribas 2002).

To investigate this in more detail, we have studied the Radius vs. Mass relation for HIP 96515 A and compared it with the predictions from evolutionary models by Baraffe et al. (1998). The adopted models correspond to a solar metallicity of $[\text{Fe}/\text{H}]=0$, and a mixing length of $\alpha=1$. The results are displayed in Fig. 9, which includes all the low-mass eclipsing binaries confirmed so far. The data has been adopted from Shkolnik et al. (2008) and references therein.

If we assume an age of 400 Myr for the triple system, the radii of HIP 96515 Aa&Ab are $\sim 15\%$ and 10% larger, respectively, than predicted by the evolutionary tracks (see Fig. 9). These discrepancies have been previously observed in similar eclipsing binaries and have been attributed to the effect of rapid rotation, magnetic activity and/or metallicity (e.g. Berger et al. 2006; Chabrier et al. 2007; López-Morales 2007).

² see <http://www.astro.umontreal.ca/~bergeron/CoolingModels>




Fig. 8. Stellar Luminosity (top panel) and surface gravity (bottom panel) vs. effective temperature. HIP 96515 Aa and Ab are represented by filled circles. The grey and black solid lines represent evolutionary tracks by Baraffe et al. (1998) and Landin et al. (2006), respectively, for 0.6 and $0.5 M_{\odot}$ stellar masses. The dotted lines represent the uncertainties in the stellar masses for the ATON models. The comparison of HIP 96515 A with the two sets of evolutionary tracks provides an age of ~ 60 Myr, that is, places the eclipsing binary members on the pre-main sequence, although with large uncertainties.

6. Results and conclusions

In this paper we have studied in detail HIP 96515, a triple system composed by a double-lined spectroscopic binary (SB2) and a co-moving white dwarf at a projected separation of $8''.6$. We have analyzed both new and archived observations of the system. Our main results can be summarized as follows:

- The analysis of multi-epoch optical photometry of HIP 96515 A has revealed that this SB2 is an eclipsing binary with almost equal-mass components: $M_{Aa}=0.59 M_{\odot}$ and $M_{Ab}=0.54 M_{\odot}$. In fact, this is the fourteenth confirmed eclipsing binary with component masses below $0.7 M_{\odot}$, which are of extreme importance to calibrate theoretical evolutionary tracks.
- Archival astrometry and new NACO/VLT data show that HIP 96515 A&B is a common proper motion pair.
- The optical spectrum of HIP 96515 B is consistent with a hot DB white dwarf. We have modeled it using evolutionary cooling sequence models, deriving a temperature, surface gravity and mass of $T_{\text{eff}} \text{ (K)}=19126$, $\log g = 8.08$,




Fig. 9. Mass-Luminosity relation for all known eclipsing binaries with masses $M < 0.7 M_{\odot}$ (see Shkolnik et al. 2008, and references therein). HIP 96515 Aa and Ab are represented by filled circles. The dotted, dashed and solid lines represent evolutionary tracks by Baraffe et al. (1998) for 50, 100 and 500 Myr, respectively.

and $M=0.65 M_{\odot}$. Empirical relations provide a progenitor mass of $3.1 \pm 0.8 M_{\odot}$, and a white dwarf cooling age of 100 ± 13 Myr. The MS lifetime of a $3.1 M_{\odot}$ star, as derived from evolutionary models, is ~ 300 Myr, resulting in a total age of ~ 40 Myr for the white dwarf.

- A direct comparison of the EB members with evolutionary tracks (Baraffe et al. 1998; Landin et al. 2006) places them on the PMS with an age of ~ 60 Myr, although with large uncertainties. We note that HIP 96515 A does not share the kinematical properties of known young moving groups in the solar vicinity.
- If HIP 96515 A&B are coeval, the system must be older than 300 Myr. The discrepancy between this age and the 60 Myr provided by PMS tracks might be explained by the different evolution of EBs, which normally show fast rotation and strong magnetic fields, compared to non-magnetic stars. Since HIP 96515 A shows fast rotation and strong X-ray emission, it can depart from the predictions of theoretical models, which place the binary on the PMS.
- If we assume an age of ~ 400 Myr for the triple system, the comparison of HIP 96515 Aa&Ab with evolutionary tracks by Baraffe et al. (1998) shows that the models underestimate the stellar radii by 15% and 10%, for the primary and the secondary, respectively. This behavior has been previously observed in similar low-mass EBs.

Finally, we note that additional RV observations are needed to better constraint the orbit and physical parameters of the eclipsing binary.

Acknowledgements. We are very grateful to I. Saviane for obtaining the NTT/EMMI spectrum presented here. NH is indebted to the Spanish *Programa Juan de la Cierva*. This research has been funded by Spanish grant MEC ESP2007-65475-C02-02 and MEC/Consolider-CSD2006-0070. NH and CT gratefully acknowledge support from ESO-DGDF 2007 program. GC acknowledges support from the Faculty of the European Space Astronomy Centre (ESAC). HB acknowledges the funding from the European Commission's Sixth Framework Program as a Marie Curie Outgoing International Fellow (MOIF-CT-2005-8389). LPRV and NRL gratefully acknowledges financial support from the Brazilian agencies CAPES, CNPq and FAPEMIG. PB is a Cottrell Scholar of the Research Corporation for Science Advancement and is supported in part by

the NSERC Canada and by the Fund FQRNT(Québec). This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France, of NASA's Astrophysics Data System Abstract Service, and of the *Washington Double Star Catalog* maintained at the U.S. Naval Observatory.

References

- Alencar, S. H. P. & Vaz, L. P. R. 1997, *A&A*, 326, 257
 Alencar, S. H. P., Vaz, L. P. R., & Nordlund, Å. 1999, *A&A*, 346, 556
 Allard, F. & Hauschildt, P. H. 1995, *ApJ*, 445, 433
 Allard, F., Hauschildt, P. H., Alexander, D. R., & Starrfield, S. 1997, *ARA&A*, 35, 137
 Baraffe, I., Chabrier, G., Allard, F., & Hauschildt, P. H. 1998, *A&A*, 337, 403
 Baraffe, I., Chabrier, G., Barman, T. S., Allard, F., & Hauschildt, P. H. 2003, *A&A*, 402, 701
 Barrado y Navascués, D., Stauffer, J. R., Song, I., & Caillault, J.-P. 1999, *ApJ*, 520, L123
 Basri, G., Marcy, G. W., & Graham, J. R. 1996, *ApJ*, 458, 600
 Beauchamp, A., Wesemael, F., & Bergeron, P. 1997, *ApJS*, 108, 559
 Beauchamp, A., Wesemael, F., Bergeron, P., Liebert, J., & Saffer, R. A. 1996, in *Astronomical Society of the Pacific Conference Series*, Vol. 96, *Hydrogen Deficient Stars*, ed. C. S. Jeffery & U. Heber, 295–+
 Berger, D. H., Gies, D. R., McAlister, H. A., et al. 2006, *ApJ*, 644, 475
 Chabrier, G., Baraffe, I., Allard, F., & Hauschildt, P. 2000, *ApJ*, 542, 464
 Chabrier, G., Gallardo, J., & Baraffe, I. 2007, *A&A*, 472, L17
 Chauvin, G., Thomson, M., Dumas, C., et al. 2003, *A&A*, 404, 157
 Claret, A. 2000, *A&A*, 363, 1081
 Claret, A. 2004, *A&A*, 424, 919
 Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, *VizieR Online Data Catalog*, 2246, 0
 D'Antona, F. & Mazzitelli, I. 1994, *ApJS*, 90, 467
 D'Antona, F. & Mazzitelli, I. 1997, *Memorie della Societa Astronomica Italiana*, 68, 807
 Devillard, N. 1997, *The Messenger*, 87, 19
 Dommanget, J. & Nys, O. 2000, *A&A*, 363, 991
 Ferrario, L., Wickramasinghe, D., Liebert, J., & Williams, K. A. 2005, *MNRAS*, 361, 1131
 Fontaine, G., Brassard, P., & Bergeron, P. 2001, *PASP*, 113, 409
 Hauschildt, P. H., Allard, F., Alexander, D. R., & Baron, E. 1997a, *ApJ*, 488, 428
 Hauschildt, P. H., Baron, E., & Allard, F. 1997b, *ApJ*, 483, 390
 Kenyon, S. J. & Hartmann, L. 1995, *ApJS*, 101, 117
 Laffer, J. & Kinman, T. D. 1965, *ApJS*, 11, 216
 Lafreniere, D., Doyon, R., Marois, C., et al. 2007, *ArXiv e-prints*, 705
 Landin, N. R., Ventura, P., D'Antona, F., Mendes, L. T. S., & Vaz, L. P. R. 2006, *A&A*, 456, 269
 Leggett, S. K., Ruiz, M. T., & Bergeron, P. 1998, *ApJ*, 497, 294
 López-Morales, M. 2007, *ApJ*, 660, 732
 Lowrance, P. J., Becklin, E. E., Schneider, G., et al. 2005, *AJ*, 130, 1845
 Masciadri, E., Mundt, R., Henning, T., Alvarez, C., & Barrado y Navascués, D. 2005, *ApJ*, 625, 1004
 McCaughrean, M. J. & Stauffer, J. R. 1994, *AJ*, 108, 1382
 Neuhäuser, R. 1997, *Science*, 276, 1363
 Paczyński, B., Szczygiel, D. M., Pilecki, B., & Pojmański, G. 2006, *MNRAS*, 368, 1311
 Perryman, M. A. C., Lindgren, L., Kovalevsky, J., et al. 1997, *A&A*, 323, L49
 Pojmanski, G. & Maciejewski, G. 2005, *Acta Astronomica*, 55, 97
 Popper, D. M. 1980, *ARA&A*, 18, 115
 Press, W. H., Flannery, B. P., & Teukolsky, S. A. 1986, *Numerical recipes. The art of scientific computing* (Cambridge: University Press, 1986)
 Rockenfelder, B., Bailer-Jones, C. A. L., & Mundt, R. 2006, *A&A*, 448, 1111
 Schmitt, J. H. M. M., Fleming, T. A., & Giampapa, M. S. 1995, *ApJ*, 450, 392
 Shkolnik, E., Liu, M. C., Reid, I. N., et al. 2008, *ApJ*, 682, 1248
 Stelzer, B. & Neuhäuser, R. 2001, *A&A*, 377, 538
 Torres, C. A. O., Quast, G. R., da Silva, L., et al. 2006, *A&A*, 460, 695
 Torres, C. A. O., Quast, G. R., de La Reza, R., et al. 2003, in *Astrophysics and Space Science Library*, Vol. 299, *Open issues in Local Star Formation*, ed. J. Lépine & J. Gregorio-Hetem, 83
 Torres, C. A. O., Quast, G. R., Melo, C. H. F., & Sterzik, M. F. 2008a, *Handbook of Star Forming Regions, Volume II; The Southern Sky*, *Astronomical Society of the Pacific, Monograph 5*, ed. Bo Reipurth
 Torres, G. & Ribas, I. 2002, *ApJ*, 567, 1140
 Torres, G., Vaz, L. P. R., & Sandberg Lacy, C. H. 2008b, *ArXiv e-prints*
 Urban, S. E., Corbin, T. E., Wycoff, G. L., et al. 1998, *AJ*, 115, 1212
 van Leeuwen, F. 2007, *A&A*, 474, 653
 Vaz, L. P. R., Andersen, J., & Claret, A. 2007, *A&A*, 469, 285
 Voges, W., Aschenbach, B., Boller, T., et al. 1999, *A&A*, 349, 389
 Wilson, R. E. 1979, *ApJ*, 234, 1054
 Wilson, R. E. 1993, in *Astronomical Society of the Pacific Conference Series*, Vol. 38, *New Frontiers in Binary Star Research*, ed. K.-C. Leung & I.-S. Nha, 91
 Wilson, R. E. & Devinney, E. J. 1971, *ApJ*, 166, 605
 Wood, M. A. 1992, *ApJ*, 386, 539
 Worley, C. E. & Douglass, G. G. 1997, *A&AS*, 125, 523
 Zahn, J.-P. 1989, *A&A*, 220, 112
 Zuckerman, B. & Song, I. 2004, *ARA&A*, 42, 685