

# Pre-nova X-ray observations of V2491 Cyg (Nova Cyg 2008b)

A. Ibarra<sup>1\*</sup>, E. Kuulkers<sup>1</sup>, J.P. Osborne<sup>2</sup>, K. Page<sup>2</sup>, J.U. Ness<sup>1</sup>, R.D. Saxton<sup>1</sup>, W. Baumgartner<sup>3,4</sup>, V. Beckmann<sup>5,6</sup>, M.F. Bode<sup>7</sup>, M. Hernanz<sup>8</sup>, K. Mukai<sup>9</sup>, M. Orío<sup>10,11</sup>, G. Sala<sup>12</sup>, S. Starrfield<sup>13</sup>, G.A. Wynn<sup>2</sup>

<sup>1</sup> ESA, European Space Astronomy Centre (ESAC), Apartado 78, 28691, Villanueva de la Cañada (Madrid), Spain  
e-mail: Aitor.Ibarra@sciops.esa.int

<sup>2</sup> The Department of Physics and Astronomy, The University of Leicester, Leicester, LE1 7RH, UK

<sup>3</sup> NASA/Goddard Space Flight Center, Code 662, Greenbelt, MD 20771, USA

<sup>4</sup> University of Maryland, Baltimore County, 1000 Hilltop Circle, Baltimore, MD 21250, USA

<sup>5</sup> ISDC Data Centre for Astrophysics, Chemin d'Écogia 16, 1290 Versoix, Switzerland

<sup>6</sup> Observatoire Astronomique de l'Université de Genève, Chemin des Maillettes 51, 1290 Sauverny, Switzerland

<sup>7</sup> Astroph. Research Institute, Liverpool John Moores Univ., Twelve Quays House, Egerton Wharf, Birkenhead CH41 1LD 7, UK

<sup>8</sup> IEEC-CSIC, Campus UAB, Facultat Ciències, C5-par, 2on, 08193 Bellaterra, Spain

<sup>9</sup> CRESST and X-ray Astrophysics Laboratory NASA/GSFC, Greenbelt, MD 20771, USA

<sup>10</sup> INAF - Osservatorio Astronomico di Padova, vicolo Osservatorio, 5, I-35122 Padova, Italy

<sup>11</sup> Department of Astronomy, University of Wisconsin, 475 N. Charter Str., Madison WI 53706, USA

<sup>12</sup> Grup d'Astronomia i Astrofísica, Dept. Física i Enginyeria Nuclear, Univ. Politècnica de Catalunya, E-08036 Barcelona, Spain

<sup>13</sup> School of Earth & Space Exploration, Arizona State University, P.O. Box 871404, AZ 85287-1404, USA

Received January 23, 2009; accepted February 23, 2009

## ABSTRACT

Classical novae are phenomena caused by explosive hydrogen burning on an accreting white dwarf. So far, only one classical nova has been identified in X-rays before the actual optical outburst occurred (V2487 Oph). The recently discovered nova, V2491 Cyg, is one of the fastest (He/N) novae observed so far. Using archival *ROSAT*, *XMM-Newton* and *Swift* data, we show that V2491 Cyg was a persistent X-ray source during its quiescent time before the optical outburst. We present the X-ray spectral characteristics and derive X-ray fluxes. The pre-outburst X-ray emission is variable, and at least in one observation it shows a very soft X-ray source.

**Key words.** X-Ray Binaries – Novae

## 1. Introduction

Classical (and recurrent) novae are the third most energetic outbursts after GRB and SNe, but are much more common than either of those types of catastrophic explosions. They take place in binary systems in which mass is transferred from a low-mass secondary star onto a white dwarf. The accreted material gradually becomes degenerate, and when temperatures become high enough and the pressure at the bottom of the accreted envelope is sufficient, a thermonuclear runaway results. Enough energy is deposited in the accreted material to eject a fraction of the envelope of the white dwarf (Bode & Evans 2008).

Novae can be classified into two observational groups: Fe II and He/N, respectively, depending on the emission lines detected in their optical spectra (Williams 1992). Novae with prominent Fe II lines evolve more slowly, have lower levels of ionization, and show P Cygni absorption components. Novae with stronger lines of He and N have larger expansion velocities and a higher level of ionization, and the lines are more flat-topped with little absorption. Only a few X-ray observations of He/N novae exist and because of their rapid evolution little is known about them.


The nova V2491 Cyg (Nova Cyg 2008b) was discovered on  $t_0=2008$  April 10.8 UT at about 7.7 mag on unfiltered CCD frames (Nakano et al. 2008). Tomov et al. (2008a) found an optical V/R/I decline rate from  $\sim 0.3$  to 0.15 mag per day between  $t_0 + 2.3$  and  $t_0 + 7.3$  days. Assuming  $t_0 + 0.6$  and  $V=7.54$

mag for the maximum (Nakano et al. 2008), the time  $t_2$  in the V band is 4.6 days, where  $t_2$  is defined as the elapsed time to decrease 2 magnitudes in its visual luminosity. This  $t_2$  value makes V2491 Cyg a very fast nova, similar to V838 Her (O'Brien et al. 1994) and V2487 Oph (Hernanz & Sala 2002). Optical post-outburst spectra showed V2491 Cyg to be a nova in its early phase of outburst (Ayani & Matsumoto 2008, Lynch et al. 2008). V2491 Cyg has been classified as a He/N nova based on the photometric and spectroscopic results (Lynch et al. 2008, Tomov et al. 2008b, Helton et al. 2008). The very fast decline and the optical spectral characteristics, such as extremely broad lines with complex profiles, and large expansion velocities ( $\sim 4000$ – $6000$  km s<sup>-1</sup>), mark V2491 Cyg as a peculiar and extremely fast nova (Ashok et al. 2008, Lynch et al. 2008, Tomov et al. 2008a, 2008b).

From optical measurements the reddening towards the source has been determined:  $E(B-V)=0.3$  (Lynch et al. 2008) and  $E(B-V)=0.43$  (Rudy et al. 2008) on  $t_0+2$  and  $t_0+7$  days, respectively. Taking the latter value and using the correlation between visual extinction,  $A_V$ , and the dust (and hydrogen) column densities, assuming there is no intrinsic absorption ( $N_H$  [cm<sup>2</sup>]/ $A_V = 1.79 \times 10^{21}$ , e.g., Predehl & Schmitt 1995), and using  $A_V = 3.1E(B-V)$ , the optical reddening corresponds to  $N_{\text{dust/H}} \approx 2.4 \times 10^{21}$  cm<sup>-2</sup>. Using the reddening of  $E(B-V)=0.43$  a distance of 10.5 kpc was estimated by Helton et al. (2008).

The object was fainter than 14 mag at  $t_0-2$  days (Nakano et al. 2008). Balman et al. (2008) did not detect the pre-nova

\* INSA (Ingeniería y Sistemas Aeroespaciales S.A.)




**Fig. 1.** *Swift*/XRT X-ray image of the V2491 Cyg field of view. The big circle corresponds to the *Swift*/BAT uncertainty and the zoomed view corresponds to the pre-nova X-ray counterpart uncertainties of Table 1. The cross corresponds to the optical coordinates.

down to  $R=18.6\pm 0.5$ , 5–9 months prior to the outburst. However, the pre-nova was detected in archival plates spanning  $\sim 16$  years showing a persistent source at  $R\approx 16.3$  (Jurđana-Sepić & Munari 2008). This indicates a dimming of the source by about 2 mag several months before the nova outburst. A search through archival X-ray data showed the presence of an X-ray source, before the nova outburst, at the nova position (Ibarra & Kuulkers 2008, Ibarra et al. 2008). We here describe in more detail the serendipitous X-ray observations performed by *ROSAT*, *XMM-Newton* and *Swift* of the pre-nova. Preliminary results have already been reported by Ibarra & Kuulkers (2008) and Ibarra et al. (2008). Since the discovery, follow-up X-ray observations have been made with *Swift* (Kuulkers et al. 2008, Page et al. 2008, Osborne et al. 2008, Page et al. in prep), *XMM-Newton* (Ness et al. 2008a, 2008b, in prep.) and *Suzaku* (Takei et al., in prep.).

## 2. X-ray observations

A search through the X-ray catalogues (Available through VizieR at the Centre de Données de Strasbourg (CDS)) showed that the field of V2491 Cyg was observed at different epochs before the nova outburst by *ROSAT*, *XMM-Newton* and *Swift*. We determine the *Swift* position by combining all the data into one single image and using the *ximage* tool. In Table 1, we list the positions and position errors of the candidate X-ray counterparts. In Fig. 1, we show a *Swift*/XRT image with all the pre-nova X-ray counterpart uncertainties. The following paragraphs describe these satellites and the relevant instruments. The results are summarised in Table 2.

The *ROSAT* satellite (Trümper et al. 1991) produced an all-sky survey (RASS) in the energy range 0.2–2.4 keV and covered about 25% of the sky during pointing observations with the PSPC camera (Voges et al. 2000). Two X-ray sources in these catalogs have coordinates close to that of V2491 Cyg, i.e., the RASS source 1RXS J194259.9+321940 and the source 2RXP J194302.0+321912 (see also Table 1 and Fig. 1). We have re-analyzed the 2RXP J194302.0+321912 observation using the HEASOFT (<http://heasarc.nasa.gov/docs/software/lheasoft/>) software suite. We filtered the *ROSAT* event list with XSELECT,



**Fig. 2.** *ROSAT*/PSPC (one) and *Swift*/XRT (three) spectra of pre-nova X-ray observations. Shown are the data points for all the observations.

using an 80'' source extraction region and a local background region.

*XMM-Newton* (Jansen et al. 2001) is performing a sensitive survey of the sky in the 0.2–12 keV energy band, when it slews between targets. A catalogue of slew point sources has been published by Saxton et al. (2008).

We report on *XMM-Newton* observations by the EPIC-pn camera (Strüder et al. 2001) operating in full-frame mode with the medium filter during a slew performed on 2006 November 15 (OBSID. 9127000003). V2491 Cyg passed through the EPIC-pn field of view at a large off-axis angle such that an effective on-axis exposure time of 3.4 s was achieved. A source with 4 photons was detected in the full (0.2–12 keV) energy band of the EPIC-pn with a significance of  $3.5\sigma$  above the very low background (0.2 cts/arcmin<sup>2</sup>) of the slew image. There were no *XMM-Newton* pointed observations of this field before outburst.

Six *Swift* observations were made of the V2491 Cyg field before the nova outburst, as part of a follow-up survey aimed at identifying the X-ray counterpart for sources discovered by the Burst Alert Telescope (BAT; Barthelmy et al. 2005). The survey (Tueller et al. 2008; Tueller et al. in prep.) pointing was towards Swift J1942.8+3220 (Sect. 3.3).

The *Swift* X-ray Telescope (XRT, 0.3–10 keV; Burrows et al. 2005) detector was operated in photon counting (PC) mode, which provides two-dimensional imaging, spectral information, and 2.5 s time resolution. All the *Swift*/XRT data have been processed through XRTPipeline (Swift\_Rel2.8), with local-background subtracted spectra binned to  $\geq 20$  cts bin<sup>-1</sup>. One of the XRT observations was too short (20 s on 2007 July 15) to be useful, and has not been included in our spectral analysis.

No optical *Swift*/UVOT data were available. This is because the UVOT telescope was operated with BLOCKED filter due to the presence of a bright star in the field of view.

## 3. Results

### 3.1. The pre-nova X-ray identification

In Table 1 we give the positions and source names of the X-ray sources we identify close to the optical position of V2491 Cyg. Based on the *Swift*/XRT observations, the pre-nova X-ray source has been designated as Swift J194302.1+321913 (Ibarra et al. 2008). The *ROSAT*/PSPC source, 2RXP J194302.0+321912, the

**Table 1.** Pre-nova X-ray counterpart source identifications, coordinates, uncertainty, and offset ( $\delta$ ) from the position of the optical nova (19h43m01.96s, +32d19'13.8"; Nakano et al. 2008).

name	position (RA,Dec; J2000.0)	error (90%)	$\delta$
1RXS J194259.9+321940	19h42m59.9s, +32d19'40.5"	28"	37"
2RXP J194302.0+321912	19h43m02.0s, +32d19'12.0"	4"	1.9"
XMMSL1 J194301.9+321911	19h43m02.0s, +32d19'10.5"	12"	3.7"
Swift J194302.1+321913	19h43m02.0s, +32d19'11.0"	3.7"	2.8"

*XMM-Newton*/EPIC-pn source XMMSL1 J194301.9+321911, and the *Swift*/XRT source Swift J194302.1+321913 are all consistent with the optical nova coordinates (see also Ibarra & Kuulkers 2008, Ibarra et al. 2008). The *ROSAT*/PSPC source 1RXS J194259.9+321940 is formally not consistent with the optical position (see Fig. 1, but due to its large uncertainty we consider that this is compatible with 2RXP J194302.0+321912. Based on the positional information, as well as the similar strength of the sources over time (see next subsection), we suggest that all the above mentioned X-ray sources are all the same source, and identify it as the X-ray counterpart to the pre-nova of V2491 Cyg.

### 3.2. Pre-nova X-ray spectral characteristics

In Table 2, we list all X-ray pre-nova observations; for *Swift*/XRT we provide the background corrected count rates in two bands, and the 68% confidence Bayesian rate errors (Kraft et al. 1991) because of the low numbers of counts. The source is clearly variable. Based on the information in the two energy bands it changes dramatically in spectral shape between observations on a time scale down to 4 days. Especially noticeable is the softness of the source on 2007 July 21 (see also Fig. 1).

The second *ROSAT* and last three *Swift* observations provide sufficient counts for a more detailed X-ray spectral analysis. We first used a power-law model to describe our data and fit them with XSPEC version 11.3. Using this model and leaving  $N_{\text{H}}$  free in the fits led to poorly constrained values, especially for  $N_{\text{H}}$ . To better constrain  $N_{\text{H}}$  we did a simultaneous fit to the four individual spectra, using the same value of  $N_{\text{H}}$  while leaving the power-law parameters independent. This resulted in a reasonable fit ( $\chi^2_{\text{red}}=1.2$  with 10 d.o.f) with  $N_{\text{H}}=2.2^{+1.3}_{-1.1} \times 10^{21} \text{ cm}^{-2}$  (90% statistical errors are quoted for all parameters). The photon index values are:  $3.6^{+0.7}_{-0.8}$  for the *ROSAT* observation and  $1.7^{+0.5}_{-0.4}$ ,  $5.2^{+1.2}_{-0.7}$ ,  $1.2 \pm 0.4$  for the last three *Swift* observations, respectively. Our value of  $N_{\text{H}}$  is compatible with that derived from the optical measurements (see Sect. 1). The softest spectrum has a very steep power-law index, compared to the other spectra. Although spectra of novae are known to show many emission lines (e.g., Mukai & Orio 2005), X-ray spectral fits using other models (e.g., the optically-thin thermal plasma model VAPEC, with abundances of C, Fe fixed at 0.3 solar and N fixed at 8 times solar, typical for novae (Nussbaumer et al. 1988, Ness et al. 2005) or Bremsstrahlung) did not provide improved fits. We also stacked all *Swift* observations (excluding the one taken 265 days before outburst; see Table 2) to get better statistics. Again, fits to the resulting spectrum show a slightly better reproduction of the data when using a power law compared to a VAPEC model. Unfortunately, the quality of the data prevents us from investigating in more detail multi-temperature emission-line models.

Using the best-fit power-law spectral fits, we derive unabsorbed X-ray (0.2–10 keV) flux values between 1 and

$30 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ , taking into account the observations from the various instruments.

### 3.3. The unidentified source Swift J1942.8+3220

It is worth considering whether V2491 Cyg might be the counterpart to the *Swift*/BAT detection Swift J1942.8+3220, whose error circle encompasses the nova position, see Fig. 1. The *Swift*/BAT survey catalog source Swift J1942.8+3220 has an estimated average flux of about  $2 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$  (14–195 keV; Tueller et al. 2008). Using public archival *INTEGRAL*/IBIS/ISGRI observations performed between March 2003 and December 2006 (for a total exposure time of about 620 ksec) we derive a  $3\sigma$  upper limit (18–50 keV) of about  $1 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$  at the position of V2491 Cyg, more or less comparable with that derived for the *Swift*/BAT source.

Apart from Swift J194302.1+321913 there is another XRT source within the  $\sim 5'$  BAT error circle of Swift J1942.8+3220 (see upper part of Fig. 1): Swift J194245.9+322411 (RA = 19h42m45.9s, Dec = +32d24'10.7", with 90% confidence error radius of 3.6"). This source has an absorbed AGN-like X-ray spectrum; a power-law fit gives  $\Gamma=1.5^{+0.3}_{-0.4}$  and  $N_{\text{H}} = (2 \pm 1) \times 10^{21} \text{ cm}^{-2}$  with an unabsorbed 0.3–10 keV flux of  $1.1 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ . Our above described *ROSAT* and *XMM-Newton* identifications are not compatible with this AGN-like XRT source. The position of this source was covered by the *XMM-Newton* Slew Survey, but no source was detected with a  $2\sigma$  upper limit of  $2 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$  (0.2–12 keV), assuming the AGN-like XRT spectrum. The AGN-like XRT source is not bright enough to account for the BAT detection, assuming it is constant, and so it is not likely to be the X-ray counterpart to the BAT source. If, on the other hand, we extrapolate the power-law fits of the *Swift*/XRT spectra of the prenova V2491 Cyg taken on 2007 July 17 and 2008 Jan 2 (i.e., when the source was hard) into the BAT band, we get fluxes which are more or less consistent with the BAT measured flux. If, however, the spectrum is assumed to be due to optically thin high-temperature thermal emission, the extrapolated flux is a few orders of magnitude lower. It remains, therefore, uncertain whether the BAT source is related to the pre-nova.

## 4. Discussion

We find a persistent, but variable, and at least on one occasion very soft X-ray source present at the position of the nova V2491 Cyg from the *ROSAT* survey era up to 3 months before the nova outburst. This is only the second nova to be detected in X-rays before eruption (after V2487 Oph, Hernanz & Sala 2002). A hard spectral component is suggested by *Swift*/BAT, but the association with the pre-nova is not secure.

V2487 Oph, also a fast nova ( $t_2 \approx 6.3$  days, see Hernanz & Sala 2002), was suggested to be a recurrent nova, because of both the rapid decline in the optical and the presence of a plateau

**Table 2.** Observation log of the *ROSAT*, *XMM-Newton* and *Swift* pre-nova observations. Apart from the exposure time the count rates in different energy bands as well as the absorbed fluxes are given (see text for more details). For the second *ROSAT* observation, the *XMM-Newton* and the first two *Swift* observations, the flux was calculated using a power-law with  $\Gamma$  between 1.3 and 5.2, and  $N_H = 2.2 \times 10^{21} \text{ cm}^{-2}$ .

Date	Days before outburst	Instrument	Exposure time (ks)	Count rate <sup>1</sup> (cts/ks)	Count rate (cts/ks) 0.3–1.0 keV	Count rate (cts/ks) 1.0–8.0 keV	0.2–10.0 keV flux absorbed ( $10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ )
1990-10-19	6383	<i>ROSAT</i> /PSPC[1RXS]	0.39	28±11	—	—	4.5 <sup>+19.5</sup> <sub>-1.8</sub>
1993-10-05	5302	<i>ROSAT</i> /PSPC[2RXP]	4.3	22±3	12±2	8±2	0.3±0.1
2006-11-15	517	<i>XMM-Newton</i> /EPIC	3.43×10 <sup>-3</sup>	1173 <sup>+613</sup> <sub>-587</sub>	—	—	2.5 <sup>+7.0</sup> <sub>-1.6</sub>
2007-05-25	322	<i>Swift</i> /XRT	1.06	21±5	1.9 <sup>+1.9</sup> <sub>-1.3</sub>	20 <sup>+5</sup> <sub>-4</sub>	5.5 <sup>+8.5</sup> <sub>-1.4</sub>
2007-06-10	306	<i>Swift</i> /XRT	0.97	41±7	11.8 <sup>+4.2</sup> <sub>-3.6</sub>	31 <sup>+7</sup> <sub>-6</sub>	10.7 <sup>+16.5</sup> <sub>-2.2</sub>
2007-07-17	269	<i>Swift</i> /XRT	3.82	23±3	3.5 <sup>+1.5</sup> <sub>-1.1</sub>	19±3	1.0±0.2
2007-07-21	265	<i>Swift</i> /XRT	4.87	37 <sup>+3</sup> <sub>-4</sub>	27.9±3.2	9±2	0.8±0.2
2008-01-02	100	<i>Swift</i> /XRT	5.65	24±2	3.9±1.0	20±2	1.6 <sup>+0.5</sup> <sub>-0.4</sub>

Note 1. Energy bands: *ROSAT*/PSPC, 0.2–2.0 keV; *XMM-Newton*/EPIC, 0.2–12 keV; *Swift*/XRT, 0.3–10 keV.

phase during the decline (Hachisu et al. 2002). Pagnotta et al. (2008) discovered a previous outburst of V2487 Oph in 1900, confirming its recurrent nature. V2491 Cyg has been suggested to be a recurrent nova also (Tomov et al. 2008b).

Assuming a distance of 10.5 kpc (Helton et al. 2008) and using the power law fit results, we derive 0.2–10 keV X-ray luminosities ranging from about  $1 \times 10^{34}$  to  $4 \times 10^{35} \text{ erg s}^{-1}$ . This is comparable to that derived for V2487 Oph, about 3 years after the outburst, i.e.,  $\approx 8 \times 10^{34} \text{ erg s}^{-1}$  (0.2–10 keV; Hernanz et al. 2008) assuming a distance of 10 kpc (but note that the distance is rather uncertain, see Hernanz & Sala 2002). It is, however, orders of magnitude higher than that seen for the recurrent nova RS Oph about 2 years after the 2006 outburst ( $\approx 5 \times 10^{31} \text{ erg s}^{-1}$ , Nelson & Orio 2008, using a distance of 1.6 kpc, Mason et al. 1987, Bode et al. 2008). This is compatible with the fact that in general fast novae appear to be brighter than slow novae, when they are at quiescence (e.g., Becker & Marshall 1981, Orio et al. 2001).

Our X-ray luminosity estimates imply inter-nova mass accretion rates in the range  $10^{-9}$  -  $10^{-8} M_{\odot} \text{ yr}^{-1}$ , for a  $1 M_{\odot}$  white dwarf. These rates are around an order of magnitude lower than those required (judged from the models of Yaron et al. 2005) to fuel the novae outbursts of RS Oph, which are recurrent on a  $\sim 20$  year period. This would imply that, even with a massive white dwarf, the recurrence timescale for novae in V2491 Cyg would be  $> \sim 100$  yrs, similar to V2487 Oph.

The soft spectra observed with *ROSAT*/PSPC and *Swift*/XRT resemble those of the post-outburst super-soft state (Page et al. 2009, in prep.), but are much fainter. This, and the fact that the *Swift*/XRT observation was taken less than a year before the nova outburst, suggests that the spectra were not obtained during a super-soft state after a previous nova outburst or accretion driven nuclear burning, in agreement with our relatively low mass transfer estimates. The *Swift*/XRT observations show that V2491 Cyg changes spectral state on at least a 4-day time scale.

Magnetic CVs in quiescence show harder spectra than their non-magnetic equivalents (e.g., Barlow et al. 2008, Landi et al. 2009). If the BAT source is the same as that seen at  $\leq 10$  keV, V2491 Cyg may be magnetic. Polars, for example, are known to change from hard to soft states as the white dwarf rotates (e.g., Heise et al. 1985), which could explain the X-ray spectral evolution we observe. However, we do not see evidence for short-term ( $\sim$ hour) orbital related variations. Moreover, in general polars are rather weak X-ray sources (see King & Watson 1987). Additionally, at  $\leq 10$  keV the spectrum is not unlike that

seen in non-magnetic CVs (e.g., Baskill et al. 2005). This brings into doubt a magnetic interpretation.

*Acknowledgements.* We acknowledge the use of observations obtained with *XMM-Newton*, an ESA science mission with instrument and contributions directly funded by ESA Member States and the USA (NASA), as well as public data from the *Swift* data archive. JO and KP acknowledge the support of the STFC; SS acknowledges partial support from NASA and NSF grants to ASU. We made use of the *XMM-Newton* archive tool, ESAC/ESA, Madrid, Spain. Also, we made use of the VizieR catalogue access tools, CDS, Strasbourg, France.

## References

- Ashok, D.P., Barnerjee, D.P.K., Joshi, V., Naik, S. 2008, CBET 1354  
Ayani, K., Matsumoto, K. 2008, CBET 1334  
Balman, S., Pekon, Y., Kiziloglu, U. 2008, ATel # 1504  
Barlow, E.J., et al. 2006, MNRAS, 372, 224  
Barthelmy, S.D., et al. 2005, SSRv, 120, 143  
Baskill, D.S., Wheatley, P.J., Osborne, J.P. 2005, MNRAS, 357, 626  
Becker, R.H., Marshall, F.E. 1981, ApJ, 244, L93  
Bode, M.F., Evans, A. 2008, in Classical Novae, CUP  
Burrows, D.N., et al. 2005, SSRv, 120, 165  
Hachisu, I. 2002, ASP Conf. Ser., 261, 629  
Heise, J., et al. 1985, A&A, 148, L14  
Helton, L.A., Woodward, C.E. 2008, CBET, 1379  
Hernanz, M., Sala, G. 2002, Science 298, 393  
Hernanz, M. 2008, in The X-ray Universe Symposium, [http://xmm.esac.esa.int/external/xmm\\_science/workshops/2008symposium/](http://xmm.esac.esa.int/external/xmm_science/workshops/2008symposium/)  
Ibarra, A., Kuulkers, E. 2008, ATel # 1473  
Ibarra, A., et al. 2008, ATel # 1478  
Jansen, F., et al. 2001, A&A, 365, L1  
Jurdana-Sepic, R., Munari, U. 2008, IBVS, 5839  
King, A.R., Watson, M.G., 1987, MNRAS, 227, 205  
Kraft, R.P., Burrows, D.N., Nousek, J.A. 1991, ApJ, 374, 344  
Landi, R., et al. 2009, MNRAS, 392, 630  
Lynch, D.K., Russell, R.W., Rudy, R.J. 2008, IAU Circ. 8935  
Kuulkers, E., et al. 2008, ATel # 1480  
Mason, K., et al. 1987, in: RS Oph (1985) and the Recurrent Nova Phenomenon, M.F. Bode (ed.), VNU Science Press (Utrecht), 167  
Mukai, K., Orio, M. 2005, ApJ, 622, 602  
Nakano, S., et al. 2008, IAU Circ. 8934  
Nelson, T., Orio, M. 2008, ATel # 1494  
Ness, J.-U., et al. 2008a, ATel # 1561  
Ness, J.-U., et al. 2008b, ATel # 1573  
Ness, J.-U. et al., 2005, MNRAS, 364, 1015  
Nussbaumer, H. et al. 1988, A&A 198, 179  
O'Brien, T.J., Lloyd, H.M., Bode, M.F. 1994, MNRAS, 271, 155  
Orio, M., Covington, J., Ögelman, H. 2001, A&A, 373, 542  
Osborne, J.P., et al. 2008, ATel # 1542  
Page, K.L., et al. 2008, ATel # 1523  
Pagnotta, A., Schaefer, B. E., Xiao, L. 2008, IAU Circ. 8951  
Predehl, P., Schmitt, J.H.M.M. 1995, A&A, 293, 889  
Rudy, R.J., Lynch, D.K. Russell, R.W. 2008, IAU Circ. 8938  
Saxton, R.D., et al. 2008, A&A, 480, 661

Strüder, L., et al. 2001, A&A, 365, 18  
Tomov, T., et al. 2008a, ATel # 1475  
Tomov, T., et al. 2008b, ATel # 1485  
Trümper, J., et al. 1991, Nat, 349, 579  
Tueller, J., et al. 2008, ApJ, 681, 113  
Voges, W., et al. 2000, IAUC 7432  
Williams, R.E. 1992, ApJ, 104, 725  
Yaron, O. et al. 2005, ApJ, 623, 398