

Clumps of material orbiting a black hole and the QPOs

Uroš Kostić, Andrej Čadež, and Andreja Gomboc

Department of Physics, Faculty of Mathematics and Physics, University of Ljubljana, Jadranska 19, 1000 Ljubljana andrej.cadez@fmf.uni-lj.si

1 Successive passages of an asteroid about a black hole

Clumps of material, the size of asteroids, may orbit a black hole and be distributed, somewhat like comets in the Kuiper belt. Some clumps are perturbed to relatively small periastron high eccentricity orbits. During each periastron passage tides do work and so change orbital parameters to make the orbit more eccentric with less orbital angular momentum and lower periastron. The periastron crossing time decreases accordingly. When the periastron touches the Roche radius, tides resonate with the fundamental quadrupole mode of the asteroid [1]. The transfer of orbital angular momentum to internal tidal modes is high and the work done by tides is large enough to heat the asteroid to high temperatures and to break loose some parts of the asteroid. Each next periastron passage has a shorter characteristic time and excites higher mechanical modes of the asteroid. The orbital energy keeps decreasing and the transferred energy is used to heat it to higher and higher temperatures and to accelerate more breaking-away parts. Soon the asteroid is split into small pieces which are sufficiently small that they are stable against tidal destruction. When the periastron reaches close to the last circular orbit, the overwhelming tidal force crushes all solids that are larger than about $0.01GM/c^2$ (here M is the mass of the black hole; $GM_{\odot}/c^2 = 1.5$ km) and produces enough work to heat the remnants to hard X-ray temperatures.

The overwhelming tidal force is very rapid with respect to the frequency of fundamental modes of the asteroid, so that the work done by tides can be as high as $0.1mc^2$, where m is the mass of the asteroid. Even a small part of this energy is more than enough to make the asteroid explode. Yet, the shock wave is still slow with respect to asteroid's relativistic speed, so that it has no time to grow large with respect to the black hole. Since the work is taken out of orbital energy, the remnants can no longer return to the far apastron, but are caught by the black hole and after a few turns about the black hole at the last circular orbit, they disappear behind its horizon. In making these few last turns they alternately move towards and away from the observer close to asteroid's orbital plane, so that the Doppler effect, aberration of light and possibly gravitational lensing make characteristic peaks (a chirp) in the light curve of this last orbit as it is seen by the observer.

2 Fourier spectra of orbiting clumps and QPOs

To investigate what kind of the observed light curve one would obtain from infalling debris of tidally disrupted asteroid, we assume that a nonrotating black hole is fed by a large number of small clumps and calculate light curves from such events and their Fourier spectra.




Fig. 1. *Left:* A typical chirp: light curve produced by a small ($r = 0.01GM/c^2$) solid body on its way to the black hole following a parabolic orbit with angular momentum $l/m = 3.999998GM/c$ which makes three turns before crossing the horizon. In this example the body's velocity, when far from the black hole, subtends an angle 24° with the line of sight and the observer is located 5° above the orbital plane. The sharp peaks in observed luminosity are the effect of gravitational lensing. The wide bumps following those peaks are due to aberration of light and Doppler shift. *Right:* The Fourier spectrum of the chirp on the left. Note that the frequency scale is in kHz, divided by the mass of the black hole in solar units; for example, a signal from a 10 solar mass black hole would have the lowest peak frequency at $\nu \sim 400\text{Hz}$. This frequency corresponds to the orbital period on the last circular orbit about the black hole.




Fig. 2. A light curve (left) and the Fourier spectrum (right) of a superposition of 24 chirps produced by 24 particles orbiting on $l/m = 3.999998GM/c$ parabolic orbits distributed randomly in time and with respect to the angle of the line of nodes. The observer is located 5° above the orbital plane.




Fig. 3. Left: A signal produced by 178 objects falling along $l/m = 3.999998GM/c$ parabolic orbits onto the black hole from random directions and at random times. Right: Fourier spectrum of this signal. The broad peaks appear at harmonics of the frequency corresponding to the period of the last circular orbit, but their amplitudes are again modulated by random factors determined by the relative timing of contributing chirps. Another random series of chirps would produce a similar, but not the same power spectrum.

3 Conclusions

Accretion of solid asteroid-like material onto a black hole can produce signals with signature similar to that of QPOs. Small solid particles are the only form of matter that can survive huge tidal forces deep in the gravitational field of a stellar mass black hole. They can enter the last circular orbit almost intact and transform the mounting tidal force into internal energy. Since the theoretical upper limit for this energy increase is $\sim 0.1mc^2$, it seems plausible that the body heats to hard X-ray temperatures when making a few last turns before plunging behind the horizon. Such a signal is modulated by Doppler shift, aberration of light and gravitational lensing and produces a characteristic chirp. In case of Schwarzschild black hole, all signals have the same characteristic frequency $1/50[c^3/GM]$, but the higher frequencies depend somewhat on the orientation of the orbit with respect to the observer. Many objects produce a signal which is very similar to that of a QPO [2, 3].

References

1. A. Gomboc, A. Čadež: *Astrophys. J.* **625**, 278 (2005)
2. G. L. Israel, et al.: *Astrophys. J. Lett.* **628**, L53 (2005)
3. J. Homan, J. M. Miller, R. Wijnands, M. van der Klis, T. Belloni, D. Steeghs, & W. H. G. Lewin: *Astrophys. J.* **623**, 383 (2005)