HOW RANDOM ARE A LEARNER'S MISTAKES ?

JOEL RATSABY

ABSTRACT. Given a random binary sequence $X^{(n)}$ of random variables, X_t , $t = 1, 2, \ldots, n$, for instance, one that is generated by a Markov source of order k^* (each state represented by k^* bits). Let β be the probability of " $X_t = 1$ " and assume it is constant with respect to t (due to stationarity). Consider a learner based on a parametric model, for instance a Markov model of order k, who trains on a sample sequence $x^{(m)}$ which is randomly drawn by the source. Test the learner's performance by giving it a sequence $x^{(n)}$ (generated by the source) and check its predictions on every bit of $x^{(n)}$. An error occurs at time t if the prediction Y_t differs from the true bit value X_t . Denote by $\xi^{(n)}$ the sequence of errors where the error bit ξ_t at time t equals 1 or 0 according to whether the event of an error occurs or not, respectively. Consider the subsequence $\xi^{(\nu)}$ of $\xi^{(n)}$ which corresponds to the errors of predicting a 0, i.e., $\xi^{(\nu)}$ consists of the bits of $\xi^{(n)}$ only at times t such that $Y_t = 0$. In this paper we compute an upper bound on the deviation of the frequency of 1s of $\xi^{(\nu)}$ from β showing dependence on k, m, v.

1. Overview

Let $\{X_t : t = 1, 2, \ldots\}$ be a sequence of Bernoulli random variables possessing the following Markovian property,

$$
P(X_t = x | X_1 = x_1, ..., X_{t-1} = x_{t-1}) = P(X_t = x | X_{t-k} = x_{t-k}, ..., X_{t-1} = x_{t-1})
$$
\n(1.1)

for some fixed k and where x_{t-k}, \ldots, x_{t-1} , and x are 0 or 1, and $t = k + 1, k + 2$, \cdots . The model is known as the k^{th} order Markov chain and we denote it by \mathcal{M}_k . Let us define by $\mathbb{S}_k = \{0,1\}^k$ the state space of \mathcal{M}_k and let $s^{(i)} \in \mathbb{S}_k$ denote the i^{th} state, $i = 0, 1, ..., 2^k - 1$. Denote by $\{S_t : t = k, k + 1, ...\}$ the sequence of random state variables where

$$
S_t := (X_{t-k+1}, X_{t-k+2}, \dots, X_t)
$$

represents the state at time t. Let $T_k = [t_{i,j}], 1 \leq i, j \leq 2^k$, be a state-transition matrix associated with model \mathcal{M}_k with elements $t_{i,j} := P(S_{t+1} = s^{(j)} | S_t = s^{(i)}),$ $0 \leq i, j \leq 2^k - 1$. Clearly, the structure of \mathcal{M}_k allows for only two possible outgoing transitions from a state S_t to the next state S_{t+1} since S_{t+1} can take only one of the two values $(X_{t-k+2}, \ldots, X_t, 0)$ or $(X_{t-k+2}, \ldots, X_t, 1)$. We call them type-0 and type-1 transitions. Using \mathcal{M}_k to produce a random sequence is done by randomly drawing a state sequence and outputting 1 or 0 for each type-1 or type-0 transition, respectively.

In this paper we consider the following estimation problem:

Estimation problem: A source produces a *data* string $x^{(m+n)} = (x_1, \ldots, x_{m+n}),$ by randomly drawing $m + n$ consecutive bits according to a Markov chain \mathcal{M}_{k^*} of order k^* . A learner (not knowing the value of k^*) estimates the parameters of another model \mathcal{M}_k based on the initial subsequence $x^{(m)} = (x_1, \ldots, x_m)$ which is

²⁰⁰⁰ Mathematics Subject Classification. 62M20, 62M05.

Key words and phrases. Prediction by Markov Models, Large deviation, Error convergence.

called *training* sequence. Afterwards, the learner uses \mathcal{M}_k to make a prediction y_t for each of the remaining bits x_{m+t} , $t = 1, 2, \ldots, n$, which form the testing sequence $x^{(n)} = (x_{m+1}, \ldots, x_{m+n})$. Denote by $\xi^{(n)} = {\{\xi_t\}}_{t=1}^n$ the corresponding binary sequence of mistakes where $\xi_t = 1$ if $y_t \neq x_{m+t}$ and is 0 otherwise. Denote by $\xi^{(\nu)} = \left\{ \xi_{i_j} \right\}_{j=1}^{\nu}, \nu \geq \ell$, a subsequence of $\xi^{(n)}$ with time instants i_j corresponding to 0-predictions, $y_{i_j} = 0, 1 \leq j \leq \nu$. Note that $\xi^{(\nu)}$ is also a subsequence of the input sequence $x^{(n)}$ hence effectively the learner acts as a selection rule Γ_d which picks certain bits $\xi^{(\nu)}$ from $x^{(n)}$ according to an algorithm that specifies its prediction rule. In this paper we compute a large deviation bound for the error sequence $\xi^{(\nu)}$.

2. Introduction

From basic theory on finite Markov chains, since the matrix $T := T_{k^*}$ is stochastic (i.e., the sum of the elements in any row equals 1) then \mathcal{M}_{k^*} has a stationary joint probability distribution

$$
\mathbb{P}\left(X^{(n)}\right) \quad := \quad P(X_1,\ldots,X_n)
$$

which is not necessarily unique. To keep the notation simple we use $\mathbb P$ to denote also any marginal distribution derived from the stationary joint distribution. For instance, $\mathbb{P}(X_1, X_2, X_3) = \sum_{(x_4,...,x_n)} \mathbb{P}(X_1, X_2, X_3, X_4 = x_4..., X_n = x_n)$. Henceforth, all random binary sequences are assumed to be drawn according to this probability distribution P. Thus for any k and ℓ satisfying $\ell > k > 1$ the probability of a string $x^{(\ell)} = (x_1, x_2, \dots, x_\ell)$ can be expressed as

$$
\mathbb{P}(X_1 = x_1, X_2 = x_2, \dots X_\ell = x_\ell) \n= \mathbb{P}(S_k = (x_1, \dots x_k)) \mathbb{P}\left(S_{k+1} = (x_2, \dots x_{k+1}) \middle| S_k = (x_1, \dots x_k)\right) \n\cdots \mathbb{P}\left(S_\ell = (x_{\ell-k+1}, \dots x_\ell) \middle| S_{\ell-1} = (x_{\ell-k}, \dots x_{\ell-1})\right).
$$
\n(2.1)

Let us denote by

$$
\beta = \mathbb{P}\left(X_t = 1\right) \tag{2.2}
$$

the stationary probability of the event $\{X_t = 1\}$ at time $t = 1, 2, \ldots$.

Data generation: We henceforth assume that the source reached stationarity and produces the data sequence $x^{(m+n)}$ with respect to \mathbb{P} .

Consider the learner's model \mathcal{M}_k . Its set of parameters are the true (unknown) probability values of transitions between states in \mathbb{S}_k where the probability values are assigned according to the source distribution P. We denote them by

$$
p_{ij} := \mathbb{P}(S_{t+1} = s^{(j)} | S_t = s^{(i)}), \quad s^{(i)}, s^{(j)} \in \mathbb{S}_k.
$$

For instance, suppose $k^* = 3$ and $k = 2$ and consider two states $s^{(i)} = (0, 1)$ and $s^{(j)} = (1, 1)$. The corresponding transition probability is

$$
p_{i,j} = P((1,1) | (0,1)) = \frac{\mathbb{P}(X_{t-1} = 0, X_t = 1, X_{t+1} = 1)}{\mathbb{P}(X_{t-1} = 0, X_t = 1)}.
$$

Based on $x^{(m)}$ the learner estimates $p_{i,j}$ by

$$
\hat{p}_{ij} = \frac{m_{i,j}}{m_i}
$$

where for a state $s^{(i)} \in \mathbb{S}_k$, m_i denotes the number of times that $s^{(i)}$ appears in $x^{(m)}$ and $m_{i,j}$ denotes the number of times there is a transition from state $s^{(i)}$ to $s^{(j)}$ in $x^{(m)}$. For instance, if $k = 3$, $x^{(m)} = 011010101$ and $s^{(i)} = 101$ then $m_i = 3$. Thus \hat{p}_{ij} are the frequency of state-transitions in $x^{(m)}$. Note that m_i , $0 \le i \le 2^k - 1$, are dependent random variables since the Markov chain may visit each state a random number of times and they must satisfy $\sum_{i=0}^{2^k-1} m_i = m - k + 1$.

After training, the learner is tested on the remaining n bits of the data $x^{(n)} =$ $x_{m+1}, x_{m+2}, \ldots, x_{m+n}$. It makes a binary prediction Y_t for X_t , $t = m+1, \ldots, n$ based on the maximum a posteriori probability which is defined as follows: suppose that the current state is $s^{(i)} \in \mathbb{S}_k$ then the prediction is

$$
\hat{d}(i) := \begin{cases} 1 & \text{if } \hat{p}(1|i) > 1 - \hat{p}(1|i) \\ 0 & \text{otherwise,} \end{cases}
$$
\n(2.3)

where $\hat{p}(1|i)$ is defined as \hat{p}_{ij} for the state $s^{(j)}$ obtained from $s^{(i)}$ by a type-1 transition, i.e., if $s^{(i)} = (x_{\ell+1}, x_{\ell+2}, \ldots, x_{\ell+k})$ then $s^{(j)} = (x_{\ell+2}, x_{\ell+3}, \ldots, x_{\ell+k}, 1)$. The corresponding true probability value is denoted by $p(1|i) = p_{ij}$. Note that [\(2.3\)](#page-2-0) may be expressed alternatively as

$$
\hat{d}(i) = \begin{cases} 1 & \text{if } \hat{p}(1|i) > \frac{1}{2} \\ 0 & \text{otherwise.} \end{cases}
$$
 (2.4)

We claim that $\hat{p}(1|i)$, $0 \leq i \leq 2^k - 1$, are independent random variables when conditioned on the vector $\underline{m} := [m_0, \ldots, m_{2^k-1}]$. We now prove the claim which will be used in Section [3.](#page-4-0) Let us denote by $\sigma^{(m)} = (\sigma_k, \ldots, \sigma_m)$, $\sigma_i \in \mathbb{S}_k$, $k \leq i \leq m$, the particular sequence of states corresponding to the sequence $x^{(m)}$. To show the dependence of $x^{(m)}$ on $\sigma^{(m)}$ we will sometimes write $x^{(m)} = x(\sigma^{(m)})$. Then by (2.1) we have

$$
\mathbb{P}\left(X_1 = x_1, \ldots, X_m = x_m\right)
$$
\n
$$
= \mathbb{P}\left(S_k = \sigma_k\right) \mathbb{P}\left(S_{k+1} = \sigma_{k+1} \middle| S_k = \sigma_k\right) \mathbb{P}\left(S_{k+2} = \sigma_{k+2} \middle| S_{k+1} = \sigma_{k+1}\right)
$$
\n
$$
\cdots \mathbb{P}\left(S_m = \sigma_m \middle| S_{m-1} = \sigma_{m-1}\right).
$$

Since at every bit there are only two types of transitions then not every sequence $\sigma^{(m)} \in (\mathbb{S}_k)^{m-k+1}$ is possible. For instance, if $k = 3$ then the state sequence $(010, 100, 001)$ is valid but $(010, 110, 001)$ is not valid. Denote by $V \subset (\mathbb{S}_k)^{m-k+1}$ the set of *valid* state sequences $\sigma^{(m)}$. We now show that if $\sigma^{(m)}$ is in V then, conditioned on <u>m</u>, any other state sequence that visits the same states as $\sigma^{(m)}$ the same number of times (perhaps in a different order) must have the same probability. For any state $s^{(i)} \in \mathbb{S}_k$ denote by $N(1|i)$ the random variable whose value is the number of type-1 transitions from state $s^{(i)}$ in a sequence of random states $S^{(m)}$ = $S_k, S_{k+1}, \ldots, S_m$. Define by $N_{\sigma^{(m)}}(1|i)$ the number of type-1 transitions from state $s^{(i)}$ in the sequence $\sigma^{(m)}$. Since all state transitions are either type-0 or type-1 then we have

$$
\mathbb{P}\left(X^{(m)} = x\left(\sigma^{(m)}\right) \middle| \underline{m}, S^{(m)} = \sigma^{(m)}, \sigma^{(m)} \in V\right)
$$
\n
$$
= \prod_{0 \le i \le 2^{k}-1} \left(p(1|i)\right)^{N_{\sigma^{(m)}}(1|i)} \left(1 - p(1|i)\right)^{m_{i} - N_{\sigma^{(m)}}(1|i)} \tag{2.5}
$$

where $p(1|i)$ was defined above. Let α be a non-negative integer parameter and define the random variable $N(i) := [N(1|i), \alpha - N(1|i)]$. Associate a conditional probability function with parameter α for the random variable $N(i)$ as

$$
\mathbb{P}\left(N(i) = \left[\ell, \alpha - \ell\right] \middle| \alpha\right) = \left(p(1|i)\right)^{\ell} \left(1 - p(1|i)\right)^{\alpha - \ell}.
$$

Then the right side of (2.5) equals

$$
\prod_{0 \le i \le 2^k - 1} \mathbb{P}(N(i) = [N_{\sigma^{(m)}}(1|i), m_i - N_{\sigma^{(m)}}(1|i)]). \tag{2.6}
$$

For a fixed value of m_i the event " $N(i) = [N_{\sigma^{(m)}}(1|i), m_i - N_{\sigma^{(m)}}(1|i)]$ " is equivalent to the event " $\hat{p}(1|i) = \frac{N_{\sigma(m)}(1|i)}{m_i}$ ". Hence alternatively, the right side of [\(2.6\)](#page-3-0) can be expressed as

$$
\prod_{0 \le i \le 2^k - 1} \mathbb{P}\left(\hat{p}(1|i) = \frac{N_{\sigma^{(m)}}(1|i)}{m_i}\right).
$$
 (2.7)

The right side of [\(2.7\)](#page-3-1) is a product of probability functions of the random variables $\hat{p}(1|i)$. So conditioned on \underline{m} and on the event that $x^{(m)}$ corresponds to a valid state sequence $\sigma^{(m)}$, the event that $x^{(m)}$ is generated by the source Markov chain \mathcal{M}_{k^*} is equivalent to the event that its corresponding state sequence $\sigma^{(m)}$ has transition frequencies $\hat{p}(1|i)$ that independently take the particular values $\frac{N_{\sigma}(m)}{m_i}$ as prescribed in $x^{(m)}$. The claim is proved. It also follows that $\hat{p}(1|i)$ is the average of independent Bernoulli trials (success taken as a type-1 transition from state $s^{(i)}$). It is distributed according to the Binomial distribution with parameters m_i and $p(1|i)$.

We now summarize the problem setting under which the main result of the paper holds.

Problem setting: Let $0 < \delta < 1$ and k, ℓ, m, n be positive integers. Let $\mathbb P$ be the stationary probability distribution based on a finite, ergodic and reversible Markov chain with probability-transition matrix T that has a second largest eigenvalue λ . All probability values are measured according to P. Denote by $\gamma =$ $(1 - \max{0, \lambda}\)/(1 + \max{0, \lambda})$. After reaching stationarity the source generates a binary sequence $X^{(n)} = X_1, X_2, \ldots, X_n$ by repeatedly drawing X_t according to $\mathbb P$. Denote by $\beta = \mathbb P(X_t = 1)$. Let $x^{(m+n)}$ be a data-sequence obtained by randomly drawing according to \mathbb{P} . Let the learner's model \mathcal{M}_k be Markov of order k, and denote by $p(1|i)$ the probability of making a type-1 transition from state $s^{(i)}$ of \mathcal{M}_k . The learner uses the first m bits, $x^{(m)}$, to estimate $p(1|i)$ by $\hat{p}(1|i)$. Let m_i denote the number of times that state $s^{(i)}$ appears in $x^{(m)}$ by $\hat{p}(1|i)$. Let m_i denote the number of times that state $s^{(i)}$ appears in $x^{(m)}$, $\sum_{0 \le i \le 2^k-1} m_i = m - k + 1$. After training, the learner's decision at state $s^{(i)}$ is to output 1 if $\hat{p}(1|i) > \frac{1}{2}$ else output 0. Denote by μ_i the probability that a Binomial random variable with parameters m_i , $p(1|i)$, is larger (or smaller) than $\frac{m_i}{2}$ given that $p(1|i)$ is smaller (or larger) than $\frac{1}{2}$, respectively. Let $\mu = \frac{1}{2^k} \sum_{0 \le i \le 2^k - 1} \mu_i$. Let $\rho^{(m)} = \frac{1}{2^k} \sum_{0 \leq i \leq 2^k-1} \exp \left\{-2m_i \left(\frac{1}{2} - p(1|i)\right)^2\right\}$. Using \mathcal{M}_k the learner is tested incrementally on the remaining *n* bits $x^{(n)} = x_{m+1}, \ldots, x_{m+n}$ of the data and predicts an output bit y_t for bit x_t in $x^{(n)}$ to be 1 if $\hat{p}(1|i) > \frac{1}{2}$, else 0. Denote by $\xi^{(n)}$ the sequence of mistakes where $\xi_t = 1$ if $y_t \neq x_t$, and $\xi_t = 0$ otherwise, $m+1 \leq t \leq m+n$. Denote by $\xi^{(\nu)} = \left\{\xi_{i,j}\right\}_{j=1}^{\nu}, \nu \geq \ell$, the subsequence of $\xi^{(n)}$ with time instants i_j corresponding to 0-predictions, $y_{i_j} = 0, 1 \le j \le \nu$. Note that $\xi^{(\nu)}$ is also a subsequence of the input sequence $x^{(n)}$ hence effectively the learner acts as a selection rule which picks certain bits $\xi^{(\nu)}$ from $x^{(n)}$.

Let

$$
\epsilon^{2}(\ell, \gamma, \delta, k, m) := \frac{1}{2\ell\gamma} \left[2^{k} \left(\sqrt{\frac{1}{2^{k-2}} \ln\left(\frac{2}{\delta}\right)} + \rho^{(m)} \right) \ln\left(\frac{e}{\left(\sqrt{\frac{1}{2^{k-2}} \ln\left(\frac{2}{\delta}\right)} + \rho^{(m)}\right)} \right) + \ln 2 + \ln\left(\frac{4}{\delta}\right) \right]
$$

and assume that the learner's model order k satisfies,

$$
k \ge 2 + 2\log_2\left(\frac{1}{(2e-1)\mu}\right) + \log_2\left(\ln\left(\frac{2}{\delta}\right)\right).
$$

We now state the main result of the paper.

Theorem 1. For any $0 < \delta < 1$, with probability at least $1-\delta$ the deviation between

 β and the frequency of 1s of the sequence $\xi^{(\nu)}$ is bounded as

$$
\left|\frac{1}{\nu}\sum_{j=1}^{\nu}\xi_j^{(\nu)}-\beta\right| \leq \epsilon(\ell,\gamma,\delta,k,m).
$$

Before presenting the proof we make the following remarks,

- (1) The effect of the training sequence length m on ϵ is as $O(\rho^{(m)})$ which is $O(e^{-m})$. As m increases the class of possible learnt models (hypothesis class) decreases in size thereby decreasing the bound ϵ on the deviation of the error sequence.
- (2) The effect of the learner's model order k is opposite of that of m . We see that $\epsilon = O(2^{k/2})$ and as k increases, the hypothesis class increases in size.
- (3) The effect of the length ℓ of the error sequence on ϵ is as $O(\frac{1}{\ell})$. Clearly, the longer the subsequence the less chance that its frequency of 1s deviate from the mean β .
- (4) The effect of the inter-dependence between the states of the source model \mathcal{M}_{k^*} on ϵ is as $O(\frac{1}{\gamma})$. As the dependence increases, γ decreases which increases the possible deviation size ϵ . As γ decreases, the bits of the sequence $X^{(n)}$ become less dependent and ϵ decreases.

3. Proof of Theorem [1](#page-4-1)

A prediction decision-rule is denoted by a binary vector

$$
d := [d(0), \dots, d(2k - 1)].
$$
\n(3.1)

Note that d describes the prediction made by the learner at each state of the model.

Assume that the length m of the training sequence $x^{(m)}$ is fixed. Let us define by d^* the Bayes optimal decision. Clearly, $d^*(i) = 1$ when $p(1|i) > \frac{1}{2}$ and $d^*(i) = 0$ otherwise, $1 \leq i \leq 2^k$. Let us define the following set for $0 \leq r \leq 2^k$,

$$
A_r^{(k)} = \left\{ d \in \{0,1\}^{2^k} : \|d - d^*\| \le r \right\},\tag{3.2}
$$

where $||d||$ denotes the l₁-norm of d. Consider the Bernoulli random variable χ_i which equals 1 if $\hat{d}(i) \neq d^*(i)$ and 0 otherwise $(\hat{d}$ is defined in (2.3)). The event that $\chi_i = 1$ occurs if $\hat{p}(1|i) > \frac{1}{2}$ and $p(1|i) \leq \frac{1}{2}$ or if $\hat{p}(1|i) \leq \frac{1}{2}$ and $p(1|i) > \frac{1}{2}$. Since $\hat{p}(1|i)$ is the average of m_i i.i.d. Bernoulli random variables each with an expected value of $p(1|i)$ then we have in the case of $p(1|i) \leq \frac{1}{2}$ that

$$
\mathbb{P}\left(\hat{p}(1|i) > \frac{1}{2}\right) = \mathbb{P}\left(\hat{p}(1|i) - p(1|i) > \frac{1}{2} - p(1|i)\right) = \mathbb{P}\left(\hat{p}(1|i) > p(1|i) + \alpha_i\right)
$$

where

$$
\alpha_i = \frac{1}{2} - p(1|i).
$$

By Chernoff's bound [\[1\]](#page-9-0) we have

$$
\mathbb{P}\left(\hat{p}(1|i) > p(1|i) + \alpha_i\right) \le \exp\left\{-2m_i\alpha_i^2\right\}.
$$

Similarly, if $p(1|i) > \frac{1}{2}$ then, denoting by $\alpha_i = p(1|i) - \frac{1}{2}$, we have

$$
\mathbb{P}\left(\hat{p}(1|i) \leq \frac{1}{2}\right) = \mathbb{P}\left(\hat{p}(1|i) - p(1|i) \leq \frac{1}{2} - p(1|i)\right)
$$

$$
= \mathbb{P}\left(\hat{p}(1|i) \leq p(1|i) - \alpha_i\right)
$$

$$
\leq \exp\left\{-2m_i\alpha_i^2\right\}.
$$

Therefore, regardless of the value of $p(1|i)$, we have

$$
\mu_i := \mathbb{P}\left(\chi_i = 1\right) \le \rho_i^{(m)}\tag{3.3}
$$

where

$$
\rho_i^{(m)}:
$$
 = $\exp \{-2m_i \alpha_i^2\}, 1 \le i \le 2^k.$

As shown in the previous section, conditioned on \underline{m} , the $\hat{p}(1|i)$ are independent. Hence $\{\chi_i\}_{i=1}^{2^k}$ are independent non-identically distributed Bernoulli random variables (known as Poisson trials).

According to $\mathbb P$ the probability of the event that $\hat d$ is not in $A_r^{(k)}$ is the same as the probability of this event conditioned on the state sequence $\sigma^{(m)}$ being valid. Hence,

$$
\mathbb{P}\left(\hat{d} \notin A_r^{(k)}\right) = \mathbb{P}\left(\hat{d} \notin A_r^{(k)} \middle| \sigma^{(m)} \in V\right)
$$

$$
= \sum_{\underline{m}} \mathbb{P}\left(\hat{d} \notin A_r^{(k)} \middle| \underline{m}, \sigma^{(m)} \in V\right) \mathbb{P}\left(\underline{m} \middle| \sigma^{(m)} \in V\right) \quad (3.4)
$$

where the sum runs over all non-negative \underline{m} that satisfy $\sum_{0 \leq i \leq 2^k-1} m_i = m-k+1$. We now bound the first factor inside the sum by a quantity which only depends on m (not on the specific vector <u>m</u>). Denote by $\hat{\chi} = \frac{1}{2^k} \sum_{i=1}^{2^k} \chi_i$, and recall the definitions $\mu := \frac{1}{2^k} \sum_{i=1}^{2^k} \mu_i$ and $\rho^{(m)} := \frac{1}{2^k} \sum_{i=1}^{2^k} \rho_i^{(m)}$ $i^{(m)}$. From [\(3.3\)](#page-5-0) it follows that $\mu \leq \rho^{(m)}$. Conditioned on \underline{m} , $\sigma^{(m)} \in V$ we have for any $\epsilon > 0$,

$$
\mathbb{P}\left(\hat{d} \notin A_r^{(k)}\right) = \mathbb{P}\left(\hat{\chi} > \frac{r}{2^k}\right)
$$

= $\mathbb{P}\left(\hat{\chi} - \mu > \frac{r}{2^k} - \mu\right)$
 $\leq \mathbb{P}\left(\hat{\chi} - \mu > \frac{r}{2^k} - \rho^{(m)}\right).$

We will use the following lemma which bounds the deviation of the average of Poisson trials from their mean.

Lemma 2. Let X_1, \ldots, X_n be independent Bernoulli random variables with $P(X_i =$ 1) = μ_i and denote by $\mu = \frac{1}{n} \sum_{i=1}^n \mu_i$. Then for any $0 < \gamma \le (2e-1)\mu$ the following bound holds:

$$
P\left(\frac{1}{n}\sum_{i=1}^{n}X_i > \mu + \gamma\right) \le e^{-n\gamma^2/4}.
$$

The proof of the lemma is based on applying Chernoff bound on the tail probability of the sum of Poisson trials (similar to Theorem 4.1 and 4.3, in [\[3](#page-9-1)]).

Substituting for γ the value $\frac{r}{2^k} - \rho^{(m)}$ in the above lemma and recalling the theorem's condition that

$$
k \ge 2 + 2\log_2\left(\frac{1}{(2e-1)\mu}\right) + \log_2\left(\ln\left(\frac{2}{\delta}\right)\right)
$$

which, with the following choice for r ,

$$
r = 2^{k} \left(\sqrt{\frac{1}{2^{k-2}} \ln\left(\frac{2}{\delta}\right)} + \rho^{(m)} \right),
$$
\n(3.5)

ensures that $\frac{r}{2^k} - \rho^{(m)} \le (2e - 1)\mu$, yields the following bound,

 $\overline{}$ $\overline{}$ \mid

$$
\mathbb{P}\left(\hat{d}\not\in A_r^{(k)}\right) \le e^{-2^{k-2}\left(\frac{r}{2^k} - \rho^{(m)}\right)^2}.
$$
\n(3.6)

Next, we estimate the cardinality of the set $A_r^{(k)}$. Without loss of generality let $d^* = [0, 0, \dots, 0]$ then we have

$$
A_r^{(k)} \Big| = \sum_{i=0}^r \binom{2^k}{i}
$$

$$
\leq \left(\frac{e2^k}{r}\right)^r.
$$
 (3.7)

Since the error subsequence $\xi^{(\nu)}$ is also a subsequence of $x^{(n)}$ then we associate a selection rule $\Gamma_d: \{0,1\}^n \to \{0,1\}^{\nu}$ which selects $\xi^{(\nu)}$ from $x^{(n)}$. Let $E_{d,\epsilon}^{(\ell)}$ denote the event that based on a given Γ_d the selected subsequence $\xi^{(\nu)}$ is of length at least ℓ and its frequency of 1s deviates from the expected value β by at least ϵ . Formally, this is defined as the large-deviation event

$$
E_{d,\epsilon}^{(\ell)} = \left\{ x^{(n)} : \xi^{(\nu)} = \Gamma_d \left(x^{(n)} \right), \, \nu \ge \ell, \, \left| \frac{\|\xi^{(\nu)}\|}{\nu} - \beta \right| > \epsilon \right\},\,
$$

where $\|\xi^{(\nu)}\|$ denotes the number of 1s in the binary sequence $\xi^{(\nu)}$ of length ν . We wish to bound from above the probability of $E_{d,\epsilon}^{(\ell)}$.

We use the following lemma which states a rate on the strong law of large numbers for a Markov Chain.

Lemma 3. [\[2\]](#page-9-2) Let Z_1, \ldots, Z_n be a finite ergodic and reversible Markov chain in stationary state with a second largest eigenvalue λ and f a function taking values in [0, 1] such that $\mathbb{E} f(Z_i) = \mu$. Denote by $\lambda_0 = \max\{0, \lambda\}$ and the stationary probability distribution \mathbb{P} . Then for all $\epsilon > 0$ such that $\mu + \epsilon < 1$, $n \geq 1$ the following bound holds:

$$
\mathbb{P}\left(\sum_{i=1}^n Z_i \ge n(\mu + \epsilon)\right) \le e^{-2n\epsilon^2\left(\frac{1-\lambda_0}{1+\lambda_0}\right)}.
$$

The lemma appears as Theorem 1 in [\[2](#page-9-2)]. We note that the bound in this lemma coincides with Hoeffding's bound and is optimal for $\lambda \geq 0$.

We now apply the lemma to the error subsequence in order to estimate the probability of the large deviation event $E_{d,\epsilon}^{(\ell)}$. Denote by $X^{(n)} = \{X_{m+i}\}_{i=1}^n$ and $Y^{(n)} = \{Y_{m+i}\}_{i=1}^n$ the sequences of random variables produced by the source according to \mathcal{M}_{k^*} and the predictions made by the learner, respectively. Let $\Xi^{(\nu)} = {\{\Xi_j\}}_{j=1}^{\nu}$ be the sequence of random variables representing the errors made when predicting zeros, i.e., $Y_{i_j} = 0$, $i_j \in \{m+1,\ldots,m+n\}$, $1 \leq j \leq \nu$. As mentioned above, $\Xi^{(\nu)}$ is a subsequence of $X^{(n)}$. Denote by $S^{(n)} = \{S_{m+i}\}_{i=1}^n$ the sequence of consecutive states of model \mathcal{M}_{k^*} that correspond to $X^{(n)}$. Let $\tilde{S}^{(\nu)}$ denote the subsequence of $S^{(n)}$ corresponding to the subsequence $\Xi^{(\nu)}$ of $X^{(n)}$.

We now apply Lemma [3](#page-6-0) to the sequence $\Xi^{(\nu)}$. In general $S^{(\nu)}$ may be split into parts each consisting of consecutive states S of $S^{(n)}$, i.e.,

$$
\tilde{S}^{(\nu)} = (\tilde{S}_1, \dots, \tilde{S}_\nu) \n= (S_{i_1}, S_{i_1+1}, \dots, S_{i_1+r_1-1}, S_{i_2}, S_{i_2+1}, \dots, S_{i_2+r_2-1}, \dots, S_{i_q}, S_{i_q+1}, \dots, S_{i_q+r_q-1}),
$$

where $\sum_{j=1}^{q} r_j = \nu$ and the parts are disjoint, i.e.,

$$
i_j + r_j \quad < \quad i_{j+1} \tag{3.8}
$$

for $1 \leq j \leq q-1$. For any state $s \in \mathbb{S}_k$ let the function $f(s)$ in Lemma [3](#page-6-0) be the value of the least significant bit of the binary-representation of s. So for \tilde{S}_i in $\tilde{S}^{(\nu)}$ we have $f(\tilde{S}_i) = \Xi_i$ and μ in the lemma equals $\mathbb{P}(\Xi_i = 1)$ which equals β by (2.2) . Let λ in the lemma be the second largest eigenvalue of the source's transition matrix T.

From Chebychev's inequality (see for instance [\[4\]](#page-10-0)), for any $t \geq 0$, the sequence $\Xi^{(\nu)}$ satisfies,

$$
\mathbb{P}\left(\sum_{j=1}^{\nu} \Xi_j > \nu(\beta + \epsilon)\right) = \mathbb{P}\left(\sum_{j=1}^{\nu} f(\tilde{S}_j) > \nu(\beta + \epsilon)\right)
$$
(3.9)

$$
\leq \exp\left\{-\nu t(\beta+\epsilon)\right\} \mathbb{E} \exp\left\{t \sum_{j=1}^{\nu} f(\tilde{S}_j)\right\}.
$$
 (3.10)

The expectation in [\(3.10\)](#page-7-0) is now expressed as

$$
\mathbb{E} \exp\left\{t\sum_{j=1}^{\nu} f(\tilde{S}_j)\right\} = \mathbb{E} \exp\left\{t\sum_{j=1}^{q-1} \sum_{k_j=0}^{r_j-1} f(S_{i_j+k_j})\right\}.
$$
 (3.11)

Recall that $S^{(\nu)}$ is a subsequence of $S^{(n)}$ so that the expectation is taken with respect to the joint probability distribution of the whole sequence $S^{(n)}$. Taking into account the states not common to $S^{(n)}$ and $\tilde{S}^{(\nu)}$ (these are at time instances when the selection rule does not select a bit from $X^{(n)}$ and recalling that $S^{(n)}$ starts at state S_{m+1} then we write,

$$
S^{(n)} = \left(S_{m+1}, \ldots, S_{i_1-1}, S_{i_1}, \ldots, S_{i_1+r_1-1}, S_{i_1+r_1}, \ldots, S_{i_2-1}, S_{i_2}, \ldots, S_{i_2+r_2-1}, S_{i_2+r_2}, \ldots, S_{i_q}, \ldots, S_{i_q+r_q-1}, \ldots, S_{m+n}\right)
$$

We can now express (3.11) explicitly as follows:

·

$$
\mathbb{E} \exp \left\{ t \sum_{j=1}^{q-1} \sum_{k_j=0}^{r_j-1} f(S_{i_j+k_j}) \right\} = \sum_{s_{m+1} \in \mathbb{S}} \mathbb{P} (s_{m+1}) \sum_{s_{m+2} \in \mathbb{S}} \mathbb{P} (s_{m+2} | s_{m+1})
$$
\n...\n
$$
\sum_{s_{i_1} = \mathbb{S}} \mathbb{P} (s_{i_1-1} | s_{i_1-2})
$$
\n
$$
\sum_{s_{i_1} \in \mathbb{S}} \mathbb{P} (s_{i_1} | s_{i_1-1}) e^{tf(s_{i_1})} \sum_{s_{i_1+1} \in \mathbb{S}} \mathbb{P} (s_{i_1+1} | s_{i_1}) e^{tf(s_{i_1+1})}
$$
\n...\n
$$
\sum_{s_{i_1+r_1-1} \in \mathbb{S}} \mathbb{P} (s_{i_1+r_1-1} | s_{i_1+r_1-2}) e^{tf(s_{i_1+r_1-1})}
$$
\n
$$
\sum_{s_{i_1+r_1} \in \mathbb{S}} \mathbb{P} (s_{i_1+r_1} | s_{i_1+r_1-1}) \cdots \sum_{s_{i_2-1} \in \mathbb{S}} \mathbb{P} (s_{i_2-1} | s_{i_2-2})
$$
\n...\n
$$
\sum_{s_{i_2} \in \mathbb{S}} \mathbb{P} (s_{i_2} | s_{i_2-1}) e^{tf(s_{i_2})} \sum_{s_{i_2+1} \in \mathbb{S}} \mathbb{P} (s_{i_2+1} | s_{i_2}) e^{tf(s_{i_2+1})}
$$
\n...\n
$$
\sum_{s_{i_2+r_2-1} \in \mathbb{S}} \mathbb{P} (s_{i_2+r_2-1} | s_{i_2+r_2-2}) e^{tf(s_{i_2+r_2-1})}
$$
\n...\n
$$
\sum_{s_{i_q} \in \mathbb{S}} \mathbb{P} (s_{i_q} | s_{i_q-1}) e^{tf(s_{i_q})} \sum_{s_{i_q+1} \in \mathbb{S}} \mathbb{P} (s_{i_q+1} | s_{i_q}) e^{tf(s_{i_q+1})}
$$
\n...\n<

$$
\sum_{s_{i_q+r_q}\in\mathbb{S}}\mathbb{P}\left(s_{i_q+r_q}\mid s_{i_q+r_q-1}\right)\cdots\sum_{s_{m+n}\in\mathbb{S}}\mathbb{P}\left(s_{m+n}\mid s_{m+n-1}\right).
$$

In the proof of Lemma [3](#page-6-0) [\[2\]](#page-9-2) the two-state case ($|\mathbb{S}| = 2$) is solved first. They show that factors of the kind of [\(3.13\)](#page-8-0) can be expressed as a product of matrices $(MD_t^2)^{r_1}$ where M denotes the 2×2 transition matrix and D_t is a diagonal matrix $\begin{pmatrix} 1 & 0 \\ 0 & \text{cm} \end{pmatrix}$ 0 $\exp\{t/2\}$. It follows immediately that a factor such as [\(3.14\)](#page-8-0) equals $(MD_0^2)^a = M^a$, $a = i_2 - i_1 - r_1$. Thus, for the two state case, the expectation (left hand side of [\(3.12\)](#page-8-0)) equals a bilinear form $\begin{bmatrix} \beta & 1-\beta \end{bmatrix} R \begin{bmatrix} 1 & 1 \end{bmatrix}$ where β is the stationary probability of the first of the two states, v' denotes transpose of the vector v and the square matrix R equals $M^{a_1} (MD_t^2)^{a_2} M^{a_3} (MD_t^2)^{a_4} \cdots M^{a_N} (MD_t^2)^{a_N}$. Since M is a stochastic matrix then each of its elements is non-negative and bounded by 1 hence we have

$$
\begin{array}{rcl}\n[\beta \quad 1 - \beta \quad]R \ [1 \quad 1 \]' & \leq & [\beta \quad 1 - \beta \quad](MD_t^2)^{a_2} (MD_t^2)^{a_4} \cdots (MD_t^2)^{a_N} \ [1 \quad 1 \]' \\
& = & [\beta \quad 1 - \beta \quad](MD_t^2)^{\nu} \ [1 \quad 1 \]'.\n\end{array} \tag{3.15}
$$

Based on the proof of [\[2](#page-9-2)], multiplying [\(3.15\)](#page-8-1) by the exponential factor $\exp\{-\nu t(\beta + \epsilon)\}\$ gives an expression which is bounded from above by $\exp\left\{-2\nu\left(\frac{1-\lambda_0}{1+\lambda_0}\right)\epsilon^2\right\}$. This holds also in the general case ($|\mathbb{S}| > 2$). Hence (3.10) is bounded from above by this exponential. We may therefore bound the probability of $E_{d,\epsilon}^{(\ell)}$, for any fixed

 $d \in \{0,1\}^{2^k}$, as follows,

$$
\mathbb{P}\left(E_{d,\epsilon}^{(\ell)}\right) = \sum_{\nu \geq \ell} \mathbb{P}\left(\left|\frac{\|\Xi^{(\nu)}\|}{\nu} - \beta\right| > \epsilon \middle| \nu\right) \mathbb{P}\left(\nu\right)
$$
\n
$$
\leq 2 \sum_{\nu \geq \ell} \exp\left\{-2\nu\gamma\epsilon^2\right\} \mathbb{P}\left(\nu\right)
$$
\n
$$
\leq 2 \exp\left\{-2\ell\gamma\epsilon^2\right\}.
$$
\n(3.16)

Denote by \hat{d} the binary vector [\(3.1\)](#page-4-2) associated with the learnt model \mathcal{M}_k (which is based on a random training sequence $x^{(m)}$). We are interested in the probability of the event $E_{\hat{j}}^{(\ell)}$ $d_{\hat{d},\epsilon}^{(\ell)}$ that after learning, the selection rule $\Gamma_{\hat{d}}$ picks a subsequence $\Xi^{(\nu)}$ from $X^{(n)}$ of length $\nu \geq \ell$ which is biased away from β by an amount greater than ϵ .

Denoting by $\overline{A}_r^{(k)}$ $r_r^{(k)}$ the complement of the set $A_r^{(k)}$ then we have

$$
\mathbb{P}\left(E_{\hat{d},\epsilon}^{(\ell)}\right) = \mathbb{P}\left(E_{\hat{d},\epsilon}^{(\ell)}\middle|\hat{d} \in A_r^{(k)}\right) \mathbb{P}\left(A_r^{(k)}\right) + \mathbb{P}\left(E_{\hat{d},\epsilon}^{(\ell)}\middle|\hat{d} \notin A_r^{(k)}\right) \mathbb{P}\left(\overline{A}_r^{(k)}\right)
$$
\n
$$
= \mathbb{P}\left(\bigcup_{d \in A_r^{(k)}} E_{d,\epsilon}^{(\ell)}\right) \mathbb{P}\left(A_r^{(k)}\right) + \mathbb{P}\left(E_{\hat{d},\epsilon}^{(\ell)}\middle|\hat{d} \notin A_r^{(k)}\right) \mathbb{P}\left(\overline{A}_r^{(k)}\right)
$$
\n
$$
\leq \mathbb{P}\left(\bigcup_{d \in A_r^{(k)}} E_{d,\epsilon}^{(\ell)}\right) + \mathbb{P}\left(\overline{A}_r^{(k)}\right)
$$
\n
$$
\leq 2 \left|A_r^{(k)}\middle|\exp\left\{-2\ell\gamma\epsilon^2\right\} + \exp\left\{-2^{k-2}\left(\frac{r}{2^k} - \rho^{(m)}\right)^2\right\}
$$
\n
$$
\leq 2 \left(\frac{e2^k}{r}\right)^r \exp\left\{-2\ell\gamma\epsilon^2\right\} + \exp\left\{-2^{k-2}\left(\frac{r}{2^k} - \rho^{(m)}\right)^2\right\} \quad (3.17)
$$

which follows from [\(3.6\)](#page-6-1), [\(3.7\)](#page-6-2) and [\(3.16\)](#page-9-3). Note that for any $0 < \delta < 1$ the choice of r in [\(3.5\)](#page-6-3) makes the second term in [\(3.17\)](#page-9-4) be no larger than $\frac{\delta}{2}$. The first term is no larger than $\frac{\delta}{2}$ if the following holds,

$$
\epsilon \le \sqrt{\frac{1}{2\ell\gamma} \left(r \ln \left(\frac{e2^k}{r} \right) + \ln 2 + \ln \left(\frac{4}{\delta} \right) \right)}.
$$
\n(3.18)

Substituting for r in [\(3.18\)](#page-9-5) the value in [\(3.5\)](#page-6-3) gives the following bound on ϵ ,

$$
\epsilon \leq \sqrt{\frac{1}{2\ell\gamma} \left(2^k \left(\sqrt{\frac{1}{2^{k-2}}\ln\left(\frac{2}{\delta}\right)} + \rho^{(m)}\right) \ln\left(\frac{e}{\left(\sqrt{\frac{1}{2^{k-2}}\ln\left(\frac{2}{\delta}\right)} + \rho^{(m)}\right)}\right) + \ln 2 + \ln\left(\frac{4}{\delta}\right)\right)}
$$

which holds with probability at least $1-\delta$. This concludes the proof of Theorem [1.](#page-4-1)

REFERENCES

- [1] H. Chernoff. A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations. Annals of Mathematical Statistics, 23:493–507, 1952.
- [2] C. A. Leon and F. Perron. Optimal hoeffding bounds for discrete reversible markov chains. Annals of Applied Probability, 14(2):958–970, 2004.
- [3] R. Motwani and P. Raghavan. Randomized Algorithms. Cambridge University Press, 1995.

[4] A. N. Shiryaev. Probability. Springer, New York, 1996.

Department of Electrical and Electronics Engineering, Ariel University Center of Samaria, Ariel 40700, ISRAEL. Email: ratsaby@ariel.ac.il