

THE COHOMOLOGY OF MOTIVIC $A(2)$

DANIEL C. ISAKSEN

ABSTRACT. Working over an algebraically closed field of characteristic zero, we compute the cohomology of the subalgebra $A(2)$ of the motivic Steenrod algebra that is generated by Sq^1 , Sq^2 , and Sq^4 . The method of calculation is a motivic version of the May spectral sequence.

Speculatively assuming that there is a “motivic modular forms” spectrum with certain properties, we use an Adams-Novikov spectral sequence to compute the homotopy of such a spectrum at the prime 2.

1. INTRODUCTION

The purpose of this article is to present some algebraic calculations that are relevant to motivic homotopy theory. Recent work [HKO] [DI] has shown that the Adams and Adams-Novikov spectral sequences are useful for computing in the motivic stable homotopy category over an algebraically closed field of characteristic zero, after completion with respect to the motivic \mathbb{F}_2 -Eilenberg-Mac Lane spectrum $H\mathbb{F}_2$ that represents motivic \mathbb{F}_2 -cohomology.

This program is built upon work of Voevodsky [V1] [V2] [V3]. Voevodsky has described the motivic \mathbb{F}_2 -cohomology of an algebraically closed field of characteristic zero, which we write as \mathbb{M}_2 . He also described the motivic \mathbb{F}_2 -Steenrod algebra A . Much like the classical Steenrod algebra A_{cl} , A is generated by elements Sq^i , subject to motivic versions of the Adem relations.

In this article, we take Voevodsky’s descriptions as given algebraic inputs, and we carry out further algebraic computations. The motivic Adams spectral sequence takes the cohomology of A , i.e., the ring $\text{Ext}_A(\mathbb{M}_2, \mathbb{M}_2)$, as input. Just as in the classical situation, it is unlikely that we will ever have a complete description of this ring. However, much can be said in low dimensions [DI].

Classically, one way to approximate difficult calculations over A_{cl} is to consider instead the subalgebra $A(2)_{cl}$ generated by Sq^1 , Sq^2 , and Sq^4 . It is possible to give an explicit but lengthy description of the cohomology of $A(2)_{cl}$ [M2] [IS]. The main purpose of this article is to carry out the motivic version of this computation. Namely, we completely describe the ring $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$.

The cohomology of $A(2)_{cl}$ is the E_2 -term of an Adams spectral sequence that converges to the homotopy of tmf at the prime 2. The original sources of this material are mostly unpublished work of Hopkins, Mahowald, and Miller [H1] [H2]. One might optimistically hope to construct a motivic spectrum mmf , defined over algebraically closed fields of characteristic zero, whose homotopy is similarly related to the cohomology of motivic $A(2)$. This is a potential homotopical application of our algebraic computations. In this article, we do not discuss any issues related to the existence of mmf .

The author was supported by NSF grant DMS0803997.

Although it is theoretically possible to compute the homotopy of tmf at the prime 2 by starting with Ext groups over $A(2)_{cl}$ and applying the Adams spectral sequence, it is difficult in practice. The motivic version of this computation is significantly more difficult. Even finding the E_3 -term of the spectral sequence requires a prohibitive amount of bookkeeping, and there are a tremendous number of exotic extensions to resolve.

Classically, a more efficient way to compute the homotopy of tmf at the prime 2 is to start with the cohomology of the elliptic curves Hopf algebroid and apply the Adams-Novikov spectral sequence. This computation is entirely described in [B], based on work of Hopkins and Mahowald [HM].

At the end of this paper, we carry out the motivic version of this computation. We make several assumptions about motivic versions of the elliptic curves Hopf algebroid and the motivic Adams-Novikov spectral sequence. Based on these assumptions, we are able to describe the homotopy of the speculative motivic spectrum mmf at the prime 2.

1.1. Organization of the paper. Section 2 gives a brief review of the algebraic objects under consideration. In Section 3, we set up a motivic May spectral sequence that converges to the cohomology of $A(2)$. We also describe the E_∞ -term of this spectral sequence. In Section 4, we compute Massey products and resolve all multiplicative extensions to give a complete description of the cohomology of $A(2)$ as a ring. Finally, Section 5 discusses an Adams-Novikov spectral sequence and describes the homotopy of the speculative motivic spectrum mmf .

1.2. Acknowledgments. We acknowledge the assistance of Robert Bruner. Section 3 relies very heavily on unpublished notes of Peter May on the cohomology of $A(2)_{cl}$ [M2]. Section 5 relies very heavily on charts of Tilman Bauer [B]. We also thank Mark Behrens, Dan Dugger, and Mike Hill for useful conversations.

2. BACKGROUND

In this section we review the basic algebraic facts about the objects under consideration. We are working in categories of bigraded objects. In a bidegree (p, q) , we shall refer to p as the topological degree and q as the weight. The terminology is motivated by the relationship between motivic cohomology and classical homotopy theory.

Definition 2.1. *The bigraded ring \mathbb{M}_2 is the polynomial ring $\mathbb{F}_2[\tau]$ on one generator τ of bidegree $(0, 1)$.*

The relevance of \mathbb{M}_2 is that it is the motivic \mathbb{F}_2 -cohomology of an algebraically closed field of characteristic zero [V1].

Definition 2.2. *The motivic Steenrod algebra A is the \mathbb{M}_2 -algebra generated by elements Sq^{2k} and Sq^{2k-1} for all $k \geq 1$, of bidegrees $(2k, k)$ and $(2k-1, k-1)$ respectively, and satisfying the following relations for $a < 2b$:*

$$Sq^a Sq^b = \sum_c \binom{b-1-c}{a-2c} \tau^? Sq^{a+b-c} Sq^c.$$

The expression $\tau^?$ stands for either 1 or τ . The distinction is easily determined by consideration of bidegrees. For example, $Sq^2 Sq^2 = \tau Sq^3 Sq^1$.

The relevance of A is that it is the ring of motivic \mathbb{F}_2 -cohomology operations over an algebraically closed field of characteristic zero [V2] [V3].

We consider \mathbb{M}_2 as an A -module, where Sq^i acts trivially on \mathbb{M}_2 for $i > 0$ and Sq^0 acts as the identity. Since \mathbb{M}_2 is concentrated in topological degree 0, this is the only possible action of A on \mathbb{M}_2 .

Definition 2.3. *The algebra $A(2)$ is the \mathbb{M}_2 -subalgebra of A generated by Sq^1 , Sq^2 , and Sq^4 .*

Remark 2.4. The algebra A has a Milnor \mathbb{M}_2 -basis consisting of elements of the form P^R , where $R = (r_1, r_2, \dots)$ ranges over all finite sequences of non-negative integers. Just as in the classical case, $A(2)$ has an \mathbb{M}_2 -basis consisting of elements of the form P^R , where $R = (r_1, r_2, r_3)$, $0 \leq r_1 \leq 7$, $0 \leq r_2 \leq 3$, and $0 \leq r_3 \leq 1$. See [DI] for more details on the motivic Milnor basis.

2.5. Ext groups. We will compute the tri-graded groups $\mathrm{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$.

We have

$$\mathrm{Ext}_{A(2)}^{0,(0,*)}(\mathbb{M}_2, \mathbb{M}_2) = \mathrm{Hom}_{A(2)}^{(0,*)}(\mathbb{M}_2, \mathbb{M}_2) = \mathbb{M}_2.$$

Here we abuse notation and write \mathbb{M}_2 where we really mean the \mathbb{M}_2 -dual $\mathrm{Hom}_{\mathbb{M}_2}(\mathbb{M}_2, \mathbb{M}_2)$ of \mathbb{M}_2 . The only important point is that now τ has bidegree $(0, -1)$. For fixed s and t , $\mathrm{Ext}_{A(2)}^{s,(t+s,*)}(\mathbb{M}_2, \mathbb{M}_2)$ is a module over $\mathrm{Ext}_{A(2)}^{0,(0,*)}(\mathbb{M}_2, \mathbb{M}_2)$. In particular, it is an \mathbb{M}_2 -module and therefore decomposes as a sum of modules of the form \mathbb{M}_2 or \mathbb{M}_2/τ^k . In Section 2.6, we will explain that the free part coincides with Ext over the classical version of $A(2)$.

2.6. Comparison with the classical Steenrod algebra. We write A_{cl} and $A(2)_{\mathrm{cl}}$ for the classical Steenrod algebra and its subalgebra generated by Sq^1 , Sq^2 , and Sq^4 .

Lemma 2.7. *There is a ring isomorphism $A[\tau^{-1}] \rightarrow A_{\mathrm{cl}} \otimes_{\mathbb{F}_2} \mathbb{M}_2[\tau^{-1}]$ that takes Sq^{2k} and Sq^{2k-1} to $\tau^{-k} \mathrm{Sq}^{2k}$ and $\tau^{-k} \mathrm{Sq}^{2k-1}$ respectively.*

In other words, the motivic Steenrod algebra and the classical Steenrod algebra are essentially the same after inverting τ . The proof of this lemma is a straightforward algebraic exercise. Similarly, there is a ring isomorphism $A(2)[\tau^{-1}] \rightarrow A(2)_{\mathrm{cl}} \otimes_{\mathbb{F}_2} \mathbb{M}_2[\tau^{-1}]$ that takes Sq^1 , Sq^2 , and Sq^4 to Sq^1 , $\tau^{-1} \mathrm{Sq}^2$, and $\tau^{-2} \mathrm{Sq}^4$ respectively.

Proposition 2.8. *There are isomorphisms of rings*

$$\mathrm{Ext}_A(\mathbb{M}_2, \mathbb{M}_2) \otimes_{\mathbb{M}_2} \mathbb{M}_2[\tau^{-1}] \cong \mathrm{Ext}_{A_{\mathrm{cl}}}(\mathbb{F}_2, \mathbb{F}_2) \otimes_{\mathbb{F}_2} \mathbb{M}_2[\tau^{-1}].$$

and

$$\mathrm{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2) \otimes_{\mathbb{M}_2} \mathbb{M}_2[\tau^{-1}] \cong \mathrm{Ext}_{A(2)_{\mathrm{cl}}}(\mathbb{F}_2, \mathbb{F}_2) \otimes_{\mathbb{F}_2} \mathbb{M}_2[\tau^{-1}].$$

The point of the proposition is that the free \mathbb{M}_2 -modules in $\mathrm{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$ correspond precisely to the free \mathbb{F}_2 -modules in $\mathrm{Ext}_{A(2)_{\mathrm{cl}}}(\mathbb{F}_2, \mathbb{F}_2)$. On the other hand, copies of \mathbb{M}_2/τ^k have no classical analogue.

The proof of Proposition 2.8 is given in [DI]. The idea is to use the isomorphism of Lemma 2.7 and the flatness of $\mathbb{M}_2 \rightarrow \mathbb{M}_2[\tau^{-1}]$.

3. THE MOTIVIC MAY SPECTRAL SEQUENCE

In this section we begin our computation of $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$ by setting up a May spectral sequence [M1] and finding the E_∞ -term of this spectral sequence. Elements of $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$ will be graded in the form (s, a, w) , where s is the stem, i.e., the topological degree minus the homological degree, a is Adams filtration, i.e., the homological degree, and w is the weight.

Let I be the two-sided \mathbb{M}_2 -ideal of $A(2)$ generated by Sq^1 , Sq^2 , and Sq^4 . Let $\text{Gr } A(2)$ denote the associated graded algebra $A(2)/I \oplus I/I^2 \oplus I^2/I^3 \oplus \dots$. Elements of $\text{Gr } A(2)$ will be graded in the form (m, t, w) , where m is the May filtration, i.e., the I -adic valuation, t is the topological degree, and w is the weight.

We will first need to compute $\text{Ext}_{\text{Gr } A(2)}(\mathbb{M}_2, \mathbb{M}_2)$. Elements of this ring will be graded in the form (m, s, a, w) , where m is the May filtration, s is the stem, i.e., the topological degree minus the homological degree, a is the Adams filtration, i.e., the homological degree, and w is the weight.

Proposition 3.1. *There is a spectral sequence*

$$E_2 = \text{Ext}_{\text{Gr } A(2)}^{(m, s, a, w)}(\mathbb{M}_2, \mathbb{M}_2) \Rightarrow \text{Ext}_{A(2)}^{(s, a, w)}(\mathbb{M}_2, \mathbb{M}_2).$$

We shall refer to this as the motivic May spectral sequence. As usual, it can be obtained by filtering the cobar complex by powers of I .

Let I_{cl} be the ideal of $A(2)_{\text{cl}}$ generated by Sq^1 , Sq^2 , and Sq^4 , and let $\text{Gr } A(2)_{\text{cl}}$ be the associated graded algebra.

Proposition 3.2.

- (a) *The tri-graded algebras $\text{Gr } A(2)$ and $\text{Gr } A(2)_{\text{cl}} \otimes_{\mathbb{F}_2} \mathbb{F}_2[\tau]$ are isomorphic.*
- (b) *The quadruply-graded rings $\text{Ext}_{\text{Gr } A(2)}(\mathbb{M}_2, \mathbb{M}_2)$ and $\text{Ext}_{\text{Gr } A(2)_{\text{cl}}}(\mathbb{F}_2, \mathbb{F}_2) \otimes_{\mathbb{F}_2} \mathbb{M}_2$ are isomorphic.*

The point is that the E_2 -terms of the motivic and classical May spectral sequences are very similar. Although the motivic and classical differentials are compatible, they are different in an important way that we shall describe later.

The proposition can be proved just as in [DI]. The main point is that the τ coefficients in the relations in A appear only on terms of higher filtration and thus do not affect the associated graded algebra.

The classical ring $\text{Ext}_{\text{Gr } A(2)_{\text{cl}}}(\mathbb{F}_2, \mathbb{F}_2)$ is computed in [M2] using the May spectral sequence. It is also computed in [IS] using a different method.

By Proposition 3.2 and the results in [M1], the ring $\text{Ext}_{\text{Gr } A(2)}(\mathbb{M}_2, \mathbb{M}_2)$ is the cohomology of the differential graded \mathbb{M}_2 -algebra $\mathbb{M}_2[h_{10}, h_{11}, h_{12}, h_{20}, h_{21}, h_{30}]$, where the differential is described in the following table. See [DI] for an explanation of the degrees.

It is relatively straightforward to compute $\text{Ext}_{\text{Gr } A(2)}(\mathbb{M}_2, \mathbb{M}_2)$ from the differentials given above. The table below lists the generators.

Proposition 3.3. *The ring $\text{Ext}_{\text{Gr } A(2)}(\mathbb{M}_2, \mathbb{M}_2)$ is generated over \mathbb{M}_2 by the elements listed in Table 2, subject to the relations*

- (1) $h_0 h_1 = 0$.
- (2) $h_1 h_2 = 0$.
- (3) $h_2 b_{20} = h_0 h_0(1)$.
- (4) $h_2 h_0(1) = h_0 b_{21}$.
- (5) $h_0(1)^2 = b_{20} b_{21} + h_1^2 b_{30}$.

TABLE 1. Differentials for computing $\text{Ext}_{\text{Gr } A(2)}(\mathbb{F}_2, \mathbb{F}_2)$

x	degree	$d(x)$
h_{10}	(1, 0, 1, 0)	0
h_{11}	(1, 1, 1, 1)	0
h_{12}	(1, 3, 1, 2)	0
h_{20}	(2, 2, 1, 1)	$h_{10}h_{11}$
h_{21}	(2, 5, 1, 3)	$h_{11}h_{12}$
h_{30}	(3, 6, 1, 3)	$h_{10}h_{21} + h_{20}h_{12}$

TABLE 2. Generators for $\text{Ext}_{\text{Gr } A(2)}(\mathbb{M}_2, \mathbb{M}_2)$

generator	degree	description in terms of h_{ij}
h_0	(1, 0, 1, 0)	h_{10}
h_1	(1, 1, 1, 1)	h_{11}
h_2	(1, 3, 1, 2)	h_{12}
b_{20}	(4, 4, 2, 2)	h_{20}^2
b_{21}	(4, 10, 2, 6)	h_{21}^2
b_{30}	(6, 12, 2, 6)	h_{30}^2
$h_0(1)$	(4, 7, 2, 4)	$h_{20}h_{21} + h_{11}h_{30}$

The proof of Proposition 3.3 is a straightforward lift of the analogous classical computation because of Proposition 3.2(b).

3.4. E_4 -term of the motivic May spectral sequence. Having described the E_2 -term of the motivic May spectral sequence, i.e., $\text{Ext}_{\text{Gr } A(2)}(\mathbb{M}_2, \mathbb{M}_2)$, we are now ready to compute the higher terms. For dimension reasons, as in the classical case, the odd differentials must vanish. In particular, $E_3 = E_4$.

The d_2 differentials on the E_2 -term are easy to analyze. They must be compatible with the d_2 differentials in the classical May spectral sequence and they must preserve the weight. The following table lists the d_2 differentials on all of our generators. From the data in this table, one can use the Leibniz rule to compute the d_2 differential on any element.

TABLE 3. d_2 differentials in the motivic May spectral sequence

x	$d_2(x)$	x	$d_2(x)$
h_0	0	b_{20}	$\tau h_1^3 + h_0^2 h_2$
h_1	0	b_{21}	h_2^3
h_2	0	b_{30}	$\tau h_1 b_{21}$
		$h_0(1)$	$h_0 h_2^2$

A straightforward computation now gives the E_4 -term of the motivic May spectral sequence. The generators are listed in the following table, and the relations are listed in the next theorem.

The notation is chosen to be compatible with the standard notation for elements in the cohomology of the classical Steenrod algebra. The elements c , d , e , and g are related to the classical elements c_0 , d_0 , e_0 , and g . The element P is related to

TABLE 4. Generators of the E_4 -term of the motivic May spectral sequence

generator	degree	description in E_2
h_0	(1, 0, 1, 0)	h_0
h_1	(1, 1, 1, 1)	h_1
h_2	(1, 3, 1, 2)	h_2
P	(8, 8, 4, 4)	b_{20}^2
c	(5, 8, 3, 5)	$h_1 h_0(1)$
u	(5, 11, 3, 7)	$h_1 b_{21}$
α	(7, 12, 3, 6)	$h_0 b_{30}$
d	(8, 14, 4, 8)	$h_1^2 b_{30} + b_{20} b_{21}$
ν	(7, 15, 3, 8)	$h_2 b_{30}$
e	(8, 17, 4, 10)	$h_0(1) b_{21}$
g	(8, 20, 4, 12)	b_{21}^2
Δ	(12, 24, 4, 12)	b_{30}^2

the Adams periodicity operator. The element Δ is related to the element in the homotopy of tmf of the same name. The elements u , α , and ν have no analogues and are given arbitrary names.

Theorem 3.5. *The E_4 -term of the motivic May spectral sequence is generated over \mathbb{M}_2 by the elements listed in Table 4, subject to the following relations:*

- (1) $h_0 h_1, h_1 h_2, h_0^2 h_2 + \tau h_1^3, h_0 h_2^2, h_2^3.$
- (2) $\tau u, \tau h_1^2 c, \tau c d, \tau c e, \tau c g.$
- (3) $h_0^2 \nu + \tau h_1 d, h_0 h_2 \nu + \tau h_1 e, h_2^2 \nu + \tau h_1 g.$
- (4) $h_2 d + h_0 e, h_2 e + h_0 g, h_2 \alpha + h_0 \nu.$
- (5) $h_0 c, h_2 c, h_0 u, h_2 u, h_1 \alpha, h_1 \nu.$
- (6) $c^2 + h_1^2 d, u^2 + h_1^2 g, cu + h_1^2 e, e^2 + dg.$
- (7) $ud + ce, ue + cg, vd + \alpha e, ve + \alpha g.$
- (8) $c\alpha, c\nu, u\alpha, u\nu.$
- (9) $\alpha^2 + h_0^2 \Delta, \alpha\nu + h_0 h_2 \Delta, \nu^2 + h_2^2 \Delta.$
- (10) $h_0^2 d + P h_2^2, h_0 \alpha d + P h_2 \nu, d^2 + h_1^4 \Delta + P g.$

Several observations can be made immediately. First, the E_4 -term contains the polynomial ring $\mathbb{M}_2[P, \Delta]$, and the E_4 -term is free as a module over $\mathbb{M}_2[P, \Delta]$. However, beware that the E_4 -term is not of the form $\mathbb{M}_2[P, \Delta] \otimes_{\mathbb{M}_2} B$, because of the relations in part (10) of the theorem.



The E_4 -term also contains the polynomial ring $\mathbb{M}_2[g]$, but the E_4 -term is not free as a module over $\mathbb{M}_2[g]$. For example, $h_0^3 g = 0$, but h_0^3 is not zero. However, the ideal generated by g is free over $\mathbb{M}_2[g]$.

Some parts of the E_4 -term are depicted in the following chart. The horizontal axis is the stem, and the vertical axis is the Adams filtration. Solid circles indicate copies of \mathbb{M}_2 , while open circles indicate copies of \mathbb{M}_2/τ . Vertical lines indicate multiplication by h_0 , lines of slope 1 indicate multiplication by h_1 , and lines of slope $\frac{1}{3}$ indicate multiplication by h_2 . Dashed lines indicate that the multiplication hits τ times a generator. For example, the relation $h_0^2 h_2 = \tau h_1^3$ occurs in the 3-stem.

Vertical arrows indicate infinite towers of copies of \mathbb{M}_2 connected by h_0 multiplications. Diagonal arrows indicate infinite towers of copies of \mathbb{M}_2/τ connected by h_1 multiplications.

For legibility, we have omitted most of the multiples of P . We have only shown a few elements in red to express multiplicative relations with elements that are not multiples of P . We have also omitted the strict multiples of Δ and Δ^2 .

The green parts of the figure consist of elements that are multiples of g , the blue parts consist of elements that are multiples of g^2 , and the magenta parts consist of elements that are multiples of g^3 . Observe that the blue part is a shifted copy of the green part. Similarly, if the figure were larger, the magenta part would be another shifted copy.



3.6. E_∞ -term of the motivic May spectral sequence. Having described the E_4 -term of the motivic May spectral sequence, we are now ready to compute the E_∞ -term.

As for the d_2 differentials, the d_4 differentials must be compatible with the differentials in the classical May spectral sequence. The only generator that supports a d_4 differential is Δ :

$$d_4(\Delta) = \tau^2 h_2 g.$$

A straightforward computation now gives the E_5 -term of the motivic May spectral sequence. By inspection, there are no higher differentials, so $E_5 = E_\infty$. The generators of E_∞ are listed in the following table, and the relations are listed in the next theorem.

Although Δ does not survive past E_4 , we shall write Δ for the Massey product operators $\langle \tau^2 g, h_2, - \rangle$ and $\langle h_2, \tau^2 g, - \rangle$.

TABLE 5. Generators of the E_∞ -term of the motivic May spectral sequence

generator	degree	generator	degree
h_0	(1, 0, 1, 0)	ν	(7, 15, 3, 8)
h_1	(1, 1, 1, 1)	e	(8, 17, 4, 10)
h_2	(1, 3, 1, 2)	g	(8, 20, 4, 12)
P	(8, 8, 4, 4)	$\Delta h_1 = \langle \tau^2 g, h_2, h_1 \rangle$	(13, 25, 5, 13)
c	(5, 8, 3, 5)	$\Delta c = \langle h_2, \tau^2 g, c \rangle$	(17, 32, 7, 17)
u	(5, 11, 3, 7)	$\Delta u = \langle h_2, \tau^2 g, u \rangle$	(17, 35, 7, 19)
α	(7, 12, 3, 6)	Δ^2	(24, 48, 8, 24)
d	(8, 14, 4, 8)		

Theorem 3.7. *The E_∞ -term of the motivic May spectral sequence is generated over \mathbb{M}_2 by the elements listed in Table 5, subject to the relations expressed in the multiplication table below, as well as the following relations:*

- (1) $h_0 h_1, h_1 h_2, h_0^2 h_2 + \tau h_1^3, h_0 h_2^2, h_2^3$.
- (2) $\tau u, \tau h_1^2 c, \tau h_1^2 \Delta c, \tau \Delta u, \tau c d, \tau c e, \tau c g, \tau d \Delta c, \tau e \Delta c, \tau g \Delta c, \tau^2 h_2 g$.
- (3) $h_0^2 \nu + \tau h_1 d, h_0 h_2 \nu + \tau h_1 e, h_2^2 \nu + \tau h_1 g, h_0 \alpha \nu + \tau h_1^2 \Delta h_1$.
- (4) $h_2 d + h_0 e, h_2 e + h_0 g, h_2 \alpha + h_0 \nu$.
- (5) $h_0 c, h_2 c, h_0 u, h_2 u, h_1 \alpha, h_1 \nu, h_0 \Delta h_1, h_2 \Delta h_1, h_0 \Delta c, h_2 \Delta c, h_0 \Delta u, h_2 \Delta u, h_0 \nu^2, h_2 \nu^2$.
- (6) $h_0^2 d + P h_2^2, h_0 \alpha d + P h_2 \nu, \alpha^2 d + P \nu^2$.
- (7) $\alpha^2 \nu + \tau d \Delta h_1, \alpha \nu^2 + \tau e \Delta h_1, \nu^3 + \tau g \Delta h_1, \alpha^4 + h_0^4 \Delta^2$.

TABLE 6. Multiplication table for E_∞

	c	u	α	d	ν	e	Δh_1	Δc	Δu
c	$h_1^2 d$	$h_1^2 e$	0	cd	0	ud	$h_1 \Delta c$	$h_1 d \Delta h_1$	$h_1 e \Delta h_1$
u		$h_1^2 g$	0	ce	0	cg	$h_1 \Delta u$	$h_1 e \Delta h_1$	$h_1 g \Delta h_1$
α			α^2	αd	$\alpha \nu$	νd	0	0	0
d				$h_1^3 \Delta h_1 + Pg$	αe	de	$d \Delta h_1$	$d \Delta c$	$e \Delta c$
ν					ν^2	αg	0	0	0
e						dg	$e \Delta h_1$	$d \Delta u$	$g \Delta c$
Δh_1							$h_1^2 \Delta^2$	$h_1 c \Delta^2$	$h_1 u \Delta^2$
Δc								$h_1^2 d \Delta^2$	$h_1^2 e \Delta^2$
Δu									$h_1^2 g \Delta^2$

Note that the E_∞ -term is free as a module over $\mathbb{M}_2[P, \Delta^2]$. It is not free as a module over $\mathbb{M}_2[g]$, but the ideal generated by g^2 is free over $\mathbb{M}_2[g]$.

Some parts of the E_∞ -term are depicted in the following chart. The horizontal axis is the stem, and the vertical axis is the Adams filtration. Solid circles indicate copies of \mathbb{M}_2 , open circles indicate copies of \mathbb{M}_2/τ , open circles with dots indicate copies of \mathbb{M}_2/τ^2 , and open boxes indicate copies of \mathbb{M}_2/τ^3 . Vertical lines indicate multiplication by h_0 , lines of slope 1 indicate multiplication by h_1 , and lines of slope $\frac{1}{3}$ indicate multiplication by h_2 . Dashed lines indicate that the multiplication hits τ times a generator.

Most of the multiples of P are not shown. A few multiples of P are shown in red to express multiplicative relations with elements that are not multiples of P . Also, multiples of Δ^2 are not shown.

Multiples of g are shown in green, multiples of g^2 are shown in blue, and multiples of g^3 are shown in magenta. If the diagram were larger, the magenta portion of the diagram would be a shifted copy of the blue portion.

Note the class $de\Delta h_1$ in the 56-stem. Classically $de\Delta h_1 = \alpha^3 g$, but motivically we only have $\alpha^3 g = \tau de\Delta h_1$. Thus $\tau de\Delta h_1$ is a multiple of g , but $de\Delta h_1$ is not a multiple of g .

4. COMPUTATION OF $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$

With the E_∞ -term of the motivic May spectral sequence in hand from the previous section, we have nearly computed $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$. It only remains to resolve some hidden extensions in the multiplicative structure.

4.1. Massey products. In this section we compute some Massey products in $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$ that will be needed to resolve the extension problems. All Massey products that we consider have zero indeterminacy.

Lemma 4.2. *In $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$,*

- (1) $\tau h_1^2 = \langle h_0, h_1, h_0 \rangle$.
- (2) $h_0 h_2 = \langle h_1, h_0, h_1 \rangle$.
- (3) $h_2^2 = \langle h_1, h_2, h_1 \rangle$.

There are several ways of understanding these formulas. The simplest is to note that these formulas are already known in $\text{Ext}_{A(2)\text{cl}}(\mathbb{F}_2, \mathbb{F}_2)$, and comparison gives the motivic formulas also. Another approach is to compute a minimal resolution for $A(2)$ in low dimensions and to compute Yoneda products and Massey products explicitly in terms of chain maps.

Lemma 4.3. *In $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$,*

- (1) $\tau d = \langle h_0, h_1, \alpha \rangle$.
- (2) $\tau e = \langle h_0, h_1, \nu \rangle$.
- (3) $\tau e = \langle h_2, h_1, \alpha \rangle$.
- (4) $\tau g = \langle h_2, h_1, \nu \rangle$.

Proof. For the first formula, we have

$$h_1 \langle h_0, h_1, \alpha \rangle = \langle h_1, h_0, h_1 \rangle \alpha = h_0 h_2 \alpha = \tau h_1 d.$$

In the relevant dimension, multiplication by h_1 is injective, so $\tau d = \langle h_0, h_1, \alpha \rangle$.

The proofs of the other formulas are similar. \square

Lemma 4.4. *In $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$,*

- (1) $c = \langle h_1, h_0, h_2^2 \rangle$.
- (2) $u = \langle h_1, h_2, h_2^2 \rangle$.
- (3) $\alpha = \langle h_0, h_1, h_2, \tau h_2^2 \rangle$.
- (4) $h_0 d = \langle \tau h_1 c, h_1, h_2 \rangle$.
- (5) $\alpha e = \langle \tau g, c, h_0 \rangle$.
- (6) $\alpha g = \langle \tau g, u, h_0 \rangle$.

Proof. The formulas are already true in the E_4 -term of the motivic May spectral sequence. They can be verified by computations with explicit cocycles. There are no extension problems to resolve in passing from E_4 to $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$. \square

We remind the reader that Δh_1 , Δc , and Δu are defined as follows.

Definition 4.5. *In $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$,*

- (1) $\Delta h_1 = \langle \tau^2 g, h_2, h_1 \rangle$.
- (2) $\Delta c = \langle h_2, \tau^2 g, c \rangle$.
- (3) $\Delta u = \langle h_2, \tau^2 g, u \rangle$.

4.6. Multiplicative Extensions.

Lemma 4.7. *In $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$, the elements $\tau\Delta u$, $\tau d\Delta c$, $\tau e\Delta c$, and $\tau g\Delta c$ are all zero.*

Proof. For the first element, note that $\tau\langle u, \tau^2 g, h_2 \rangle \subseteq \langle 0, \tau^2 g, h_2 \rangle = 0$.

For the second element,

$$\langle h_2, \tau^2 g, c \rangle \tau d = h_2 \langle \tau^2 g, c, \tau d \rangle,$$

which equals zero since $\langle \tau^2 g, c, \tau d \rangle$ is zero for dimension reasons. The argument for the last two elements is similar. \square

Lemma 4.8. *In $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$, we have*

- (1) $d\Delta u = e\Delta c$.
- (2) $e\Delta u = g\Delta c$.

Proof. We know from E_∞ that $e\Delta u$ equals either $g\Delta c$ or $g\Delta c + \tau\alpha g^2$. We have already shown that $\tau \cdot g\Delta c$ and $\tau \cdot e\Delta u$ are both zero, but $\tau^3\alpha g^2$ is non-zero. Therefore, $e\Delta u = g\Delta c$.

The proof of the second formula is similar. \square

Lemma 4.9. *In $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$, we have the following formulas:*

- (1) $h_0\Delta c = \tau h_0\alpha g$.
- (2) $h_0\Delta u = \tau h_0\nu g$.
- (3) $h_2\Delta c = \tau h_0\nu g$.
- (4) $h_2\Delta u = \tau h_2\nu g$.

Proof. For the first formula,

$$h_0\langle c, \tau^2 g, h_2 \rangle = \langle h_0, c, \tau g \rangle \tau h_2 = \tau h_2\alpha e = \tau h_0\alpha g.$$

The proof of the second formula is similar.

The third and fourth formulas follow easily by multiplying the first and second formulas by h_2 , using that multiplication by h_0 is injective in the relevant dimensions. \square

Lemma 4.10. *In $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$, we have the following formulas:*

- (1) $\alpha c = \tau h_0^2 g$.
- (2) $\nu c = \tau h_0 h_2 g$.
- (3) $\alpha u = \tau h_0 h_2 g$.
- (4) $\nu u = \tau h_2^2 g$.
- (5) $\alpha\Delta h_1 = \tau^3 e g$.
- (6) $\nu\Delta h_1 = \tau^3 g^2$.

Proof. For the first formula,

$$\alpha\langle h_1, h_0, h_2^2 \rangle = \langle \alpha, h_1, h_0 \rangle h_2^2 = \tau d h_2^2 = \tau h_0^2 g.$$

The other calculations are similar. \square

Lemma 4.11. *In $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$, the products $\alpha\Delta c$, $\alpha\Delta u$, $\nu\Delta c$, and $\nu\Delta u$ are all zero.*

Proof. For the first product,

$$\alpha\Delta c = \langle h_0, h_1, h_2, \tau h_2^2 \rangle \Delta c \subseteq \langle h_0, h_1, h_2, \tau^3 h_1 e g \rangle = \langle h_0, h_1, h_2, 0 \rangle = 0.$$

The second product vanishes for similar reasons.

For the third product,

$$\tau\nu\Delta c = \langle h_2, \tau^2 g, c \rangle \tau\nu = h_2 \langle \tau^2 g, c, \tau\nu \rangle,$$

which is zero because $\langle \tau^2 g, c, \tau\nu \rangle$ is zero for dimension reasons. Therefore, $\nu\Delta c$ does not equal $\tau\alpha\nu g$ since $\tau^2\alpha\nu g$ is not zero. The fourth product vanishes for similar reasons. \square

Lemma 4.12. *In $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$, we have the following formulas:*

- (1) $(\Delta h_1)^2 = h_1^2 \Delta^2 + \tau^2 \nu^2 g.$
- (2) $\Delta h_1 \cdot \Delta c = h_1 c \Delta^2.$
- (3) $\Delta h_1 \cdot \Delta u = h_1 u \Delta^2.$
- (4) $(\Delta c)^2 = h_1^2 d \Delta^2.$
- (5) $\Delta c \cdot \Delta u = h_1^2 e \Delta^2.$
- (6) $(\Delta u)^2 = h_1^2 g \Delta^2.$

Proof. For the first formula, we know from E_∞ that $(\Delta h_1)^2$ equals either $h_1^2 \Delta^2$ or $h_1^2 \Delta^2 + \tau^2 \nu^2 g$. We have already shown that $\nu(\Delta h_1)^2 = \tau^3 g^2 \Delta h_1$ is non-zero, but $\nu \cdot h_1^2 \Delta^2$ is zero. It follows that $(\Delta h_1)^2 = h_1^2 \Delta^2 + \nu^2 g$.

For the second formula, we know from E_∞ that $\Delta h_1 \cdot \Delta c$ equals either $h_1 c \Delta^2$ or $h_1 c \Delta^2 + \tau^4 e g^2$. We have already shown that $\tau g \cdot \Delta h_1 \Delta c$ and $\tau g \cdot h_1 c \Delta^2$ are both zero, but $\tau g \cdot \tau^4 e g^3$ is non-zero.

For the third formula, we know from E_∞ that $\Delta h_1 \cdot \Delta u$ equals $h_1 u \Delta^2$ or $h_1 u \Delta^2 + \tau^4 g^3$. We have already shown that $\tau \cdot \Delta h_1 \Delta u$ and $\tau \cdot h_1 u \Delta^2$ are zero, but $\tau \cdot \tau^4 g^3$ is non-zero.

For the fourth formula, we know from E_∞ that $(\Delta c)^2$ equals $h_1^2 d \Delta^2$ or $h_1^2 d \Delta^2 + \tau^2 \alpha^2 g^2$. Observe that

$$\tau \cdot (\Delta c)^2 = \langle h_2, \tau^2 g, c \rangle \tau \Delta c = h_2 \langle \tau^2 g, c, \tau \Delta c \rangle,$$

which must be zero because $\langle \tau^2 g, c, \tau \Delta c \rangle$ is zero for dimension reasons. Also, $\tau \cdot h_1^2 d \Delta^2$ is zero, but $\tau \cdot \tau^2 \alpha^2 g^2$ is non-zero.

For the fifth formula, we know from E_∞ that $\Delta c \cdot \Delta u$ equals $h_1^2 e \Delta^2$ or $h_1^2 e \Delta^2 + \tau^2 \alpha \nu g^2$. We already know that $\tau \cdot (\Delta c \cdot \Delta u)$ and $\tau \cdot h_1^2 e \Delta^2$ are zero, but $\tau \cdot \tau^2 \alpha \nu g^2$ is non-zero.

For the sixth formula, we know from E_∞ that $(\Delta u)^2$ equals $h_1^2 g \Delta^2$ or $h_1^2 g \Delta^2 + \tau^2 \nu^2 g^2$. We already know that $\tau \cdot (\Delta u)^2$ and $\tau \cdot h_1^2 g \Delta^2$ are zero, but $\tau \cdot \tau^2 \nu^2 g^2$ is non-zero. \square

Lemma 4.13. *In $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$, we have $\tau h_1^2 \Delta c = \tau^2 h_0 d g$ but $h_1^2 \Delta c$ does not equal $\tau h_0 d g$.*

Proof. From E_∞ , we know that $h_1 \cdot h_1^2 \Delta c$ is non-zero, but $h_1 \cdot \tau h_0 d g$ is zero. Therefore, $h_1^2 \Delta c$ and $\tau h_0 d g$ cannot be equal.

From E_∞ , we also know that $h_1 \Delta c$ equals $c \Delta h_1$. Therefore,

$$\tau h_1^2 \Delta c = \tau h_1 c \Delta h_1 = \tau h_1 c \langle h_1, h_2, \tau^2 g \rangle = \langle \tau h_1 c, h_1, h_2 \rangle \tau^2 g.$$

It follows that $\tau h_1^2 \Delta c = \tau^2 h_0 d g$. \square

Lemma 4.14. *In $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$, we have $\alpha^4 = h_0^4 \Delta^2 + \tau^4 P g^2$.*

Proof. From E_∞ , we know that α^4 equals either $h_0^4\Delta^2$ or $h_0^4\Delta^2 + \tau^4Pg^2$. We have already shown that $\Delta h_1 \cdot \alpha^4$ equals

$$\alpha^3 \cdot \tau^3 eg = \tau^3 \alpha^2 e \cdot \nu e = \tau^4 d \Delta h_1 \cdot dg = \tau^4 g \Delta h_1 (Pg + h_1^3 \Delta h_1) = \tau^4 Pg^2 \Delta h_1.$$

However, $\Delta h_1 \cdot h_0^4 \Delta^2$ equals zero. It follows that α^4 must equal $h_0^4 \Delta^2 + \tau^4 Pg^2$. \square

4.15. Ring structure of $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$. Finally, we assemble the calculations made in the previous sections to compute $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$ as a ring. The generators are listed in the following table. All possible multiplicative extensions have been resolved in the previous section.

TABLE 7. Generators of $\text{Ext}_{A(2)}$

generator	degree	generator	degree
h_0	(0, 1, 0)	ν	(15, 3, 8)
h_1	(1, 1, 1)	e	(17, 4, 10)
h_2	(3, 1, 2)	g	(20, 4, 12)
P	(8, 4, 4)	Δh_1	(25, 5, 13)
c	(8, 3, 5)	Δc	(32, 7, 17)
u	(11, 3, 7)	Δu	(35, 7, 19)
α	(12, 3, 6)	Δ^2	(48, 8, 24)
d	(14, 4, 8)		

Theorem 4.16. *The ring $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$ is generated over \mathbb{M}_2 by the elements listed in Table 7, subject to the relations expressed in the multiplication table below, as well as the following relations:*

- (1) $h_0 h_1, h_1 h_2, h_0^2 h_2 + \tau h_1^3, h_0 h_2^2, h_2^3$.
- (2) $\tau u, \tau h_1^2 c, \tau h_1^2 \Delta c + \tau^2 h_0 d g, \tau \Delta u, \tau c d, \tau c e, \tau c g, \tau d \Delta c, \tau e \Delta c, \tau g \Delta c, \tau^2 h_2 g$.
- (3) $h_0^2 \nu + \tau h_1 d, h_0 h_2 \nu + \tau h_1 e, h_2^2 \nu + \tau h_1 g, h_0 \alpha \nu + \tau h_1^2 \Delta h_1$.
- (4) $h_2 d + h_0 e, h_2 e + h_0 g, h_2 \alpha + h_0 \nu$.
- (5) $h_0 c, h_2 c, h_0 u, h_2 u, h_1 \alpha, h_1 \nu, h_0 \Delta h_1, h_2 \Delta h_1, h_0 \Delta c + \tau h_0 \alpha g, h_2 \Delta c + \tau h_0 \nu g, h_0 \Delta u + \tau h_0 \nu g, h_2 \Delta u + \tau h_2 \nu g, h_0 \nu^2, h_2 \nu^2$.
- (6) $h_0^2 d + P h_2^2, h_0 \alpha d + P h_2 \nu, \alpha^2 d + P \nu^2$.
- (7) $\alpha^2 \nu + \tau d \Delta h_1, \alpha \nu^2 + \tau e \Delta h_1, \nu^3 + \tau g \Delta h_1, \alpha^4 + h_0^4 \Delta^2 + \tau^4 P g^2$.

TABLE 8. Multiplication table for $\text{Ext}_{A(2)}$

	c	u	α	d	ν
c	$h_1^2 d$	$h_1^2 e$	$\tau h_0^2 g$	cd	$\tau h_0 h_2 g$
u		$h_1^2 g$	$\tau h_0 h_2 g$	ce	$\tau h_2^2 g$
α			α^2	αd	$\alpha \nu$
d				$h_1^3 \Delta h_1 + Pg$	αe
ν					ν^2

	e	Δh_1	Δc	Δu
c	ud	$h_1 \Delta c$	$h_1 d \Delta h_1$	$h_1 e \Delta h_1$
u	cg	$h_1 \Delta u$	$h_1 e \Delta h_1$	$h_1 g \Delta h_1$
α	νd	$\tau^3 e g$	0	0
d	de	$d \Delta h_1$	$d \Delta c$	$e \Delta c$
ν	αg	$\tau^3 g^2$	0	0
e	dg	$e \Delta h_1$	$d \Delta u$	$g \Delta c$
Δh_1		$h_1^2 \Delta^2 + \tau^2 \nu^2 g$	$h_1 c \Delta^2$	$h_1 u \Delta^2$
Δc			$h_1^2 d \Delta^2$	$h_1^2 e \Delta^2$
Δu				$h_1^2 g \Delta^2$

Some parts of $\text{Ext}_{A(2)}(\mathbb{M}_2, \mathbb{M}_2)$ are depicted in the following chart. The notation is the same as in the chart for the E_∞ -term of the motivic May spectral sequence.

The most interesting difference occurs with Δc . Note that Δc supports exotic multiplications by h_0 and h_2 . Note also that $\tau h_1^2 \Delta c$ is no longer zero; this is an exotic τ -extension. In fact, $h_1^2 \Delta c$ is the sum $(h_1^2 \Delta c + \tau h_0 dg) + \tau h_0 dg$, where the first term is killed by τ and the second is killed by no power of τ .



Similarly to Δc , the classes Δu , $d \Delta c$, and $e \Delta c$ all support exotic multiplications by h_0 and h_2 .

4.17. An Adams spectral sequence? At this point, it is natural to wonder whether the cohomology of $A(2)$ is the E_2 -term of an Adams spectral sequence that converges to the homotopy of some motivic spectrum that is analogous to tmf .

Assuming that such a spectral sequence exists, it is possible to determine the d_2 -differentials by comparison to calculations over the full motivic Steenrod algebra A and by comparison to calculations over the classical $A(2)_{cl}$.

For example, comparison to the full motivic Steenrod algebra implies that $d_2(e) = h_1^2 d$ [DI]. Then the relation $ud = ce$ implies that $d_2(u) = h_1^2 c$. Similar kinds of arguments allow one to compute the entire d_2 differential.

We have not pursued this computation any further because there is a simpler way of making the same speculative calculation that we present in the following section.



5. ADAMS-NOVIKOV SPECTRAL SEQUENCE FOR “MOTIVIC MODULAR FORMS”

In this section, we make a leap of faith and assume that there exists a motivic spectrum mmf , called “motivic modular forms”, defined over $\text{Spec } \mathbb{C}$. We assume that the topological realization of this motivic spectrum is the classical spectrum tmf . We also assume that the homotopy of mmf can be computed by an Adams-Novikov spectral sequence whose E_2 -term is the cohomology of a version of the elliptic curves Hopf algebroid.

Recall that the elliptic curves Hopf algebroid localized at the prime 2 has the form $(\tilde{A}_{\text{cl}}, \tilde{\Gamma}_{\text{cl}})$, where \tilde{A}_{cl} is the ring $\mathbb{Z}_{(2)}[a_1, a_3, a_4, a_6]$ and $\tilde{\Gamma}_{\text{cl}} = \tilde{A}_{\text{cl}}[s, t]$ [B].

We write $\mathbb{M}_{(2)}$ for the ring $\mathbb{Z}_{(2)}[\tau]$, i.e., the 2-local motivic cohomology of a point.

Definition 5.1. *The motivic elliptic curves Hopf algebroid localized at 2 is $(\tilde{A}, \tilde{\Gamma})$, where $\tilde{A} = \mathbb{M}_{(2)}[a_1, a_3, a_4, a_6]$ and $\tilde{\Gamma} = \tilde{A}[s, t]$. The bidegree of a_i is $(2i, i)$, while the bidegrees of s and t are $(2, 1)$ and $(6, 3)$ respectively.*

The structure maps of $(\tilde{A}, \tilde{\Gamma})$ are defined to be compatible with the classical structure maps.

One may compute the cohomology of $(\tilde{A}, \tilde{\Gamma})$ as in [B, Section 7]. It turns out that the weights introduce no new complications. The answer is essentially the same as the classical answer, as shown in the chart on page 30 of [B]. In fact, the motivic answer is equal to the classical answer tensored over $\mathbb{Z}_{(2)}$ with $\mathbb{M}_{(2)}$. Also, the generators are assigned weights, but these are easy to determine. At location (s, t) of the chart, the weight of the generator is $\frac{s+t}{2}$.

We now have the E_2 -term of a (speculative) spectral sequence for computing the homotopy groups of mmf at the prime 2. Next we consider differentials and compute the E_∞ -term.

We assume that the topological realization of mmf is tmf . This implies that the motivic differentials are algebraically compatible with the classical differentials, i.e., they are equal after inverting τ . Using this fact, it turns out that the motivic differentials are entirely determined by the classical differentials. However, one must be careful with the weights. We describe this issue next.

The first classical differential is a d_3 that hits h_1^4 . The input to this differential has weight 3 (since it lies in the 5-stem and has filtration 1), but h_1^4 has weight 4. It follows that d_3 hits τh_1^4 in the motivic calculation, not h_1^4 as in the classical calculation. As a result, instead of finding that $h_1^4 = 0$, we have that h_1^k is non-zero for all $k \geq 0$ but $\tau h_1^k = 0$ for $k \geq 4$.

The same situation occurs for the h_1 -multiples of many of the classes in filtration 0. Classically, d_3 -differentials tell us that these classes are killed by h_1^3 . Motivically, these classes are not killed by h_1^k for any k , but they are killed by τh_1^3 .

In order to simplify our discussion and make our diagrams legible, from here on we shall ignore these classes in filtration 0 and their h_1^k -multiples without further mention.

Classically, the next differential is $d_5(\Delta) = h_2g$. Since Δ has weight 12 while h_2g has weight 14, the motivic differential is $d_5(\Delta) = \tau^2 h_2g$. We observe that h_2g survives the motivic calculation, but $\tau^2 h_2g = 0$.

The entire calculation proceeds similarly. Elementary bookkeeping with the weights shows that a non-zero d_{2k+1} -differential hits τ^k times a generator. In other words, non-zero d_{2k+1} -differentials produce classes that are killed by τ^k .

The results of the full analysis are shown in the following tables. These are the E_∞ -term of a (speculative) spectral sequence converging to the 2-complete homotopy groups of the (speculative) motivic spectrum mmf .

Boxes represent copies of $\mathbb{Z}_2[\tau]$. Solid dots represent copies of $\mathbb{Z}/2[\tau]$. A number k represents $\mathbb{Z}/2[\tau]/\tau^k$. A horizontal row of symbols at a single location represents extensions by 2. For example, at location $(3, 1)$ there is a copy of $\mathbb{Z}/4[\tau]$, and at location $(23, 5)$ there is a copy of $\mathbb{Z}/4[\tau]/\tau^2$. More interestingly, at location $(20, 4)$ there is a copy of $\mathbb{Z}/8[\tau]$; at location $(40, 8)$ there is a copy of $\mathbb{Z}/8[\tau]/4\tau^2$; and at location $(120, 24)$ there is a copy of $\mathbb{Z}/8[\tau]/(\tau^{11}, 2\tau^6, 4\tau^2)$.

Observe that if we ignore the numbers and just consider the boxes and solid dots, we obtain the classical picture. This expresses the principle that the classical calculation is recovered by inverting τ .

The blue lines show extensions by η and ν that are detected in the E_∞ -term. They take generators to generators in the predictable way. The blue arrows of slope 1 indicate infinite sequences of elements that are connected by h_1 -multiplications and are killed by τ . We have not shown them explicitly in order to make the diagrams legible.

The red lines show exotic extensions by 2, η , and ν that are not detected in E_∞ . Beware that these exotic extensions do not take generators to generators. The first example occurs in the 3-stem, where $4\nu = \tau\eta^3$. The next example is $\nu e[25, 1] = \tau^2\epsilon\bar{\kappa}$. As always, the required power of τ is easily determined by the weights. Here we write $e[s, t]$ for the homotopy element that is represented by the generator in stem s and filtration t . Note that ϵ and $\bar{\kappa}$ are names for $e[8, 2]$ and $e[20, 4]$.

The motivic calculation shows many more exotic extensions than the classical calculation. However, all of the motivic extensions are easily implied by the classical ones. We remark that the charts in [B] inadvertently failed to indicate the exotic extension $\eta e[104, 2] = e[105, 17]$.

Example 5.2. Note that $\pi_{0,*}mmf$ is $\mathbb{Z}_2[\tau]$. Hence for any k , $\pi_{k,*}mmf$ is a $\mathbb{Z}_2[\tau]$ -module.

Consider $\pi_{120,*}mmf$. This module has 6 free generators of weight 60. These arise in filtration 0; only one is shown on the chart.

In addition, $\pi_{120,*}mmf$ contains many copies of $\mathbb{Z}/2[\tau]/\tau$; these arise as h_1 -multiples and are not shown explicitly on the chart.


Finally, $\pi_{120,*}mmf$ contains a copy of $\mathbb{Z}/8[\tau]/(\tau^{11}, 2\tau^6, 4\tau^2)$, generated by $\bar{\kappa}^6$ with weight 72.


Example 5.3. The $\mathbb{Z}_2[\tau]$ -module $\pi_{170,*}mmf$ contains a copy of


$$\mathbb{Z}/8[x, y, z]/(\tau^2x, \tau^6y, \tau^{11}z, 2x = \tau^4y, 2y = \tau^6z)$$


where the generators x , y , and z have weights 88, 92, and 98 respectively.

Remark 5.4. In [B, Proposition 8.4], the classical relation $\epsilon\kappa = \eta^2\bar{\kappa}$ is computed by consideration of $\eta^3\bar{\kappa}$. Here κ is a name for $e[14, 2]$. However, the argument is not quite so simple, since this classical product is zero. Motivically, this complication does not occur. We can show that $\epsilon\kappa = \tau\eta^2\bar{\kappa}$ by considering $\tau\eta^3\bar{\kappa}$, which is non-zero. (Note that $\tau^3\eta^3\bar{\kappa}$ is zero, but this does not interfere with the computation.)









REFERENCES

- [B] T. Bauer, *Computation of the homotopy of the spectrum tmf* , Geometry & Topology Monographs **13** (2008) 1140.
- [DI] D. Dugger and D. C. Isaksen, *The motivic Adams spectral sequence*, preprint (2009).
- [H1] M. J. Hopkins, *Topological modular forms, the Witten genus, and the theorem of the cube*, Proc. International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994), 554–565, Birkhäuser, 1995.
- [H2] M. J. Hopkins, *Algebraic topology and modular forms*, Proc. International Congress of Mathematicians, Vol. I (Beijing, 2002), 291–317, Higher Ed. Press, 2002.
- [HM] M. J. Hopkins and M. Mahowald, *From elliptic curves to homotopy theory*, preprint, 1998.
- [HKO] P. Hu, I. Kriz, and K. Ormsby, *Some remarks on motivic homotopy theory over algebraically closed fields*, preprint, 2008.
- [IS] N. Shimada and A. Iwai, *On the cohomology of some Hopf algebras*, Nagoya Math. J. **30** (1967) 103–111.
- [M1] J. P. May, *The cohomology of restricted Lie algebras and of Hopf algebras; application to the Steenrod algebra*, Ph.D. dissertation, Princeton, 1964.
- [M2] J. P. May, *Cohomology of $A(2)$* , unpublished notes.
- [V1] V. Voevodsky, *Motivic cohomology with $\mathbb{Z}/2$ -coefficients*, Publ. Math. Inst. Hautes Études Sci. **98** (2003) 59–104.
- [V2] V. Voevodsky, *Reduced power operations in motivic cohomology*, Publ. Math. Inst. Hautes Études Sci. **98** (2003) 1–57.
- [V3] V. Voevodsky, *Motivic Eilenberg-Mac Lane spaces*, preprint, 2007.

DEPARTMENT OF MATHEMATICS, WAYNE STATE UNIVERSITY, DETROIT, MI 48202
E-mail address: isaksen@math.wayne.edu