Comments on "Modelling the gap size distribution of parked cars"

M. Girardi

Universidade Federal do Pampa - Caixa Postal 7, 96412-420, Bagé, Rio Grande do Sul, Brazil.

Abstract

In this Comment we discuss some points concerning the modeling of parked cars proposed in the article by Rawal and Rodgers, Physica A (2005). We also introduce another approach to this problem which leads to a better description of the empirical data collected by the authors.

PACS: 05.20.-y; 02.50.Cw; 80.20.Mj

Key words: Parked cars; Simulations.

Rawal and Rodgers [1] presented in their article two different models to describe parked cars in a central area of London. They measured the distances (gaps) between about 500 cars in four streets, and observed that the gap distribution have minima for small and large gaps. Figure 1 exhibits a reproduction of the collected data.

In the first proposed model (model A), cars of length 1 are placed randomly on an infinite linear road if the vacancy is greater than or equal to $1 + \varepsilon$, with $\varepsilon \ge 0$. The extra space ε mimics the need to manoeuvre. For the full car road, the gap distribution which they found has a plateau for gaps lower than ε , decreasing monotonically for gaps of sizes lower then $1 + \varepsilon$. These results are very different from the empirical one.

In the second model (model B), a car of length 1 can park and stay in its position with probability p, and park, driving to the nearest car, with probability 1 - p. In this case, the distance y from the nearest car is chosen with probability f(y). Two functional forms for f(y) were tested to reproduce the empirical data, and the authors argued that f(y) = 6y(1 - y) fits the data very well.

Email address: mauricio.girardi@unipampa.edu.br (M. Girardi).

Three points about the above article deserve some discussion:

- (1) In their first model, the idea to introduce of an extra space to manoeuvre was good, and seems to be an important ingredient to any model for parked cars. However, it was not included in their second model.
- (2) In model B, the functional form of f(y) had no justification or explanation, and the authors gave no argument to why they used it.
- (3) The results obtained from the second model are qualitatively different from the empirical ones, since the gap distribution does not have the convex curvature for large gaps.

In this way, we propose a model where the extra space ε is maintained, while another dynamics for car parking is chosen. Here, a car is displaced between the two other cars with a Gaussian probability $p(x) \propto \exp\left(-\frac{(x-x_0)^2}{\sigma^2}\right)$, where xis the distance from the center x_0 of the vacancy. This dynamics resembles the drivers desire to leave a secure gap between front and rear cars (the maximum probability is when the car stays at the center of the vacancy), and their ability to measure this distance, given by the standard deviation σ .

Figure 2 shows simulation results [2,3] of the present model for various values of ε and σ . In figure 2a, σ is equal to gap/10, indicating the case where drivers are very skillful, and the cars are placed almost at center of the gap. The opposite case ($\sigma = gap/0.1$) is shown in figure 2d, where the cars are placed at random in the gap, as in model A of ref. [1]. As can be seen, changing the value of ε does not modify qualitatively the gap distribution, except for $\varepsilon = 0$ where the distribution is monotonically decreasing. The interesting results are exhibited in figures 2b and 2c for $\sigma = gap/4$ and $\sigma = gap/2$ respectively. Here the curves resemble the empirical data, with lower densities for small gaps (non zero densities), a maximum at a typical gap size, and a convex curvature of the distribution for large gaps. This contrasts with Fig. 9 of ref. [1], which is concave for large gaps.

Concluding, we found a model for parked cars which agrees qualitatively with empirical data for full car roads. The essential ingredients are the early employed extra space for manoeuvre, and a dynamics where the driver's skill to park his car keeping equal gaps from rear and front cars is represented by the standard deviation in the Gaussian probability to put the car in the center of the vacancy.

Acknowledgements

We would like to thank the financial support of the Brazilian agencies FAPESP and CNPq.

Figures

1 - Gap distribution between parked cars in the central area of London (same as Fig1. of ref. [1]).

2 - Gap size distribution for (a) $\sigma = gap/10$, (b) $\sigma = gap/4$, (c) $\sigma = gap/2$ and $\sigma = gap/0.1$. In each plot, from left to right, we have $\varepsilon = 0, 1, 3, 5, 7$ and 9.

References

- [1] S. Rawal and G. J. Rodgers, Physica A 346, 621 (2005).
- [2] M. E. Newman and G. T. Barkema, Monte Carlo Methods in Statistical Physics (Clarendom Press, Oxford, 1999).
- [3] W. H. Press et al, "Numerical Recipes in Fortran 77: The Art of Scientific Computing" (Cambridge Press, Cambridge, 1992).




Fig. 1 - Girardi




Fig. 2 - Girardi

Gap size