

ON THE HEREDITARY PROXIMITY TO ℓ_1

S.A. ARGYROS, A. MANOUSSAKIS, AND A.M. PELCZAR

ABSTRACT. In the first part of the paper we present and discuss concepts of local and asymptotic hereditary proximity to ℓ_1 . The second part is devoted to a complete separation of the hereditary local proximity to ℓ_1 from the asymptotic one. More precisely for every countable ordinal ξ we construct a separable reflexive space \mathfrak{X}_ξ such that every infinite dimensional subspace of it has Bourgain ℓ_1 -index greater than ω^ξ and the space itself has no ℓ_1 -spreading model. We also present a reflexive HI space admitting no ℓ_p as a spreading model.

1. INTRODUCTION

Concepts of proximity to a classical ℓ_p (or c_0) space play a significant role to the understanding of the structure of a Banach space. They are categorized as follows:

The first one is the global proximity to ℓ_p which simply means that ℓ_p is isomorphic to a subspace Y of X . The local proximity which occurs more frequently, due to J.L. Krivine's theorem [19], is measured through the Bourgain ℓ_p -index [10]. The last concept is the asymptotic proximity that varies from A. Brunel-L. Sucheston ℓ_p -spreading models, [11], to the asymptotic ℓ_p spaces. The latter class of Banach spaces appeared after B.S. Tsirelson space [28] that answered in negative the famous Banach's problem by showing that global proximity to some ℓ_p is not always possible.

It is easy to see that the global proximity to ℓ_p is the strongest one followed by the asymptotic one. The local proximity is the weakest among them. It is also known that the three classes are separated for each ℓ_p . Namely there are spaces with arbitrarily large local proximity to ℓ_p but no asymptotic one and similarly for the asymptotic and the global ones. The present paper is mainly devoted to the separation of the local and asymptotic proximity to ℓ_1 when the first one is hereditarily large. In particular our work is motivated by a result of the third named author stated as follows.

Theorem. ([25]) *Let X be a separable Banach space and ξ be a countable ordinal. If X is boundedly distortable and has hereditary Bourgain ℓ_1 -index greater than ω^ξ then it is saturated by asymptotic ℓ_1^ξ spaces.*

Let's recall that the hereditary Bourgain ℓ_p -index of a Banach space X is the minimum of Bourgain ℓ_p -index of its subspaces. In the sequel by the ℓ_p -index we will mean the Bourgain ℓ_p -index.

In view of the above theorem it is natural to ask how critical is the bounded distortion of X for the final conclusion. It is also worth adding that heredity assumptions for the local proximity to ℓ_1 could yield large asymptotic one. In this direction we prove the following.

Proposition. *Let (e_n) be a Schauder basis of a Banach space X such that the Bourgain ℓ_1 -tree supported by any subsequence of $(e_n)_{n \in \mathbb{N}}$ has order greater than ω^ξ . Then there exists a subsequence generating an ℓ_1^ξ -spreading model.*

2000 *Mathematics Subject Classification.* 46B20, 46B15, 03E10, 05A17.

Key words and phrases. Bourgain ℓ_1 -index, ℓ_1 -spreading model, attractors method.

Our aim is to show that large hereditary ℓ_1 -structure in a Banach space X does not imply in general any asymptotic one. More precisely the main goal at the present paper is to prove the next

Theorem. *For every countable ordinal ξ there exists a separable reflexive space \mathfrak{X}_ξ with the hereditary ℓ_1 -index greater than ω^ξ such that \mathfrak{X}_ξ does not admit an ℓ_1 -spreading model. Moreover the dual \mathfrak{X}_ξ^* has hereditary c_0 -index greater than ω^ξ and does not admit c_0 as a spreading model.*

Our approach in constructing the space \mathfrak{X}_ξ is based on mixed Tsirelson extensions of a ground set G_ξ using the method of attractors. The latter appeared in [8] and is extensively used in [3]. The method of attractors has two separated steps that we are about to describe.

In the first step an auxiliary space is constructed that partially solves the required problem. In our case for a given countable ordinal ξ we construct a ground set G_ξ such that the resulting space X_{G_ξ} is reflexive, has a Schauder basis $(e_n)_{n \in \mathbb{N}}$ and satisfies the following properties.

- i) The space X_{G_ξ} does not have an ℓ_1 -spreading model.
- ii) For every $L \in [\mathbb{N}]$ the ℓ_1 -index of the subspace $\langle (e_n)_{n \in L} \rangle$ is greater than ω^ξ .

The above stated proposition together with property i) of the space X_{G_ξ} indicate that property ii) requires special attention. Namely for every subsequence $(e_n)_{n \in L}$ of the basis the ℓ_1 -index of the corresponding subspace should be greater than ω^ξ but the ℓ_1 -tree supported by that subsequence should be of small height. To achieve those requirements we include in the set G_ξ a c_0 -tree generated by functionals of the form $m_{2^j-1}^{-2} \sum_{k \in B} e_k^*$, $\#B \leq n_{2^j-1}$ and the tree order induced by an appropriate coding function.

The proof that X_{G_ξ} does not have an ℓ_1 -spreading model heavily relies on combinatorics in particular on Ramsey theory.

The next step is to define the space X_ξ . For this purpose we proceed to a mixed Tsirelson extension K_ξ of the set G_ξ including also the attracting functionals. The space X_ξ is the completion of $(c_{00}(\mathbb{N}), \|\cdot\|_{K_\xi})$ where $\|\cdot\|_{K_\xi}$ is the norm induced by the set K_ξ .

For our approach the attracting functionals are of significant importance. They are the tool for transferring to every block subspace of X_ξ^* a large c_0 -subtree from the set G_ξ . Thus we are able to show that the hereditary c_0 index of X_ξ^* is greater than ω^ξ . This does not yield that the corresponding ℓ_1 -index of X_ξ is also greater than ω^ξ . Therefore we need one more step, namely the desired space \mathfrak{X}_ξ is a quotient X_ξ/X_ξ^L where $X_\xi^L = \langle (e_n)_{n \in L} \rangle$ with L a suitable subset of \mathbb{N} . Both spaces X_ξ, \mathfrak{X}_ξ do not have an ℓ_1 spreading model.

We proceed now to describe how the paper is organized.

Section 2 is devoted to preliminary notions and results.

In Section 3 we discuss different concepts of local and asymptotic proximity to ℓ_1 . More precisely we introduce the hereditary strategic ℓ_1 -index of a Banach space X denoted as $I_{hs}(X)$. This index is formulated in terms of Gowers game and Schreier families $\mathcal{S}_\xi, \xi < \omega_1$. A non-hereditary version of the aforementioned index, related to the asymptotic structures defined in [21], is also presented. The $I_{hs}(X)$ is essentially equivalent to the hereditary block ℓ_1 -index, denoted as $I_{hb}(X)$, in the following manner: First we show that for every countable ordinal ξ if $I_{hs}(X) > \xi$ then $I_{hb}(X) > \omega^\xi$. In the opposite direction Gowers dichotomy [15] yields that if $I_{hb}(X) > \omega^\xi$ then there exists a closed subspace Y of X with $I_{hs}(Y) > \xi$. Furthermore we examine the relation between the concepts of non-hereditary ℓ_1 -proximity. We finish Section 3 by showing strong correlation between the above notions in a subsequence setting, i.e. with block sequences replaced by subsequences of a fixed basis. The main result in this part is Proposition 3.7 which has been mentioned before.

With Section 4 we start dealing with our final goal namely the space \mathfrak{X}_ξ . Thus in this section for a given countable ordinal ξ we define a ground set G_ξ which serves as

a norming set for the aforementioned auxiliary space X_{G_ξ} . The set G_ξ includes a rich c_0 -tree of height greater than ω^ξ . This tree is defined as follows. First G_ξ contains all functionals of the form $m_{2^{j-1}}^{-2} \sum_{k \in B} e_k^*$ with $\#B \leq n_{2^{j-1}}$ where $(n_j)_{j \in \mathbb{N}}$, $(m_j)_{j \in \mathbb{N}}$ are appropriate increasing sequences of natural numbers. Next using a coding function σ , in a similar manner as in the classical work of B. Maurey and H. Rosenthal [22], we define a well-founded tree of σ -special sequences (f_1, \dots, f_d) ordered by the initial segment inclusion. Each f_l is of the form $m_{2^{j_l-1}}^{-2} \sum_{k \in B_l} e_k^*$, $\#B_l \leq n_{2^{j_l-1}}$. Then we include into G_ξ all $\sum_{l=1}^d \pm f_l$ where $(f_l)_{l=1}^d$ is a σ -special sequence.

A second ingredient of G_ξ is coming from James tree-like spaces [17] (see also [1], Chapter 13 or [3]). Namely we include all rational ℓ_2 -convex combinations of functionals $\sum_{l=1}^d \pm f_l$ with disjoint weights. We denote by $X_{G_\xi} = \overline{(c_{00}(\mathbb{N}), \|\cdot\|_{G_\xi})}$ the space with the set G_ξ as the norming set.

In Section 5 we present the basic properties of the space X_{G_ξ} . Namely it is reflexive, has a Schauder basis $(e_n)_{n \in \mathbb{N}}$, is ℓ_2 -saturated and for every $L \in [\mathbb{N}]$ the ℓ_1 -index of the subspace $\langle (e_n)_{n \in L} \rangle$ is greater than ω^ξ . Also we show a dual result. Namely for every $(e_n^*)_{n \in L}$ in the dual $X_{G_\xi}^*$ the c_0 -index of $\langle (e_n^*)_{n \in L} \rangle$ is similarly large.

In Section 6 it is shown that the space X_{G_ξ} does not have an ℓ_1 -spreading model. This is the most involved part of the study of X_{G_ξ} . This result is critical as with some small additional effort yields that the ultimate space \mathfrak{X}_ξ shares the same property. The difference of the space X_{G_ξ} from the earlier examples of spaces with no ℓ_p -spreading model, (i.e. [24]) is that X_{G_ξ} has a rich local ℓ_1 structure. Therefore the proof requires new tools which are of combinatorial nature. The main part of the proof is given by Proposition 6.3.

In Section 7 we define the set K_ξ which is the norming set of the intermediate space X_ξ . The ingredients of the set K_ξ are the following:

i) The set G_ξ is included into K_ξ . In particular K_ξ is a mixed Tsirelson extension of G_ξ .

ii) For the aforementioned sequences $(m_j)_j, (n_j)_j$ the set K_ξ is closed under the even operations. Namely it contains all $f = m_{2^j}^{-1} \sum_{i=1}^{n_{2^j}} f_i$ with $f_1 < \dots < f_{n_{2^j}}$ in K_ξ .

iii) For the odd operations $(\mathcal{A}_{n_{2^{j-1}}}, m_{2^{j-1}}^{-1})$ K_ξ includes the attracting functionals. Those are functionals of the form $f = m_{2^{j-1}}^{-1} \sum_{i=1}^{n_{2^{j-1}}/2} (f_{2i-1} + e_{l_{2i}}^*)$ with $f_1 < e_{l_2}^* < f_3 < e_{l_4}^* < \dots$ and the whole sequence is selected with the aid of a coding function.

iv) The set K_ξ contains all rational ℓ_2 -convex combinations of its weighted functionals with disjoint weights and also it is a rationally convex set.

The set K_ξ induces a norm on $c_{00}(\mathbb{N})$ and the space X_ξ is its completion. The attracting functionals are responsible for carrying structure from the set G_ξ to block subspaces of X_ξ^* . To make more transparent the role of the attracting functionals let us first notice, that each attracting functional f consists of two parts. Namely $f = g_1 + g_2$ where $g_1 = m_{2^{j-1}}^{-1} (f_1 + f_3 + \dots)$ and $g_2 = m_{2^{j-1}}^{-1} (e_{l_2}^* + e_{l_4}^* + \dots)$. The key result for applying the method of attractors is the following

Lemma. *For every block subspace Y of X_ξ^* , for every $j_0 \in \mathbb{N}$ and every $\epsilon > 0$ there exists an attracting functional $f = m_{2^{j_0-1}}^{-1} \sum_{i=1}^{n_{2^{j_0-1}}/2} (f_{2i-1} + e_{l_{2i}}^*)$ such that writing $f = g_1 + g_2$ we have $\text{dist}(g_1, Y) < \epsilon$ and*

$$(1) \quad \|g_1\| \geq c m_{2^{j_0-1}}$$

where $c \in (0, 1)$ is a universal constant.

Granting (1) we proceed as follows. We observe that $\|f\| \leq 1$ as $f \in K_\xi$ and also (1) yields $\|m_{2^{j_0-1}}^{-1} g_1\| \geq c$. Therefore the functional $m_{2^{j_0-1}}^{-1} g_1$ has norm bounded from below by c and also

$$(2) \quad \|m_{2^{j_0-1}}^{-1} g_1 - (-m_{2^{j_0-1}}^{-1} g_2)\| \leq m_{2^{j_0-1}}^{-1}.$$

We recall that $(-m_{2j_0-1}^{-1}g_2) \in G_\xi$ and in fact is a component of the c_0 -tree structure included in G_ξ . Therefore inequality (2) permit us to transfer into an arbitrary block subspace of X_ξ^* a c_0 -tree structure of height greater than ω^ξ . It is worth pointing out that the method of attractors offers a considerable reduction to the complexity of the proofs for properties of the ultimate space. For example in the case of the space \mathfrak{X}_ξ the proof that the space does not have an ℓ_1 -spreading model is essentially given for the auxiliary space X_{G_ξ} where the norming set G_ξ is simpler than the set K_ξ .

Sections 8,9,10 are rather technical and include the necessary estimations for proving the aforementioned inequality (1). This part is closely related to the well known estimations in mixed Tsirelson and Hereditarily Indecomposable spaces. The additional complexity of this part, compared to the previous similar results, arises from property *iv*) of the norming set K_ξ and the local c_0 -structure of the set G_ξ . A consequence of the new estimations is the following theorem.

Theorem. *There exists a reflexive HI space X admitting no ℓ_p , $1 \leq p < \infty$, or c_0 as a spreading model.*

This result is presented in Section 11 which also includes a general result concerning spaces with no ℓ_p (or c_0) as a spreading model (Theorem 11.3). E. Odell and Th. Schlumprecht, [24], have presented the first example of a Banach space with no ℓ_p (or c_0) as a spreading model. The aforementioned result provides an alternative proof of the latter property of E. Odell and Th. Schlumprecht example and also yields that the space X_ξ does not have an ℓ_1 -spreading model.

In Section 12 we show that the space X_ξ^* has hereditary c_0 -index greater than ω^ξ . This result does not allow us to conclude that the hereditary ℓ_1 -index of X_ξ is also greater than ω^ξ . It is worth pointing out that in general the existence of local or asymptotic c_0 -structure in X^* yields that the corresponding ℓ_1 will occur on X . For example it is easy to see that if X^* has a c_0 -spreading model then X will have a corresponding ℓ_1 . This fact seems not to remain valid for the hereditary local or asymptotic structure as X_ξ indicates.

In Section 13 we make the final step in defining the space $\mathfrak{X}_\xi = X_\xi/X_\xi^L$ where $X_\xi^L = \langle (e_n)_{n \in L} \rangle$ with a suitable subset L of \mathbb{N} . The space \mathfrak{X}_ξ has a Schauder basis and $I_{hb}(\mathfrak{X}_\xi) > \omega^\xi$. The space \mathfrak{X}_ξ does not have an ℓ_1 -spreading model since it is quotient of X_ξ which satisfies the same property. Actually both spaces X_ξ and \mathfrak{X}_ξ does not have any ℓ_p as a spreading model.

2. PRELIMINARIES

We start by recalling some basic definitions and standard notation. Let X be a Banach space with a basis (e_i) . Given any basic sequence (x_n) by $\langle (x_n)_n \rangle$ we denote the closed vector subspace spanned by (x_n) . The *support* of a vector $x = \sum_i x_i e_i$ is the set $\text{supp } x = \{i \in \mathbb{N} : x_i \neq 0\}$, we define the range x of a vector $x \in X$ as the smallest interval in \mathbb{N} containing support of x . Given any $x = \sum_i x_i e_i$ and finite $E \subset \mathbb{N}$ put $E x = \sum_{i \in E} x_i e_i$. We write $x < y$ for vectors $x, y \in X$, if $\text{maxsupp}(x) < \text{minsupp}(y)$. A *block sequence* is any sequence $(x_i) \subset X$ satisfying $x_1 < x_2 < \dots$, a *block subspace* of X - any closed subspace spanned by an infinite block sequence. A *tail subspace* of X is any subspace of the form $\langle (e_n)_{n \geq n_0} \rangle$ for some $n_0 \in \mathbb{N}$.

Let $(x_n)_{n \in \mathbb{N}}$ be a sequence and $(\epsilon_n)_{n \in \mathbb{N}}$ be a sequence of positive numbers. We say that $(y_n)_{n \in \mathbb{N}}$ is (ϵ_n) -close to $(x_n)_{n \in \mathbb{N}}$ if $\|x_n - y_n\| < \epsilon_n$ for every $n \in \mathbb{N}$.

Given infinite $M \subset \mathbb{N}$ by $[M]$ we denote the family of all infinite subsets of M , by $[M]^{< \infty}$ - the family of all finite subsets of M . By $[M]^n$, $n \in \mathbb{N}$, we denote all finite subsets of M of cardinality n . A family \mathcal{F} of finite subsets of \mathbb{N} is *regular*, if it is *hereditary*, i.e. for any $G \subset F$, $F \in \mathcal{F}$ also $G \in \mathcal{F}$, *spreading*, i.e. for any integers

$n_1 < \dots < n_k$ and $m_1 < \dots < m_k$ with $n_i \leq m_i$, $i = 1, \dots, k$, if $(n_1, \dots, n_k) \in \mathcal{F}$ then also $(m_1, \dots, m_k) \in \mathcal{F}$, and *compact* in the product topology of $2^{\mathbb{N}}$.

The families \mathcal{A}_n , $n \in \mathbb{N}$, are defined by the following formula:

$$\mathcal{A}_n = \{F \subset \mathbb{N} : \#F \leq n\}, \quad n \in \mathbb{N}.$$

Define the *generalized Schreier families* $(\mathcal{S}_\xi)_{\xi < \omega_1}$ of finite subsets of \mathbb{N} by the transfinite induction [2]:

$$\mathcal{S}_0 = \{\{n\} : n \in \mathbb{N}\} \cup \{\emptyset\}$$

$$\mathcal{S}_{\xi+1} = \mathcal{S}_1[\mathcal{S}_\xi] = \{F_1 \cup \dots \cup F_m : m \in \mathbb{N}, F_1, \dots, F_m \in \mathcal{S}_\xi, m \leq F_1 < \dots < F_m\}$$

for any $\xi < \omega_1$. If ξ is a limit ordinal, choose $\xi_n \nearrow \xi$ and set

$$\mathcal{S}_\xi = \{F : F \in \mathcal{S}_{\xi_n} \text{ and } n \leq F \text{ for some } n \in \mathbb{N}\}.$$

It is well known that the families \mathcal{A}_n , $n \in \mathbb{N}$, \mathcal{S}_ξ , $\xi < \omega_1$, are regular families of finite subsets of \mathbb{N} .

Let us recall that a set $F \in \mathcal{S}_\xi$ is called a maximal set if there is no $G \in \mathcal{S}_\xi$ such that $F \subseteq G$. In [14] it is proved that F is \mathcal{S}_ξ -maximal if and only if there is no $k \in \mathbb{N}$ with $F < k$ and $F \cup \{k\} \in \mathcal{S}_\xi$.

Definition 2.1. Let \mathcal{F} be one of the families \mathcal{A}_n , $n \in \mathbb{N}$, \mathcal{S}_ξ , $\xi < \omega_1$, and $\theta \in (0, 1)$.

1) A finite sequence (f_1, \dots, f_k) in $c_{00}(\mathbb{N})$ is said to be \mathcal{F} -admissible if

$$\text{supp}(f_1) < \dots < \text{supp}(f_k) \text{ and } \{\min(f_1), \dots, \min(f_k)\} \in \mathcal{F}$$

2) The (\mathcal{F}, θ) -operation on $c_{00}(\mathbb{N})$ is the operation which assigns to each \mathcal{F} -admissible sequence $f_1 < \dots < f_d$ the vector $\theta(f_1 + \dots + f_d)$.

Throughout this paper by a *tree on a set X* we mean a subset \mathcal{T} of $\bigcup_{n=1}^{\infty} X^n$ such that $(x_1, \dots, x_k) \in \mathcal{T}$ whenever $(x_1, \dots, x_k, x_{k+1}) \in \mathcal{T}$, $k \in \mathbb{N}$, ordered by the initial segment inclusion. A tree \mathcal{T} is *well-founded*, if there is no infinite sequence $(x_i) \subset X$ with $(x_1, \dots, x_k) \in \mathcal{T}$ for any $k \in \mathbb{N}$. Given a tree \mathcal{T} on X put

$$D(\mathcal{T}) = \{(x_1, \dots, x_k) : (x_1, \dots, x_k, x) \in \mathcal{T} \text{ for some } x \in X\}.$$

Inductively define trees $D^\alpha(\mathcal{T})$: $D^0(\mathcal{T}) = \mathcal{T}$, $D^{\alpha+1}(\mathcal{T}) = D(D^\alpha(\mathcal{T}))$ for α ordinal and $D^\alpha(\mathcal{T}) = \bigcap_{\xi < \alpha} D^\xi(\mathcal{T})$ for α limit ordinal. The *order* of a well-founded tree \mathcal{T} is given by $o(\mathcal{T}) = \inf\{\alpha : D^\alpha(\mathcal{T}) = \emptyset\}$.

Let \mathcal{F} be a countable family of finite subset of \mathbb{N} endowed with the topology of the pointwise topology. For $\alpha < \omega_1$, we set $\mathcal{F}^{\alpha+1} = \{F \in \mathcal{F} : F \text{ a limit point of } \mathcal{F}^\alpha\}$ and for α limit ordinal $\mathcal{F}^\alpha = \bigcap_{\beta < \alpha} \mathcal{F}^\beta$. The Cantor-Bendixson index of \mathcal{F} , denoted by $CB(\mathcal{F})$, is defined as the least α for which $\mathcal{F}^\alpha = \emptyset$.

Let \mathcal{T} be a countable tree on \mathbb{N} . Then \mathcal{T} defines the family

$$\mathcal{F}_{\mathcal{T}} = \{F \in [\mathbb{N}]^{<\omega} : \text{there exists } t \in \mathcal{T} \text{ with } F \text{ a subset of the range of } t\}.$$

It follows that the family $\mathcal{F}_{\mathcal{T}}$ is hereditary. Conversely given \mathcal{F} a countable family of finite subsets of \mathbb{N} with each $F \in \mathcal{F}$ we associate the finite strictly increasing sequence t_F of integers with range equal to the set F . We set

$$\mathcal{T}_{\mathcal{F}} = \{t \in \bigcup_n [\mathbb{N}]^n : \text{there exists } F \in \mathcal{F} \text{ such that } t \text{ is initial segment of } t_F\}.$$

From the above definitions it follows that $F \in \mathcal{F}^1$ if and only if the corresponding node in the tree has infinitely many immediate successors. It follows $CB(\mathcal{F}) \leq o(\mathcal{T}_{\mathcal{F}})$. In the case of the Schreier families (c.f [2]) it follows that

$$CB(\mathcal{S}_\xi) = o(\mathcal{T}_{\mathcal{S}_\xi}) = \omega^\xi + 1.$$

For unexplained notions and notations we refer the reader to [20].

3. CONCEPTS OF PROXIMITY TO ℓ_1

In this section we introduce the hereditary strategic ℓ_1 -index (Def. 3.3) and show that it is essentially equivalent to the notion of the hereditary Bourgain ℓ_1 -index (Prop. 3.4). We discuss also non-hereditary concepts of proximity to ℓ_1 and show strong relation between these notions in the sequence setting (Prop. 3.7).

Definition 3.1. *Let X be a Banach space with a basis.*

A tree \mathcal{T} on X is an ℓ_1 -tree on X with constant $C \geq 1$, if any $(x_1, \dots, x_k) \in \mathcal{T}$ is a normalized sequence C -equivalent to the unit vector basis of ℓ_1^k . Let

$$I(X, C) = \sup\{o(\mathcal{T}) : \mathcal{T} \text{ - } \ell_1\text{-tree on } X \text{ with constant } C\}, \quad C \geq 1$$

The (Bourgain) ℓ_1 -index of X is defined by $I(X) = \sup\{I(X, C) : C \geq 1\}$.

The block (Bourgain) ℓ_1 -index $I_b(X)$, is defined analogously, using ℓ_1 -trees consisting only of block sequences.

It follows by [10] that for a separable Banach space X the ℓ_1 -index is a countable ordinal if and only if X does not contain ℓ_1 . In this case the ℓ_1 -index is of the form ω^ξ for some $\xi < \omega_1$ and also it is greater than $I_b(X, C)$ for any $C \geq 1$, and the same holds for the block ℓ_1 -index [18]. Recall that if $I(X) \geq \omega^\omega$, then $I(X) = I_b(X)$, if $I(X) = \omega^{n+1}$, then $I_b(X) = \omega^n$ or $I_b(X) = \omega^{n+1}$ [18]. For more information on the block ℓ_1 -index see [18].

The hereditary (Bourgain) ℓ_1 -index is defined as $I_h(X) = \min\{I(Y) : Y \text{ is subspace of } X\}$. The hereditary block (Bourgain) ℓ_1 -index, denoted as $I_{hb}(X)$, is defined similarly taking block Bourgain ℓ_1 -indices of block subspaces.

Next we introduce the following notion.

Definition 3.2. *Let X be a Banach space with basis, $\xi < \omega_1$, $C \geq 1$ and consider an \mathcal{S}_ξ -game between the players S and V defined as follows:*

in the i -th move player S chooses a block subspace X_i of X and player V picks a normalized block vector $x_i \in X_i$.

We say that player V wins, if the resulting sequence $(x_i)_{i=1}^k$ is C -equivalent to the standard basis of ℓ_1^k and maximal \mathcal{S}_ξ -admissible (i.e. $\{\text{minsupp } x_i : i = 1, \dots, k\}$ is \mathcal{S}_ξ -maximal). We say that V has a winning strategy, if V wins the game for any possible choice of the player S .

Definition 3.3. *Let X be a Banach space with a basis. We define the hereditary strategic ℓ_1 -index of X by the formula*

$$I_{hs}(X) = \sup\{\xi < \omega_1 : \text{for any } \zeta < \xi \text{ there is } C \geq 1 \text{ such that} \\ V \text{ has a winning strategy in } \mathcal{S}_\zeta \text{ - game with constant } C\}.$$

If we consider the game above where S chooses only tail subspaces instead of arbitrary block subspaces, then we define the strategic ℓ_1 -index of X denoted by $I_s(X)$.

In non-hereditary case for $\xi = 1$ we obtain Definition 2.1. [21].

The next result describes the relations between the above introduced notions.

Proposition 3.4. *Let X be a Banach space with a basis, $\xi < \omega_1$.*

If $I_{hs}(X) > \xi$ then $I_{hb}(X) > \omega^\xi$.

If $I_{hb}(X) > \omega^\xi$ then $I_{hs}(Y_0) > \xi$ for some block subspace Y_0 of X .

Proof. We can assume that X does not contain a copy of ℓ_1 . Notice that for any $\xi < \omega_1$ we have $I_{hs}(X) > \xi$ if and only if player V has a winning strategy in \mathcal{S}_ξ -game with some constant $C \geq 1$.

Assume first that player V has a winning strategy in the \mathcal{S}_ξ -game (with some constant C). Fix a block subspace Y . Then player V in particular has a winning strategy in producing maximal \mathcal{S}_ξ -admissible sequences C -equivalent to the unit vector basis of ℓ_1

in the \mathcal{S}_ξ^Y -game, in which player S chooses at each step some $m_k \in \mathbb{N}$ and V chooses $x_k \in Y$ with $x_k > m_k$ (i.e. player S chooses only tail subspaces of Y).

Take the tree \mathcal{T} of all block sequences produced by player V in all \mathcal{S}_ξ^Y -games for all block subspaces Y , with all possible moves of player S , i.e. all block sequences (x_1, \dots, x_k) produced by player V at some point in all \mathcal{S}_ξ^Y -games according to his winning strategy. It follows that \mathcal{T} is an ℓ_1 -tree with a constant C . We show now that $o(\mathcal{T}) \geq o(\mathcal{S}_\xi) = \omega^\xi + 1$ which implies that $I_b(Y) > \omega^\xi$.

For any $(m_1, \dots, m_k) \in \mathcal{S}_\xi$ by $\mathcal{T}_{m_1, \dots, m_k}$ denote the set of all block sequences of length k produced by V according to his winning strategy in all \mathcal{S}_ξ^Y -games, where S has chosen in his first k moves m_1, \dots, m_k .

By induction we show that for any $\alpha < \omega_1$ we have

$$D^\alpha(\mathcal{T}) \supset \cup \{ \mathcal{T}_{m_1, \dots, m_k} : (m_1, \dots, m_k) \in D^\alpha(\mathcal{S}_\xi) \}.$$

Indeed, if $\alpha = \beta + 1$ (in particular if $\beta = 0$), then by the inductive assumption (or by the definition of \mathcal{T} in case of $\beta = 0$) we have

$$\begin{aligned} D(D^\beta \mathcal{T}) &\supset \cup \{ D(\mathcal{T}_{m_1, \dots, m_k}) : (m_1, \dots, m_k) \in D^\beta(\mathcal{S}_\xi) \} \\ &= \cup \{ \mathcal{T}_{m_1, \dots, m_{k-1}} : (m_1, \dots, m_k) \in D^\beta(\mathcal{S}_\xi) \} = \cup \{ \mathcal{T}_{m_1, \dots, m_l} : (m_1, \dots, m_l) \in D^\alpha(\mathcal{S}_\xi) \} \end{aligned}$$

For any limit $\alpha < \omega_1$ by the inductive assumption we have

$$\begin{aligned} \cap_{\beta < \alpha} D^\beta \mathcal{T} &\supset \cap_{\beta < \alpha} \cup \{ \mathcal{T}_{m_1, \dots, m_k} : (m_1, \dots, m_k) \in D^\beta(\mathcal{S}_\xi) \} \\ &\supset \cup \{ \mathcal{T}_{m_1, \dots, m_k} : (m_1, \dots, m_k) \in D^\alpha(\mathcal{S}_\xi) \}. \end{aligned}$$

It follows that $o(\mathcal{T}) \geq o(\mathcal{S}_\xi)$, which ends the proof of the first implication.

Assume now that for any block subspace Y of X we have $I_b(Y) > \omega^\xi$. We will show now that in any block subspace Y there is a block subspace Y_0 and $C \geq 1$ such that in any subspace $W \subset Y_0$ there is a maximal \mathcal{S}_ξ -admissible block sequence (x_1, \dots, x_k) C -equivalent to the unit vector basis of ℓ_1^k . Then by Gowers dichotomy for games for families of finite block sequences [15] in some block subspace of Y player V has a winning strategy for producing maximal \mathcal{S}_ξ -admissible block sequences $2C$ -equivalent to unit vector basis of suitable finite dimensional spaces ℓ_1 , which will prove that $I_{hs}(X) > \xi$.

First notice that there is some block subspace Y_0 and universal constant C such that for any block subspace Z of Y_0 there is an ℓ_1 -tree \mathcal{T}_Z on Z with constant C and $o(\mathcal{T}_Z) > \omega^\xi + 1$. Since we deal with ℓ_1 -trees, we can assume that they are hereditary trees, i.e. for any ℓ_1 -tree \mathcal{R} , any $(x_i)_{i \in F} \in \mathcal{R}$ and any $G \subset F$ we have also $(x_i)_{i \in G} \in \mathcal{R}$.

Indeed, otherwise we can produce a decreasing sequence of block subspaces (Y_n) such that in each Y_n there is no ℓ_1 -tree with constant n and order greater than $\omega^\xi + 1$. Then the diagonal subspace does not contain any ℓ_1 -tree of order greater than $\omega^\xi + 1$, a contradiction with the assumption, since the block ℓ_1 -index of a Banach space is a limit ordinal.

Take Y_0 and the constant C as above, for any block subspace Z of Y_0 pick an ℓ_1 -tree \mathcal{T}_Z with constant C and $o(\mathcal{T}_Z) > \omega^\xi + 1$. Let $\mathcal{T} = \bigcup \{ \mathcal{T}_Z : Z \subset Y_0 \}$. Then \mathcal{T} is also an hereditary ℓ_1 -tree (with constant C).

Pick any block subspace $W = \langle (w_n)_n \rangle$ of Y_0 . Let $L = \{ \text{minsupp } w_n : n \in \mathbb{N} \}$ and

$$\mathcal{F} = \{ (\text{minsupp } x_1, \dots, \text{minsupp } x_k) \subset L : (x_1, \dots, x_k) \in \mathcal{T} \cap W^k, k \in \mathbb{N} \}$$

Notice that the tree \mathcal{F} is hereditary and well-founded. Indeed, assume that there is $(n_i)_{i \in \mathbb{N}}$ such that $(n_1, \dots, n_k) \in \mathcal{F}$ for any $k \in \mathbb{N}$. Thus for any $k \in \mathbb{N}$ there is $(x_1^k, \dots, x_k^k) \in \mathcal{T}$ with $n_1 \leq x_1^k < n_2 \leq x_2^k < \dots < n_k \leq x_k^k$. By compactness argument we can assume that for some block sequence (x_i) we have $x_i^k \rightarrow x_i$ in X as $k \rightarrow \infty$. Since each sequence (x_1^k, \dots, x_k^k) is C -equivalent to the unit vector basis of ℓ_1^k , thus $(x_i)_{i \in \mathbb{N}}$ is equivalent to the unit vector basis of ℓ_1 , a contradiction with the assumption from the beginning of the proof.

By I. Gasparis' dichotomy [14] there is some infinite $M \subset L$ such that either $\mathcal{S}_\xi \cap [M]^{<\infty} \subset \mathcal{F}$ or $\mathcal{F} \cap [M]^{<\infty} \subset \mathcal{S}_\xi$.

We show that the first case holds by the following

Claim. *Given any tree \mathcal{R} on X let*

$$\mathcal{F}_\mathcal{R} = \{(\text{minsupp } x_1, \dots, \text{minsupp } x_k) : (x_1, \dots, x_k) \in \mathcal{R}\}$$

Then if trees \mathcal{R} and $\mathcal{F}_\mathcal{R}$ are well-founded it follows that $o(\mathcal{R}) \leq o(\mathcal{F}_\mathcal{R})$.

To prove the claim it is enough to show by induction that $\mathcal{F}_{D^\alpha(\mathcal{R})} \subset D^\alpha(\mathcal{F}_\mathcal{R})$ for any $\alpha < \omega_1$.

Let now $Z = \langle (w_n)_{\text{minsupp } w_n \in M} \rangle$. By definition of \mathcal{T} and \mathcal{F} we have that $\mathcal{F} \cap [M]^{<\infty} \supset \mathcal{F}_{\mathcal{T}_Z}$, therefore by the above Claim

$$o(\mathcal{F} \cap [M]^{<\infty}) \geq o(\mathcal{T}_Z) > \omega^\xi + 1 = o(\mathcal{S}_\xi)$$

hence $\mathcal{S}_\xi \cap [M]^{<\infty} \subset \mathcal{F}$ must hold.

Take now any (n_1, \dots, n_k) maximal in $\mathcal{S}_\xi \cap [M]^{<\infty} \subset \mathcal{F}$ and the corresponding $(x_1, \dots, x_k) \in \mathcal{T} \cap W^k$ with $\text{minsupp } x_i = n_i$, $i = 1, \dots, k$. It is clear that (x_1, \dots, x_k) is maximal \mathcal{S}_ξ -admissible. Therefore we picked in W a maximal \mathcal{S}_ξ -admissible block sequence C -equivalent to the unit vector basis of some finite dimensional ℓ_1 , which by previous remarks ends the proof. \square

Remark 3.5. *The non-hereditary case the above proposition takes the following form for a Banach space X with a basis:*

If $I_s(X) > \xi$, then $I_b(X) > \omega^\xi$.

If $I_b(\langle (e_n)_{n \in M} \rangle) > \omega^\xi$ for any $M \in [\mathbb{N}]$, then $I_s(X) > \xi$.

The proof goes along the same scheme as above, with passing to subspaces spanned by subsequences of the basis instead of block subspaces.

A stronger representation of ℓ_1 in a Banach space is described by the following notions:

Definition 3.6. *Let X be a Banach space with a basis and $\xi < \omega_1$ be a countable ordinal.*

A normalized basic sequence $(x_i)_{i \in \mathbb{N}} \subset X$ generates an ℓ_1^ξ -spreading model, $\xi < \omega_1$, with constant $C \geq 1$, if for any $F \in \mathcal{S}_\xi$ the sequence $(x_i)_{i \in F}$ is C -equivalent to the unit vector basis of $\ell_1^{\#F}$.

We say that (x_i) generates an ℓ_1 -spreading model if the above property holds for $\xi = 1$.

The space X is ℓ_1^ξ -asymptotic, if any \mathcal{S}_ξ -admissible sequence $(x_i)_{i=1}^n$ is C -equivalent to the unit vector basis of ℓ_1^n .

The space X is said to be ℓ_1 -asymptotic if the above property holds for $\xi = 1$.

The asymptotic ℓ_1 spaces were introduced in [23], ℓ_1 -spreading models of higher order were studied in [6],[4].

Replacing ℓ_1 by c_0 in the above definitions we obtain the (Bourgain) c_0 -index, a c_0^ξ -spreading model and an asymptotic c_0 space.

The structures described in the above definitions give us an hierarchy of the representation of ℓ_1 in a Banach space X with a basis, not containing ℓ_1 . For a countable ordinal $\xi < \omega_1$ we consider the following four "local" structures:

- A) The space X is ℓ_1^ξ asymptotic.
- B) The space X contains a sequence generating an ℓ_1^ξ -spreading model.
- C) The strategic ℓ_1 -index of X is greater than ξ .
- D) The block ℓ_1 -index of X is greater than ω^ξ .

Let us observe that the following implications hold for the above structures

$$A) \Rightarrow B) \Rightarrow C) \Rightarrow D)$$

The reverse implications are not true.

First observe that for every $\xi < \omega_1$ the Tsirelson spaces $T[\mathcal{S}_\xi, \theta]$, $\theta \in (0, 1)$, are examples of reflexive Banach space which are ℓ_1^ξ -asymptotic however they do not contain ℓ_1 .

The Schreier space X_ξ , $\xi < \omega_1$, is an example of Banach space with basis for which every subsequence of the basis generates an ℓ_1^ξ -spreading model and it does not contain an asymptotic ℓ_1^ζ subspace for any $\zeta \leq \xi$ since, as it is well known, X_ξ is c_0 -saturated.

In the next section of the paper for every $\xi < \omega_1$ we provide an example of Banach space X_{G_ξ} with strategic ℓ_1 -index greater than ξ yet X_{G_ξ} does not contain sequence generating an ℓ_1 -spreading model. More precisely the space X_{G_ξ} has basis such that in the game, where S chooses subspaces spanned by subsequences of the basis, player V has a winning strategy. Therefore if a space X has strategic ℓ_1 -index greater than ξ , then it does not follow that X contains a sequences generating an ℓ_1 -spreading model.

A richer asymptotic ℓ_1^ξ structure than the one described in property C) would be provided by a winning strategy of player V in the following modification of the \mathcal{S}_ξ -game: player S chooses subsequences of the basis instead of block subspaces and player V picks vectors from subsequences chosen by player S . In such a case every subsequence of the basis would admit ℓ_1 -tree (i.e. formed by elements of this subsequence) of order ω^ξ . Let us observe the following

Proposition 3.7. *Let (e_n) be a basis of X such that any subsequence $(e_n)_{n \in L}$ admit an ℓ_1 -tree $\mathcal{T}_L \subset \{(e_n)_{n \in F} : F \in [L]^{<\infty}\}$ of order ω^ξ , $\xi < \omega_1$. Then some subsequence of $(e_n)_{n \in \mathbb{N}}$ generates an ℓ_1^ξ -spreading model.*

Proof. Assume first that any subsequence $(e_n)_{n \in L}$ of the basis admits an ℓ_1 -tree of order greater than $\omega^\xi + 1$.

Repeating the reasoning and notation from the proof of Prop. 3.4 we can assume that there is some subsequence $(e_n)_{n \in L_0}$ and an hereditary ℓ_1 -tree \mathcal{T} formed by elements of this subsequence such that for any $M \subset L_0$ the order of $\mathcal{F}_\mathcal{T} \cap [M]^{<\infty}$ is greater than $\omega^\xi + 1$.

As before by I.Gasparis' dichotomy [14] pick an infinite $M \subset \mathbb{N}$ such that either $\mathcal{S}_\xi \cap [M]^{<\infty} \subset \mathcal{F}_\mathcal{T}$ or $\mathcal{F}_\mathcal{T} \cap [M]^{<\infty} \subset \mathcal{S}_\xi$. By the above remark the first case holds. Notice that if $M = (m_i)$, then for any $F \in \mathcal{S}_\xi$ we have $(m_i)_{i \in F} \in \mathcal{S}_\xi \cap [M]^{<\infty} \subset \mathcal{F}_\mathcal{T}$, which implies that $(e_n)_{n \in M}$ generates an ℓ_1^ξ -spreading model.

Now notice that if (e_n) admits an ℓ_1 -tree of order ω^ξ , then it also admit an ℓ_1 -tree of order greater than $\omega^\xi + 1$ (maybe with worse constant).

It follows by repeating the reasoning in Lemma 5.7 [18] in case of the block ℓ_1 -index defined not by using all block ℓ_1 -trees but only ℓ_1 -trees consisting of finite subsequences of (e_n) . Therefore we can reduce the case where all subsequences admit ℓ_1 -trees of order ω^ξ to the case of order greater than $\omega^\xi + 1$, which was treated above and hence we finish the proof of the observation. \square

We end this section with the following observation regarding the non-hereditary indices. Assuming only $I_b(X) > \omega^\xi$ does not imply $I_s(X) > 1$, as it shown by the following example.

Let X be an ℓ_2 -direct sum of ℓ_1^n 's, i.e. $X = (\bigoplus_{n=1}^{\infty} \ell_1^n)_{\ell_2}$.

It is clear that $I_b(X) > \omega$, since $\sup\{o(\mathcal{T}) : \mathcal{T} \text{ } \ell_1\text{-tree with constant } 1\} = \omega$.

On the other hand $I_s(X) = 1$. Indeed, consider \mathcal{S}_1 -game with player S choosing tail subspaces and let in the first move player S pick a tail subspace $(\bigoplus_{n=n_0}^{\infty} \ell_1^n)_{\ell_2}$. In the $(i+1)$ -th move player S chooses tail subspaces after the support of vector x_i picked by V in the i -th move, i.e. if in the i -th move player V picks a vector $x_i \in (\bigoplus_{n=1}^{n_i} \ell_1^n)_{\ell_2}$, then in the $(i+1)$ -th move player S chooses tail subspace $(\bigoplus_{n=n_i+1}^{\infty} \ell_1^n)_{\ell_2}$.

Therefore player S forces the sequence (x_1, \dots, x_k) produced by player V to be 1-equivalent to the unit vector basis of ℓ_2^k , hence there is no universal constant C for which

player V has a winning strategy for producing \mathcal{S}_1 -admissible block sequences C -equivalent to the unit vector basis of suitable finite dimensional ℓ_1 .

4. DEFINITION OF THE GROUND SET G_ξ

In this section we define for any $\xi < \omega_1$ ground sets G'_ξ and G_ξ for the auxiliary spaces $Y_{G'_\xi}$ and $X_{G'_\xi}$ respectively, in particular we introduce the tree of G_ξ -special functionals, defined with the use of special coding function σ_1 , and related notions. We show that the space $Y_{G'_\xi}$ is c_0 -saturated (Prop. 4.7).

We recall that a subset G of $c_{00}(\mathbb{N})$ is said to be a ground set if:

- (1) G is symmetric and $\{e_n^* : n \in \mathbb{N}\}$ is contained in G .
- (2) $\|\phi\|_\infty \leq 1$ and $\phi(n) \in \mathbb{Q}$ for $\phi \in G$.
- (3) G is closed under the restriction of its elements to intervals of \mathbb{N} .

A ground set G induces a norm $\|\cdot\|_G$ in $c_{00}(\mathbb{N})$ defined by

$$\|x\|_G = \sup\{\phi(x) : \phi \in G\} \text{ and we set } X_G = \overline{(c_{00}(\mathbb{N}), \|\cdot\|_G)}.$$

It is easy to see that the natural basis of $c_{00}(\mathbb{N})$ is a Schauder basis of the space X_G . In the opposite for every Banach space with a Schauder basis $(x_n)_{n \in \mathbb{N}}$ there exists a ground set G such that the natural correspondence $e_n \rightarrow x_n$ is extended to an isomorphism between X_G and X .

We pass now to define the ground set G_ξ . Fix $\xi < \omega_1$. We choose two strictly increasing sequences $(n_j)_j, (m_j)_j$ of positive integers, such that

- (i) $m_1 = 2^5$ and $m_{j+1} = m_j^5$
- (ii) $n_1 = 2^6$ and $n_{j+1} = (2n_j)^{s_j}$ where $2^{s_j} = m_{j+1}^3$.

Let us observe, for later use, that $260m_{2j}^4 \leq n_{2j-1}$ for $j \geq 2$. Set $G_0 = \{\pm e_n^* : n \in \mathbb{N}\}$

$$G_1^j = \left\{ \frac{1}{m_{2j-1}^2} \sum_{i \in E} f_i : f_i \in G_0, (f_i)_{i \in E}, \#E \leq n_{2j-1} \right\}, j \in \mathbb{N}.$$

Finally we set $G_1 = \cup_{j \in \mathbb{N}} G_1^j$.

Notation 4.1. For every $f \in G_1^j$ we define the weight of f as $w(f) = m_{2j-1}^2$ and the index as $\text{ind}(f) = j$. The elements of G_1 are called functionals of type I.

We consider the following set

$$\mathcal{U} = \{(f_1, \dots, f_d) : f_1 < \dots < f_d, f_i \in G_1^{j_i}, j_i < j_{i+1} \text{ for all } i < d \in \mathbb{N}\}.$$

Let $\mathbb{N} = M_1 \cup M_2$ where M_1, M_2 are infinite disjoint sets. Let also $\sigma_1 : \mathcal{U} \rightarrow M_2$ be a 1-1 function (i.e. a coding function) such that $\sigma_1(f_1, \dots, f_{i+1}) > \sigma_1(f_1, \dots, f_i)$ for every $i \in \mathbb{N}$.

Definition 4.2. A special sequence is an element (f_1, \dots, f_d) of \mathcal{U} satisfying the following:

- 1) $(\text{minsupp } f_i)_{i=1}^d \in \mathcal{S}_\xi$.
- 2) $f_1 \in G_1^{j_1}, j_1 \in M_1, f_{i+1} \in G_1^{\sigma_1(f_1, \dots, f_i)}$ for every $i = 1, \dots, d-1$.

We define

$$G_{sp} = \left\{ E \sum_{i=1}^d \epsilon_i f_i : (f_1, \dots, f_d) \text{ is a special sequence, } \epsilon_i \in \{-1, 1\}, E \text{ interval of } \mathbb{N} \right\}$$

For any element $\phi = E \sum_{i=1}^d \epsilon_i f_i$ of G_{sp} we set $\text{ind}(\phi) = \{\text{ind}(f_i) : E f_i \neq \emptyset, i \leq d\}$. We define also

$$G_{\ell_2} = \left\{ \sum_{i=1}^d a_i \phi_i : d \in \mathbb{N}, \sum_{i=1}^d a_i^2 \leq 1, (\phi_i)_{i=1}^d \subset G_{sp} \cup G_1, (\text{ind } \phi_i)_{i=1}^d \text{ pairwise disjoint} \right\}.$$

Definition 4.3. The ground set G_ξ is defined to be the set

$$G_\xi = G_0 \cup G_1 \cup G_{sp} \cup G_{\ell_2}.$$

Remarks 4.4. 1) The set G_ξ is symmetric and closed under the restriction of its elements on intervals of \mathbb{N} .

2) The injectivity of the coding function σ_1 yields that the set of the special sequences has a tree structure i.e. if $(f_1, \dots, f_d), (g_1, \dots, g_n)$ are two special sequences then either $f_i \neq g_j$ for all i, j or there exists $i_0 \leq \min\{d, n\}$ such that $f_i = g_i$ for all $i < i_0$ and $f_i \neq g_j$ for all $i, j \geq i_0$, in particular $w(f_i) \neq w(g_j)$ for all $i, j > i_0$.

We shall call every special sequence also a segment of the tree of the special sequences. The elements of G_{sp} are called G_ξ -special functionals.

Notation 4.5. For every segment $s = (f_1, \dots, f_d)$ of the tree of the special sequences we set $F(s) = \{\sum_{i=1}^d \epsilon_i f_i : \epsilon_i \in \{-1, 1\}\}$ and for $f \in F(s)$ we set $\text{ind}(f) = \text{ind}(s)$.

If $V = \{s_1, \dots, s_d\}$ is a set of segments we set

$$\bar{V} = \left\{ \sum_{i=1}^d \lambda_i \phi_i : \sum_{i=1}^d \lambda_i^2 \leq 1, \phi_i \in F(s_i) \text{ and } \text{ind}(\phi_i) \cap \text{ind}(\phi_j) = \emptyset \text{ for all } 1 \leq i \neq j \leq d \right\}$$

It readily follows that \bar{V} is symmetric.

We set $G'_\xi = G_0 \cup G_1 \cup G_{sp} \cup \{0\}$.

Lemma 4.6. The set G'_ξ is a closed subset of $[0, 1]^{<\omega}$ in the pointwise topology.

Proof. Let $(\phi_n)_{n \in \mathbb{N}}$ be a sequence of elements of G'_ξ such that $\lim_n \phi_n = \phi$ pointwise. It is clear that if $\phi_n \in G_0$ for infinitely many n 's then $\phi \in G_0 \cup \{0\}$.

Also if $\phi_n \in G_1$ for all but finitely many n 's, then if $\{\phi_n : n \in \mathbb{N}\} \cap G_1^j \neq \emptyset$ for infinitely many j 's then $\phi = 0$. Otherwise there exists j_0 such that $\phi_n \in G_1^{j_0}$ for all but finitely many n 's. Since $\mathcal{A}_{n_{2j_0-1}}$ is regular we get $\phi \in G_1^{j_0}$.

Finally assume that for all but finitely many n 's there exists a segment $s_n = (f_1^n, \dots, f_d^n)$ of the tree of the special sequences such that $\phi_n = E_n \sum_{i=1}^d \epsilon_i^n f_i^n \in F(s_n)$.

If $\sup_n \min E_n = +\infty$, then $\phi = 0$. Otherwise we can assume that for all but finitely many n 's we have $E_n = [k, K)$, for some $k \in \mathbb{N}$ and $K \in \mathbb{N} \cup \{+\infty\}$.

Since the family \mathcal{S}_ξ is regular we get $\{m_i\}_{i=1}^d \in \mathcal{S}_\xi$ such that

$$m_1 \leq f_1^n < m_2 \leq f_2^n \cdots < m_d \leq f_d^n \quad (\text{and } \text{minsupp } f_{d+1}^n \rightarrow \infty)$$

Let $i_0 = \min\{i \leq d : \text{there is no } n \in \mathbb{N} \text{ such that } f_i^n = f_i^m \text{ for all } m \geq n\}$. By the reasoning above in case $(\phi_n) \subset G_1$ we can assume that for any $i = 1, \dots, d$ we have $f_i^n \rightarrow f_i$ for some $f_i \in G_1 \cup \{0\}$. Also we may assume that $\epsilon_i^n = \epsilon_i$ for all $i \leq d$ and $n \in \mathbb{N}$.

If $\limsup_n w(f_{i_0}^n) = +\infty$ then $f_{i_0} = 0$. Also since σ_1 is 1-1 we get $\limsup w(f_{i_0+j}^n) = +\infty$ and therefore $f_{i_0+j} = 0$ for all $j = 1, \dots, d - i_0$.

If $w(f_{i_0}^n) = m_{2j_0-1}^2$ for all but finitely many n 's, since the set $G_1^{j_0}$ is compact we get that $f_{i_0}^n \rightarrow f_{i_0} \in G_1^{j_0}$. Again since σ_1 is 1-1 we get $\limsup w(f_{i_0+j}^n) = +\infty$ and therefore $f_{i_0+j} = 0$ for all $j = 1, \dots, d - i_0$.

It follows that (f_1, \dots, f_d) is a special sequence and therefore $\sum_{i=1}^d \epsilon_i f_i^n \rightarrow \sum_{i=1}^d \epsilon_i f_i$ is a G_ξ -special functional. \square

We define $X_{G_\xi} = \overline{(c_{00}(\mathbb{N}), \|\cdot\|_{G_\xi})}$ and $Y_{G_\xi} = \overline{(c_{00}(\mathbb{N}), \|\cdot\|_{G'_\xi})}$.

Let us observe that since the sets G_ξ, G'_ξ are symmetric and closed under projections on intervals it follows that $(e_n)_{n \in \mathbb{N}}$ is a bimonotone basis for the spaces X_{G_ξ} and Y_{G_ξ} .

Proposition 4.7. Y_{G_ξ} is c_0 -saturated.

Proof. The set G'_ξ is countable by the construction and compact by Lemma 4.6. Since G'_ξ is norming set of Y_{G_ξ} we conclude that Y_{G_ξ} is isometric to a subspace of $C(G'_\xi)$ and hence is c_0 -saturated, see [26] (see also [5] and [1], Theorem 4.5). \square

We end this section with the following observations regarding the norming set G_ξ .

From the definition of the norming set G_ξ it follows that for any finite sequence $(\phi_i) \subset G_1 \cup G_{sp}$ with pairwise disjoint index sets and any $(a_i) \in \ell_2$ we have $\|\sum_i a_i \phi_i\|_{G_\xi}^* \leq (\sum_i a_i^2)^{1/2}$. Therefore for any infinite sequence $(\phi_i)_i \subset G_1 \cup G_{sp}$ with pairwise disjoint index sets and any $(a_i)_i \in \ell_2$ the series $\sum_i a_i \phi_i$ is convergent in norm and

$$\left\| \sum_{i=1}^{\infty} a_i \phi_i \right\|_{G_\xi}^* \leq \left(\sum_{i=1}^{\infty} a_i^2 \right)^{1/2}.$$

Notice also that since the set G_ξ is a norming set for X_{G_ξ} we obtain that $B_{X_{G_\xi}^*} = \overline{\text{conv}(G_\xi)^{w^*}}$. Also it is not hard to show that $\overline{G_\xi}^{w^*} = G_\xi \cup \mathcal{F}$ where

$$\mathcal{F} = \left\{ \sum_{i=1}^{\infty} a_i \phi_i : (a_i)_{i \in \mathbb{N}} \in B_{\ell_2}, \phi_i \in G_1 \cup G_{sp}, \text{ind}(\phi_i) \cap \text{ind}(\phi_j) = \emptyset \text{ for all } i \neq j \right\}.$$

We omit the proof since we shall not make use of this result.

5. BASIC PROPERTIES OF THE SPACE X_{G_ξ}

In this section we shall prove that the space X_{G_ξ} has the following properties

- 1) It is reflexive.
- 2) For every subsequence $(e_n)_{n \in M}$ of the basis the subspace $\langle (e_n)_{n \in M} \rangle$ has ℓ_1 -index greater than ω^ξ .
- 3) Every subspace of X_{G_ξ} contains ℓ_2 .

In order to prove the theorem we shall need the following definition

Definition 5.1. Let $(x_n)_n$ be a bounded block sequence in X_{G_ξ} and $\epsilon > 0$. We say that $(x_n)_n$ is ϵ -separated if for every $\phi \in G_1 = \cup_{j \in \mathbb{N}} G_1^j$

$$\#\{n : |\phi(x_n)| \geq \epsilon\} \leq 1.$$

In addition, we say that $(x_n)_n$ is separated if for every $L \in [\mathbb{N}]$ and $\epsilon > 0$ there exists an $M \in [L]$ such that $(x_n)_{n \in M}$ is ϵ -separated.

Concerning the separated sequences the following holds:

Proposition 5.2. Let $(x_n)_{n \in \mathbb{N}}$ be a separated sequence in X_{G_ξ} with $\|x_n\|_{G_\xi} \leq 1$. Then for all $m \in \mathbb{N}$ there is $L \in [\mathbb{N}]$ such that for all $g \in G_{\ell_2}$

$$\#\{n \in L : |g(x_n)| \geq m^{-1}\} \leq 65m^2.$$

The proof of the above results follows the arguments of Lemma 3.12 and Proposition 3.14 in [3] where we refer for the proofs.

Lemma 5.3. Let $(x_n)_{n \in \mathbb{N}}$ be a bounded block sequence. Setting $y_n = \frac{1}{n} \sum_{i \in F_n} x_i$ where $\#F_n = n$ and $F_n < F_{n+1}$ we get that $(y_n)_{n \in \mathbb{N}}$ is separated.

Proof. Let $\epsilon > 0$, $L \in [\mathbb{N}]$ and assume that $\|x_n\|_{G_\xi} \leq C$ for any n . Pick inductively sequences $(j_i), (l_i) \subset L$ such that

$$1) \epsilon m_{2j_{i-1}}^2 > C \# \text{range } y_{l_i} \quad \text{and} \quad 2) \epsilon l_i > C n_{2j_{i-1}-1}.$$

Take now any $\phi \in G_1^j$ and l_i . If $j \geq j_i$, then by (1) $|\phi(y_{l_i})| < \epsilon$. If $j < j_{i-1}$, then by (2) also $|\phi(y_{l_i})| < \epsilon$. Therefore $\#\{i : |\phi(y_{l_i})| \geq \epsilon\} \leq 1$ and hence setting $M = (l_i)_i$ we end the proof that $(y_l)_{l \in M}$ is ϵ -separated. \square

Combining the above lemma with Proposition 5.2 and the choice of (m_j) , (n_j) we obtain the following.

Proposition 5.4. *Let $(x_n)_{n \in \mathbb{N}}$ be a bounded block sequence. Assume that $y_n = \frac{1}{n} \sum_{i \in F_n} x_i$, where $\#F_n = n$ and $F_n < F_{n+1}$, satisfy $\|y_n\| \leq 1$. Then for any $j \geq 2$ there is an infinite $L \subset \mathbb{N}$ such that for any $g \in G_\xi$*

$$\#\{n \in L : |g(y_n)| \geq 2m_{2j}^{-2}\} \leq n_{2j-1}$$

As corollary of the above proposition we obtain that the basis is shrinking.

Corollary 5.5. *Every bounded block sequence in X_{G_ξ} is weakly null.*

Proof. Assume that there exist a normalized block sequence $(x_n)_{n \in \mathbb{N}}$, $x^* \in X_{G_\xi}^*$ of norm one and $\epsilon > 0$ such that $x^*(x_n) > \epsilon$ for all n .

Let $j \in \mathbb{N}$, $j \geq 2$ such that $1/2m_{2j}^2 < \epsilon/4$. By Proposition 5.4 setting $y_n = \frac{1}{\#F_n} \sum_{i \in F_n} x_n$, where $\#F_n = n$ and $F_n < F_{n+1}$ for all $n \in \mathbb{N}$, we may assume that for all $g \in G_\xi$ it holds that

$$(3) \quad \#\{n : |g(y_n)| \geq 2m_{2j}^{-2}\} \leq n_{2j-1} \Rightarrow |g(\frac{1}{n_{2j}} \sum_{i=1}^{n_{2j}} y_n)| \leq \frac{n_{2j-1}}{n_{2j}} + \frac{1}{2m_{2j}^2} \leq \frac{1}{m_{2j}^2} < \frac{\epsilon}{2}.$$

This yields a contradiction since G_ξ is a norming set for X_{G_ξ} . \square

We prove now the reflexivity of X_{G_ξ} .

Theorem 5.6. *The space X_{G_ξ} is reflexive.*

Proof. We show that the basis is shrinking and boundedly complete. Corollary 5.5 yields that the basis $(e_n)_{n \in \mathbb{N}}$ is shrinking. We prove that $(e_n)_{n \in \mathbb{N}}$ is also boundedly complete.

On the contrary assume that $\sup_{n \in \mathbb{N}} \|\sum_{i=1}^n a_i e_i\|_{G_\xi} \leq 1$ and there exist $\epsilon_0 > 0$, successive intervals $F_1 < F_2 < \dots$ of \mathbb{N} such that $\|\sum_{i \in F_n} a_i e_i\|_{G_\xi} > \epsilon_0$.

For every n choose $g_n \in G_\xi$ such that $g_n(\sum_{i \in F_n} a_i e_i) \geq \epsilon_0$ and $\text{range } g_n \subset F_n$.

We distinguish the following cases

Case 1. $g_n \in G_0$ for infinitely many n 's.

Then for $j \in \mathbb{N}$ and $M > \max F_{n_{2j-1}}$ and we get

$$\|\sum_{i=1}^M a_i e_i\|_{G_\xi} \geq \frac{1}{m_{2j-1}^2} \sum_{n=1}^{n_{2j-1}} g_n(\sum_{i=1}^M a_i e_i) \geq \frac{\epsilon_0 n_{2j-1}}{m_{2j-1}^2},$$

a contradiction for large j .

We state the next three cases.

Case 2. $g_n \in G_1$ for all but finitely many n 's.

Case 3. $g_n \in G_{sp}$ for all but finitely many n 's.

Case 4. $g_n \in G_{\ell_2}$ for all but finitely many n 's.

The proofs of these three cases follow the same argument hence we shall give only the proof of *Case 4*.

If we have that infinitely many g_n 's have pairwise disjoint index sets then for suitable $n \in \mathbb{N}$ and $A \in \mathcal{A}_n$ the functional $g = \frac{1}{\sqrt{n}} \sum_{i \in A} g_i \in G_{\ell_2}$ will give us a contradiction.

Assume that only finitely many g_n 's have pairwise disjoint index sets.

Let $g_n = \sum_{i \in D_n} c_i \phi_i$ where $(\phi_i)_{i \in D_n}$ have pairwise disjoint index sets. For every $j \in \mathbb{N}$ we set $\phi_i = \phi_{i,j}^1 + \phi_{i,j}^2$ where $\text{ind}(\phi_{i,j}^1) \subset \{1, \dots, j\}$ and $\text{ind}(\phi_{i,j}^2) \subset \{j+1, j+2, \dots\}$.

Then $g_n = g_{n,j}^1 + g_{n,j}^2 = \sum_{i \in D_{n,1}} c_i \phi_{i,j}^1 + \sum_{i \in D_{n,2}} c_i \phi_{i,j}^2$. Let us observe that $\#D_{n,1} \leq j$.

We distinguish the following two subcases.

Subcase 4a. There exists $j_0 \in \mathbb{N}$ such that for all but finitely many n 's,

$$|g_{n,j_0}^1(\sum_{i \in F_n} a_i e_i)| \geq \epsilon_0/2.$$

Since $\text{ind}(\phi_{i,j_0}^1) \subset \{1, \dots, j_0\}$ it follows that $\#\text{supp } \phi_{i,j_0}^1 \leq \sum_{i \leq j_0} n_{2i-1} = n_0$ and

$$|\phi_{i,j_0}^1(\sum_{i \in F_n} a_i e_i)| \leq n_0 \|\phi_{i,j_0}^1\|_\infty \max_{i \in F_n} |a_i| \leq \frac{n_0}{m_1^2} \max_{i \in F_n} |a_n|.$$

It follows

$$\epsilon_0/2 \leq |\sum_{i \in D_{n,1}} c_i \phi_{i,j_0}^1(\sum_{i \in F_n} a_i e_i)| \leq \sum_{i \in D_{n,1}} |c_i| \frac{n_0}{m_1^2} \max_{i \in F_n} |a_n| \leq j_0 \frac{n_0}{m_1^2} \max_{i \in F_n} |a_i|.$$

From the above relation as in *Case 1* we derive a contradiction.

Subcase 4b. For every $j, m \in \mathbb{N}$ there exists $n \in \mathbb{N}$, $n > m$ with

$$|g_{n,j}^1(\sum_{i \in F_n} a_i e_i)| \leq \epsilon_0/2.$$

Then we choose inductively an increasing sequence $(n_i)_{i \in \mathbb{N}}$ such that

$$|g_{n_i, j_{i-1}}^1(\sum_{l \in F_{n_i}} a_l e_l)| \leq \epsilon_0/2 \text{ where } j_{i-1} = \max\{\text{ind}(g_{n_k}) : k \leq i-1\}.$$

It follows that $|g_{n_i, j_{i-1}}^2(\sum_{l \in F_{n_i}} a_l e_l)| \geq \epsilon_0/2$ and the functionals $g_{n_i, j_{i-1}}^2$ have pairwise disjoint index sets.

Setting $g = \frac{1}{\sqrt{n}} \sum_{i \in A} g_{n_i, j_{i-1}}^2$ for suitable n and $A \in \mathcal{A}_n$ we derive a contradiction. \square

From Theorem 5.6 and Proposition 4.7 we obtain the following

Corollary 5.7. *The identity map $Id : X_{G_\xi} \rightarrow Y_{G_\xi}$ is strictly singular.*

We prove now that every subsequence of the basis generates a subspace with ℓ_1 -index greater than ω^ξ .

Proposition 5.8. *For every $M \in [\mathbb{N}]$ the subspace $\langle (e_n)_{n \in M} \rangle$ has ℓ_1 -index greater than ω^ξ .*

Proof. It is not hard to see that for every $j \in \mathbb{N}$ the following holds

$$(4) \quad \frac{1}{m_{2^j-1}^2} \leq \left\| \frac{1}{n_{2^j-1}} \sum_{i=1}^{n_{2^j-1}} e_{k_i} \right\|_{G_\xi} \leq \frac{2}{m_{2^j-1}^2}.$$

Let (f_1, \dots, f_d) be a special sequence where $f_i = \frac{1}{m_{2^{j_i-1}}} \sum_{k \in F_i} e_k^*$ with $\#F = n_{2^{j_i-1}}$ and $F \subset M$.

For every $i \leq d$ take the vector $x_i = \frac{m_{2^{j_i-1}}}{n_{2^{j_i-1}}} \sum_{k \in F_i} e_k$. It follows that $1 \leq \|x_i\|_{G_\xi} \leq 2$ and

$$\left\| \sum_{i=1}^d a_i x_i \right\|_{G_\xi} \geq \sum_{i=1}^d \epsilon_i f_i \left(\sum_{i=1}^d a_i x_i \right) = \sum_{i=1}^d |a_i|.$$

Since for every $F \in \mathcal{S}_\xi$ we have a special sequence $(f_i)_{i \in F}$ it follows that the subspace $\langle (e_n)_{n \in M} \rangle$ contains an ℓ_1 -tree with constant 1 of order ω^ξ . It follows that the ℓ_1 -index of the subspace $\langle (e_n)_{n \in M} \rangle$ is greater than ω^ξ . \square

From (4) it follows that for every $M \in [\mathbb{N}]$ the c_0 -index of the subspace generated by the subsequence $(e_n^*)_{n \in M}$ of the basis of $X_{G_\xi}^*$ is greater than ω^ξ . Indeed, take an G_ξ -special sequence $(f_i)_{i=1}^d$ supported in the set M such that each $f_i = \frac{1}{m_{2^{j_i-1}}} \sum_{k \in B} e_k^*$ with $\#B = n_{2^{j_i-1}}$. From (4) we get $\|f_i\| \geq 1/2$ for all $i \leq d$ and hence

$$1/2 \leq \left\| \sum_{i=1}^d \pm f_i \right\| \leq 1.$$

From the above inequality easily follows that the c_0 -index of $\langle (e_n^*)_{n \in M} \rangle$ is greater than ω^ξ .

We show now the the space X_{G_ξ} is ℓ_2 -saturated.

Theorem 5.9. *For any subspace Y of X_{G_ϵ} and any $\epsilon > 0$ there is a subspace of Y which is $(1 + \epsilon)$ isomorphic to ℓ_2 .*

The proof of the theorem follows the arguments of Lemma B.13 and Theorem B.14 from [3] and we omit it.

6. THE SPACE X_{G_ϵ} DOES NOT HAVE ℓ_1 AS A SPREADING MODEL.

In this section we prove that X_{G_ϵ} does not contain a sequence generating an ℓ_1 -spreading model. The basic tool for the proof is Proposition 6.3 which is a combinatorial result. We begin with the following lemmas.

Lemma 6.1. *For any $x \in c_{00}(\mathbb{N})$ and $\epsilon > 0$ there is $j_0 = j_0(x, \epsilon) \in \mathbb{N}$ such that for any $g \in G_{\ell_2}$ with $\text{ind}(g) \cap \{1, \dots, j_0\} = \emptyset$ we have $|g(x)| < \epsilon$*

Proof. Let $D = \|x\|_{\ell_1}$ and take j_0 so that $\sum_{j=j_0+1}^{\infty} m_{2^{j-1}}^{-4} < (\epsilon/D)^2$. Take now any $g \in G_{\ell_2}$, $g = \sum_i a_i \phi_i$ with $\text{ind}(g) \cap \{1, \dots, j_0\} = \emptyset$ and compute

$$\begin{aligned} |g(x)| &\leq \left(\sum_i a_i^2\right)^{1/2} \left(\sum_i |\phi_i(x)|^2\right)^{1/2} \leq \left(\sum_i \|\phi_i\|_\infty^2 \|x\|_{\ell_1}^2\right)^{1/2} \\ &\leq D \left(\sum_{j=j_0+1}^{\infty} m_{2^{j-1}}^{-4}\right)^{1/2} < \epsilon. \end{aligned}$$

□

Lemma 6.2. *Let $(x_n)_{n \in \mathbb{N}}$ be a normalized block sequence and $\epsilon > 0$. There exist $j_1 \in \mathbb{N}$ and $L \in [\mathbb{N}]$ such that for every f of type I with $\text{ind}(f) \geq j_1$ it holds that $|f(x_n)| > \epsilon$ for at most one $n \in L$.*

Proof. We set $\mathcal{D}_1 = \{(n, m) : \text{exists } f \text{ of type I with } |f(x_n)| \geq \epsilon \text{ and } |f(x_m)| \geq \epsilon\}$.

If there exists $L \in [\mathbb{N}]$ with $[L]^2 \cap \mathcal{D}_1 = \emptyset$ the proof is complete. Otherwise by Ramsey theorem there exists L_1 such that $[L_1]^2 \subset \mathcal{D}_1$. Let $l_1 = \min L_1$ and let j_1 be such that $|f(x_{l_1})| < \epsilon$ for all f of type I with $\text{ind}(f) \geq j_1$.

We set

$$\mathcal{D}_2 = \{(n, m) \in [L_1]^2 : \text{exists } f \text{ of type I with } \text{ind}(f) \geq j_1, |f(x_n)| \geq \epsilon \text{ and } |f(x_m)| \geq \epsilon\}$$

If there exists $L_2 \in [L_1]$ with $[L_2]^2 \cap \mathcal{D}_2 = \emptyset$ the proof is complete. Otherwise by Ramsey theorem there exists L_2 such that $[L_2]^2 \subset \mathcal{D}_2$. Let $l_2 = \min L_2 > l_1$ and let j_2 be such that $|f(x_{l_2})| < \epsilon$ for all f of type I with $\text{ind}(f) \geq j_2$.

If the conclusion does not hold, choosing k_0 such the $\epsilon\sqrt{k_0} > 1$, following the above arguments we get $L_1 \supset L_2 \supset \dots \supset L_{k_0+1}$ such that $[L_i]^2 \subset \mathcal{D}_i$, where

$$\mathcal{D}_i = \{(n, m) \in [L_{i-1}]^2 : \text{exists } f \text{ of type I with } \text{ind}(f) \geq j_{i-1}, |f(x_n)| \geq \epsilon \text{ and } |f(x_m)| \geq \epsilon\}$$

and $|f(x_{\min L_i})| < \epsilon$ for all f of type I with $\text{ind}(f) \geq j_i$.

Setting $l_i = \min L_i$ we get $(l_i, l_{k_0+1}) \in \mathcal{D}_i$ for every $i = 1, \dots, k_0$ and therefore there exists f_i of type I with $\text{ind}(f_i) \in [j_{i-1}, j_i)$ such that $|f_i(x_{l_{k_0+1}})| \geq \epsilon$.

It follows $\frac{1}{\sqrt{k_0}} \sum_{i=1}^{k_0} \epsilon_i f_i(x_{k_0+1}) \geq \epsilon\sqrt{k_0} > 1$, a contradiction. □

Proposition 6.3. *Let $0 < \epsilon < 10^{-10}$ and $(x_n)_{n \in \mathbb{N}}$ be a normalized block basis such that for all $n_1 < n_2 < n_3$ there exists $\phi \in G_{\ell_2}$ such that*

$$(5) \quad |\phi(x_{n_i})| \geq 1 - \epsilon \text{ for all } i=1,2,3.$$

Then there exists $j_1 \in \mathbb{N}$ and $L \in [\mathbb{N}]$ such that for all $n \in L$ there exists $\phi_n = \sum_{i \in B_n} \lambda_{i,n} \phi_{i,n}$ with $\text{ind}(\phi_{i,n}) \subset \{1, \dots, j_1\}$ for all $i \in B_n$ and $\sum_{i \in B_n} \lambda_{i,n}^2 \leq 1$ such that

$$\phi_n(x_n) \geq 0.75.$$

Proof. From Lemma 6.1 we get $j_0 \in \mathbb{N}$ such that

$$(6) \quad \forall \phi = \sum_{i=1}^d \lambda_i \phi_i \text{ with } \text{ind}(\phi_i) \cap \{1, \dots, j_0\} = \emptyset \text{ and } \sum_{i \leq d} \lambda_i^2 \leq 1 \Rightarrow |\phi(x_1)| < \epsilon$$

Let $\delta > 0$ such that $\delta j_0 < \epsilon$. From Lemma 6.2 we get $M \in [\mathbb{N}]$ and $j_1 \in \mathbb{N}$ such that

$$(7) \quad \text{for every } f \text{ of type I with } \text{ind}(f) > j_1, |f(x_n)| > \delta \text{ for at most one } n \in M.$$

Let $M_1 = \{1\} \cup M$. After reordering we may assume that $M_1 = \mathbb{N}$.

Let $n \in \mathbb{N}$ and for every $1 < k < n$ consider the triple $(1, k, n) \in [\mathbb{N}]^3$. Let $\phi_{k,n} = \sum_{s \in S_{k,n}} c_s \phi_s \in G_{\ell_2}$, where $\phi_s \in F(s)$, be the functional we obtain from (5) for x_1, x_k and x_n . We set

$$J_{k,n} = \{s \in S_{k,n} : \text{ind}(\phi_s) > j_0\} \text{ and } G_{k,n} \text{ the complement of } J_{k,n}.$$

From (6) we get $|\sum_{s \in J_{k,n}} c_s \phi_s(x_1)| < 10^{-10}$ and therefore

$$(8) \quad \left(\sum_{s \in G_{k,n}} c_s^2 \right)^{1/2} \geq \left| \sum_{s \in G_{k,n}} c_s \phi_s(x_1) \right| > 1 - 2/10^{10}.$$

The disjointness of the index sets of the segments $s \in S_{k,n}$ yields that $\#G_{k,n} \leq j_0$ for every k .

By (8) we get $\sum_{s \in G_{k,n}} c_s^2 \geq (1 - 2/10^{10})^2$ and therefore

$$(9) \quad \sum_{s \in J_{k,n}} c_s^2 \leq 1 - (1 - 2/10^{10})^2 < 4/10^{10} \Rightarrow \left\| \sum_{s \in J_{k,n}} c_s \phi_s \right\|_{G_\xi}^* \leq 2/10^5.$$

From (5) and (9) it follows that for $r = k, n$

$$(10) \quad \left| \sum_{s \in G_{k,n}} c_s \phi_s(x_r) \right| \geq 1 - 10^{-10} - 2/10^5 > 1 - 3/10^5.$$

For every $s \in G_{k,n}$, if $\phi_s = \sum_{i \in V_s} \epsilon_i f_{s,i}$ let $f_{s,k,n}$ be its first node with the property $\text{range}(f_{s,i}) \cap \text{range}(x_n) \neq \emptyset$. Set

$$\begin{aligned} G_{k,n}^0 &= \{s \in G_{k,n} : \text{ind}(f_{s,k,n}) \leq j_1\}, \\ G_{k,n}^1 &= \{s \in G_{k,n} : \text{ind}(f_{s,k,n}) > j_1 \text{ and } |f_{s,k,n}(x_n)| \leq \delta\}, \\ G_{k,n}^2 &= \{s \in G_{k,n} \setminus G_{k,n}^1 : \text{ind}(f_{s,k,n}) > j_1 \text{ and } |f_{s,k,n}(x_k)| \leq \delta\}. \end{aligned}$$

From (7) it follows that for every $s \notin (G_{k,n}^0 \cup G_{k,n}^1)$ it holds that $|f_{s,k,n}(x_k)| \leq \delta$ and hence the sets $G_{k,n}^i$, $i = 0, 1, 2$ give a partition of $G_{k,n}$.

For every $1 < k < n$ we set

$$(11) \quad H_{k,n} = \begin{cases} G_{k,n}^0 & \text{if } \left| \sum_{s \in G_{k,n}^0} c_s \phi_s(x_k) \right| \geq 0.75 \quad (**), \\ G_{k,n}^1 & \text{if } (**) \text{ fails and } \sum_{s \in G_{k,n}^1} c_s^2 \geq \sum_{s \in G_{k,n}^2} c_s^2, \\ G_{k,n}^2 & \text{otherwise.} \end{cases}$$

By Ramsey theorem passing to an infinite subset N of \mathbb{N} we may assume that there exists $i \in \{0, 1, 2\}$ such that for all $1 < k < n \in N$, $H_{k,n} = G_{k,n}^i$. Without loss of generality we may assume that $N = \mathbb{N}$.

If $H_{k,n} = G_{k,n}^0$ for all $k < n$ then the proof is complete. Next we show that the cases $H_{k,n} = G_{k,n}^1$ or $H_{k,n} = G_{k,n}^2$ are not possible.

Claim. Assume that either $H_{k,n} = G_{k,n}^1$ for all $k < n$ or $H_{k,n} = G_{k,n}^2$ for all $k < n$.

Then there exists $N \in \mathbb{N}$ such that for all $n \in \mathbb{N}$ there exists a family U_n of segments such that:

- 1) $\#U_n \leq N$.
- 2) For every $1 < k < n$, $\sup\{\phi(x_k) : \phi \in \overline{U}_n\} \geq \delta$.

Let's see first how the conclusion of the claim makes impossible $H_{k,n} = G_{k,n}^1$ or $H_{k,n} = G_{k,n}^2$.

Let $U_n = \{s_{1,n}, \dots, s_{d_n,n}\}$ with $d_n \leq N$ for every $n > 1$. From 2) we obtain that for every $1 < k < n$ there exist $\phi_k \in \overline{U}_n$ such that

$$\phi_k(x_k) \geq \delta.$$

By the compactness of the set G'_ξ passing to an infinite subset of \mathbb{N} we get that there exist finite segments s_1, \dots, s_N such that $w^* - \lim_{n \rightarrow \infty} s_{i,n} = s_i$ for every $i \leq N$.

Let $k \in \mathbb{N}$ such that $\text{supp}(x_k) > \max\{\text{maxsupp } s_i : i \leq N\}$. Then since $w^* - \lim_{n \rightarrow \infty} s_{i,n} = s_i$ we obtain that there exists $n_0 > k$ such that $\text{supp } s_{i,n_0} \cap \text{supp}(x_k) = \emptyset$ for all $i \leq N$, a contradiction.

We proceed now to the proof of the Claim.

Assume that $H_{k,n} \neq G_{k,n}^0$. It follows that $|\sum_{s \in G_{k,n}^0} c_s \phi_s(x_k)| < 0.75$ and from (10) we get

$$(12) \quad \left| \sum_{s \in G_{k,n} \setminus G_{k,n}^0} c_s \phi_s(x_k) \right| \geq 1 - 3/10^5 - 0.75 = 0.25 - 3/10^5.$$

As in (8)-(9) we get that

$$(13) \quad 2 \sum_{s \in H_{k,n}} c_s^2 \geq \sum_{s \in G_{k,n} \setminus G_{k,n}^0} c_s^2 \geq (0.25 - 3/10^5)^2 \Rightarrow \sum_{s \in H_{k,n}} c_s^2 > 0.031.$$

From (13) we obtain $\sum_{s \in G_{k,n} \setminus H_{k,n}} c_s^2 \leq 0.97 \Rightarrow \|\sum_{s \in G_{k,n} \setminus H_{k,n}} c_s \phi_s\|_{G'_\xi}^* < 0.985$ and therefore for $l = k, n$ the following holds:

$$(14) \quad \left| \sum_{s \in H_{k,n}} c_s \phi_s(x_l) \right| \geq \left| \sum_{s \in G_{k,n}} c_s \phi_s(x_l) \right| - \left| \sum_{s \in G_{k,n} \setminus H_{k,n}} c_s \phi_s(x_l) \right| \geq 1 - 3/10^5 - 0.985 = 0.0149.$$

Assume now that $H_{k,n} = G_{k,n}^1$ for all $k < n$.

For every $s \in G_{k,n}^1$ we set $\tilde{s} = s|_{[1, \text{maxsupp}(f_{s,k,n})]}$.

Using finite induction we define sets $U_{2,n} \subset U_{3,n} \subset \dots \subset U_{n-1,n}$ as follows:

Set $U_{2,n} = \{\tilde{s} : s \in G_{2,n}^1\}$. Let $k = 3, \dots, n-1$ and assume that the set $U_{k-1,n}$ has been defined.

If there exists $\phi = \sum_i c_i \phi_i \in \overline{U}_{k-1,n}$, see Notation 4.5, with $|\phi(x_k)| \geq \delta$ we set $U_{k,n} = U_{k-1,n}$.

Assume that

$$(15) \quad \text{for all } \phi = \sum_i c_i \phi_i \in \overline{U}_{k-1,n} \text{ it holds that } |\phi(x_k)| < \delta.$$

For the functional $\phi_{k,n} = \sum_{s \in G_{k,n}^1} c_s \phi_s$ we set

$$E_{k,n} = \{s \in G_{k,n}^1 : \tilde{s} \in U_{k-1,n}\}.$$

Notice that the segments $\tilde{s}, s \in G_{k,n}^1 \setminus E_{k,n}$, are different from the segments in $U_{k-1,n}$ and therefore the functionals $\phi_s^1 = \phi_s|_{(\text{maxsupp}(f_{s,k,n}), +\infty)}$, $s \in G_{k,n}^1 \setminus E_{k,n}$, have disjoint index sets from the functionals in $\overline{U}_{k-1,n}$.

From (15) we get

$$(16) \quad \left| \sum_{s \in E_{k,n}} c_s \phi_{\tilde{s}}(x_k) \right| = \left| \sum_{s \in E_{k,n}} c_s \phi_s(x_k) \right| < \delta.$$

Combining the above inequality with (10) we get

$$(17) \quad \left(\sum_{s \in G_{k,n} \setminus E_{k,n}} c_s^2 \right)^{1/2} \geq \left| \sum_{s \in G_{k,n} \setminus E_{k,n}} c_s \phi_s(x_k) \right| > 1 - 3/10^5 - \delta > 1 - 4/10^5.$$

while from (14)

$$(18) \quad \left| \sum_{s \in G_{k,n}^1 \setminus E_{k,n}} c_s \phi_{\tilde{s}}(x_k) \right| > 0.0149 - \delta > 0.0148.$$

From (17) following the arguments we used to get (9) it follows

$$(19) \quad \left(\sum_{s \in E_{k,n}} c_s^2 \right)^{1/2} \leq (1 - (1 - 4/10^5)^2)^{1/2} \leq 0.009.$$

From the definition of the set $H_{k,n}$, using that $|f_{s,k,n}(x_n)| < \delta$, and (19) we obtain

$$\begin{aligned} \left| \sum_{s \in G_{k,n}^1 \setminus E_{k,n}} c_s \phi_s^1(x_n) \right| &\geq \left| \sum_{s \in G_{k,n}^1} c_s \phi_s(x_n) \right| - \sum_{s \in G_{k,n}^1} |c_s f_{s,k,n}(x_n)| - \left| \sum_{s \in E_{k,n}} c_s \phi_s(x_n) \right| \\ &\geq 0.0149 - j_0 \delta - 0.009 > 0.005. \end{aligned}$$

Set $U_{k,n} = U_{k-1,n} \cup \{\tilde{s} : s \in G_{k,n}^1 \setminus E_{k,n}\}$.

Note that in this case $\#(U_{k,n} \setminus U_{k-1,n}) \leq j_0$ and from (18) we obtain that for the functional $\phi_k = \sum_{s \in G_{k,n}^1 \setminus E_{k,n}} c_s \phi_{\tilde{s}} \in \overline{U}_{k,n}$ it holds that $\phi_k(x_k) \geq 0.0148$.

Let $U_{n-1,n}$ be the set of the segments we get after we complete the above procedure for $k = 2, \dots, n-1$.

We show now that there is no n with $\#U_{n-1,n} > j_0(2 + 1/\delta_0^2)$, $\delta_0 = 0.005$.

Indeed assume that there exists n such that $\#U_{n-1,n} > j_0(2 + 1/\delta_0^2)$. Since $\#U_{2,n} \leq j_0$ and $\#(U_{k,n} \setminus U_{k-1,n}) \leq j_0$ it follows that $U_{k-1,n} \not\subseteq U_{k,n}$ holds for at least $t = 1 + 1/\delta_0^2$ different k 's. For every such k we get a functional $\phi_k^1 = \sum_{s \in G_{k,n}^1 \setminus E_{k,n}} c_s \phi_s^1$ such that $\phi_k^1(x_n) > \delta_0$ and the functionals ϕ_k^1 have pairwise disjoint index sets. It follows that

$$\frac{1}{\sqrt{t}} \sum_k \phi_k^1(x_n) \geq \sqrt{t} \delta_0 > 1,$$

a contradiction.

Case 2. $H_{k,n} = G_{k,n}^2$ for every $1 < k < n$.

For every $s \in G_{k,n}^2$ we set $\tilde{s} = s|_{[1, \text{minsupp } f_{s,k,n}]}$.

Using finite induction we define sets $U_{2,n} \subset U_{3,n} \subset \dots \subset U_{n-1,n}$ as follows:

Set $U_{2,n} = \{\tilde{s} : s \in G_{2,n}^2\}$. Let $k = 3, \dots, n-1$ and assume that $U_{k-1,n}$ has been defined.

If there exists $\phi \in \overline{U}_{k-1,n}$ such that $|\phi(x_k)| > \delta$ we set $U_{k,n} = U_{k-1,n}$.

Assume that

$$(20) \quad \text{for all } \phi = \sum_i c_i \phi_i \text{ ind } \overline{U}_{k-1,n} \text{ it holds that } |\phi(x_k)| < \delta.$$

Set

$$E_{k,n} = \{s \in G_{k,n}^2 : \tilde{s} \in U_{k-1,n}\}.$$

We get that the segments $\tilde{s}, s \in G_{k,n}^2 \setminus E_{k,n}$ are different from the segments in $U_{k-1,n}$ and therefore the functionals $\phi_s^2 = \phi_{s|_{[\text{minsupp } f_{s,k,n}, +\infty]}}$ have disjoint index set from the functionals in $\overline{U}_{k-1,n}$.

As in the previous case, see (17),(19), we obtain that $(\sum_{s \in E_{k,n}} c_s^2)^{1/2} < 0.009$.

From (14) and (20) we obtain

$$(21) \quad \begin{aligned} \left| \sum_{s \in G_{k,n}^2 \setminus E_{k,n}} c_s \phi_{\tilde{s}}(x_k) \right| &\geq \left| \sum_{s \in G_{k,n}^2} c_s \phi_s(x_k) \right| - \sum_{s \in G_{k,n}^2} |c_s f_{s,k,n}(x_k)| - \left| \sum_{s \in E_{k,n}} c_s \phi_{\tilde{s}}(x_k) \right| \\ &= 0.0149 - j_0 \delta - \delta > 0.0148 \end{aligned}$$

Also we get

$$\begin{aligned} \left| \sum_{s \in G_{k,n}^2 \setminus E_{k,n}} c_s \phi_s^2(x_n) \right| &\geq \left| \sum_{s \in G_{k,n}^2} c_s \phi_s(x_n) \right| - \left| \sum_{s \in E_{k,n}} c_s \phi_s(x_n) \right| \\ &\geq 0.0149 - 0.009 > 0.005 \end{aligned}$$

We set $U_{k,n} = U_{k-1,n} \cup \{\tilde{s} : s \in G_{k,n}^2 \setminus E_{k,n}\}$. Note that in this case $\#(U_{k,n} \setminus U_{k-1,n}) \leq j_0$ and from (21) we get that for the functional $\phi_k = \sum_{s \in G_{k,n}^2 \setminus E_{k,n}} c_s \phi_{\tilde{s}} \in \overline{U_{k,n}}$ it holds that $\phi_k(x_k) \geq 0.0148$.

Let $U_{n-1,n}$ be the set of the segments we obtain after we complete the above procedure. Assuming that there exists n such that $\#U_{n-1,n} > j_0(2 + 1/\delta_0^2)$ as in the case where $H_{k,n} = G_{k,n}^1$ we get a contradiction. \square

Theorem 6.4. *The space X_{G_ϵ} does not contain a normalized sequence generating an ℓ_1 -spreading model.*

In the proof of the theorem we shall use the following well known fact.

Fact 6.5. *Let X be a Banach space and $(x_n)_n$ be a normalized sequence generating an ℓ_1 -spreading model with constant c . Then for every $\epsilon > 0$ there exists a normalized sequence $(y_n)_{n \in \mathbb{N}}$ generating an ℓ_1 -spreading model with constant $(1 - \epsilon)^{-1}$. In particular $(y_n)_n$ can be chosen such that each y_n is a normalized convex combination of $(x_n)_{n \in \mathbb{N}}$.*

Proof of Theorem 6.4. Let $(x_n)_{n \in \mathbb{N}}$ be a sequence which generates an ℓ_1 -spreading model with constant c . Since the space X_{G_ϵ} does not contain ℓ_1 by standard arguments passing to a subsequence and taking the differences we may assume that $(x_n)_{n \in \mathbb{N}}$ is a block sequence and generates a spreading model with constant $(1 - \epsilon)^{-1}$, $\epsilon < 10^{-10}$.

By proposition 6.3 we get that there exist $j_1 \in \mathbb{N}$ and $L \in [\mathbb{N}]$ such that for all $n \in L$ there exists $\phi_n \in G_{\ell_2}$ with index set contained in $\{1, \dots, j_1\}$ such that $\phi_n(x_n) \geq 0.75$.

It follows that for every $n \in L$ and every $\phi \in G_{\ell_2}$ with $\text{ind}(\phi) \subset \{j_1 + 1, \dots\}$ it holds that $|\phi(x_n)| < 0.75$.

Indeed if there exists such ϕ with $|\phi(x_n)| \geq 0.75$ then the functionals ϕ and ϕ_n have disjoint index sets and hence $\psi = \frac{\phi + \phi_n}{\sqrt{2}} \in K$. It follows that $\psi(x_n) \geq \frac{0.75 + 0.75}{\sqrt{2}} > 1$, a contradiction.

Let $n_0 \in \mathbb{N}$ with $n_0 > n_{j_1}$. Let $x = \frac{1}{n_0} \sum_{i=1}^{n_0} x_{l_i}$ with $n_0 \leq l_1 < \dots < l_{n_0} \in L$.

Since $(x_n)_{n \in \mathbb{N}}$ is assumed to generate an ℓ_1 -spreading model with constant $(1 - \epsilon)^{-1}$ it follows that $\|x\|_{G_\epsilon} \geq 1 - \epsilon$. Let $\phi \in K$ be a functional which norms x . It readily follows that $\phi = \sum_{i=1}^d \lambda_i \phi_i \in G_{\ell_2}$.

For every i let ϕ_i^1 be the part of ϕ_i with the index set contained in $\{1, \dots, j_1\}$ and ϕ_i^2 be the part with index set contained in $\{j_1 + 1, j_1 + 2, \dots\}$. Let $\phi_1 = \sum_{i=1}^d \lambda_i \phi_i^1$ and $\phi_2 = \sum_{i=1}^d \lambda_i \phi_i^2$.

By the above note we get that $\phi_2(x_{l_i}) \leq 0.75$ for all i and hence $\phi_2(x) \leq 0.75$.

Also for every f of type I with $\text{ind}(f) \leq j_1$ it follows that $|f(x)| \leq \frac{\#\text{supp}(f)}{n_0} \leq w(f)^{-1}$. Hence

$$|\phi^1(x)| \leq \sum_{i=1}^d \sum_{k \in \text{ind}(\phi_i^1)} m_k^{-1} \leq \sum_{k=1}^{\infty} m_k^{-1} < 0.1.$$

Therefore $|\phi(x)| \leq |\phi_1(x)| + |\phi_2(x)| < 0.85$, a contradiction. \square

7. THE SPACE X_ξ AS AN EXTENSION OF THE SPACE X_{G_ξ}

In this section we define the space X_ξ by its norming set K_ξ . We introduce the key ingredient for the definition of K_ξ - attractor sequences (Def. 7.1), and the basic tool in estimating the norm in X_ξ - a tree-analysis of a functional from K_ξ (Def. 7.3).

Let $(\Lambda_i)_{i \in \mathbb{N}}$ be a partition of \mathbb{N} into pairwise disjoint infinite sets. Let \mathbb{Q}_s denote the set of all finite sequences (f_1, \dots, f_d) such that for all i , $f_i \in c_{00}(\mathbb{N})$, $f_i \neq 0$, $f_i(n) \in \mathbb{Q}$ for all $n \in \mathbb{N}$ and $f_1 < f_2 < \dots < f_d$.

We fix a partition N_1, N_2 of \mathbb{N} . Let $\sigma : \mathbb{Q}_s \rightarrow N_2$ be an injective function such that

$$m_{2\sigma(f_1, \dots, f_d)}^{1/2} > \max\{|f_i(e_l)|^{-1} : l \in \text{supp } f_i, i \leq d\} \max \text{supp } f_d.$$

Such an injective function exists since the set \mathbb{Q}_s is countable.

Let K_ξ be the minimal subset of $c_{00}(\mathbb{N})$ satisfying the following conditions

- (1) K_ξ is symmetric i.e. if $f \in K_\xi$ then $-f \in K_\xi$, K_ξ is closed under the restriction of its elements to intervals of \mathbb{N} and $G_\xi \subset K_\xi$.
- (2) K_ξ is closed under $(\mathcal{A}_{n_{2j}}, m_{2j}^{-1})$ -operations.
- (3) K_ξ is closed under $(\mathcal{A}_{n_{2j-1}}, m_{2j-1}^{-1})$ -operations on attractor sequences.
- (4) K_ξ is closed under the operation $\sum_{i \in A} \lambda_i f_i$ whenever
 - (a) f_i is the result of an $(\mathcal{A}_{n_{j_i}}, m_{j_i}^{-1})$ -operation and $n_{j_i} \neq n_{j_k}$ for every $i \neq k \in A$
 - (b) $(\lambda_i)_{i \in A} \in B_{\ell_2} \cap [\mathbb{Q}]^{<\infty}$.
- (5) K_ξ is rationally convex.

In order to complete the definition of the set K_ξ we have to define the attractor sequences.

Definition 7.1. *A finite sequence (f_1, \dots, f_d) is said to be a n_{2j-1} -attractor sequence provided that*

- (1) $(f_1, \dots, f_d) \in \mathbb{Q}_s$ and $f_i \in K_\xi$ for all $i = 1, \dots, d \leq n_{2j-1}$.
- (2) The functional f_1 is the result of an $(\mathcal{A}_{n_{2j_1}}, m_{2j_1}^{-1})$ operation on a family of functionals of K_ξ for some $j_1 \in N_1$ with $m_{2j_1} > n_{2j_1}^3$. Also for every $i = 2, \dots, n_{2j-1}/2$, f_{2i-1} is the result of an $(\mathcal{A}_{n_{2\sigma(f_1, \dots, f_{2i-2})}}, m_{2\sigma(f_1, \dots, f_{2i-2})}^{-1})$ operation on a family of functionals of K_ξ .
- (3) $f_{2i} = e_{\lambda_{2i}^*}$ for some $\lambda_{2i} \in \Lambda_{\sigma(f_1, \dots, f_{2i-1})}$.

We say that $f \in K_\xi$ is of type I if it is a result of some $(\mathcal{A}_{n_j}, m_j^{-1})$ operation. In this case we set $w(f) = m_j$ and $\text{ind}(f) = j$.

We say that $f \in K_\xi$ is of type II if $f = \sum_{i=1}^n \lambda_i f_i$ for some $(f_i)_{i=1}^n \subset K_\xi$ with $w(f_i) \neq w(f_j)$ for all $i \neq j$ and $\sum_{i=1}^n \lambda_i^2 \leq 1$.

We say that f is of type III if it is a rational combination of elements of K_ξ .

Remarks 7.2. *a) Using the partition of $\mathbb{N} = \cup_i \Lambda_i$ it follows that the set of the attractor sequences has a tree structure i.e. if $(f_i)_{i=1}^{n_{2j-1}}$, $(g_i)_{i=1}^{n_{2k-1}}$ are two attractor sequences either $g_i \neq f_r$ for every i, r or there exists $i_0 \leq \min\{n_{2j-1}, n_{2k-1}\}$ such that $f_i = g_i$ for all $i \leq i_0 - 1$ and $f_i \neq g_r$ for all $i, r \geq i_0$. In particular $w(f_{2i-1}) \neq w(g_{2r-1})$ for all $2i - 1, 2r - 1 > i_0$ and $f_{2i} \neq g_{2r}$ for all $2i, 2r > i_0$.*

b) Note that in the definition of the norming set when we take functionals $\sum_{i=1}^n \lambda_i f_i$ with $\sum_{i=1}^n \lambda_i^2 \leq 1$ we require $w(f_i) \neq w(f_j)$ but do not require $\text{supp}(f_i) \cap \text{supp}(f_j) = \emptyset$. This is will be essential in order to get that no sequence generates an ℓ_1 -spreading model.

Definition 7.3. *[The tree T_f of a functional $f \in K_\xi$] Let $f \in K_\xi$. By a tree of f we mean a finite family $T_f = (f_\alpha)_{\alpha \in \mathcal{A}}$ indexed by a finite tree \mathcal{A} with a unique root $0 \in \mathcal{A}$ such that the following holds*

- (1) $f_0 = f$ and $f_\alpha \in K_\xi$ for all $\alpha \in \mathcal{A}$.
- (2) An $\alpha \in \mathcal{A}$ is maximal if and only if $f_\alpha \in G_\xi$.
- (3) For every $\alpha \in \mathcal{A}$ not maximal, denoting by S_α the set of the immediate successors of α one of the following holds
 - (a) There exists $j \in \mathbb{N}$ such that $\#S_\alpha \leq n_j$ and f_α is the result of the $(\mathcal{A}_{n_j}, m_j^{-1})$ -operation in the set $\{f_\beta : \beta \in S_\alpha\}$. In this case we say that the weight of f is $w(f_\alpha) = m_j$.

- (b) $f_\alpha = \sum_{\beta \in S_\alpha} \lambda_\beta f_\beta$ where $\sum_{\beta \in S_\alpha} \lambda_\beta^2 \leq 1$ the functionals f_β are of type I and have different weights.
- (c) f_α is a rational convex combination of the elements f_β , $\beta \in S_\alpha$. Moreover for every $\beta \in S_\alpha$ $\text{range } f_\beta \subset \text{range } f_\alpha$

The order $o(f_\alpha)$ for each $\alpha \in \mathcal{A}$ is also defined by backward induction as follows.

If $f_\alpha \in G_\xi$ then $o(f_\alpha) = 1$, otherwise $o(f_\alpha) = 1 + \max\{o(f_\beta) : \beta \in S_\alpha\}$.

The order $o(T_f)$ of the aforementioned tree is defined to be equal to $o(f_0)$.

Definition 7.4. The order $o(f)$ of an $f \in K_\xi$, is defined as

$$o(f) = \min\{o(T_f) : T_f \text{ is a tree of } f\}.$$

Every functional $f \in K_\xi$ admits a tree analysis not necessarily unique.

Definition 7.5. We define $X_\xi = \overline{(c_{00}(\mathbb{N}), \|\cdot\|_{K_\xi})}$ and we denote the norm $\|\cdot\|_{K_\xi}$ by $\|\cdot\|$.

Definition 7.6. Let $k \in \mathbb{N}$. A vector $x \in c_{00}(\mathbb{N})$ is said to be a $C - \ell_1^k$ average if there exists $x_1 < \dots < x_k$, $\|x_i\| \leq C\|x\|$ and $x = \frac{1}{k} \sum_{i=1}^k x_i$. Moreover, if $\|x\| = 1$ then x is called a normalized $C - \ell_1^k$ average.

Lemma 7.7 ([27], (or [16], Lemma 4)). Let $j \geq 1$, x be a $C - \ell_1^{n_j}$ average. Then for every $n \leq n_{j-1}$ and every $E_1 < \dots < E_n$, we have that

$$\sum_{i=1}^n \|E_i x\| \leq C(1 + \frac{2n}{n_j}) < \frac{3}{2}C.$$

Proposition 7.8. For every block sequence $(y_\ell)_\ell \subset X_\xi$, $\epsilon > 0$ and every $k \geq m_2$ there exists $x \in \langle (y_\ell)_\ell \rangle$ which is a normalized $2 - \ell_1^k$ average.

In particular for every $k \in \mathbb{N}$ there exists $n \in \mathbb{N}$ such that for every finite block subsequence $(y_{\ell_i})_{i=1}^n$ of $(y_\ell)_\ell$ there exists $(z_i)_{i=1}^k$, a normalized block sequence of $(y_{\ell_i})_{i=1}^n$ such that $\|z_1 + \dots + z_k\| \geq k/2$.

Lemma II.22 [7] give us the existence of such average.

Proposition 7.9. X_ξ is a strictly singular extension of X_{G_ξ} , i.e. the identity map $X_{G_\xi} \rightarrow X_\xi$ is strictly singular.

Proof. Let Y be a block subspace of X_ξ and $j \in \mathbb{N}$. By the Proposition 7.8 we obtain a sequence $(y_n)_{n \in \mathbb{N}} \subset Y$ of normalized $\ell_1^{k_n}$ averages and by Proposition 5.4 for any fixed $j > 1$ we get $L \in [\mathbb{N}]$ such that for every $g \in G_\xi$

$$\{n \in L : |g(y_n)| \geq m_{2j}^{-2}\} \leq n_{2j-1}.$$

It follows that

$$\left\| \frac{1}{n_{2j}} \sum_{i=1}^{n_{2j}} y_i \right\|_{G_\xi} \leq \frac{n_{2j-1}}{n_{2j}} + \frac{1}{m_{2j}^2} \leq \frac{2}{m_{2j}^2}$$

while $\left\| \frac{1}{n_{2j}} \sum_{i=1}^{n_{2j}} y_i \right\| \geq \frac{1}{m_{2j}}$. Since j was arbitrarily chosen we have a strictly singular extension. \square

8. THE AUXILIARY SPACE

Now we define for any $j_0 > 1$ the auxiliary space $X_{W_{j_0}}$ and discuss construction and relation between its norming sets: W_{j_0} and W'_{j_0} . Next in Subsection 8.1 we estimate the supremum norm of any norming functional from W'_{j_0} (Lemma 8.2) and its action on a suitable average of $(e_n)_n$ of length n_{j_0} (Lemma 8.10).

Let $j_0 > 1$ be fixed. We set $C_{j_0} = \{\sum_{i \in F} \pm e_i^* : F \in \mathcal{A}_{n_{j_0-1}}\}$. Let W_{j_0} be the minimal subset of $c_{00}(\mathbb{N})$ satisfying the following conditions

- 1) W_{j_0} contains the set C_{j_0} .

- 2) W_{j_0} is closed under $(\mathcal{A}_{2n_j}, m_j^{-1})$ -operations for every $j \in \mathbb{N}$.
- 3) W_{j_0} is closed under the operation $\sum_{i \in A} \lambda_i f_i + \sum_{i \in B} \lambda_i e_{t_i}$ whenever $A \cap B = \emptyset$, $t_i \neq t_j$ for all $i \neq j \in B$, f_i is the result of an $(\mathcal{A}_{2n_{j_i}}, m_{j_i})$ -operation, $m_{j_i} \neq m_{j_k}$ for all $i \neq k \in A$ and $\sum_{i \in A \cup B} \lambda_i^2 \leq 1$.
- 4) W_{j_0} is rationally convex.

The auxiliary space is the space $X_{W_{j_0}} = \overline{(c_{00}(\mathbb{N}), \|\cdot\|_{W_{j_0}})}$.

For every functional $f \in W_{j_0}$ which is the result of an $(\mathcal{A}_{2n_j}, m_j^{-1})$ -operation we set $w(f) = m_j$ and we say that f is of type I. Every functional of the form $\sum_{i \in A} \lambda_i f_i + \sum_{i \in B} \lambda_i e_{n_i} \in W_{j_0}$ is called of type II. Every functional which is a convex combination of elements of W_{j_0} is called of type III.

Consider the minimal set W'_{j_0} which satisfies Properties 1)-3) in the definition of W_{j_0} . Then following the arguments of Lemma 3.15 [9], every $f \in W_{j_0}$ of type I with $w(f) = m_j$ can be written as a convex combination $\sum_i \lambda_i f_i$ of functionals from the set W'_{j_0} with $w(f_i) = m_j$ for every i . Hence in order to get an upper estimate for functionals of type I from the set W_{j_0} it is enough to get an upper estimate for the functionals from the set W'_{j_0} .

An alternative definition of the norming set W'_{j_0} is the following:

Set $W_0 = C_{j_0}$ and assume that W_{n-1} has been defined.

We set

$$W_n^1 = \bigcup_j \left\{ m_j^{-1} \sum_{i=1}^d f_i : (f_i)_{i=1}^d \text{ is } \mathcal{A}_{2n_j}\text{-admissible subset of } W_{n-1} \right\}$$

and

$$W_n^2 = \left\{ \sum_{i \in A} \lambda_i f_i + \sum_{i \in B} \lambda_i e_{t_i} : \sum_{i \in A \cup B} \lambda_i^2 \leq 1, t_i \neq t_j \text{ for all } i \neq j \in B, A \cap B = \emptyset, \right. \\ \left. f_i \in W_{n-1} \text{ is of type I for all } i \in A \text{ with } w(f_i) \neq w(f_j) \text{ for all } i \neq j \in A \right\}.$$

We set $W_n = W_n^1 \cup W_n^2$. Then $W'_{j_0} = \bigcup_n W_n$.

Every $f \in W'_{j_0}$ admits a tree analysis $(f_\alpha)_{\alpha \in \mathcal{T}}$ satisfying properties similar to Definition 7.3.

Remarks 8.1. Let $f \in W'_{j_0}$ and $(f_\alpha)_{\alpha \in \mathcal{T}}$ be a tree analysis of f .

a) If α is a terminal node then $f_\alpha \in C_{j_0}$. In particular if $\alpha \in S_\beta$ and f_β is of type II then $f_\alpha = \pm e_k^*$ for some k .

b) If α, β are different maximal nodes it is not necessary true that $\text{supp}(f_\alpha) \cap \text{supp}(f_\beta) = \emptyset$.

c) A consequence of b) is that it does not hold that $\|f\|_\infty \leq 1$ for every $f \in W'_{j_0}$. We show in Lemma 8.2 that there is a constant $c_1 > 1$ such that $\|f\|_\infty \leq c_1$ for every $f \in W'_{j_0}$.

8.1. Estimates on the auxiliary space. Set $c_0 = \sum_{j=1}^\infty m_j^{-2}$ and $c_1 = (\sum_{n=0}^\infty c_0^n)^{1/2}$.

Lemma 8.2. For every $f \in W'_{j_0}$ it holds that $|f(e_t)| \leq \begin{cases} \frac{c_1}{w(f)} & \text{if } f \text{ is of type I} \\ c_1 & \text{if } f \text{ is of type II.} \end{cases}$

Proof. We prove by induction that for every $f \in W_n$ the following holds:

$$(22) \quad |f(e_t)| \leq u_f (1 + c_0 + \cdots + c_0^n)^{1/2}$$

where $u_f = w(f)^{-1}$ if f is of type I and $u_f = 1$ otherwise.

Assume that for every $f \in W_{n-1}$ it holds that $|f(e_t)| \leq u_f (1 + c_0 + \cdots + c_0^{n-1})^{1/2}$ which is certainly true for $n = 1$.

If $f = w(f)^{-1} \sum_{i=1}^d f_i \in W_n^1$ then there exists unique $i \leq d$ such that $t \in \text{supp}(f_i)$. It follows from the inductive hypothesis that

$$|f(e_t)| \leq \frac{1}{w(f)} |f_i(e_t)| \leq \frac{1}{w(f)} (1 + c_0 + \dots + c_0^{n-1})^{1/2}.$$

If $f = \sum_{i \in A} \lambda_i f_i + \sum_{i \in B} \lambda_i e_i^* \in W_n^2$ then either $t \notin B$ or there exists unique $i_t \in B$ such that $e_t = e_{i_t}$. Assume that the latter holds. Setting $A_t = \{i \in A : t \in \text{supp} f_i\}$ from the inductive hypothesis and Hölder inequality we get

$$\begin{aligned} |f(e_t)| &\leq \sum_{i \in A_t} |\lambda_i f_i(e_t)| + |\lambda_{i_t}| \leq (1 + \sum_{i \in A_t} w(f_i)^{-2} (1 + c_0 + \dots + c_0^{n-1}))^{1/2} \\ &\leq (1 + (1 + c_0 + \dots + c_0^{n-1}) \sum_j m_j^{-2})^{1/2} = (1 + c_0 + \dots + c_0^n)^{1/2}. \end{aligned}$$

This proves (22) and completes the proof of the lemma. \square

Recall that in the definition of the space we have chosen $n_j = (2n_{j-1})^{s_{j-1}}$ where $2^{s_{j-1}} = m_j^3$.

Lemma 8.3. *Let $f \in W'_{j_0}$ such that $|f(e_t)| > \frac{3}{m_{j_0}}$ for every $t \in \text{supp}(f)$. Then*

$$\#\text{supp}(f) \leq (2n_{j_0-1})^{s_{j_0-1}-2}.$$

This lemma is the key ingredient of our evaluations in the auxiliary space. Its proof is given in four steps. As usually we shall consider the tree analysis $(f_\alpha)_{\alpha \in \mathcal{T}}$ of a functional f in W'_{j_0} . In the first two steps we shall show that we may assume that the height of the tree is less or equal to $s_{j_0-1} - 3$, and for all functionals f_α of type I in a tree analysis of f it holds that $w(f_\alpha) < m_{j_0}$. This will enable us to use an argument similar to Lemma A4 in [3] and get the result.

Proof. Let $(f_\alpha)_{\alpha \in \mathcal{T}}$ be a tree analysis of the functional f . For every $t \in \text{supp}(f)$ we set

$$D_1 = \{\alpha \in \mathcal{T} : \alpha \text{ is a minimal node with } f_\alpha \text{ of type I and } w(f_\alpha) \geq m_{j_0}\}.$$

Let us observe that the set D_1 consists of incomparable nodes. We set

$$M_1 = \{\gamma \in \mathcal{T} : \gamma \text{ is a maximal node and there exists } \alpha \in D_1 \text{ with } \alpha \prec \gamma\}$$

and also

$$M_2 = \{\gamma \in \mathcal{T} : \gamma \text{ is a maximal node and } \gamma \notin M_1\}.$$

The definition of M_2 yields that for every $\gamma \in M_2$ and every $\alpha \prec \gamma$ with f_α of type I, $w(f_\alpha) < m_{j_0}$.

We denote by f_1 the functional that we get following backwards the tree analysis of f with maximal nodes the nodes of M_1 and f_2 the corresponding one with maximal nodes those of M_2 . Observe that $f = f_1 + f_2$.

We also set $\mathcal{T}_1 = \{\beta : \text{there exists } \alpha \in D_1 \text{ with } \beta \preceq \alpha\}$ which is a complete subtree of the tree \mathcal{T} and $(g_\alpha)_{\alpha \in \mathcal{T}_1}$ the tree analysis of the functional f_1 . Let us observe that every maximal node α of \mathcal{T}_1 belongs to the set D_1 and $g_\alpha = f_\alpha$.

Sublemma 8.4. *For the functional f_1 , its tree analysis $(g_\alpha)_{\alpha \in \mathcal{T}_1}$ and $t \in \mathbb{N}$ the following hold:*

- (1) for every $\alpha \in D_1$ we have $|g_\alpha(e_t)| \leq \frac{c_1}{m_{j_0}}$.
- (2) for every non-maximal node $\alpha \in \mathcal{T}_1$ for which g_α is of type I we have

$$|g_\alpha(e_t)| < \frac{9c_1}{8w(g_\alpha)m_{j_0}}.$$

- (3) For all $\alpha \in \mathcal{T}_1$ non-maximal with $g_\alpha = \sum_{\beta \in S_\alpha} \lambda_\beta g_\beta$ of type II we have

$$|g_\alpha(e_t)| \leq \frac{9c_1}{8m_{j_0}}.$$

Proof. (1) follows from Lemma 8.2 and the definition of the set M_1 .

(2) and (3) are proved inductively. Let $\alpha \in \mathcal{T}_1$ and assume the result for all $\alpha \prec \beta$.

If $g_\alpha = w(g_\alpha)^{-1} \sum_{\beta \in S_\alpha} g_\beta$ it is clear that (2) holds.

We pass now to prove (3). Assume that $g_\alpha = \sum_{\beta \in S_\alpha} \lambda_\beta g_\beta$. Then $|g_\alpha(e_t)| = |\sum_{\beta \in S_\alpha} \lambda_\beta g_\beta(e_t)|$.

If $\beta \in S_\alpha$ is not maximal then g_β is of type I and the inductive assumptions yields

$$(23) \quad |g_\beta(e_t)| \leq \frac{9c_1}{8w(g_\beta)m_{j_0}}.$$

Let $\beta \in S_\alpha$ be a maximal node, namely $\beta \in D_1$. From the definition of the set M_1 it follows that g_β is of type I and $w(g_\beta) = m_{j_0+k}$ for some $k \geq 0$. Lemma 8.2 yields

$$(24) \quad |g_\beta(e_t)| \leq \frac{c_1}{w(g_\beta)} (\leq \frac{1}{m_{j_0} m_{j_0}^k} \text{ if } k > 0).$$

We also observe that there exists at most one $\beta \in S_\alpha$ with $w(g_\beta) = m_{j_0}$. Without loss of generality we assume that there does exist $\beta_0 \in S_\alpha$ with $w(g_{\beta_0}) = m_{j_0}$.

Set $S_\alpha^1 = \{\beta \in S_\alpha : g_\beta \text{ of type I with } w(g_\beta) < m_{j_0}\}$,

$S_\alpha^2 = \{\beta \in S_\alpha : g_\beta \text{ of type I with } w(g_\beta) > m_{j_0}\} \cup \{\beta \in S_\alpha : \beta \text{ is maximal}\}$.

It follows from (23), (24) and the fact that $w(g_\beta) \neq w(g_\gamma)$ for every $\beta \neq \gamma \in S_\alpha$, that

$$\begin{aligned} |g_\alpha(e_t)| &\leq \sum_{\beta \in S_\alpha^1} |\lambda_\beta g_\beta(e_t)| + |\lambda_{\beta_0} g_{\beta_0}(e_t)| + \sum_{\beta \in S_\alpha^2} |\lambda_\beta g_\beta(e_t)| \\ &\leq |\lambda_{\beta_0} g_{\beta_0}(e_t)| + \sum_{\beta \in S_\alpha^1} \frac{2c_1 |\lambda_\beta|}{m_{j_0} w(g_\beta)} + \sum_{\beta \in S_\alpha^2} \frac{c_1 |\lambda_\beta|}{w(g_\beta)} \\ &\leq |g_{\beta_0}(e_t)| + \sum_{i < j_0} \frac{2c_1}{m_{j_0} m_i} + \sum_{k \geq 1} \frac{1}{m_{j_0}^{k+1}} \leq \frac{c_1}{m_{j_0}} + \frac{4c_1}{m_{j_0} m_1} \leq \frac{9c_1}{8m_{j_0}}. \end{aligned}$$

□

Remark 8.5. Let $(f_\alpha)_{\alpha \in \mathcal{T}}$ be a tree analysis of f_1 such that for all f_α of type I it holds that $w(f_\alpha) \neq m_{j_0}$. The proof of the above sublemma yields that in this case $|f_1(e_t)| \leq \frac{9c_1}{8m_{j_0}^2}$.

Let us observe that, since $f = f_1 + f_2$, the previous sublemma yields that $\text{supp}(f_2) = \text{supp}(f)$. So we may assume that $f = f_2$. Let $(f_\alpha)_{\alpha \in \mathcal{T}}$ be the tree analysis of f with all maximal nodes belonging to M_2 . For $\beta \in \mathcal{T}$ we denote by

$$|\beta| = \#\{\alpha \in \mathcal{T} : \alpha \prec \beta\}$$

the order of β .

We set $d_{j_0} = s_{j_0-1} - 3$ and we observe that $c_1 2^{-d_{j_0}} < m_{j_0}^{-2}$. We also set

$$D = \{\alpha : \alpha \text{ is a maximal node of } \mathcal{T} \text{ with } |\alpha| \leq d_{j_0} \text{ and if } \alpha \in S_\beta,$$

$$f_\beta \text{ is of type II then } |\lambda_\alpha| \geq m_{j_0}^{-1}\}.$$

With the next two sublemmas we show that $\text{supp}(f) = \cup_{\alpha \in D} \text{supp}(f_\alpha)$.

Sublemma 8.6. Let $t \in \text{supp}(f)$ be such that there is no maximal node β with $|\beta| \leq d_{j_0}$ and $t \in \text{supp}(f_\beta)$. Then it holds $|f(e_t)| \leq c_1 2^{-d_{j_0}}$.

Proof. Let $t \in \text{supp}(f)$. By induction we shall show the following: for every $\gamma \in \mathcal{T}$ such that $|\gamma| = d_{j_0} - j$, $j = 0, 1, \dots, d_{j_0}$ we have that

$$(25) \quad |f_\gamma(e_t)| \leq c_1 2^{-j} u_\gamma,$$

where $u_\gamma = m_{r-1}^{-1}$ if f_γ is of type I and $w(f_\gamma) = m_r$, (we make the convention that $m_0 = 2^{-2}$), $u_\gamma = 1$ if f_γ is of type II.

First we observe that Lemma 8.2 yields $|f_\beta(e_t)| \leq c_1$ for every maximal node $\beta \in \mathcal{T}$.

Assume that the result holds for all $\delta \succ \gamma$ with $|\delta| = d_{j_0} - (j - 1)$ and $|\gamma| = d_{j_0} - j$.

If $f_\gamma = m_r^{-1} \sum_{\delta \in S_\gamma} f_\delta$ is of type I then since there exists unique $\delta \in S_\gamma$ with $t \in \text{supp } f_\delta$, from the inductive hypothesis we obtain

$$|f_\gamma(e_t)| \leq m_r^{-1} \frac{c_1}{2^{j-1}} \leq \frac{c_1}{m_{r-1} 2^j}.$$

If $f_\gamma = \sum_{\delta \in S_\gamma} \lambda_\delta f_\delta$ is of type II then each f_δ is of type I. This follows from our assumption that every δ with $|\delta| \leq d_{j_0}$ is not a maximal node of the tree \mathcal{T} .

Hence setting $A_t = \{\delta : t \in \text{supp } f_\delta\}$ we get

$$|f_\gamma(e_t)| \leq \sum_{\delta \in A_t} |\lambda_\delta f_\delta(e_t)| \leq \sum_i \frac{1}{m_{i-1}} \frac{c_1}{2^{j-1}} \leq \frac{c_1}{2^j}.$$

□

By the above sublemma we may assume

$$\text{supp}(f) = \cup \{\text{supp}(f_\alpha) : \alpha \text{ maximal node with } |\alpha| \leq d_{j_0}\}.$$

Sublemma 8.7. *Let $t \in \text{supp}(f)$ be such that for every maximal node $\alpha \in \mathcal{T}$ with $t \in \text{supp}(f_\alpha)$, we have $|\alpha| \leq d_{j_0}$ and $\alpha \in S_\beta$ with f_β is of type II and $|\lambda_\alpha| \leq m_{j_0}^{-1}$. Then*

$$|f(e_t)| < \frac{3}{2m_{j_0}}$$

Proof. We prove by induction on $j = 0, \dots, d_{j_0}$ that if $|\gamma| = d_{j_0} - j$ then

$$|f_\gamma(e_t)| < \begin{cases} 3/2m_{j_0}, & \text{if } f_\gamma \text{ is of type II} \\ 1/2^2 m_{j_0} m_{r-1} & \text{if } f_\gamma \text{ is of type I and } w(f_\gamma) = m_r. \end{cases}$$

For the predecessors of maximal nodes we have it by assumption. If f_γ is of type I, then only one of its successors have e_t in its support and by the inductive hypothesis

$$|f_\gamma(e_t)| < \frac{3}{2m_{j_0} m_r} \leq \frac{1}{2^2 m_{j_0} m_{r-1}}$$

If f_γ is of type II and $|f_\gamma| = d_{j_0} - j$, let $f_\gamma = \sum_{\beta \in S_\gamma} \lambda_\beta f_\beta$. Set $A_t = \{\beta \in S_\gamma : t \in \text{supp}(f_\beta)\}$. There exists at most one $f_{\beta_0} = e_t^*$ and all other f_β are of type I with different weights. The inductive hypothesis and Hölder inequality yield

$$|f_\gamma(e_t)| \leq \sum_{\beta \in A_t, \beta \neq \beta_0} |\lambda_\beta| \frac{1}{2^2 m_{j_0} m_{\beta-1}} + |\lambda_{\beta_0}| e_t^*(e_t) < \frac{1}{2m_{j_0}} + \frac{1}{m_{j_0}} \leq \frac{3}{2m_{j_0}}$$

and the induction is finished. □

The above sublemmas for the tree analysis $(f_\alpha)_{\alpha \in \mathcal{T}}$ of f with all maximal nodes belonging to M_2 yields that

$$(26) \quad \text{supp}(f) = \cup \{\text{supp}(f_\alpha) : \alpha \text{ is a maximal node of } \mathcal{T} \text{ with } |\alpha| \leq d_{j_0} \text{ and} \\ \text{if } \alpha \in S_\beta, f_\beta \text{ is of type II then } |\lambda_\alpha| \geq m_{j_0}^{-1}\}$$

Sublemma 8.8. *Let $f \in W'_{j_0}$ with an tree analysis $(f_\alpha)_{\alpha \in \mathcal{T}}$ such that*

- 1) *the height of \mathcal{T} is less or equal to d_{j_0} ,*
- 2) *if f_α is of type I then $w(f_\alpha) < m_{j_0}$ holds,*
- 3) *if α is a maximal node, $\alpha \in S_\beta$, f_β is of type II then $|\lambda_\alpha| \geq m_{j_0}^{-1}$.*

Then

$$\# \text{supp}(f) \leq (2n_{j_0-1})^{d_{j_0}+1}.$$

Proof. Inductively we show that for every α with $|\alpha| = d_{j_0} - j$, $j = 0, 1, \dots, d_{j_0}$ holds the following

$$(27) \quad \#\text{supp}(f_\alpha) \leq (2n_{j_0-1})^{j+1}.$$

If f_α is a terminal node it is clear that (27) holds.

Assume that the result holds for all α of order $d_{j_0}, \dots, d_{j_0} - (j-1)$ and let $|\alpha| = d_{j_0} - j$.

If $f_\alpha = w(f_\alpha)^{-1} \sum_{\beta \in S_\alpha} f_\beta$ is of type I, then we get $w(f_\alpha) \leq m_{j_0-1}$ and hence $\#S_\alpha \leq 2n_{j_0-1}$. Using the inductive hypothesis we obtain

$$\#\text{supp}(f_\alpha) \leq 2n_{j_0-1} \max\{\#\text{supp}(f_\beta) : \beta \in S_\alpha\} \leq (2n_{j_0-1})^{j+1}.$$

If $f_\alpha = \sum_{\beta \in S_\alpha} \lambda_\beta f_\beta$, setting $S_\alpha^1 = \{\beta \in S_\alpha : \beta \text{ maximal}\}$ from assumption 3) we get $|\lambda_\beta| \geq m_{j_0}^{-1}$ for all $\beta \in S_\alpha^1$ and therefore $\#S_\alpha^1 \leq m_{j_0}^2$.

Note also that for every $\beta \in S_\alpha \setminus S_\alpha^1$ it holds that $w(f_\beta) < m_{j_0}$ and hence $\#(S_\alpha \setminus S_\alpha^1) \leq j_0 - 1$. Hence by the inductive hypothesis we get

$$\begin{aligned} \#\text{supp}(f_\alpha) &\leq (j_0 - 1) \max\{\#\text{supp}(f_\beta) : \beta \in S_\alpha \setminus S_\alpha^1\} + m_{j_0}^2 \\ &\leq (j_0 - 1)(2n_{j_0-1})^j + m_{j_0}^2 \leq (2n_{j_0-1})^{j+1}. \end{aligned}$$

□

Proof of Lemma 8.3 completed. The desired inequality follows from (26) and Sublemma 8.8. □

Remark 8.9. Take $f \in W'_{j_0}$ such that for every $t \in \text{supp}(f)$, $|f(e_t)| > 3m_{j_0}^{-2}$ and every f_α in the tree analysis $(f_\alpha)_{\alpha \in \mathcal{T}}$ of f satisfies $w(f_\alpha) \neq m_{j_0}$. Then Remark 8.5, Sublemma 8.6 and 8.8 yields that $\#\text{supp}(f) \leq (2n_{j_0-1})^{s_{j_0-1}-2}$.

Lemma 8.10. Let $f \in W'_{j_0}$ be of type I. Then

$$(28) \quad \left| f\left(\frac{1}{n_{j_0}} \sum_{i=1}^{n_{j_0}} e_t\right) \right| \leq \begin{cases} \frac{4}{w(f)m_{j_0}}, & \text{if } w(f) < m_{j_0} \\ \frac{e_1}{w(f)}, & \text{if } w(f) \geq m_{j_0}. \end{cases}$$

If moreover we assume that there exists a tree analysis $(f_\alpha)_{\alpha \in \mathcal{T}}$ of f , such that $w(f_\alpha) \neq m_{j_0}$ for every $\alpha \in \mathcal{T}$, we have that

$$(29) \quad \left| f\left(\frac{1}{n_{j_0}} \sum_{i=1}^{n_{j_0}} e_{k_i}\right) \right| \leq \begin{cases} \frac{4}{m_i m_{j_0}^2} & \text{if } w(f) = m_i < m_{j_0} \\ \frac{e_1}{m_i} & \text{if } w(f) = m_i > m_{j_0} \end{cases}$$

Proof. If $w(f) \geq m_{j_0}$ Lemma 8.2 yields the result.

Let $f = w(f)^{-1} \sum_{i=1}^d f_i$ with $w(f) < m_{j_0}$. We set

$$A_1 = \{t : t \in \text{supp}(f_i) \text{ for some } i \leq d \text{ and } f_i(e_t) \leq 3m_{j_0}^{-1}\}.$$

It readily follows that

$$(30) \quad |f(e_t)| \leq \frac{3}{w(f)m_{j_0}} \text{ for every } t \in A_1.$$

We set $B_i = \text{supp}(f_i) \setminus A_1$. Lemma 8.3 yields that

$$\#\text{supp}(B_i) \leq (2n_{j_0-1})^{s_{j_0-1}-2}.$$

It follows that $\#\{t : t \notin A_1\} \leq 2n_{j_0-1}(2n_{j_0-1})^{s_{j_0-1}-2} = (2n_{j_0-1})^{s_{j_0-1}-1}$ and therefore

$$(31) \quad \left| f\left(\sum_{t \notin A_1} \frac{e_t}{n_{j_0}}\right) \right| \leq \frac{(2n_{j_0-1})^{s_{j_0-1}-1}}{n_{j_0}} < m_{j_0}^{-2}.$$

From (30) and (31) we get the result.

For the proof of (29) we consider the set

$$A_1 = \{t : t \in \text{supp}(f_i) \text{ for some } i \leq d \text{ and } f_i(t) \leq 3m_{j_0}^{-2}\}.$$

and using Remark 8.9 we obtain again the relation (31) for the new set A_1 . \square

9. THE BASIC INEQUALITY AND ITS CONSEQUENCES

In this section we introduce the notion of rapidly increasing sequences (RIS) and prove the Basic Inequality (Prop. 9.3), which by results from the previous section provides estimation on the norm of suitable averages of RIS (Prop. 9.4). In particular we obtain reflexivity of X_ξ .

Definition 9.1. *Let $\epsilon > 0$. A block sequence (x_k) in X_ξ is said to be a (C, ϵ) -rapidly increasing sequence (RIS), if there exists a strictly increasing sequence (j_k) of positive integers such that*

- a) $\|x_k\| \leq 1$ for all k .
- b) $m_{2j_1}^{-1/2} < \epsilon$ and $\#(\text{range}(x_k))m_{2j_{k+1}}^{-1/2} < \epsilon$ for all $k \geq 1$
- c) For every $k = 1, 2, \dots$ and every $f \in K_\xi$ of type I with $w(f) < m_{2j_k}$ we have that $|f(x_k)| \leq \frac{C}{w(f)}$.

Remark 9.2. *From Proposition 7.8 and Lemma 7.7 we get that for every $\epsilon > 0$ any block subspace contains a $(3, \epsilon)$ -RIS $(x_k)_k$ where x_k is a normalized $2 - \ell_1^{m_{j_k}}$ average with $(2j_k)_k$ satisfying condition b) of the above definition.*

Proposition 9.3 (Basic Inequality). *Let $\epsilon > 0$, $1 < j_0 \in \mathbb{N}$, $(x_k)_{k \in \mathbb{N}}$ be a (C, ϵ) -RIS in X_ξ with the associated sequence $(j_k)_k$. Assume that for every $g \in G_\xi$ the set $I_g = \{k : |g(x_k)| \geq \epsilon\}$ has cardinality at most n_{j_0-1} .*

Let $(c_k)_k$ be a sequence of scalars. Then for every $f \in K_\xi$ and every interval I there exists a functional $g \in W_{j_0}$ such that

$$(32) \quad |f(\sum_{k \in I} c_k x_k)| \leq C(g(\sum_{k \in I} |c_k| e_k) + \epsilon_f \sum_{k \in I} |c_k|), \quad \epsilon_f \leq \epsilon.$$

Moreover, if f is the result of an $(\mathcal{A}_{n_j}, m_j^{-1})$ -operation then $g = e_r^$ or 0 or g is the result of an $(\mathcal{A}_{2n_j}, m_j^{-1})$ operation and $\epsilon_f \leq \epsilon w(f)^{-1/2}$.*

If we additionally assume that for every $f \in K_\xi$ with $w(f) = m_{j_0}$, for every interval J it holds that

$$(33) \quad |f(\sum_{k \in J} c_k x_k)| \leq C(\max_{k \in J} |c_k f(x_k)| + \epsilon m_{j_0}^{-1/2} \sum_{k \in J} |c_k|),$$

then we may select the functional g to have a tree analysis $(g_\alpha)_\alpha$ with $w(g_\alpha) \neq m_{j_0}$ for all $\alpha \in \mathcal{A}$.

Proof. We shall treat the case that there exists j_0 satisfying (33). We also assume that $\text{range}(x_k) \cap \text{range}(f) \neq \emptyset$ for every $k \in I$. We proceed by induction of the order $o(f)$ of the functional f .

Let $o(f) = 1$. Then we have that $f \in G_\xi$ and from the assumptions the set $R_f = \{k \in I : |f(x_k)| \geq \epsilon\}$ has cardinality at most n_{j_0-1} . We set $g_f = \sum_{k \in R_f} e_k^*$ and $\epsilon_f = \epsilon$. It readily follows

$$|f(\sum_{k \in I} c_k x_k)| \leq g_f(\sum_{k \in I} |c_k| e_k) + \epsilon_f \sum_{k \in I} |c_k|.$$

Suppose now that the result holds for every functional in K_ξ with order less than q and consider $f \in K_\xi$ with $o(f) = q$.

We consider the following three cases.

Case 1. f is of type I and $w(f) = m_{j_0}$. We choose $k_0 \in I$ with $|c_{k_0}f(x_{k_0})| = \max_{k \in I} |c_k f(x_k)|$ and we set $g_f = |f(x_{k_0})|e_{k_0}^*$. Then from our assumption (33) it follows

$$\begin{aligned} |f(\sum_{k \in I} c_k x_k)| &\leq C \left(\max_{k \in I} |c_k f(x_k)| + \epsilon m_{j_0}^{-1/2} \sum_{k \in I} |c_k| \right) \\ &\leq C \left(g(\sum_{k \in I} |c_k| e_k) + \epsilon m_{j_0}^{-1/2} \sum_{k \in I} |c_k| \right). \end{aligned}$$

Case 2. f is of type I and $w(f) \neq m_{j_0}$.

Then $f = m_j^{-1} \sum_{i=1}^d f_i$ with $j \neq j_0$ and $d \leq n_j$. We consider the following three subcases.

Subcase 2a. $w(f) < m_{2j_k}$ for all $k \in I$.

For every $i \leq d$ we set

$$I_i = \{k \in I : \text{range}(x_k) \cap \text{range}(f_i) \neq \emptyset \text{ and } \text{range}(x_k) \cap \text{range}(f_{i'}) = \emptyset \text{ for all } i' \neq i\}.$$

We also set $I_0 = \{k \in I : \text{range}(x_k) \cap \text{range}(f_i) \neq \emptyset \text{ for at least two } i \in \{1, \dots, d\}\}$. We observe that $\#I_0 \leq d$. Condition c) in the definition of the RIS yields

$$(34) \quad |f(x_k)| \leq \frac{C}{w(f)} \text{ for every } k \in I_0.$$

For every $i \leq d$ we have that I_i is a subinterval of I , hence our inductive assumption yields that there exists $g_{f_i} \in W_{j_0}$ with $\text{supp}(g_{f_i}) \subset I_i$ such that

$$(35) \quad |f_i(\sum_{k \in I_i} c_k x_k)| \leq C(g_{f_i}(\sum_{k \in I_i} |c_k| e_k) + \epsilon_{f_i} \sum_{k \in I_i} |c_k|).$$

The family $\{I_1, \dots, I_d\} \cup \{\{k\} : k \in I_0\}$ consists of pairwise disjoint intervals and has cardinality less than or equal to $2d \leq 2n_j$. We set

$$g_f = \frac{1}{w(f)} \left(\sum_{i=1}^d g_{f_i} + \sum_{k \in I_0} e_k^* \right).$$

Then $g_f \in W_{j_0}$, $\text{supp } g_f \subset I$, while from (34), (35) we obtain

$$\begin{aligned} |f(\sum_{k \in I} c_k x_k)| &\leq \sum_{k \in I_0} |c_k| |f(x_k)| + \frac{1}{w(f)} \sum_{i=1}^d C \left(g_{f_i}(\sum_{k \in I_i} |c_k| e_k) + \epsilon \sum_{k \in I_i} |c_k| \right) \\ (36) \quad &\leq C \left(g_f(\sum_{k \in I} |c_k| e_k) + \epsilon_f \sum_{k \in I} |c_k| \right), \text{ where } \epsilon_f = \epsilon w(f)^{-1}. \end{aligned}$$

Subcase 2b. $m_{2j_{k_0}} \leq w(f) < m_{2j_{k_0+1}}$ for some $k_0 \in I$.

From condition b) in the definition of RIS we get

$$(37) \quad |f(x_k)| \leq \epsilon w(f)^{-1/2} = \epsilon_f \text{ for all } k \in I \text{ with } k < k_0.$$

Using that $w(f) \geq m_{2j_1}$ and condition c) we get

$$(38) \quad |f(x_k)| \leq C w(f)^{-1} \leq C \epsilon w(f)^{-1/2} = C \epsilon_f \text{ for every } k_0 < k \in I.$$

Thus setting $g_f = |f(x_{k_0})|e_{k_0}^*$ from (37),(38) we get

$$\begin{aligned} (39) \quad |f(\sum_{k \in I} c_k x_k)| &\leq |c_{k_0} f(x_{k_0})| + \sum_{k \in I \setminus \{k_0\}} |c_k f(x_k)| \\ &\leq |c_{k_0} f(x_{k_0})| + C \epsilon_f \sum_{k \in I \setminus \{k_0\}} |c_k| \leq C \left(g_f(\sum_{k \in I} |c_k| e_k) + \epsilon_f \sum_{k \in I} |c_k| \right). \end{aligned}$$

Subcase 2c. $m_{2j_{k+1}} \leq w(f)$ for all $k \in I$.

In this case as in (37), $|f(x_k)| \leq \epsilon w(f)^{-1/2} = \epsilon_f$ for all $k \in I$, and we set $g_f = 0$. It follows easily that (32) holds.

Case 3. f is of type II, i.e. $f = \sum_{i=1}^d \lambda_i f_i$.

For the given interval I the inductive assumption associates to each f_i a functional g_{f_i} satisfying (32).

We note that there exists at most one $i \in \{1, \dots, d\}$ with $w(f_i) = m_{j_0}$. Without loss of generality we assume that there does exist such an i , denoted by i_0 . From *Case 1* we have $g_{f_{i_0}} = |f(x_{k_0})|e_{k_0}^*$ for some $k_0 \in I$.

For every $1 \leq i \leq d$ we set $C_{f_i} = \{k \in I : \text{range}(f_i) \cap \text{range}(x_k) \neq \emptyset\}$.

We partition the set $M = \{1, \dots, d\} \setminus \{i_0\}$ as follows:

$$L_0 = \{i \in M : w(f_i) < m_{2j_k} \text{ for all } k \in C_{f_i}\}$$

and for $k = 1, \dots$, we set

$$L_k = \{i \in M : k \in C_{f_i} \text{ and } m_{2j_k} \leq w(f_i) < m_{2j_{k+1}}\}.$$

We enlarge L_{k_0} by adding i_0 to its elements. Note that if $i \notin L_0 \cup \cup_k L_k$ then $g_{f_i} = 0$. We set

$$g_f = \sum_{i=1}^d |\lambda_i| g_{f_i} = \sum_{i \in L_0} |\lambda_i| g_{f_i} + \sum_k \left(\sum_{i \in L_k} |\lambda_i f_i(x_k)| \right) e_k^*.$$

We show that $g_f \in W_{j_0}$. Since for every $i \in L_0$ it holds that $w(g_{f_i}) = w(f_i)$ we have that the functionals $g_{f_i}, i \in L_0$, have different weights. Also since the sets L_k are pairwise disjoint it follows

$$\sum_{i \in L_0} \lambda_i^2 + \sum_k \left(\sum_{i \in L_k} |\lambda_i f_i(x_k)| \right)^2 \leq \sum_{i \in L_0} \lambda_i^2 + \sum_k \left(\sum_{i \in L_k} \lambda_i^2 \|x_k\| \right) \leq \sum_{i=1}^d \lambda_i^2 \leq 1,$$

hence $g_f \in W_{j_0}$.

We show that (32) holds. First we observe that since for all i , $\epsilon_{f_i} \leq \epsilon w(f_i)^{-1/2}$ it follows that $\sum_{i \leq d} |\lambda_i| \epsilon_{f_i} \leq \epsilon$. Also,

$$\begin{aligned} |f(\sum_{k \in I} c_k x_k)| &\leq \sum_{i=1}^d |\lambda_i f_i(\sum_{k \in C_{f_i}} c_k x_k)| \leq \sum_{i=1}^d C |\lambda_i| (g_{f_i}(\sum_{k \in C_{f_i}} |c_k| e_k) + \epsilon_{f_i} \sum_{k \in C_{f_i}} |c_k|) \\ &\leq C \left(\sum_{i \in L_0} |\lambda_i| g_{f_i} + \sum_k \left(\sum_{i \in L_k} |\lambda_i f_i(x_k)| \right) e_k^* \right) \left(\sum_{k \in I} |c_k| e_k \right) + C \epsilon \sum_{k \in I} |c_k| \\ &= C g_f(\sum_{k \in I} |c_k| e_k) + C \epsilon \sum_{k \in I} |c_k|. \end{aligned}$$

Case 4. $f = \sum_{i=1}^d r_i f_i$, where $(r_i)_{i=1}^d \subset \mathbb{Q}$, is a rational convex combination.

As in the previous case for every $i = 1, \dots, d$ we set

$$I_i = \{k \in I : \text{range}(f_i) \cap \text{range}(x_k) \neq \emptyset\}.$$

Take suitable (g_{f_i}) by the inductive hypothesis. Setting $g_f = \sum_{i=1}^d r_i g_{f_i}$ we get

$$|f(\sum_{k \in I} c_k x_k)| \leq C (g_f(\sum_{k \in I} |c_k| e_k) + \epsilon \sum_{k \in I} |c_k|)$$

□

Proposition 9.4. *Let $\epsilon > 0$, $1 < j \in \mathbb{N}$, $(x_k)_{k \in \mathbb{N}}$ be a (C, ϵ) -RIS in X_ξ with the associated sequence $(j_k)_k$ and $j < j_1$. Assume that for every $g \in G_\xi$ the set $I_g = \{k : |g(x_k)| \geq 2m_j^{-2}\}$ has cardinality at most n_{j-1} . Then*

a) If $\varepsilon \leq \frac{2}{m_j^2}$ then for every $f \in K_\xi$ of type I

$$\left| f\left(\frac{1}{n_j} \sum_{k=1}^{n_j} x_k\right) \right| \leq \begin{cases} \frac{5C}{m_j w(f)}, & \text{if } w(f) < m_j \\ \frac{c_1 C}{w(f)} + \frac{2C}{m_j^2}, & \text{if } w(f) \geq m_j. \end{cases}$$

In particular $\left\| \frac{1}{n_j} \sum_{k=1}^{n_j} x_k \right\| \leq \frac{3C}{m_j}$.

b) If moreover for $j_0 = j$ the additional assumption of the Basic Inequality is fulfilled (Proposition 9.3 (33)), then for a linear combination $\frac{1}{n_j} \sum_{i=1}^{n_j} b_i x_i$, where $|b_i| \leq 1$, we have

$$\left\| \frac{1}{n_j} \sum_{i=1}^{n_j} b_i x_i \right\| \leq \frac{3C}{m_j^2}.$$

Proof. The proof is an application of the Basic Inequality and Lemma 8.10. \square

Proposition 9.5. *The space X_ξ is reflexive.*

The reflexivity of X_ξ is consequence of Proposition 9.4 following standard arguments, see [3],[8].

Remark 9.6. *The reflexivity of the space X_ξ yields that K_ξ is norm-dense subset of $B_{X_\xi^*}$. Indeed since K_ξ is norming set it follows that $\text{conv}(K_\xi)$ is w^* -dense. The reflexivity of X_ξ implies that $\text{conv}(K_\xi)$ is w -dense and hence $\|\cdot\|$ -dense. Since K_ξ is rationally convex we get that K_ξ is in fact norm-dense subset of $B_{X_\xi^*}$.*

10. EXACT PAIRS AND ATTRACTING SEQUENCES

Now we define exact pairs and show saturation of X_ξ by them. Next we introduce the crucial notion of attracting sequences and estimate the norms of averages of elements forming these sequences: vectors (Corollary 10.5) and functionals (Corollary 10.6).

Definition 10.1. *A pair (x, f) with $x \in X_\xi$ and $f \in K_\xi$ is said to be a $(C, 2j)$ -exact pair, $C \geq 1$, $j \in \mathbb{N}$ if the following conditions are satisfied*

- 1) $f(x) = 1$ and $\text{range } f = \text{range } x$.
- 2) f is of type I and $w(f) = m_{2j}$.
- 3) $1 \leq \|x\| \leq 3C$, $\|x\|_\infty \leq m_{2j}^{-2}$ and for every g of type I with $w(g) < m_{2j}$ it holds that $|g(x)| \leq \frac{5C}{w(g)}$ while for g of type I with $w(g) > m_{2j}$, $|g(x)| \leq 5Cm_{2j}^{-1}$.

Proposition 10.2. *Let $j \in \mathbb{N}$ and Y be a block subspace of X_ξ . There exists a $(3, 2j)$ -exact pair with $x \in Y$.*

Proof. Let Y be a block subspace of X_ξ and $j \in \mathbb{N}$. By Remark 9.2, Proposition 5.4 we can choose for $\epsilon \leq 2m_j^{-2}$ a $(3, \epsilon)$ -RIS $(x_k)_{k=1}^{n_{2j}}$ satisfying the assumptions of Proposition 9.4. It follows that

$$\left\| \frac{m_{2j}}{n_{2j}} \sum_{i=1}^{n_{2j}} x_i \right\| \leq 9.$$

Choosing $f_i \in K_\xi$ such that $f_i(x_i) = 1$ and $\text{range}(f_i) \subset \text{range}(x_i)$ we have that $f = m_{2j}^{-1} \sum_{i=1}^{n_{2j}} f_i \in K_\xi$ and $f\left(\frac{m_{2j}}{n_{2j}} \sum_{i=1}^{n_{2j}} x_i\right) = 1$. Setting $x = E \frac{m_{2j}}{n_{2j}} \sum_{i=1}^{n_{2j}} x_i$, where $E = \text{range}(f)$, Proposition 9.4 a) yields that (x, f) is a $(3, 2j)$ -exact pair. \square

Definition 10.3. *A double sequence $(x_k, f_k)_{k=1}^{n_{2j-1}}$ is called a $(C, 2j-1)$ attracting sequence, if there is a sequence $(j_k)_{k=1}^{n_{2j-1}}$ such that*

- (1) $(f_k)_{k=1}^{n_{2j-1}}$ is a $(2j-1)$ -attractor sequence with $w(f_{2k-1}) = m_{j_{2k-1}}$ and $f_{2k} = e_{l_{2k}}^*$ where $l_{2k} \in \Lambda_{j_{2k}}$ for all $k \leq n_{2j-1}/2$,
- (2) $x_{2k} = e_{l_{2k}}$ for all $k \leq n_{2j-1}/2$,

- (3) (x_{2k-1}, f_{2k-1}) is a (C, j_{2k-1}) exact pair.
(4) $j_k = 2\sigma(f_1, \dots, f_{k-1})$ for any $k \leq n_{2j-1}$

Remark 10.4. If $(x_k, f_k)_{k=1}^{n_{2j-1}}$ is a $(C, 2j-1)$ -attracting sequence, then $(x_k/(3C))_{k=1}^{n_{2j-1}}$ is a $(5/3, n_{2j-1})$ RIS. Indeed,

$$\#(\text{range } x_k) \frac{1}{m_{j_{k+1}}^{1/2}} = \#(\text{range } f_k) \frac{1}{m_{2\sigma(f_1, \dots, f_k)}^{1/2}} < \min\{\|f_i\|_\infty, i \leq k\} \leq m_{j_1}^{-1/2} \leq n_{2j-1}^{-1}$$

by the condition on σ . Condition (c) in definition of RIS is satisfied thanks to $x_{2k} = e_{l_{2k}}$ and the fact that (x_{2k-1}, f_{2k-1}) is a (C, j_{2k-1}) exact pair.

Corollary 10.5. Let $(x_k, f_k)_{k=1}^{n_{2j-1}}$ be a $(C, 2j-1)$ attracting sequence with associated sequence $(j_k)_k$, with $\|x_{2k-1}\|_{G_\xi} \leq m_{2j-1}^{-2}$ for every $k \leq n_{2j-1}/2$. Then

$$(40) \quad \frac{1}{2m_{2j-1}^2} \leq \left\| \frac{1}{n_{2j-1}} \sum_{k=1}^{n_{2j-1}} (-1)^{k+1} x_k \right\| \leq \frac{15C}{m_{2j-1}^2}$$

Proof. To see the lower estimate in (40) note that $\tilde{f} = m_{2j-1}^{-2} \sum_{k=1}^{n_{2j-1}/2} f_{2k} \in G_\xi \subset K_\xi$. Hence

$$\left\| \frac{1}{n_{2j-1}} \sum_{k=1}^{n_{2j-1}} (-1)^{k+1} x_k \right\| \geq |\tilde{f}(\frac{1}{n_{2j-1}} \sum_{k=1}^{n_{2j-1}} (-1)^{k+1} x_k)| = m_{2j-1}^{-2}/2.$$

The upper estimation in (40) follows from Proposition 9.4 b) for $b_k = (-1)^k$ and $j_0 = 2j-1$ after we show that

- K1) for every $g \in G_\xi$ it holds that $\#\{k : |g(x_k)| \geq 2m_{2j-1}^{-2}\} \leq n_{2j-2}$
K2) The additional property of the Basic Inequality holds for the sequence $(x_k/(3C))_{k=1}^{n_{2j-1}}$ with constant $5/3$.

To see K1) note that from the assumption we have that $\|x_{2k-1}\|_{G_\xi} \leq m_{2j-1}^{-2}$. Take now any $g \in G_1^r = \{m_{2r-1}^{-2} \sum_{i \in F} \pm e_i^* : F \in \mathcal{A}_{n_{2r-1}}\}$. If $r \geq j$ then $|g(e_{l_{2k}})| \leq m_{2j-1}^{-2}$ while if $r < j$ it holds that $\#\{l_{2k} : |g(e_{l_{2k}})| \geq m_{2j-1}^{-2}\} \leq n_{2r-1}$.

From the definition of the G_ξ -special functionals $g = E \sum_{r=1}^d g_r$ we obtain

$$|g(e_{l_{2k}})| \geq 2m_{2j-1}^{-2} \Rightarrow l_{2k} \in \text{supp } g_r \text{ with } g_r \in \cup_{i \leq j-1} G_1^i$$

and therefore

$$\#\{l_{2k} : |g(e_{l_{2k}})| \geq 2m_{2j-1}^{-2}\} \leq \sum_{r < j} n_{2r-1} \leq n_{2j-2}.$$

Finally let $y^* = \sum_{k=1}^d a_k y_k^* \in G_{\ell_2}$. For every $k = 1, \dots, d$ let $y_k^* = y_{k,1}^* + y_{k,2}^*$ with $\text{ind}(y_{k,1}^*) \subset \{1, \dots, j\}$ and $\text{ind}(y_{k,2}^*) \subset \{j+1, j+2, \dots\}$. So we may write $y^* = \sum_{k=1}^d a_k y_{k,1}^* + \sum_{k=1}^d a_k y_{k,2}^*$.

Since $\sum_{i \geq j} m_{2i-1}^{-2} < \frac{3}{2m_{2j-1}^2}$ and the sets $\text{ind}(y_k^*)_k$ are pairwise disjoint in order $|y^*(e_{l_{2i}})| \geq 2/m_{2j-1}^2$ it must hold that $l_{2i} \in \text{supp } y_{k,1}^*$ for some $k \leq d$. As in the previous case we get

$$\#\{l_{2i} : |y^*(e_{l_{2i}})| \geq 2m_{2j-1}^{-2}\} \leq \sum_{r < j} n_{2r-1} \leq n_{2j-2}.$$

To see K2) we have to show that for any $(2j-1)$ -attractor functional g and for any interval $J \subset \{1, \dots, n_{2j-1}\}$ we have

$$|g(\sum_{k \in J} (-1)^k x_k)| \leq 5C(\max |g(x_k)| + m_{2j-1}^{-4} \#J).$$

Let $g = m_{2j-1}^{-1} \sum_{i=1}^d g_i$, $d \leq n_{2j-1}$. If $g = f = m_{2j-1}^{-1} \sum_{i=1}^{n_{2j-1}} f_i$ then $|g(\sum_k (-1)^k x_k)| = 0$. Otherwise let $i_0 = \min\{i \leq d : f_i \neq g_i\}$. Then, by definition of the attracting sequence and since J is an interval, we have

$$(41) \quad \left| \sum_{i=1}^{i_0-1} g_i \left(\sum_{k \in J} (-1)^k x_k \right) \right| \leq 3 \max_{i \leq i_0-1} |g_i(x_i)|.$$

By Remark 7.2a) for any $2i-1 > i_0, 2k-1 > i_0$ we have $w(g_{2i-1}) \neq w(f_{2k-1})$ and for any $2i > i_0, 2k > i_0$ we have $g_{2i} \neq f_{2k}$. Notice $g_{2i}(x_{2k}) = 0$ for any $2i > i_0$ and any $2k \in J$. By the definition of the attractor sequence we get for any $2k > i_0$

$$(42) \quad \left| \sum_{i \geq i_0} g_i(x_{2k}) \right| \leq \max_i \|g_{2i-1}\|_\infty \leq m_{j_1}^{-1} \leq m_{2j-1}^{-4}.$$

Now by the definition of the exact sequence for any $2i-1 \geq i_0$ and any $2k-1 > i_0$ we have

$$|g_{2i-1}(x_{2k-1})| \leq 5C \max\{w(g_{2i-1})^{-1}, m_{j_{2k-1}}^{-1}\} \leq 5C n_{2j-1}^{-2}.$$

Hence using that $\|x_{2k-1}\|_\infty \leq m_{j_{2k-1}}^{-2}$ we obtain

$$(43) \quad \left| \sum_{i \geq i_0} g_i(x_{2k-1}) \right| \leq 5C n_{2j-1} (n_{2j-1}^{-2} + \|x_{2k-1}\|_\infty) \leq 5C m_{2j-1}^{-4}.$$

From (41), (42), (43) we get

$$\begin{aligned} |g(\sum_{k \in J} (-1)^k x_k)| &= \left| \frac{1}{m_{2j-1}} \left(\sum_{i=1}^{i_0-1} g_i + \sum_{i \geq i_0} g_i \right) \left(\sum_{k \in J} (-1)^k x_k \right) \right| \\ &\leq 3 \max_{i < i_0} |g(x_i)| + |g(x_{i_0})| + 5C \sum_{k \in J} m_{2j-1}^{-4} \leq 5C (\max_{k \in J} |g(x_k)| + \frac{\#J}{m_{2j-1}^4}) \end{aligned}$$

which ends the proof of K2) and thus the whole proof. \square

Corollary 10.6. *Let $(x_i, f_i)_{i=1}^{n_{2j-1}}$ be a $(C, 2j-1)$ -attracting sequence of length n_{2j-1} satisfying the assumption of Corollary 10.5. Set*

$$\phi = m_{2j-1}^{-2} \sum_{i=1}^{n_{2j-1}/2} f_{2i-1}, \quad \psi = m_{2j-1}^{-2} \sum_{i=1}^{n_{2j-1}/2} f_{2i}$$

Then

$$\frac{1}{30C} \leq \|\psi\| \leq 1, \quad \|\phi + \psi\| \leq m_{2j-1}^{-1}.$$

Proof. Notice that $m_{2j-1}(\phi + \psi) \in K_\xi$ hence the second inequality holds. To prove the first, from Corollary 10.5 we have $\|\frac{1}{n_{2j-1}} \sum_{k=1}^{n_{2j-1}} (-1)^{k+1} x_k\| \leq \frac{15C}{m_{2j-1}^2}$ and therefore

$$\|\psi\| \geq \psi \left(\frac{m_{2j-1}^2}{15C n_{2j-1}} \sum_{i=1}^{n_{2j-1}} (-1)^{i+1} x_i \right) = \frac{m_{2j-1}^2}{15C n_{2j-1}} \frac{1}{m_{2j-1}^2} \sum_{i=1}^{n_{2j-1}/2} f_{2i}(x_{2i}) = \frac{1}{30C}.$$

\square

11. SPACES WITH NO ℓ_p AS A SPREADING MODEL

In this section we show that the space X_ξ does not admit c_0 or $\ell_p, 1 \leq p < \infty$, as a spreading model. Actually we show that this holds for a wider class of Banach spaces which describe now.

Let G be a ground set. Let \mathcal{W}_G denote the smallest subset of $c_{00}(\mathbb{N})$ which

- (1) is symmetric, closed under the projections of its elements on intervals of \mathbb{N} and $G \subset \mathcal{W}_G$.

- (2) for every $j \in \mathbb{N}$ is closed under the (A_{n_j}, m_j^{-1}) operation.
- (3) whenever $(f_i)_{i=1}^d$ is the result of an $(A_{n_{j_k}}, m_{j_k}^{-1})$ -operation with $n_{j_k} \neq n_{j_m}$ for $k \neq m$, then $\sum_{i=1}^d \lambda_i f_i \in \mathcal{W}_G$ for all $(\lambda_i)_{i=1}^d \in B_{\ell_2} \cap [\mathbb{Q}]^{<\infty}$.
- (4) is rationally convex.

Definition 11.1. A subset D_G of \mathcal{W}_G is said to be an extension of G if:

- (i) The set D_G is symmetric, closed under the projections of its elements on intervals of \mathbb{N} and $G \subset D_G$.
- (ii) For any $j \in \mathbb{N}$ we have that D_G is closed under the $(A_{n_{2j}}, m_{2j}^{-1})$ -operation.
- (iii) Whenever $(f_i)_{i=1}^d$ is the result of an $(A_{n_{j_k}}, m_{j_k}^{-1})$ -operation with $n_{j_k} \neq n_{j_m}$ for $k \neq m$, then $\sum_{i=1}^d \lambda_i f_i \in D_G$ for all $(\lambda_i)_{i=1}^d \in B_{\ell_2} \cap [\mathbb{Q}]^{<\infty}$.
- iv) It is rationally convex.

Definition 11.2. Let D_G be an extension subset of \mathcal{W}_G . We define $\mathcal{Y}_{D_G} = \overline{(c_{00}(\mathbb{N}), \|\cdot\|_{D_G})}$.

We prove now the following theorem

Theorem 11.3. Let G be a ground set such that the corresponding space X_G does not admit ℓ_1 as a spreading model. Then for every extension D_G the space \mathcal{Y}_{D_G} does not admit any ℓ_p or c_0 as a spreading model.

Remark 11.4. The first example of a Banach space X with no ℓ_p as a spreading model was given by E. Odell and Th. Schlumprecht [24]. The spaces we consider in Theorem 11.3 are extensions of their example. In particular the space \mathcal{Y}_{D_G} is similar to their example when $G = \{\pm e_n^* : n \in \mathbb{N}\}$ and $D_G = \mathcal{W}_G$. Our proof provides also an alternative proof of their result.

Proof of the theorem. First we note that Proposition 7.8 holds for the space \mathcal{Y}_{D_G} . It follows that only ℓ_1 is finitely block representable in \mathcal{Y}_{D_G} and hence \mathcal{Y}_{D_G} does not admit c_0 or ℓ_p , $p > 1$, as a spreading model. We prove now that the space \mathcal{Y}_{D_G} does not contain a normalized sequence generating an ℓ_1 -spreading model.

Since the space X_G does not admit ℓ_1 as a spreading model Erdos-Magidor theorem [12], yields that for every bounded sequence $(x_n)_{n \in \mathbb{N}}$ and every $\epsilon > 0$ we can choose a block sequence $(y_n)_{n \in \mathbb{N}}$ of $(x_n)_{n \in \mathbb{N}}$ where each $y_n = \sum_{k \in F_n} x_k / n_0$, $\#F_n = n_0$, such that $\|y_n\|_G < \epsilon$.

Also by the Fact 6.5 if a sequence $(x_n)_{n \in \mathbb{N}}$ generates an ℓ_1 -spreading model with constant c then for every $\epsilon > 0$ there exists a block sequence $(y_n)_{n \in \mathbb{N}}$ of $(x_n)_{n \in \mathbb{N}}$ generating an ℓ_1 -spreading model with constant $(1 - \epsilon)^{-1}$. So assuming that a normalized block sequence (y_n) generates an ℓ_1 -spreading model with constant C , passing to suitable block sequence $(z_n)_{n \in \mathbb{N}}$ of $(y_n)_{n \in \mathbb{N}}$ we may assume that

- A) $(z_n)_{n \in \mathbb{N}}$ generates an ℓ_1 -spreading model with constant $(1 - \epsilon)^{-1}$.
- B) $\|z_n\|_G < \epsilon$ for every $n \in \mathbb{N}$.

We shall need also the following lemmas

Lemma 11.5. Let $x \in \mathcal{Y}_{D_G}$. Then for every $\epsilon > 0$ there exists $j_0 \in \mathbb{N}$ such that for every $\phi \in D_G$ of type II with $\text{ind}(\phi) > j_0$ it holds that $|\phi(x)| < \epsilon$.

The proof is similar to the proof of Lemma 6.1 and we omit it.

Lemma 11.6. Assume that $(z_n)_{n \in \mathbb{N}}$ satisfies A) and B) for $\epsilon = 10^{-3}$. Then there exists $j_0 \in \mathbb{N}$ such that for all $n \in \mathbb{N}$ there exists ϕ of type II with $\text{ind}(\phi) \leq j_0$ such that $|\phi(z_n)| \geq 0.9$.

Proof. Let $\phi \in D_G$ be such that $\phi(z_2 + z_n) \geq 1.998$. It follows that $|\phi(z_2)| \geq 0.998$ and $|\phi(z_n)| \geq 0.998$.

By B) we get that $\phi = \sum_{i=1}^d \lambda_i f_i$, where each f_i is a weighted functional i.e. is a result of an $(A_{n_{j_i}}, m_{j_i}^{-1})$ -operation and weights of $(f_i)_{i=1}^d$ are different. Let $j_0 \in \mathbb{N}$ be the

number we obtain from Lemma 11.5 for z_2 . Setting $A = \{i \leq d : w(f_i) \leq m_{j_0}\}$ and B its complement we get $|\sum_{i \in B} \lambda_i f_i(z_2)| < 0.001$. Therefore

$$\left(\sum_{i \in A} \lambda_i^2\right)^{1/2} \geq \left|\sum_{i \in A} \lambda_i f_i(z_2)\right| \geq 0.997.$$

It follows that

$$\left(\sum_{i \in B} \lambda_i^2\right)^{1/2} \leq (1 - 0.997^2)^{1/2} \Rightarrow \left\|\sum_{i \in B} \lambda_i f_i\right\| \leq 0.09$$

Hence $|\sum_{i \in A} \lambda_i f_i(z_n)| \geq 0.9$. □

By the above lemma we get that

(44) for every ϕ of type II with $\text{ind}(\phi) > j_0$ it holds that $|\phi(z_n)| < 0.6$ for all $n \in \mathbb{N}$.

Indeed, assume that there exists ϕ_2 of type II with $\text{ind}(\phi_2) > j_0$ and $|\phi_2(z_n)| \geq 0.6$. Then by the previous lemma we get ϕ_1 of type II with $\text{ind}(\phi_1) \leq j_0$ such that $\phi_1(z_n) \geq 0.9$. It follows that $\phi = \frac{1}{\sqrt{2}}(\phi_1 + \phi_2) \in D_G$ and hence $\|z_n\| \geq \phi(z_n) \geq \frac{0.9+0.6}{\sqrt{2}} > 1$, a contradiction.

Consider now the vector $u = \frac{1}{n_{j_0+1}} \sum_{i=1}^{n_{j_0+1}} z_{n_{j_0+1}+i}$. Since we have that $(z_n)_{n \in \mathbb{N}}$ generates an ℓ_1 -spreading model with constant 0.999^{-1} , there exists $\phi \in D_G$ such that $\phi(u) \geq 0.999$. Since $\|z_n\|_G \leq 10^{-3}$ for all $n \in \mathbb{N}$ we get that $\phi = \sum_{i=1}^d \lambda_i f_i$ where each f_i is a weighted functional and their weights are different. Set

$$R_1 = \{i : w(f_i) \leq m_{j_0}\} \text{ and } R_2 = \{i : w(f_i) > m_{j_0}\}.$$

By (44) for every n we obtain $|\sum_{i \in R_2} \lambda_i f_i(z_n)| \leq 0.6$ and hence

$$(45) \quad \left|\sum_{i \in R_2} \lambda_i f_i(u)\right| \leq 0.6.$$

On the other hand if $i \in R_1$ by Lemma 7.7 we get $|f_i(u)| \leq \frac{2}{w(f_i)}$.

Combining the above inequality with (45) we get

$$0.999 \leq |\phi(u)| \leq \sum_{i \in R_1} \frac{2|\lambda_i|}{w(f_i)} + 0.6 < 0.8,$$

a contradiction. □

Corollary 11.7. *The space X_ξ does not admit any ℓ_p (or c_0) as a spreading model.*

The abstraction of the properties of the set D_G enable us to derive also the following

Theorem 11.8. *There exists a reflexive Hereditarily Indecomposable Banach space \mathfrak{X}_{HI} with no ℓ_p , $1 \leq p < \infty$, or c_0 as a spreading model.*

Sketch of the proof. First we shall define the norming set K_{HI} of the space \mathfrak{X}_{HI} . The set K_{HI} is the smallest subset of $c_{00}(\mathbb{N})$ satisfying the following conditions

- (1) K_{HI} is symmetric i.e. if $f \in K_{HI}$ then $-f \in K_{HI}$, it is closed under the restriction of its elements to intervals of \mathbb{N} and $\{e_n^* : n \in \mathbb{N}\} \subset K_{HI}$.
- (2) K_{HI} is closed under $(\mathcal{A}_{n_{2j}}, m_{2j}^{-1})$ -operations.
- (3) K_{HI} is closed under $(\mathcal{A}_{n_{2j-1}}, m_{2j-1}^{-1})$ -operations on special sequences.
- (4) K_{HI} is closed under the operation $\sum_{i \in A} \lambda_i f_i$ whenever
 - (a) f_i is the result of an $(\mathcal{A}_{n_{j_i}}, m_{j_i}^{-1})$ -operation and $n_{j_i} \neq n_{j_k}$ for every $i \neq k \in A$
 - (b) $(\lambda_i)_{i \in A} \in B_{\ell_2} \cap [\mathbb{Q}]^{<\infty}$.
- (5) K_{HI} is rationally convex.

To complete the definition of K_{HI} we must define the special sequences. An n_{2j-1} -special sequence $(f_i)_{i=1}^{n_{2j-1}}$ is defined as the n_{2j-1} -attractor sequence (Def. 7.1) with the exception that the functionals f_{2i} are the results of an $(\mathcal{A}_{n_{2\sigma}(f_1, \dots, f_{2i-1})}, m_{2\sigma}^{-1}(f_1, \dots, f_{2i-1}))$ -operation of functionals of K_{HI} instead of $(e_{\lambda_{2i}}^*)$ we use in the attractor sequence. Observe that as a ground set we take the set $\{e_n^* : n \in \mathbb{N}\}$. The space \mathfrak{X}_{HI} is the completion of $(c_{00}(\mathbb{N}), \|\cdot\|_{K_{HI}})$. In order to show that \mathfrak{X}_{HI} is HI space we follow the method initiated in [16] and extended in [8],[3],[7]. Namely first we observe that Proposition 7.8 holds for the space \mathfrak{X}_{HI} hence there exist seminormalized ℓ_1 averages in every block subspace. Next we consider rapidly increasing sequence (RIS) of ℓ_1 averages, Def. 9.1 and we observe that Basic Inequality, Prop.9.3, holds also for the space \mathfrak{X}_{HI} . It follows that the estimations of Prop. 9.4 holds for RIS in the space \mathfrak{X}_{HI} . This enable us to consider the exact pairs, Def. 10.1. and also the dependent sequences. A (C, n_{2j-1}) -dependent sequence $(x_k, f_k)_{k=1}^{n_{2j-1}}$, $j \in \mathbb{N}$, is defined as the attractor sequence, Def. 10.3, with the exception that for all $k \leq n_{2j-1}$, (x_k, f_k) is an exact pair. From the Basic Inequality and the estimations on RIS we obtain that estimations of Corollary 10.5 also holds for the dependent sequences. The estimations in Corollary 10.5 easily yields that \mathfrak{X}_{HI} is indeed HI space.

Since the ground set of the space X_{HI} is the set $G = \{\pm e_n^* : n \in \mathbb{N}\}$, Theorem 11.3 yields that the space \mathfrak{X}_{HI} does not admit any ℓ_p or c_0 as a spreading model. \square

12. THE c_0 -INDEX OF THE DUAL OF X_ξ

We show now that every subspace of the dual space X_ξ^* has c_0 -index greater than ω^ξ and does not contain a sequence generating a c_0 -spreading model. As we have observe in Remark 9.6 the set K_ξ is a norm-dense subset of $B_{X_\xi^*}$. So proving that every block subspace generated by a block sequence of elements of K_ξ has c_0 -index greater than ω^ξ we get that the same holds for all block subspaces of X_ξ^* . Hence in the sequel we will assume that the block subspaces are generated by block sequences of K_ξ .

We shall need the dual result to Proposition 7.8.

Proposition 12.1. *For every $j \in \mathbb{N}$ every block subspace Y of X_ξ^* contains a $2 - c_0^{n_{2j}}$ average i.e. there exists a block sequence $(x_i^*)_{i=1}^{n_{2j}}$ in Y such that $\|x_i^*\| \geq 2^{-1}$ for every $i \leq n_{2j}$ and $2^{-1} \leq \|\sum_{i=1}^{n_{2j}} x_i^*\| \leq 1$.*

For the proof see Lemma 5.4 in [8].

Proposition 12.2. *a) The c_0 -index of every subspace Y of X_ξ^* is greater than ω^ξ .
b) The dual space X_ξ^* of X_ξ does not contain a normalized basic sequence generating a c_0 -spreading model.*

Proof. a) It is enough to prove the result for the block subspaces. Let Y be a block subspace of X_ξ^* . By Proposition 12.1 for every $j \in \mathbb{N}$ we choose x_i^* , $i \in \mathbb{N}$ such that x_i^* is a $2 - c_0^{n_{2j}}$ average. Let $x_i^* = \sum_{t=1}^{n_{2j}} x_{i,t}^*$ and $x_i = \sum_{t=1}^{n_{2j}} \frac{x_{i,t}}{n_{2j}}$ where $x_{i,t}^*(x_{i,t}) \geq 2^{-1}$ and $\|x_{i,t}\| = 1$.

Let $1 < j \in \mathbb{N}$. Passing to a subsequence we may assume that $(x_i)_{i=1}^{n_{2j}}$ is a $(3, 1/n_{2j})$ -RIS and by Proposition 5.4 for every $g \in G_\xi$ holds

$$\#\{n \in \mathbb{N} : |g(x_n)| \geq \frac{2}{m_{2j}^4}\} \leq 20m_{2j}^4 \leq n_{2j-1}$$

Set $\check{y} = \frac{m_{2j}}{n_{2j}} \sum_{i=1}^{n_{2j}} x_i$ and $y^* = m_{2j}^{-1} \sum_{i=1}^{n_{2j}} x_i^*$.

Then setting $y := \lambda \check{y}$ for some $\lambda \in [1, 2]$, Proposition 9.4 yields that (y, y^*) is a $(6, 2j)$ -exact pair.

Note that for every $g \in G_\xi$, setting $A = \{i : |g(x_i)| \geq \frac{2}{m_{2j}^2}\}$ we get

$$(46) \quad |g(y)| \leq \frac{2m_{2j}}{n_{2j}} \left(\sum_{i \in A} |g(x_i)| + \sum_{i \notin A} |g(x_i)| \right) \leq \frac{2m_{2j}}{n_{2j}} (n_{2j-1} + \frac{2n_{2j}}{m_{2j}^2}) \leq \frac{6}{m_{2j}}$$

It follows that for every $j \in \mathbb{N}$ and every block subspace we have a $(6, 2j)$ -exact pair (y, y^*) with $y^* \in Y$ and $\|y\|_{G_\xi} \leq 6/m_{2j}$.

Therefore we are able to construct for every $j \in \mathbb{N}$ a n_{2j-1} -attractor sequence $(y_i, y_i^*)_{i=1}^{n_{2j-1}}$ with (y_{2i-1}, y_{2i-1}^*) - a $(6, 2j_{2i-1})$ -exact pair, $y_{2i-1}^* \in Y^*$ and $\|y_{2i-1}\|_{G_\xi} < m_{2j-1}^{-2}$ for every i . From Corollary 10.5 we get

$$\frac{1}{2m_{2j-1}^2} \leq \left\| \frac{1}{n_{2j-1}} \sum_{i=1}^{n_{2j-1}} (-1)^i y_i \right\| \leq \frac{90}{m_{2j-1}^2}.$$

Setting $z_j^* = m_{2j-1}^{-2} \sum_{i=1}^{n_{2j-1}/2} y_{2i-1}^*$ and $w_j^* = -m_{2j-1}^{-2} \sum_{i=1}^{n_{2j-1}/2} y_{2i}^*$ we get that

$$(47) \quad \|z_j^*\|, \|w_j^*\| \geq \frac{1}{180}, \text{ and } \|z_j^* - w_j^*\| \leq \frac{1}{m_{2j-1}}.$$

Let $F \in \mathcal{S}_\xi$ and $(w_j^*)_{j \in F}$ be a G_ξ -special sequence such that w_j^* satisfies (47) for an appropriate z_j^* and $\text{minsupp } w_j^* > j$ for every $j \in F$.

From the definition of the ground set G_ξ we have $\sum_{j \in F} \epsilon_j w_j^* \in G_\xi \subset B_{X_\xi^*}$ and hence

$$\left\| \sum_{j \in F} \epsilon_j z_j^* \right\| \leq \left\| \sum_{j \in F} \epsilon_j w_j^* \right\| + \left\| \sum_{j \in F} \epsilon_j z_j^* - \epsilon_j w_j^* \right\| \leq 2.$$

Notice that the tree consisting of sequences $(z_j^*)_{j \in F}$, with $F \in \mathcal{S}_\xi$, obtained in the way described above has order greater or equal $o(\mathcal{S}_\xi)$.

b) Notice that if there is a normalized basic sequence in X_ξ^* generating a c_0 -spreading model, then we get also a block sequence (f_n) generating a c_0 -spreading model, and then a normalized block sequence $(x_n) \subset X_\xi$ with $f_i(x_j) = \delta_{i,j}$ generates an ℓ_1 -spreading model in X_ξ , a contradiction with Corollary 11.7. \square

13. THE SPACE \mathfrak{X}_ξ

In this section we define our final space \mathfrak{X}_ξ as a suitable quotient X_ξ/X_L and show the desired properties of \mathfrak{X}_ξ (Corollaries 13.5 and 13.6). In order to this we show for any subspace $Y \subset X_\xi$ with Y/X_L infinite dimensional the existence of an ℓ_1 average and a norming functional with controlled behavior with respect to Y/X_L and X_L^\perp (Lemma 13.2) and consequently the existence of a suitable attracting sequence (Corollary 13.4).

For an infinite subset L of \mathbb{N} we set $X_L = \overline{\langle (e_n)_{n \in L} \rangle}$, where $(e_n)_{n \in \mathbb{N}}$ is the basis of X_ξ . We shall prove that for every $L \in [\mathbb{N}]$ with $\mathbb{N} \setminus L$ also infinite and satisfying $\#(L \cap \Lambda_i) = \infty$ for every $i \in \mathbb{N}$, the quotient space $\mathfrak{X}_\xi = X_\xi/X_L$, does not contain a (normalized) sequence generating an ℓ_1 -spreading model but every of its subspaces has ℓ_1 -index greater than ω^ξ . Let $Q : X_\xi \rightarrow \mathfrak{X}_\xi$ denotes the quotient map.

Let $\mathbb{N} = L \cup M$, where $M \cap L = \emptyset$. First we prove the following

Proposition 13.1. *The sequence $(Q(e_n))_{n \in M}$ is a basis for the quotient \mathfrak{X}_ξ .*

Proof. For every $x = \sum_{n \in \mathbb{N}} a_n e_n \in X_\xi$ we get $Q(x) = \sum_{n \in M} a_n Q(e_n)$. Let $M = (m_i)$ and $j < n$. By the bimonotonicity of $(e_n)_{n \in \mathbb{N}}$ we have that $\|\sum_{i=1}^n a_{m_i} Q(e_{m_i})\| = \|\sum_{i=m_1}^{m_n} a_i e_i\|$ for some $a_i, i \in [m_1, m_n]$. Using again that $(e_n)_{n \in \mathbb{N}}$ is bimonotone, for every $j < n$ we obtain

$$\left\| \sum_{i=m_1}^{m_j} a_i e_i \right\| \leq \left\| \sum_{i=m_1}^{m_n} a_i e_i \right\| = \left\| \sum_{i=1}^n a_{m_i} Q(e_{m_i}) \right\|$$

and therefore $\|\sum_{i=1}^j a_{m_i} Q(e_{m_i})\| \leq \|\sum_{i=m_1}^{m_j} a_i e_i\| \leq \|\sum_{i=1}^n a_{m_i} Q(e_{m_i})\|$. \square

We shall need the following result which is analogous to Lemma 11 [13].

Lemma 13.2. *Let $N, m \in \mathbb{N}$, $\epsilon \in (0, 1/4)$ and Y subspace of X_ξ such that the quotient Y/X_L is infinite dimensional. Then there exist a $2 - \ell_1^N$ average $x \in \{e_i : i \geq m\}$ and $f \in K_\xi$ such that*

$$f(x) \geq 1/2, \quad \text{minsupp } f \geq \text{minsupp } x, \quad \text{dist}(Q(x), Y/X_L) < \epsilon \quad \text{and} \quad \text{dist}(f, X_L^\perp) < \epsilon.$$

Proof. Let $(\epsilon_n)_{n \in \mathbb{N}}$ be positive numbers with $\sum_n \epsilon_n < \epsilon$. Let $(\hat{y}_n)_{n \in \mathbb{N}}$ be a normalized w -null sequence in Y/X_L . Passing to a subsequence we may assume that $(\hat{y}_n)_{n \in \mathbb{N}}$ is (ϵ_n) -close to a normalized block sequence $(\hat{x}_n)_{n \in \mathbb{N}} \in \mathfrak{X}_\xi$ and there exists $x_n \in X_\xi$ such that

$$Q(x_n) = \hat{x}_n, \quad \|x_n\| = \|\hat{x}_n\| \quad \text{and} \quad \text{range}(x_n) = [\text{minsupp}(\hat{x}_n), \text{maxsupp}(\hat{x}_n)].$$

Let also $y_n^* \in \mathfrak{X}_\xi^* = X_L^\perp$ such that $y_n^*(x_n) = 1 = \|\hat{x}_n\|$, $\|y_n^*\| = 1$ for every $n \in \mathbb{N}$. Since $(e_n^*)_{n \in \mathbb{N}}$ is bimonotone basis of X_ξ^* , setting $x_n^* = E_n y_n^*$, where $E_n = \text{range}(x_n)$, we get that $(x_n^*)_{n \in \mathbb{N}}$ is a block sequence

$$\|x_n^*\| \leq \|y_n^*\|, \quad x_n^*(x_n) = y_n^*(x_n) \quad \text{and} \quad x_n^* \in X_L^\perp$$

Let $k, j \in \mathbb{N}$ be such that $2^k > m_{2j}^2$ and $N^k \leq n_{2j}$. Such j, k exist, since by definition $n_{2j} = (2n_{2j-1})^{s_{2j-1}}$ and $2^{s_{2j-1}} = m_{2j}^3$. Hence if $N \leq m_{2j-1}$ setting $k = s_{2j-1}$ we get $N^k \leq n_{2j}$ and $2^{-k} < m_{2j}^{-2}$. We set

$$A_1 = \left\{ L \in [\mathbb{N}] : L = (l_i)_{i \in \mathbb{N}} : \left\| \frac{1}{N} \sum_{i=1}^N \hat{x}_{l_i} \right\|_{\mathfrak{X}_\xi} > 1/2 \right\}.$$

By Ramsey theorem we may find an $L \in [\mathbb{N}]$ such that either $[L] \subset A_1$ or $[L] \cap A_1 = \emptyset$.

Assume first that $[L] \subset A_1$. We may assume that $x_{\min L} \geq m$. Since $\hat{x}_n = Q(x_n)$ for every $n \in \mathbb{N}$ it follows that

$$\left\| \frac{1}{N} \sum_{i=1}^N x_{l_i} \right\| \geq \left\| \frac{1}{N} \sum_{i=1}^N \hat{x}_{l_i} \right\|_{\mathfrak{X}_\xi} > 1/2.$$

Let $x = \frac{1}{N} \sum_{i=1}^N x_{l_i}$. Take $g \in \mathfrak{X}_\xi^* = X_L^\perp$ such that $g(\frac{1}{N} \sum_{i=1}^N x_{l_i}) > 1/2$. As before we may assume that $\text{minsupp } x = \text{minsupp } g$.

Setting $\hat{x} = Q(\frac{1}{N} \sum_{i=1}^N x_{l_i})$ we get

$$\text{dist}(\hat{x}, Y/X_L) \leq \left\| \frac{1}{N} \sum_{i=1}^N (\hat{x}_{l_i} - \hat{y}_{l_i}) \right\|_{\mathfrak{X}_\xi} \leq \frac{1}{N} \sum_{i=1}^N \|\hat{x}_{l_i} - \hat{y}_{l_i}\|_{\mathfrak{X}_\xi} < \sum_i \epsilon_{l_i} \leq \epsilon.$$

Since $B_{X_\xi^*} = \overline{K_\xi}^{w^*}$ we choose $f \in K_\xi$ with

$$f(x) \geq 1/2, \quad \text{minsupp } f \geq \text{minsupp } x \quad \text{and} \quad \|f - g\| < \epsilon.$$

This is possible by taking an interval B such that $\|g - Bg\| < \epsilon/2$ and using that $\text{minsupp } x = \text{minsupp } g$ and $B_{X_\xi^*} = \overline{K_\xi}^{w^*}$.

Assume now that $[L] \cap A_1 = \emptyset$. Set $\hat{y}_n^{(1)} = \frac{1}{N} \sum_{i \in F_n} \hat{x}_{l_i}$ where $\#F_i = N$, $F_n < F_{n+1}$ for every n and $\cup_i F_i = L$.

Passing to a subsequence we may assume that there exists $a_1 \geq 2$ such that $\hat{x}_n^{(2)} = a_1 \frac{1}{N} \sum_{i \in F_n} \hat{x}_i^{(1)}$ is a normalized sequence in Y/Z . We may again apply Ramsey theorem defining A_2 as before. If we get some L with $[L] \subset A_2$ the proof finishes as before.

Assume that in none of the first k steps we get $L \in [\mathbb{N}]$ with $[L] \subset A_k$. Then there exist $a_1, a_2, \dots, a_k \geq 2$ and $l_1 < l_2 < \dots < l_{N^k}$ in \mathbb{N} such that the vector

$$\hat{y} = a_1 a_2 \cdots a_k \frac{1}{N^k} \sum_{i=1}^{N^k} \hat{x}_{l_i}$$

satisfies $\|\widehat{y}\|_{\mathfrak{X}_\xi} = 1$.

The functional $y^* = \frac{1}{m_{2j}} \sum_{i=1}^{N^k} x_{i_i}^* \in X_L^\perp$ satisfies $\|y^*\| \leq 1$. Therefore

$$1 = \|\widehat{y}\|_{\mathfrak{X}_\xi} \geq y^*(\widehat{y}) = \frac{2^k N^k}{m_{2j} N^k} = \frac{2^k}{m_{2j}}$$

which contradicts our choice of k and j . \square

Corollary 13.3. *Let Y/X_L be an infinitely dimensional subspace of the quotient X_ξ/X_L . For every $j \in \mathbb{N}$, $\epsilon > 0$ there exists a $(6, 2j)$ -exact pair (y, f) such that*

- (1) $\text{dist}(Q(y), Y/X_L) < \epsilon$ and $\text{dist}(f, X_L^\perp) < \epsilon$,
- (2) $\|y\|_{G_\xi} < 6m_{2j}^{-1}$.

Proof. Let $j \in \mathbb{N}$ and $(\epsilon_i)_i \subset (0, 1)$ such that $\sum_i \epsilon_i < \epsilon \leq n_{2j}^{-1}$. Using Lemma 13.2 inductively we choose $(z_i, f_i)_{i=1}^{n_{2j}} \in (c_{00}(\mathbb{N}) \times K_\xi)$ such that

- a) z_i is a $2 - \ell_1^{n_{2j}}$ average and $(z_i)_{i=1}^{n_{2j}}$ is a $(3, \epsilon)$ -RIS.
- b) $\text{dist}(Q(z_i), Y/X_L) < \epsilon_i$ and $\text{dist}(f_i, X_L^\perp) < \epsilon_i$ for every i .
- c) $f_i(z_i) \geq 1/2$.
- d) $\text{range}(z_i) \cup \text{range}(f_i) \subset \text{range}(z_{i+1}) \cup \text{range}(f_{i+1})$ for every i .

By Proposition 5.4 we can assume that for every $g \in G_\xi$ the set $\{i \in \mathbb{N} : |g(z_i)| \geq 2m_j^{-2}\}$ has cardinality at most n_{j-1} . Setting $z = \frac{m_{2j}}{n_{2j}} \sum_{i=1}^{n_{2j}} z_i$ and $f = m_{2j}^{-1} \sum_{i=1}^{n_{2j}} f_i$ we have $f(z) \geq 1/2$. Taking $y := \lambda z$ for some $\lambda \in [1, 2]$ we get from Proposition 9.4 that (y, f) is a $(6, 2j)$ -exact pair. It is easy to see that the exact pair (y, f) satisfies the requirements (1). The argument of (46) yields that (2) also holds. \square

Corollary 13.4. *For every $j \in \mathbb{N}$, any Y/X_L infinitely dimensional subspace of the quotient X_ξ/X_L and any $\epsilon > 0$ there exists a $(6, 2j - 1)$ -attracting sequence $(y_i, f_i)_{i=1}^{n_{2j-1}}$ with the associated sequence $(j_i)_i$ such that*

- A) $\text{supp } f_{2i} = \text{supp } y_{2i} \subset L$,
- B) For every $i \leq n_{2j-1}$, (y_{2i-1}, f_{2i-1}) is a $(6, j_{2i-1})$ -exact pair and $\|y_{2i-1}\|_{G_\xi} < m_{2j-1}^{-2}$,
- C) $\text{dist}(Q(\sum_i y_{2i-1}), Y/X_L) < \epsilon$,
- D) There exists $g \in X_L^\perp$ with

$$\|g - \frac{1}{m_{2j-1}} \sum_{i=1}^{n_{2j-1}/2} f_{2i-1}\| \leq m_{2j-1}^{-2} \quad \text{and} \quad \|g - \frac{1}{m_{2j-1}} \sum_{i=1}^{n_{2j-1}/2} f_{2i}\| \leq 2m_{2j-1}^{-1}.$$

Proof. Using Corollary 13.3 and assumption on the set L we can choose an attracting sequence $(y_i, f_i)_{i=1}^{n_{2j-1}}$ such that A) holds and for every $i \leq n_{2j-1}/2$ the couple (y_{2i-1}, f_{2i-1}) is a $(6, j_{2i-1})$ -exact pair satisfying the conclusion of Corollary 13.3 for $\epsilon_i = \epsilon 2^{-i}$, i.e.

- (1) $\text{dist}(Q(y_{2i-1}), Y/X_L) < \epsilon_i$ and $\text{dist}(f_{2i-1}, X_L^\perp) < \epsilon_i$
- (2) $\|y_{2i-1}\|_{G_\xi} < 6m_{j_{2i-1}}^{-1} < m_{2j-1}^{-2}$.

For every $i \leq n_{2j-1}/2$ choose $g_{2i-1} \in X_L^\perp$ with $\|f_{2i-1} - g_{2i-1}\| < \epsilon 2^{-i}$ and let $g = m_{2j-1}^{-2} \sum_{i=1}^{n_{2j-1}/2} g_{2i-1}$.

By Corollaries 10.5, 10.6 the chosen vectors and functionals satisfy the desired conditions. \square

Corollary 13.5. *Every subspace of the quotient \mathfrak{X}_ξ has ℓ_1 -index greater than ω^ξ .*

Proof. By Corollary 13.4 we can choose for every subspace Y/X_L of the quotient \mathfrak{X}_ξ and any j an attracting sequence $(y_l^j, f_l^j)_{l=1}^{n_{2j-1}}$ and $g_j \in X_L^\perp$ such that for all l , $\|y_l^j\|_G < m_{2j-1}^{-2}$

and for

$$z_j^* = m_{2^{j-1}}^{-2} \sum_{l=1}^{n_{2^{j-1}}/2} f_{2l}^j \in G_1, \quad f_j = m_{2^{j-1}}^{-2} \sum_{l=1}^{n_{2^{j-1}}/2} f_{2l-1}^j, \quad u_j = \frac{m_{2^{j-1}}^2}{n_{2^{j-1}}} \sum_{l=1}^{n_{2^{j-1}}/2} y_{2l-1}^j$$

we have

$$\|g_j - z_j^*\| < 2m_{2^{j-1}}^{-1} \text{ and } \|g_j - f_j\| \leq m_{2^{j-1}}^{-2} \text{ and } \text{dist}(Q(u_j), Y/X_L) < 16^{-j}.$$

In particular we get $\|g_j\| \leq 2$. Setting $\hat{u}_j = Q(u_j) = Q\left(\frac{m_{2^{j-1}}^2}{n_{2^{j-1}}} \sum_{l=1}^{n_{2^{j-1}}/2} (-1)^{j+1} y_l^j\right)$ we get by Corollary 10.5

$$\begin{aligned} \frac{1}{8} &\leq \frac{1}{4} - m_{2^{j-1}}^{-2}/2 \leq f_j(u_j)/2 - \|f_j - g_j\|/2 \leq g_j(u_j)/2 \leq \|\hat{u}_j\|_{\mathfrak{X}_\xi} \\ &\leq \left\| \frac{m_{2^{j-1}}^2}{n_{2^{j-1}}} \sum_{l=1}^{n_{2^{j-1}}/2} (-1)^{j+1} y_l^j \right\| \leq 90. \end{aligned}$$

For any $F \in \mathcal{S}_\xi$ we pick $(z_j^*)_{j \in F}$, $(g_j)_{j \in F}$, $(u_j)_{j \in F}$ so that

- (1) z_j^* , g_j , u_j are picked as above for any $j \in F$,
- (2) $\text{minsupp}(z_j) > j$ for any $j \in F$,
- (3) $(z_j^*)_{j \in F}$ is a G_ξ -special sequence.

Then we have that $\|\sum_{j \in F} \epsilon_j z_j^*\| \leq 1$ for every $\epsilon_j \in \{-1, 1\}$. Since $\|g_j - z_j^*\| \leq 2^{-j}$ for every $j \in F$ it follows that $\|\sum_{j \in F} \epsilon_j g_j\| \leq 2$ for every $\epsilon_j = \pm 1$. Therefore

$$\left\| \sum_{j \in F} a_j \hat{u}_j \right\|_{\mathfrak{X}_\xi} \geq \frac{1}{2} \sum_{j \in F} \text{sgn}(a_j) g_j \left(\sum_{j \in F} a_j u_j \right) \geq \frac{1}{2} \sum_j \frac{1}{8} |a_j| \geq \frac{1}{16} \sum |a_j|.$$

Since $\text{dist}(\hat{u}_j, Y/X_L) < 16^{-j}$ for any j , by the above procedure we can obtain an ℓ_1 -tree in Y/X_L with order greater or equal $o(\mathcal{S}_\xi)$, hence the ℓ_1 -index of Y/X_L is greater than ω^ξ . \square

Corollary 13.6. *The space $\mathfrak{X}_\xi = X_\xi/X_L$ does not contain a normalized basic sequence generating an ℓ_1 -spreading model.*

Proof. If a normalized basic sequence $(\hat{x}_n)_{n \in \mathbb{N}} \subset \mathfrak{X}_\xi$ generates an ℓ_1 -spreading model in \mathfrak{X}_ξ , we may assume that it is a block sequence. Then we get a block sequence in X_ξ which generates an ℓ_1 -spreading model, a contradiction by Corollary 11.7. \square

Gathering the results of this and previous sections we get the following

Theorem 13.7. *For every countable ordinal ξ there exists a separable reflexive Banach space \mathfrak{X}_ξ with the hereditary Bourgain ℓ_1 -index greater than ω^ξ such that \mathfrak{X}_ξ does not admit an ℓ_1 -spreading model. Moreover the dual \mathfrak{X}_ξ^* has hereditary c_0 -index greater than ω^ξ and does not admit a c_0 -spreading model.*

Remark 13.8. *The results presented above concerning indices of the dual space X_ξ^* and the quotient space \mathfrak{X}_ξ suggest the following problem:*

Assume that X is a reflexive Banach space such that every subspace of the dual has c_0 -index greater than ω^ξ . Does there exist a subspace Y such that every subspace of the quotient has X/Y has Bourgain ℓ_1 -index greater than ω^ξ ?

REFERENCES

- [1] F. Albiac and N.J. Kalton, *Topics in Banach space theory*, Graduate Texts in Mathematics 233, Springer, New York, 2006.
- [2] D. Alspach and S.A. Argyros, *Complexity of weakly null sequences*, Diss. Math. 321 (1992), 1-44.

- [3] S.A. Argyros, A. D. Arvanitakis and A. Tolia, *Saturated extensions, the attractors method and Hereditarily James Tree Spaces*, London Math. Soc. Lecture Note Ser., 337, Cambridge Univ. Press, Cambridge, 2006.
- [4] S.A. Argyros and I.Gasparis, *Unconditional structures of weakly null sequences*, Trans. Amer. Math. Soc. 353(5), 2019-2058, 2001.
- [5] S.A. Argyros and V. Kanellopoulos, *Determining c_0 in $C(K)$ spaces*, Fund. Math. 187 (2005), 61–93.
- [6] S. A. Argyros, S. Mercourakis and A. Tsarpalias, *Convex unconditionality and summability of weakly null sequences*, Israel J. Math. 107 (1998), 157–193.
- [7] S.A. Argyros and S.Todorcevic, *Ramsey methods in analysis. Advanced Courses in Mathematics*. CRM Barcelona. Birkhäuser Verlag, Basel, 2005.
- [8] S.A. Argyros and A. Tolia, *Indecomposability and unconditionality in duality*, Geom. Funct. Anal. 14 (2004), no. 2, 247–282.
- [9] S.A. Argyros and A. Tolia, *Methods in the theory of hereditarily indecomposable Banach spaces*, Mem. Amer. Math. Soc. 170 (2004), no. 806.
- [10] J. Bourgain, *On convergent sequences of continuous functions*, Bull. Soc. Math. Belg. Sér. B 32 (1980), 235–249.
- [11] A. Brunel and L.Sucheston, *On B -convex Banach spaces*, Math. Systems Theory, 7 (1974), no. 4, 294-299.
- [12] P. Erdős and M. Magidor, *A note on regular methods of summability and the Banach-Saks property*, Proc. Amer. Math. Soc., 59 (1976), no. 2, 232–234.
- [13] V. Ferenczi, *Quotient hereditarily indecomposable Banach spaces*, Canadian J. Math. 51 (1999), no.3, 566–584.
- [14] I. Gasparis, *A dichotomy theorem for subsets of the power set of the natural numbers*, Proc. Amer. Math. Soc. 129 (2001), no. 3, 759–764.
- [15] W.T. Gowers *An infinite Ramsey theorem and some Banach-space dichotomies*, Ann. of Math. (2), 156 (2002), no. 3, 797-833.
- [16] W.T. Gowers and B. Maurey, *The unconditional basic sequence problem*, J. Amer. Math. Soc., 6 (1993), no.4, 851–874.
- [17] R.C. James, *A separable somewhat reflexive space with nonseparable dual*, Bull. Amer. Math. Soc., 80 (1974), 738–743.
- [18] R. Judd and E. Odell, *Concerning Bourgain’s ℓ_1 -index of a Banach space*, Israel J. Math. 108 (1998), 145–171.
- [19] J.L. Krivine, *Sous espaces de dimension finie des espaces de Banach réticulés*, Ann. of Math. (2) 104 (1976), 1–29.
- [20] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces I*, Springer-Verlag 92, 1977.
- [21] B. Maurey, V.D. Milman and N. Tomczak-Jaegermann, *Asymptotic Infinite-Dimensional theory of Banach spaces*, Geometric aspects of functional analysis (Israel, 1992–1994), 149–175, Oper. Theory Adv. Appl., 77, Birkhäuser, Basel, 1995.
- [22] B. Maurey and H. Rosenthal, *Normalized weakly null sequence with no unconditionall subsequence*, Studia Math. 61 (1977), 77–98.
- [23] V. Milman and N. Tomczak-Jaegermann, *Asymptotic ℓ_p spaces and bounded distortion*, Banach spaces (Merida 1992), 173–195, Contemp. Math. 144, Amer. Math. Soc., Providence, RI, 1993.
- [24] E. Odell and Th. Schlumprecht, *On the richness of the set of p 's in Krivine’s theorem*, Geometric aspects of functional analysis (Israel, 1992–1994), 177–198, Oper. Theory Adv. Appl., 77, Birkhäuser, Basel, 1995.
- [25] A.M. Pelczar, *Note on distortion and Bourgain ℓ_1 -index*, Studia Math. 190 (2009), no. 2, 147–161.
- [26] A. Pełczyński and Z. Semadeni, *Spaces of continuous functions III, Spaces $C(\Omega)$ for Ω without perfect subsets*, Studia Math. 18 (1959), 211–222.
- [27] Th. Schlumprecht, *An arbitrarily distortable Banach space*, Israel J. Math. 76 (1991), 81–95.
- [28] B. S. Tsirelson, *Not every Banach space contains ℓ_p or c_0* , Funct. Anal. Appl. 8 (1974), 138–141.

DEPARTMENT OF MATHEMATICS, NATIONAL TECHNICAL UNIVERSITY OF ATHENS, ATHENS 15780, GREECE

E-mail address: sargyros@math.ntua.gr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AEGEAN, KARLOVASI, GREECE

E-mail address: amanouss@aegean.gr

INSTITUTE OF MATHEMATICS, JAGIELLONIAN UNIVERSITY, ŁOJASIEWICZA 6, 30-348 KRAKÓW, POLAND

E-mail address: anna.pelczar@im.uj.edu.pl