# Hitting all maximum cliques with a stable set using lopsided independent transversals

Andrew D. King<sup>\*</sup>

November 2, 2018

#### Abstract

Rabern recently proved that any graph with  $\omega \geq \frac{3}{4}(\Delta+1)$  contains a stable set meeting all maximum cliques. We strengthen this result, proving that such a stable set exists for any graph with  $\omega > \frac{2}{3}(\Delta +$ 1). This is tight, i.e. the inequality in the statement must be strict. The proof relies on finding an independent transversal in a graph partitioned into vertex sets of unequal size.

**Keywords:** maximum clique, stable set, graph colouring, independent transversal, independent system of representatives.

### **1** Introduction and motivation

When colouring a graph G, we often desire a stable set S meeting every maximum clique. For example, finding such a set S efficiently is the key to colouring perfect graphs in polynomial time [11]. Proving the existence of Shas also been very useful in attacking Reed's  $\omega$ ,  $\Delta$ , and  $\chi$  conjecture<sup>1</sup>:

Conjecture 1 (Reed [10]). For any graph G,  $\chi(G) \leq \lfloor \frac{1}{2}(\Delta(G) + 1 + \omega(G)) \rfloor$ .

<sup>\*</sup>Department of Industrial Engineering and Operations Research, Columbia University, New York. Email: andrew.d.king@gmail.com. Research supported by an NSERC Postdoctoral Fellowship.

 $<sup>{}^{1}\</sup>omega$ ,  $\Delta$ , and  $\chi$  denote the clique number, maximum degree, and chromatic number of a graph, respectively.

No minimum counterexample G to this conjecture has such a stable set S. For if it did, it would contain a maximal stable set S' meeting every maximum clique, and we would have  $\lceil \frac{1}{2}(\Delta(G-S')+1+\omega(G-S'))\rceil+1 \leq \lceil \frac{1}{2}(\Delta(G)+1+\omega(G))\rceil$ . Since S' is a stable set,  $\chi(G) \leq \chi(G-S')+1$ , contradicting the minimality of G.

Thus such a stable set S is highly desirable when attacking Reed's conjecture for a hereditary class of graphs. The proof of Reed's conjecture for line graphs [7] exemplifies the general approach: If the maximum degree and clique number are far apart, a combination of previously known results suffices. If they are not far apart, we can use the structure of line graphs to prove the existence of a stable set S meeting all maximum cliques.

Rabern [9] recently proved that if the maximum degree and clique number are close enough, we need not consider the structure of the graph class at all:

**Theorem 2** (Rabern). If a graph G satisfies  $\omega(G) \geq \frac{3}{4}(\Delta(G) + 1)$ , then G contains a stable set meeting all maximum cliques.

Here we prove the best possible theorem of this type:

**Theorem 3.** If a graph G satisfies  $\omega(G) > \frac{2}{3}(\Delta(G) + 1)$ , then G contains a stable set S meeting every maximum clique.

To see that this is best possible, let  $G_k$  be the graph obtained by substituting every vertex of a 5-cycle with a clique of size k. Then  $\omega(G_k) = 2k = \frac{2}{3}(\Delta(G_k) + 1)$ , and no stable set meets every maximum clique. To prove Theorem 3 we apply Rabern's approach with a stronger final step. Rabern applies Haxell's theorem [4], which can be stated as follows:

**Theorem 4** (Haxell). For a positive integer k, let G be a graph with vertices partitioned into r cliques of size  $\geq 2k$ . If every vertex has at most k neighbours outside its own clique, then G contains a stable set of size r.

To prove our theorem we need to deal with a graph that has been partitioned into cliques of unequal size. We use the following extension of Theorem 4:

**Theorem 5.** For a positive integer k, let G be a graph with vertices partitioned into cliques  $V_1, \ldots V_r$ . If for every i and every  $v \in V_i$ , v has at most  $\min\{k, |V_i| - k\}$  neighbours outside  $V_i$ , then G contains a stable set of size r.

Although this is not at all obvious, it is a straightforward consequence of observations made by Aharoni, Berger, and Ziv about the proof of Theorem 4 [1].

#### 2 Hitting the maximum cliques

To prove Theorem 3 we must investigate intersections of maximum cliques. Given a graph G and the set  $\mathcal{C}$  of maximum cliques in G, we define the *clique* graph  $G(\mathcal{C})$  as follows. The vertices of  $G(\mathcal{C})$  are the cliques of  $\mathcal{C}$ , and two vertices of  $G(\mathcal{C})$  are adjacent if their corresponding cliques in G intersect. For a connected component  $G(\mathcal{C}_i)$  of  $G(\mathcal{C})$ , let  $D_i \subseteq V(G)$  and  $F_i \subseteq V(G)$ denote the union and the mutual intersection of the cliques of  $\mathcal{C}_i$  respectively, i.e.  $D_i = \bigcup_{C \in \mathcal{C}_i} C$  and  $F_i = \bigcap_{C \in \mathcal{C}_i} C$ .

The proof uses three intermediate results. The first, due to Hajnal [2] (also see [9]), tells us that for each component of  $G(\mathcal{C})$ ,  $|D_i| + |F_i|$  is large:

**Lemma 6** (Hajnal). Let G be a graph and  $C_1, \ldots, C_r$  be a collection of maximum cliques in G. Then

$$\left| \bigcap_{i \leq r} C_i \right| + \left| \bigcup_{i \leq r} C_i \right| \geq 2\omega(G).$$

The second is due to Kostochka [8] (proven in English in [9]). It tells us that if  $\omega(G)$  is sufficiently close to  $\Delta(G) + 1$ , then  $|F_i|$  is large:

**Lemma 7** (Kostochka). Let G be a graph with  $\omega(G) > \frac{2}{3}(\Delta(G) + 1)$  and let  $\mathcal{C}$  be the set of maximum cliques in G. Then for each connected component  $G(\mathcal{C}_i)$  of  $G(\mathcal{C})$ ,

$$\left| \bigcap_{C \in \mathcal{C}_i} C \right| \ge 2\omega(G) - (\Delta(G) + 1).$$

The third intermediate result is Theorem 5. Combining them to prove Theorem 3 is a simple matter.

Proof of Theorem 3. Let  $\mathcal{C}$  be the set of maximum cliques of G, and denote the connected components of  $G(\mathcal{C})$  by  $G(\mathcal{C}_1), \ldots, G(\mathcal{C}_r)$ . For each  $\mathcal{C}_i$ , let  $F_i = \bigcap_{C \in \mathcal{C}_i} C$  and let  $D_i = \bigcup_{C \in \mathcal{C}_i} C$ . It suffices to prove the existence of a stable set S in G intersecting each clique  $F_i$ .

Lemma 7 tells us that  $|F_i| > \frac{1}{3}(\Delta(G) + 1)$ . Consider a vertex  $v \in F_i$ , noting that v is universal in  $G[D_i]$ . By Lemma 6, we know that  $|F_i| + |D_i| > \frac{4}{3}(\Delta(G) + 1)$ . Therefore  $\Delta(G) + 1 - |D_i| < |F_i| - \frac{1}{3}(\Delta(G) + 1)$ , so v has fewer than  $|F_i| - \frac{1}{3}(\Delta(G) + 1)$  neighbours in  $\bigcup_{j \neq i} F_i$ . Furthermore v certainly has fewer than  $\frac{1}{3}(\Delta(G) + 1)$  neighbours in  $\bigcup_{j \neq i} F_i$ . Now let H be the subgraph of G induced on  $\cup_i F_i$ , and let  $k = \frac{1}{3}(\Delta(G)+1)$ . Clearly the cliques  $F_1, \ldots, F_r$  partition V(H). A vertex  $v \in F_i$  has at most  $\min\{k, |F_i| - k\}$  neighbours outside  $F_i$ . Therefore by Theorem 5, H contains a stable set S of size r. This set S intersects each  $F_i$ , and consequently it intersects every clique in  $\mathcal{C}$ , proving the theorem.  $\Box$ 

It remains to prove Theorem 5. We do this in the next section.

#### **3** Independent transversals with lopsided sets

Suppose we are given a finite graph whose vertices are partitioned into stable sets  $V_1, \ldots, V_r$ . An *independent system of representatives* or *ISR* of  $(V_1, \ldots, V_r)$  is a stable set of size r in G intersecting each  $V_i$  exactly once. A *partial ISR*, then, is simply a stable set in G intersecting no  $V_i$  more than once. ISR's are intimately related to both the strong chromatic number [6] and list colourings [5].

A totally dominating set D is a set of vertices such that every vertex of G has a neighbour in D, including the vertices of D. Given  $J \subseteq [m]$ , we use  $V_J$  to denote  $(V_i \mid i \in J)$ . Given  $X \subseteq V(G)$ , we use I(X) to denote the set of partitions intersected by X, i.e.  $I(X) = \{i \in [r] \mid V_i \cap X \neq \emptyset\}$ . For an induced subgraph H of G, we implicitly consider H to inherit the partitioning of G.

To prove our lopsided existence condition for ISR's, we use a consequence of Haxell's proof of Theorem 4 [4] pointed out (and proved explicitly) by Aharoni, Berger, and Ziv [1]. Actually we prove a slight strengthening of their result:

**Lemma 8.** Let  $x_1$  be a vertex in  $V_r$ , and suppose  $G[V_{[r-1]}]$  has an ISR. Suppose there is no  $J \subseteq [r-1]$  and  $D \subseteq V_J \cup \{x_1\}$  totally dominating  $V_J \cup \{x_1\}$  with the following properties:

- 1. D is the union of disjoint stable sets X and Y.
- 2. Y is a (not necessarily proper) partial ISR for  $V_J$ . Thus  $|Y| \leq |J|$ .
- 3. Every vertex in Y has exactly one neighbour in X. Thus  $|X| \leq |Y|$ .
- 4. X contains  $x_1$ .

Then G has an ISR containing  $\{x_1\}$ .

Proof. Let G be a minimum counterexample; we can assume  $G = G[V_{[r-1]} \cup \{x_1\}]$ . Furthermore, r > 1 otherwise the lemma is trivial. Let  $R_1$  be an ISR of  $G[V_{[r-1]}]$  chosen such that the set  $Y'_1 = Y_1 = R_1 \cap N(x_1)$  has minimum size. We know that  $R_1$  exists because  $G[V_{[r-1]}]$  has at least one ISR, and we know that  $Y'_1$  is nonempty because G does not have an ISR. Now let  $X_1 = \{x_1\}$  and let  $D_1 = X_1 \cup Y_1$ .

We now construct an infinite sequence of partial ISRs  $Y_1 \subset Y_2 \subset \ldots$ , which contradicts the fact that G is finite. Let i > 1, and suppose we have sets  $\{R_j, Y_j, X_j \mid 1 \leq j < i\}$  such that:

- $X_j$  is a stable set consisting of distinct vertices  $\{x_1, \ldots, x_j\}$ . For j > 1,  $x_j$  is a vertex in  $G[V_{I(Y_{j-1})}]$  with no neighbour in  $X_{j-1} \cup Y_{j-1}$ .
- $R_j$  is an ISR of  $G[V_{[r-1]}]$  such that for every  $1 \le \ell < j, R_j \cap N(X_\ell) = Y_\ell$ . Subject to that,  $R_j$  is chosen so that  $Y'_j = R_j \cap N(x_j)$  is minimum. For  $1 \le j < i, Y'_j$  is nonempty.
- $Y_j = \bigcup_{i=1}^j Y'_j$ .

To find  $x_i, Y'_i$ , and  $R_i$ , we proceed as follows.

- 1. Let  $x_i$  be any vertex in  $G[V_{I(Y_{i-1})}]$  with no neighbour in  $X_{i-1} \cup Y_{i-1}$ . We know that  $x_i$  exists, otherwise the set  $D_{i-1} = X_{i-1} \cup Y_{i-1}$  would be a total dominating set for  $G[V_{I(Y_{i-1})} \cup \{x_1\}]$ , contradicting the fact that G is a counterexample.
- 2. Let  $R_i$  be an ISR of  $G[V_{[r-1]}]$  chosen so that for all  $1 \leq j < i, R_i \cap N(x_j) = R_j \cap N(x_j) = Y'_j$ . Subject to that, choose  $R_i$  so that  $Y'_i = R_i \cap N(x_i)$  is minimum. We know that  $R_i$  exists because  $R_{i-1}$  is a possible candidate for the ISR.
- 3. It remains to show that  $Y'_i$  is nonempty, i.e. that  $Y_i \neq Y_{i-1}$ . Suppose  $Y'_i = \emptyset$ . We will show that this contradicts our choice of  $R_j$  for the unique j < i such that  $x_i \in V_{I(Y'_j)}$ . Let y be the unique vertex in  $R_i \cap V_{I(x_i)}$ . Construct  $R'_j$  from  $R_i$  by removing y and inserting  $x_i$ . Now for every  $\ell$  such that  $1 \leq \ell < j$ ,  $R'_j \cap N(x_\ell) = Y'_\ell = R_j \cap N(x_\ell)$ . For j,  $R'_i \cap N(x_i) = (R_j \cap N(x_i)) \setminus \{y\}$ , a contradiction. Thus  $Y'_i$  is nonempty.
- 4. Set  $X_i = X_{i-1} \cup \{x_i\}$  and  $Y_i = Y_{i-1} \cup Y'_i$ .

This choice of  $X_i$ ,  $R_i$ , and  $Y_i$  sets up the conditions so that we can repeat our argument indefinitely for increasing i, a contradiction since G is finite.  $\Box$ 

The lemma easily implies Theorem 3.5 in [1], and allows us to prove a strengthening of Theorem 5:

**Theorem 9.** Let k be a positive integer and let G be a graph partitioned into stable sets  $(V_1, \ldots, V_r)$ . If for each  $i \in [r]$ , each vertex in  $V_i$  has degree at most  $\min\{k, |V_i| - k\}$ , then for any vertex v, G has an ISR containing v.

Proof. Suppose G is a minimum counterexample for a given value of k. Clearly we can assume each  $V_i$  has size greater than k, and that  $G[V_J]$  has an ISR for all  $J \subset [r]$ . Take v such that G does not have an ISR containing v; we can assume  $v \in V_r$ . By Lemma 8, there is some  $J \subseteq [r-1]$  and a set  $D \subseteq V_J \cup \{v\}$  totally dominating  $V_J \cup \{v\}$  such that (i) D is the union of disjoint stable sets X and Y, (ii) Y is a partial ISR of  $V_J$ , (iii)  $|X| \leq |Y| \leq |J|$ , and (iv)  $v \in X$ .

Since *D* totally dominates  $V_J \cup \{v\}$ , the sum of degrees of vertices in *D* must be greater than the number of vertices in  $V_J$ . That is,  $\sum_{v \in D} d(v) > \sum_{i \in J} |V_i|$ . Clearly  $\sum_{v \in X} d(v) \le k \cdot |J|$  and  $\sum_{v \in Y} d(v) \le \sum_{i \in J} (|V_i| - k)$ , so  $\sum_{v \in D} d(v) \le \sum_{i \in J} |V_i|$ , contradicting the fact that *D* is totally dominating. This proves the theorem.

This extends Haxell's theorem by bounding the difference between the degree of a vertex and the size of its partition. One might hope that bounding the ratio of these by  $\frac{1}{2}$  is enough, but it is not: Given  $V_1$  of size four and  $V_2, \ldots, V_5$  of size two, in which each vertex of  $V_1$  dominates one of the smaller sets, there exists no ISR [3]. Lemma 8 cannot imply such a result because in the totally dominating set  $D = X \cup Y$ , we have no control over the average degree of a vertex in X – it may be k. So while we know that the average degree of a vertex in Y behaves nicely with respect to the average partition size, the same is not necessarily true of X. Thus Theorem 9 gives a lopsided existence condition that is not only a useful consequence of Lemma 8, but also a natural one.

# 4 Acknowledgements

The author is grateful to Landon Rabern and Penny Haxell for helpful discussions, and to Robert Himmelmann for pointing out an error in an earlier version of the paper.

## References

- R. Aharoni, E. Berger, and R. Ziv. Independent systems of representatives in weighted graphs. *Combinatorica*, 27(3):253–267, 2007.
- [2] A. Hajnal. A theorem on k-saturated graphs. *Canadian J. Math.*, 17:720–724, 1965.
- [3] P. E. Haxell. Personal communication.
- [4] P. E. Haxell. A condition for matchability in hypergraphs. Graphs and Combinatorics, 11(3):245–248, 1995.
- [5] P. E. Haxell. A note on vertex list colouring. *Combinatorics, Probability* and Computing, 10(04):345–347, 2001.
- [6] P. E. Haxell. On the strong chromatic number. Combinatorics, Probability and Computing, 13(06):857–865, 2004.
- [7] A. D. King, B. A. Reed, and A. Vetta. An upper bound for the chromatic number of line graphs. *Eur. J. Comb.*, 28(8):2182–2187, 2007.
- [8] A. V. Kostochka. Degree, density, and chromatic number of graphs. Metody Diskret. Analiz. (In Russian), 35:45–70, 1980.
- [9] L. Rabern. On hitting all maximum cliques with an independent set. Arxiv preprint arXiv:0907.3705, 2009. Submitted.
- [10] B. A. Reed.  $\omega$ ,  $\Delta$ , and  $\chi$ . J. Graph Theory, 27:177–212, 1998.
- [11] B. A. Reed. A gentle introduction to semi-definite programming. In J. L. Ramírez Alfonsín and B. A. Reed, editors, *Perfect Graphs*, chapter 11. Wiley, 2001.