On the quantum Rényi relative entropies and related capacity formulas

Milán Mosonyi and Fumio Hiai

Abstract—Following Csiszár's approach in classical information theory, we show that the quantum α -relative entropies with parameter $\alpha \in (0,1)$ can be represented as generalized cutoff rates, and hence provide a direct operational interpretation to the quantum α -relative entropies. We also show that various generalizations of the Holevo capacity, defined in terms of the α -relative entropies, coincide for the parameter range $\alpha \in (0,2]$, and show an upper bound on the one-shot ε -capacity of a classical-quantum channel in terms of these capacities.

Index Terms—Rényi relative entropies, Hoeffding distances, generalized cutoff rates, quantum channels, α -capacities, one-shot capacities.

I. INTRODUCTION

N information theory, it is convenient to measure the distance of states (probability distributions in the classical, and density operators in the quantum case) with measures that do not satisfy the axioms of a metric. In a broad sense, a *statistical distance* is a function taking non-negative values on pairs of states, that satisfies some convexity properties in its arguments and which cannot increase when its arguments are subjected to a stochastic operation. Probably the most popular statistical distance, for a good reason, is the *relative entropy* S, defined for density operators ρ , σ as

$$S\left(\rho \left| \right|\sigma\right) := \begin{cases} \operatorname{Tr}\rho(\log\rho - \log\sigma), & \text{if } \operatorname{supp}\rho \leq \operatorname{supp}\sigma, \\ +\infty, & \text{otherwise}. \end{cases}$$

While various generalizations of the relative entropy, leading to statistical distances in the above sense, are easy to define, they are not equally important, and the relevant ones are those that appear in answers to natural statistical problems, or in other terms, those that admit an operational interpretation.

The operational interpretation of the relative entropy is given in the problem of *asymptotic binary state discrimination*, where one is provided with several identical copies of a quantum system and the knowledge that the state of the system

The work of M. Mosonyi was supported by the Hungarian Research Grant OTKA T068258, the JSPS Japan-Hungary Joint Project, and by the Centre for Quantum Technologies, which is funded by the Singapore Ministry of Education and the National Research Foundation as part of the Research Centres of Excellence program. Part of this work was done when M. Mosonyi was a Scientific Researcher at the Fields Institute during the Thematic Program on Mathematics in Quantum Information. The work of F. Hiai was supported by the Grant-in-Aid for Scientific Research (C)21540208 and by the JSPS Japan-Hungary Joint Project.

M. Mosonyi is with the Centre for Quantum Technologies, National University of Singapore, 117543 Singapore and the Mathematical Institute, Budapest University of Technology and Economics Budapest, 1111 Hungary (e-mail: milan.mosonyi@gmail.com).

F. Hiai is with the Graduate School of Information Sciences, Tohoku University, Sendai, 980-8579 Japan (e-mail: hiai@math.is.tohoku.ac.jp).

is either ρ (null hypothesis) or σ (alternative hypothesis), where ρ and σ are density operators on the system's Hilbert space \mathcal{H} , and one's goal is to make a good guess for the true state of the system, based on measurement results on the copies. It is easy to see that the most general inference scheme, based on measurements on n copies, can be described by a binary positive operator valued measurement (T, I-T), where $T \in \mathcal{B}(\mathcal{H}^{\otimes n}), 0 < T < I$, and the guess is ρ if the outcome corresponding to T occurs, and σ otherwise. The probability of a wrong guess is $\alpha_n(T) := \operatorname{Tr} \rho^{\otimes n}(I - T)$ if the true state is ρ (error probability of the first kind) and $\beta_n(T) := \operatorname{Tr} \sigma^{\otimes n} T$ if the true state is σ (error probability of the second kind). Unless the two states have orthogonal supports, there is a trade-off between the two error probabilities, and it is not possible to find a measurement that makes both error probabilities equal to zero. As it turns out, if we require the error probabilities of the first kind to go to zero asymptotically then, under an optimal sequence of measurements, the error probabilities of the second kind decay exponentially, and the decay rate is given by $S(\rho || \sigma)$ [1], [2]. On the other hand, if we impose the stronger condition that the error probabilities of the first kind go to zero asymptotically as $\alpha_n \sim 2^{-nr}$ for some r > 0then, under an optimal sequence of measurements, the error probabilities of the second kind decay as $\beta_n \sim 2^{-nH_r(\rho || \sigma)}$, where $H_r(\rho || \sigma)$ is the *Hoeffding distance* of ρ and σ with parameter r [3]–[6].

The Hoeffding distances can be obtained as a certain transform of the α -relative entropies that were defined by Rényi, based on purely axiomatic considerations [7]. While the above state discrimination result relates Rényi's α -relative entropies to statistical distances with operational interpretation, a direct operational interpretation of the Rényi relative entropies was missing for a long time. This gap was filled in the classical case by Csiszár [8], who defined the operational notion of cutoff rates and showed that the α -relative entropies arise as cutoff rates in state discrimination problems. In Section III we follow Csiszár's approach to show that the α -relative entropies can be given the same operational interpretation in the quantum case, at least for the parameter range $\alpha \in (0,1)$.

Given a state shared by several parties, and a statistical distance D, the D-distance of the state from the set of uncorrelated states yields a measure of correlations among the parties. For instance, a popular measure of quantum correlations is the *relative entropy of entanglement* [9], which is the relative entropy distance of a multipartite quantum state from the set of separable (i.e., only classically correlated) states. Similarly, a measure of the total amount of correlations between parties A and B sharing a bipartite quantum state ρ_{AB} , can be defined

by the D-distance of ρ_{AB} from the set of product states,

$$I_D(A:B \mid \rho_{AB}) := \inf_{\sigma_A \in \mathcal{S}(\mathcal{H}_A), \sigma_B \in \mathcal{S}(\mathcal{H}_B)} D\left(\rho_{AB} \mid\mid \sigma_A \otimes \sigma_B\right),$$

where $\mathcal{S}(\mathcal{H}_A)$ and $\mathcal{S}(\mathcal{H}_B)$ denote the state spaces of parties A and B, respectively. When the statistical distance is the relative entropy S, there is a unique product state closest to ρ_{AB} , which is the product $\rho_A \otimes \rho_B$ of the marginals of ρ_{AB} , and we have the identities

$$I_{S}(A:B \mid \rho_{AB}) = S(\rho_{AB} \mid\mid \rho_{A} \otimes \rho_{B})$$

$$= \inf_{\sigma_{A} \in \mathcal{S}(\mathcal{H}_{A})} S(\rho_{AB} \mid\mid \sigma_{A} \otimes \rho_{B})$$

$$= \inf_{\sigma_{B} \in \mathcal{S}(\mathcal{H}_{B})} S(\rho_{AB} \mid\mid \rho_{A} \otimes \sigma_{B}).$$
(1)

These identities, however, are not valid any longer if S is replaced with some other statistical distance D, and one may wonder which formula gives the "right" measure of correlations, i.e., which one admits an operational interpretation. When D is an α -relative entropy or a Hoeffding distance, an operational interpretation can be obtained for $D(\rho_{AB} || \rho_A \otimes$ ρ_B) in the setting of discriminating ρ_{AB} from $\rho_A \otimes \rho_B$, as described above. It seems, however, that when D is an α relative entropy and the aim is to measure correlations between the input and the output of a stochastic communication channel then it is the last formula in (1) (with S replaced with an α relative entropy) that yields a natural operational interpretation, as we will see below.

By a classical-quantum communication channel (or simply a channel) we mean a map $W: \mathcal{X} \to \mathcal{S}(\mathcal{H})$, where \mathcal{X} is a set and \mathcal{H} is a Hilbert space, which we assume to be finitedimensional. Note that there is no restriction on the cardinality of \mathcal{X} , and this formulation encompasses both the case of classical channels (i.e., when the range of W is commutative) and the standard formalism for quantum channels (i.e., when \mathcal{X} is the state space of an input Hilbert space and W is a completely positive trace-preserving map). A "lifting" of the channel can be defined by $\hat{W}: \mathcal{X} \to \mathcal{S}(\mathcal{H}_{\mathcal{X}} \otimes \mathcal{H}), \quad \hat{W}: x \mapsto \delta_x \otimes W_x$, where $\mathcal{H}_{\mathcal{X}}$ is some auxiliary Hilbert space with dimension equal to the cardinality of \mathcal{X} , and $\delta_x := |e_x\rangle\langle e_x|$ for some orthonormal system $\{e_x\}_{x\in\mathcal{X}}$ in $\mathcal{H}_{\mathcal{X}}$. The expectation value of \hat{W} with respect to a finitely supported probability measure $p \in$ $\mathcal{M}_f(\mathcal{X})$ is a classical-quantum state $\mathbb{E}_p \hat{W} = \sum_x p(x) \delta_x \otimes W_x$ on the joint system of the input and the output of the channel, and its marginals are given by $\operatorname{Tr}_{\mathcal{H}} \mathbb{E}_p \hat{W} = \hat{p} := \sum_x p(x) \delta_x$ and $\operatorname{Tr}_{l^2(\mathcal{X})} \mathbb{E}_p \hat{W} = \mathbb{E}_p W = \sum_x p(x) W_x$. The amount of correlations between the input and the output in the state $\mathbb{E}_p \hat{W}$, as measured by the relative entropy, can be written in various equivalent ways:

$$I_{S}(p; W)$$

$$:= S\left(\mathbb{E}_{p} \hat{W} || \hat{p} \otimes \mathbb{E}_{p} W\right) = \inf_{\sigma \in S(\mathcal{H})} S\left(\mathbb{E}_{p} \hat{W} || \hat{p} \otimes \sigma\right) \quad (2)$$

$$= \sum_{x} p(x) S\left(W_{x} || \mathbb{E}_{p} W\right) = \inf_{\sigma \in \mathcal{S}(\mathcal{H})} \sum_{x} p(x) S\left(W_{x} || \sigma\right) \quad (3)$$

$$= S(\mathbb{E}_p W) - \sum p(x)S(W_x). \tag{4}$$

The Holevo-Schumacher-Westmoreland theorem [10], [11] shows that the asymptotic information transmission capacity of a channel, under the assumption of product encoding, is given by the Holevo capacity

$$\chi_S^*(W) := \sup_{p \in \mathcal{M}_f(\mathcal{X})} I_S(p; W), \tag{5}$$

2

which is the maximal amount of correlation that can be created between the classical input and the quantum output in a classical-quantum state of the form $\mathbb{E}_p W$, $p \in \mathcal{M}_f(W)$. A geometric interpretation of the Holevo capacity was given in [12], where it was shown that the Holevo capacity of a channel W is equal to the relative entropy radius $R_S(\operatorname{ran} W)$ of its range, where the *D*-radius of a subset $\Sigma \subset \mathcal{S}(\mathcal{H})$ for a statistical distance D is defined as

$$R_D(\Sigma) := \inf_{\sigma \in \mathcal{S}(\mathcal{H})} \sup_{\rho \in \Sigma} D(\rho || \sigma). \tag{6}$$

Not so suprisingly, the identities in (2)–(4) do not hold for a general statistical distance D, and one may define various formal generalizations of the Holevo capacity. Here we will be interested in the quantities

$$\chi_{D,0}^*(W) := \sup_{p \in \mathcal{M}_f(\mathcal{X})} D(\mathbb{E}_p \hat{W} || \hat{p} \otimes \mathbb{E}_p W), \tag{7}$$

$$\chi_{D,1}^*(W) := \sup_{p \in \mathcal{M}_f(\mathcal{X})} \inf_{\sigma \in \mathcal{S}(\mathcal{H})} D(\mathbb{E}_p \hat{W} || \hat{p} \otimes \sigma), \tag{8}$$

$$\chi_{D,2}^{*}(W) := \sup_{p \in \mathcal{M}_{f}(\mathcal{X})} \inf_{\sigma \in \mathcal{S}(\mathcal{H})} \sum_{x \in \mathcal{X}} p(x) D(W_{x} || \sigma), \quad (9)$$

$$R_{D}(\operatorname{ran} W) := \inf_{\sigma \in \mathcal{S}(\mathcal{H})} \sup_{x \in \mathcal{X}} D(W_{x} || \sigma). \quad (10)$$

$$R_D(\operatorname{ran} W) := \inf_{\sigma \in \mathcal{S}(\mathcal{H})} \sup_{x \in \mathcal{X}} D(W_x \mid\mid \sigma). \tag{10}$$

The capacities $\chi_{D,1}^*(W), \chi_{D,2}^*(W)$ and $R_D(\operatorname{ran} W)$ were shown to be equal in [8] when the channel is classical and D is an α -relative entropy S_{α} with arbitrary non-negative parameter α , and in [13], the identity $\chi_{S_{\alpha},1}^*(W) = R_{S_{\alpha}}(\operatorname{ran} W)$ was shown for quantum channels and $\alpha \in (1, +\infty)$. In Section IV we follow the approach of [8] to show that $\chi_{D,1}^*(W) =$ $\chi_{D,2}^*(W) = R_D(\operatorname{ran} W)$ for classical-quantum channels when D is an α -relative entropy with parameter $\alpha \in (0, 2]$.

The Holevo-Schumacher-Westmoreland theorem identifies the Holevo capacity (5) as the optimal rate of information transmission through the channel in an asymptotic scenario, under the assumption that the noise described by the channel occurs independently at consecutive uses of the channel (memoryless channel). However, in practical applications one can use a channel only finitely many times, and the memoryless condition might not always be realistic, either. Hence, it is desirable to have bounds on the information transmission capacity of a channel for finitely many uses. For a given threshold $\varepsilon > 0$, the one-shot ε -capacity of the channel is the maximal number of bits that can be transmitted by one single use of the channel, with an average error not exceeding ε . Note that finitely many (possibly correlated) uses of a channel can be described as the action of one single channel acting on sequences of inputs, and hence the study of one-shot capacities addresses the generalization of coding theorems in the direction of finitely many uses and possibly correlated channels at the same time. In [14] a lower bound on the one-shot ε -capacity of an arbitrary classical-quantum channel

W was given in terms of the Rényi capacities $\chi_{S_{\alpha},0}^*(W)$ with parameter $\alpha \in [0,1)$. This bound was shown to be asymptotically optimal in the sense of yielding the Holevo capacity as a lower bound in the asymptotic limit, but no upper bound of similar form has been known up till now. In Section V we show an upper bound on the one-shot ε -capacity in terms of the Rényi capacities $\chi_{S_{\alpha},1}^*(W)$ with parameter $\alpha > 1$ that is again asymptotically optimal in the above sense. It remains an open question whether the capacities $\chi_{S_{\alpha},0}^*(W)$ and $\chi_{S_{\alpha},1}^*(W)$ are equal for a given α . To the best of our knowledge, the answer to this question is unknown even in the classical case.

II. PRELIMINARIES ON THE RÉNYI RELATIVE ENTROPIES

Let \mathcal{H} be a finite-dimensional Hilbert space with d := $\dim \mathcal{H}$. We will use the notations $\mathcal{B}(\mathcal{H})_+$ and $\mathcal{B}(\mathcal{H})_{++}$ to denote the positive semidefinite and the strictly positive definite operators on \mathcal{H} , respectively. Similarly, we denote the set of density operators (positive semidefinite operators with unit trace) by $S(\mathcal{H})$, and use the notation $S(\mathcal{H})_{++}$ for the set of invertible density operators. We will use the conventions $0^{\alpha} := 0, \ \alpha \in \mathbb{R}, \ \text{and} \ \log 0 := -\infty, \ \log +\infty := +\infty.$ By the former, powers of a positive semidefinite operator are only taken on its support, i.e., if the spectral decomposition of an $A \in \mathcal{B}(\mathcal{H})_+$ is $A = \sum_k a_k P_k$, where all $a_k > 0$, then $A^{\alpha} := \sum_k a_k^{\alpha} P_k$ for all $\alpha \in \mathbb{R}$. In particular, A^0 is the projection onto the support of A.

Following [15], we define for every $\alpha \in [0, +\infty) \setminus \{1\}$ the α -quasi-relative entropy of an $A \in \mathcal{B}(\mathcal{H})_+$ with respect to a $B \in \mathcal{B}(\mathcal{H})_+$ as

$$\begin{aligned} Q_{\alpha}\left(A \mid\mid B\right) \\ := \begin{cases} \operatorname{sign}(\alpha - 1)\operatorname{Tr} A^{\alpha}B^{1 - \alpha}, & \operatorname{supp} A \leq \operatorname{supp} B \\ & \text{or } \alpha \in [0, 1), \\ +\infty, & \text{otherwise.} \end{cases} \end{aligned}$$

The Rényi α -relative entropy of A with respect to B is then defined as

$$S_{\alpha}(A || B) := \frac{1}{\alpha - 1} \log \operatorname{sign}(\alpha - 1) Q_{\alpha}(A || B).$$

Note that $S_{\alpha}(A||B) = +\infty$ if supp $A \perp \text{supp } B$, or if supp $A \nleq \text{supp } B \text{ and } \alpha > 1$. In all other cases, $S_{\alpha}(A || B)$ is a finite number, given by $S_{\alpha}(A || B) = \frac{1}{\alpha - 1} \log \operatorname{Tr} A^{\alpha} B^{1 - \alpha}$. Note that for $\alpha \in (0,1)$, we have

$$S_{1-\alpha}(A \parallel B) = \frac{1-\alpha}{\alpha} S_{\alpha}(B \parallel A). \tag{11}$$

It is easy to see that if $\operatorname{Tr} A = 1$ then

$$S_1(A || B) := \lim_{\alpha \to 1} S_{\alpha}(A || B) = S(A || B)$$

where S(A||B) is the *relative entropy*

$$S\left(A \mid\mid B\right) := \begin{cases} \operatorname{Tr} A(\log A - \log B), & \operatorname{supp} A \leq \operatorname{supp} B, \\ +\infty, & \text{otherwise}. \end{cases}$$

Operator monotonicity of the function $x \mapsto x^{1-\alpha}, x > 0$. for $\alpha \in [0,1]$ yields that

$$\begin{aligned} Q_{\alpha}\left(A \mid\mid B+C\right) &\leq Q_{\alpha}\left(A \mid\mid B\right) \quad \text{and} \\ S_{\alpha}\left(A \mid\mid B+C\right) &\leq S_{\alpha}\left(A \mid\mid B\right) \end{aligned}$$

for any $A, B, C \in \mathcal{B}(\mathcal{H})_+$ and $\alpha \in [0, 1]$, and the same holds for $\alpha > 1$ if B and C commute. In particular, for fixed $A, B \in$ $\mathcal{B}(\mathcal{H})_+$, the maps $0 < \varepsilon \mapsto Q_{\alpha}(A || B + \varepsilon I)$ and $0 < \varepsilon \mapsto$ $S_{\alpha}(A || B + \varepsilon I)$ are monotonic decreasing, and it is easy to see that, for any $\alpha \in [0, +\infty)$,

$$Q_{\alpha}(A || B) = \sup_{\varepsilon > 0} Q_{\alpha}(A || B + \varepsilon I), \qquad (12)$$

$$S_{\alpha}(A || B) = \sup_{\varepsilon > 0} S_{\alpha}(A || B + \varepsilon I). \qquad (13)$$

$$S_{\alpha}(A || B) = \sup_{\varepsilon > 0} S_{\alpha}(A || B + \varepsilon I). \tag{13}$$

For $\alpha \in [0,2] \setminus \{1\}$, the α -quasi-relative entropies have the monotonicity property [15]–[17]

$$Q_{\alpha}\left(\Phi(A) \mid\mid \Phi(B)\right) \le Q_{\alpha}\left(A \mid\mid B\right), \quad A, B \in \mathcal{B}(\mathcal{H})_{+}, \quad (14)$$

where Φ is any completely positive trace-preserving (CPTP) map on $\mathcal{B}(\mathcal{H})$. As a consequence, the α -quasi-relative entropies are jointly convex in their arguments for $\alpha \in [0, 2] \setminus$ $\{1\}$:

$$Q_{\alpha}\left(\sum_{i} p_{i} A_{i} \mid\mid \sum_{i} p_{i} B_{i}\right) \leq \sum_{i} p_{i} Q_{\alpha}\left(A_{i} \mid\mid B_{i}\right), \quad (15)$$

where $A_i, B_i \in \mathcal{B}(\mathcal{H})_+$, and $\{p_i\}$ is a finite probability distribution [15], [18], [19].

The monotonicity property (14) of the α -quasi-relative entropies yields that, for any CPTP map Φ on $\mathcal{B}(\mathcal{H})$ and $\alpha \in [0,2],$

$$S_{\alpha}\left(\Phi(A) \mid\mid \Phi(B)\right) \leq S_{\alpha}\left(A \mid\mid B\right), \quad A, B \in \mathcal{B}(\mathcal{H})_{+}.$$

Convexity of the function $\frac{1}{\alpha-1}\log$ for $\alpha\in[0,1)$ yields, by (15), that for $\alpha \in [0, 1]$,

$$S_{\alpha}\left(\sum_{i} p_{i} A_{i} \mid\mid \sum_{i} p_{i} B_{i}\right) \leq \sum_{i} p_{i} S_{\alpha}\left(A_{i} \mid\mid B_{i}\right) \quad (16)$$

for any finite probability distribution $\{p_i\}$ and $A_i, B_i \in$ $\mathcal{B}(\mathcal{H})_+$. Note that the joint convexity (15) of the α -quasirelative entropies for $\alpha \in (1,2]$ is not inherited by the corresponding Rényi relative entropies, as $\frac{1}{\alpha-1}\log$ is not convex for $\alpha > 1$; for a counterexample, see e.g. [20]. Actually, the example of [20] shows that the Rényi relative entropies are not even convex in their first argument for $\alpha > 1$. However, we have the following:

Theorem II.1. For a fixed $A \in \mathcal{B}(\mathcal{H})_+$, the map $B \mapsto$ $S_{\alpha}(A || B)$ is convex on $\mathcal{B}(\mathcal{H})_+$ for every $\alpha \in [0, 2]$.

Proof: For $\alpha \in [0,1]$, the assertion is a weaker version of (16), and hence for the rest we assume that $\alpha \in (1, 2]$. Let $A, B_1, B_2 \in \mathcal{B}(\mathcal{H})_+$; it suffices to show that

$$S_{\alpha}\left(A \mid\mid \eta(B_{1} + \varepsilon I) + (1 - \eta)(B_{2} + \varepsilon I)\right)$$

$$\leq \eta S_{\alpha}\left(A \mid\mid B_{1} + \varepsilon I\right) + (1 - \eta)S_{\alpha}\left(A \mid\mid B_{2} + \varepsilon I\right) \quad (17)$$

holds for every $\eta \in (0,1)$. Taking the limit $\varepsilon \searrow 0$ will then give the desired convexity inequality. Note that (17) is equivalent to

$$\log \omega (\eta(B_1 + \varepsilon I) + (1 - \eta)(B_2 + \varepsilon I)^{1 - \alpha})$$

$$\leq \eta \log \omega ((B_1 + \varepsilon I)^{1 - \alpha}) + (1 - \eta) \log \omega ((B_2 + \varepsilon I)^{1 - \alpha}),$$

where $\omega(X) := \operatorname{Tr} A^{\alpha}X, X \in \mathcal{B}(\mathcal{H})$, is a positive linear functional on $\mathcal{B}(\mathcal{H})$. Proposition 1.1 in [21] states that the functional $X \mapsto \log \omega(f(X)), X \in \mathcal{B}(\mathcal{H})_{++}$, is convex whenever ω is a positive linear functional and f is a nonnegative operator monotone decreasing function on $(0, +\infty)$. Applying this to the ω above and $f(x) := x^{1-\alpha}, \, x > 0$, the assertion follows.

By computing its second derivative, it is easy to see that the function $\alpha \mapsto \log \operatorname{Tr} A^{\alpha} B^{1-\alpha}$, $\alpha \in \mathbb{R}$, is convex on \mathbb{R} for any fixed $A, B \in \mathcal{B}(\mathcal{H})_+$, which yields by a simple computation the following:

Lemma II.2. If $\operatorname{Tr} A \leq 1$ then the function $\alpha \mapsto S_{\alpha}(A \mid\mid B)$ is monotonically increasing on [0,1) and on $(1,+\infty)$. Moreover, if $\operatorname{Tr} A = 1$ then $\alpha \mapsto S_{\alpha}(A \mid\mid B)$ is monotonically increasing on $[0,+\infty)$.

Proposition II.3. Assume that $\operatorname{Tr} A \leq 1$ and $\operatorname{Tr} B \leq 1$. For $\alpha \in (0,1)$, $S_{\alpha}(A \mid\mid B) \geq 0$ with equality if and only if A=B and $\operatorname{Tr} A=1$. If A is a density operator and $\operatorname{Tr} B \leq 1$ then, for all $\alpha \in [1, +\infty)$, $S_{\alpha}(A \mid\mid B) \geq 0$, and $S_{\alpha}(A \mid\mid B) = 0$ if and only if A=B. Moreover, if both A and B are density operators then the Csiszár-Pinsker inequality

$$S_{\alpha}(A || B) \ge \frac{1}{2} ||A - B||_{1}^{2}$$

holds for all $\alpha > 1$.

Proof: Assume first that $\alpha \in [0,1)$. Then, by Hölder's inequality,

$$\operatorname{Tr} A^{\alpha} B^{1-\alpha} \leq (\operatorname{Tr} A)^{\alpha} (\operatorname{Tr} B)^{1-\alpha} \leq 1,$$

from which $S_{\alpha}\left(A \mid\mid B\right) = \frac{1}{\alpha-1}\log\operatorname{Tr}A^{\alpha}B^{1-\alpha} \geq 0$. Obviously, $S_{\alpha}\left(A \mid\mid B\right) = 0$ if and only if $\operatorname{Tr}A^{\alpha}B^{1-\alpha} = 1$. By the above, this is true if and only if $\operatorname{Tr}A = \operatorname{Tr}B = 1$, and Hölder's inequality holds with equality. The latter condition yields that $B = \lambda A$ for some $\lambda \geq 0$, and $\operatorname{Tr}A = \operatorname{Tr}B$ yields $\lambda = 1$. Lemma II.2 yields the assertion on strict positivity for $\alpha \geq 1$ when A is a density operator. The Csiszár-Pinsker inequality holds for $\alpha = 1$ (cf. Theorem 3.1 in [22]) and hence, by Lemma II.2, for all $\alpha > 1$.

For a density operator $\rho \in \mathcal{S}(\mathcal{H})$, its *Rényi* α -entropy for $\alpha \in [0, +\infty)$ is

$$S_{\alpha}(\rho) := \log d - S_{\alpha}(\rho || (1/d)I).$$

For $\alpha \neq 1$ we have $S_{\alpha}(\rho) = \frac{1}{1-\alpha} \log \operatorname{Tr} \rho^{\alpha}$, which is easily seen to be non-negative, and $S_{\alpha}(\rho \mid |(1/d)I) \geq 0$ yields that

$$0 \le S_{\alpha}(\rho) \le \log d, \qquad \alpha \in [0, +\infty).$$
 (18)

The *Hoeffding distance* of states $\rho, \sigma \in \mathcal{S}(\mathcal{H})$ with parameter $r \geq 0$ is defined as

$$H_{r}(\rho || \sigma) := \sup_{0 \leq \alpha < 1} \left\{ \frac{-\alpha r}{1 - \alpha} + S_{\alpha}(\rho || \sigma) \right\}$$

$$= \sup_{0 \leq \alpha < 1} \frac{-\alpha r - \psi(\alpha)}{1 - \alpha} = \sup_{s \geq 0} \{-sr - \tilde{\psi}(s)\},$$
(19)

where

$$\psi(\alpha) := \log \operatorname{Tr} \rho^{\alpha} \sigma^{1-\alpha}, \quad \alpha \in \mathbb{R},$$

$$\tilde{\psi}(s) := (1+s)\psi\left(s/(1+s)\right), \quad s > -1. \tag{20}$$

Convexity of ψ yields the convexity of $\tilde{\psi}$, and a simple computation shows that $\psi(0) + \psi'(0) = \tilde{\psi}'(0) \leq \lim_{s \to \infty} \tilde{\psi}'(s) = \psi(1) \leq 0$. Hence,

$$H_r(\rho || \sigma) = \begin{cases} -\tilde{\psi}(0) = -\psi(0), & -r \le \psi(0) + \psi'(0), \\ +\infty, & -r > \psi(1). \end{cases}$$

The function $r \mapsto H_r(\rho || \sigma)$ is the Legendre-Fenchel transform (up to the sign of the variable) of $\tilde{\psi}$ on $[0, +\infty)$ and hence it is convex on $[0, +\infty)$. Using the bipolar theorem for convex functions [23, Proposition 4.1], we get

$$S_{\alpha}\left(\rho \mid\mid \sigma\right) = -\sup_{r>0} \left\{ \frac{-r\alpha}{1-\alpha} - H_r\left(\rho \mid\mid \sigma\right) \right\}, \quad 0 \le \alpha < 1.$$

That is, the Rényi relative entropies with parameter in [0,1) and the Hoeffding distances with parameter $r\geq 0$ mutually determine each other. Note that $r\mapsto H_r\left(\rho\,||\,\sigma\right)$ is monotonic decreasing, and

$$S_{0}\left(\rho \mid\mid \sigma\right) = \lim_{r \to \infty} H_{r}\left(\rho \mid\mid \sigma\right) \leq H_{0}\left(\rho \mid\mid \sigma\right) = S_{1}\left(\rho \mid\mid \sigma\right).$$

Finally, the max-relative entropy of $A, B \in \mathcal{S}(\mathcal{H})_+$ was defined in [24] as $S_{\max}(A \mid\mid B) := \inf\{\gamma: A \leq 2^{\gamma}B\}$. One can easily see that if A and B commute then $S_{\max}(A \mid\mid B) = S_{\infty}(A \mid\mid B) := \lim_{\alpha \to \infty} S_{\alpha}(A \mid\mid B)$, but for non-commuting A and B, $S_{\max}(A \mid\mid B) < S_{\infty}(A \mid\mid B)$ might happen [14]. In general, $S_2(A \mid\mid B) \leq S_{\max}(A \mid\mid B) \leq S_{\infty}(A \mid\mid B)$ [25], [26].

III. CUTOFF RATES FOR QUANTUM STATE DISCRIMINATION

Consider the asymptotic binary state discrimination problem with null hypothesis ρ and alternative hypothesis σ , as described in the Introduction. We will consider the scenario where the error probability of the second kind is minimized under an exponential constraint on the error probability of the first kind; the quantity of interest in this case is

$$\beta_{n,r} := \min\{\beta_n(T) \mid T \in \mathcal{B}(\mathcal{H}^{\otimes n}), \ 0 \le T \le I,$$
 and $\alpha_n(T) < 2^{-nr}\},$

where r is some fixed positive number. In general, there is no closed formula to express $\beta_{n,r}$ or the optimal measurement in terms of ρ and σ for a finite n, but it becomes possible in the limit of large n. We define the *Hoeffding exponents* for a parameter r>0 as

$$\underline{h}_{r}\left(\rho \mid\mid \sigma\right) := \inf_{\{T_{n}\}} \left\{ \liminf_{n \to \infty} (1/n) \log \beta_{n}(T_{n}) \mid \\ \limsup_{n \to \infty} (1/n) \log \alpha_{n}(T_{n}) < -r \right\},$$

$$\overline{h}_{r}\left(\rho \mid\mid \sigma\right) := \inf_{\{T_{n}\}} \left\{ \limsup_{n \to \infty} (1/n) \log \beta_{n}(T_{n}) \mid \\ \limsup_{n \to \infty} (1/n) \log \alpha_{n}(T_{n}) < -r \right\},$$

$$h_{r}\left(\rho \mid\mid \sigma\right) := \inf_{\{T_{n}\}} \left\{ \lim_{n \to \infty} (1/n) \log \beta_{n}(T_{n}) \mid \\ \limsup_{n \to \infty} (1/n) \log \alpha_{n}(T_{n}) < -r \right\}.$$

It is easy to see that

$$\underline{h}_{r}(\rho || \sigma) \leq \liminf_{n \to \infty} \frac{1}{n} \log \beta_{n,r}
\leq \limsup_{n \to \infty} \frac{1}{n} \log \beta_{n,r} \leq \overline{h}_{r}(\rho || \sigma).$$

Moreover, as it was shown in [3]-[6], we have

$$\underline{h}_{r}(\rho \mid\mid \sigma) = \overline{h}_{r}(\rho \mid\mid \sigma) = h_{r}(\rho \mid\mid \sigma) = -H_{r}(\rho \mid\mid \sigma), \quad (21)$$

where $H_r(\rho || \sigma)$ is the Hoeffding distance defined in (19), and hence, the limit $\lim_{n\to\infty} \frac{1}{n} \log \beta_{n,r}$ exists and

$$\lim_{n \to \infty} \frac{1}{n} \log \beta_{n,r} = -H_r \left(\rho \mid\mid \sigma \right).$$

Note that while the above result gives the exact value of the optimal exponential decay rate for every r, the evaluation of $H_r\left(\rho\mid\mid\sigma\right)$ is a non-trivial task even for one single r. Indeed, there is no closed formula known for the Hoeffding distance in general, and, as the definition (19) shows, in order to compute $H_r\left(\rho\mid\mid\sigma\right)$, one has to know in principle all the Rényi relative entropies $S_\alpha\left(\rho\mid\mid\sigma\right)$ for every $\alpha\in(0,1)$, and solve an optimization problem. It is thus natural to look for simple approximants of the function $r\mapsto H_r\left(\rho\mid\mid\sigma\right)$ for given ρ and σ . Following [8], for a $\kappa<0$ we define the *generalized* κ -cutoff rate $C_\kappa\left(\rho\mid\mid\sigma\right)$ as the supremum of all $r_0\geq0$ that satisfy

$$\overline{h}_r(\rho || \sigma) \le \kappa(r_0 - r), \qquad r \ge 0.$$
 (22)

That is, we are looking for a linear approximation of $r\mapsto H_r\left(\rho\,||\,\sigma\right)$ which is optimal among all the linear functions with a given slope. Note that (22) gives a restriction only for $r\le r_0$, as otherwise the right-hand side is non-negative and the inequality holds trivially. That is, one can ensure an exponential decay rate at least as fast as given in the right-hand side of (22) whenever $r< r_0:=C_\kappa\left(\rho\,||\,\sigma\right)$. Moreover, as the following Theorem shows, the cutoff rate is easy to evaluate, as it is equal to a Rényi relative entropy with a given parameter depending on κ .

Theorem III.1. For every $\kappa < 0$,

$$C_{\kappa}\left(\rho \mid\mid \sigma\right) = \frac{1}{|\kappa|} S_{\frac{|\kappa|}{1+|\kappa|}}\left(\rho \mid\mid \sigma\right) = S_{\frac{1}{1+|\kappa|}}\left(\sigma \mid\mid \rho\right). \tag{23}$$

Proof: If supp $\rho \perp$ supp σ then all the quantities in (23) are $+\infty$ and the assertion holds trivially. Hence, for the rest we assume that supp ρ is not orthogonal to supp σ . Note that the second identity follows from (11). Let $\kappa < 0$ be fixed. By (21), our goal is to determine the largest r_0 such that

$$-|\kappa|r + |\kappa|r_0 \le -\overline{h}_r(\rho||\sigma) = H_r(\rho||\sigma), \qquad r \ge 0.$$

By (19), $H_r(\rho||\sigma) \ge -|\kappa|r - \tilde{\psi}(|\kappa|)$ for every $r \ge 0$, where $\tilde{\psi}$ is given in (20). On the other hand, for $r_{\kappa} := -\tilde{\psi}'(|\kappa|)$ we have $\tilde{\psi}(s) \ge \tilde{\psi}(|\kappa|) + (s - |\kappa|)\tilde{\psi}'(|\kappa|)$, $s \ge 0$, due to the convexity of $\tilde{\psi}$ and hence,

$$H_{r_{\kappa}}(\rho \mid\mid \sigma) = \sup_{s \geq 0} \{s\tilde{\psi}'(|\kappa|) - \tilde{\psi}(s)\} = |\kappa|\tilde{\psi}'(|\kappa|) - \tilde{\psi}(|\kappa|)$$
$$= -|\kappa|r_{\kappa} - \tilde{\psi}(|\kappa|).$$

Therefore,

$$C_{\kappa} \left(\rho \mid\mid \sigma \right) = -\frac{1}{|\kappa|} \tilde{\psi}(|\kappa|) = -\frac{1+|\kappa|}{|\kappa|} \psi \left(\frac{|\kappa|}{1+|\kappa|} \right)$$
$$= \frac{1}{|\kappa|} S_{\frac{|\kappa|}{1+|\kappa|}} \left(\rho \mid\mid \sigma \right).$$

The following Corollary is immediate from Theorem III.1, and gives an operational interpretation of the Rényi relative entropies with parameter between 0 and 1:

Corollary III.2. For every $\rho, \sigma \in \mathcal{S}(\mathcal{H})$ and every $\alpha \in (0,1)$,

$$S_{\alpha}\left(\rho\mid\mid\sigma\right) = \frac{\alpha}{1-\alpha} C_{\frac{\alpha}{\alpha-1}}\left(\rho\mid\mid\sigma\right) = C_{\frac{\alpha-1}{\alpha}}\left(\sigma\mid\mid\rho\right).$$

In the above, we considered the scenario where the consecutive trials are independent and identically distributed, and hence the state describing the outcome probabilities of n trials is a state of the form $\rho^{\otimes n}$ or $\sigma^{\otimes n}$. In a more general scenario, that encompasses correlated trials, one can consider a sequence of Hilbert spaces $\vec{\mathcal{H}} := \{\mathcal{H}_n\}_{n\in\mathbb{N}}$ and two sequences of states $\vec{\rho} := \{\rho_n\}_{n\in\mathbb{N}}$ and $\vec{\sigma} := \{\sigma_n\}_{n\in\mathbb{N}}$. The goal is again to analyze the asymptotic performance of a decision scheme for deciding between ρ_n and σ_n for each $n \in \mathbb{N}$. The error probabilities α_n and β_n can be defined in the same way as above, and in analogy with the above problem, the limit $\lim_{n\to\infty} (1/c(n)) \log \beta_{n,r}$ can be considered, where $c: \mathbb{N} \to \mathbb{N}$ is some monotonically increasing function such that $\lim_{n\to\infty} c(n) = +\infty$. The following was shown in [6]:

Theorem III.3. Assume that the limit $\psi(\alpha) := \lim_{n\to\infty} \frac{1}{c(n)} (\alpha-1) S_{\alpha} (\rho_n || \sigma_n)$ exists for all $\alpha \in [0,1)$ and the convergence is uniform on [0,1). Assume, moreover, that ψ is differentiable on (0,1). Then,

$$\lim_{n \to \infty} \frac{1}{c(n)} \log \beta_{n,r} = -\lim_{n \to \infty} \frac{1}{c(n)} H_{c(n)r} \left(\rho_n \mid\mid \sigma_n \right)$$
$$=: -H_r \left(\vec{\rho} \mid\mid \vec{\sigma} \right).$$

Moreover, $H_r\left(\vec{\rho} \mid\mid \vec{\sigma}\right) = \sup_{0 \leq \alpha < 1} \left\{ \frac{-\alpha r}{1-\alpha} + \frac{\psi(\alpha)}{\alpha - 1} \right\}$, where

$$\frac{\psi(\alpha)}{\alpha - 1} = S_{\alpha}\left(\vec{\rho} \mid\mid \vec{\sigma}\right) := \lim_{n \to \infty} \frac{1}{c(n)} S_{\alpha}\left(\rho_n \mid\mid \sigma_n\right).$$

A particular example that satisfies the conditions of Theorem III.3 is the case where ρ_n and σ_n are the n-step restrictions of classical ergodic Markov chains with finite state-space [6]. Physically motivated examples can be obtained by considering ρ_n and σ_n to be finite-block restrictions of temperature states of non-interacting fermionic and bosonic systems on cubic lattices [27], [28].

The cutoff rates $C_{\kappa}(\vec{\rho}||\vec{\sigma})$ can again be defined in the same way as in (22) (with the scale 1/n replaced with 1/c(n) in the definition of $\overline{h}_r(\vec{\rho}||\vec{\sigma})$). The same argument as in the proof of Theorem III.1 leads to the following:

Theorem III.4. Under the assumptions of Theorem III.3, we have

$$C_{\kappa}\left(\vec{\rho}\,||\,\vec{\sigma}\right) = \frac{1}{|\kappa|} S_{\frac{|\kappa|}{1+|\kappa|}}\left(\vec{\rho}\,||\,\vec{\sigma}\right) = S_{\frac{1}{1+|\kappa|}}\left(\vec{\sigma}\,||\,\vec{\rho}\right)$$

for every $\kappa < 0$, or equivalently, for every $\alpha \in (0,1)$,

$$S_{\alpha}\left(\vec{\rho}\,||\,\vec{\sigma}\right) = \frac{\alpha}{1-\alpha}C_{\frac{\alpha}{\alpha-1}}\left(\vec{\rho}\,||\,\vec{\sigma}\right) = C_{\frac{\alpha-1}{\alpha}}\left(\vec{\sigma}\,||\,\vec{\rho}\right).$$

IV. EQUIVALENCE OF CAPACITIES

Let $W: \mathcal{X} \to \mathcal{S}(\mathcal{H})$ be a classical-quantum channel as in the Introduction. Our aim in this section is to show that the capacities defined in (8)–(10) are equal to each other when $D=S_{\alpha}$ is a Rényi relative entropy with parameter $\alpha \in (0,2]$. We will assume that $\operatorname{ran} W$ is compact in $\mathcal{S}(\mathcal{H})$. This assumption is satisfied when W is a CPTP map on the state space of an input Hilbert space as well as when \mathcal{X} is a finite set.

Note that $\mathcal{S}(\mathcal{H})$ is a compact convex subset of the Euclidean space $B(\mathcal{H})_{sa}$ (with the Hilbert-Schmidt norm). Let \mathcal{K} be a compact subset of $\mathcal{S}(\mathcal{H})$ and $\mathcal{M}(\mathcal{K})$ be the set of all Borel probability measures on \mathcal{K} . Let $C_{\mathbb{R}}(\mathcal{K})$ be the real Banach space of all real continuous functions on \mathcal{K} with the sup-norm; then $\mathcal{M}(\mathcal{K})$ is identified with a w*-compact convex subset of the dual Banach space $C_{\mathbb{R}}(\mathcal{K})^*$. We also introduce the subset $\mathcal{M}_f(\mathcal{K})$ of $\mathcal{M}(\mathcal{K})$, consisting of finitely supported measures. For every $\alpha \in (0,2] \setminus \{1\}$ and $\varepsilon \geq 0$, define the functions

$$f_{\alpha,\varepsilon}(p,\sigma) := \int_{\mathcal{K}} S_{\alpha}(\rho \| \sigma + \varepsilon I) \, dp(\rho),$$

$$g_{\alpha,\varepsilon}(p,\sigma) := \int_{\mathcal{K}} Q_{\alpha}(\rho \| \sigma + \varepsilon I) \, dp(\rho).$$

 $f_{\alpha,\varepsilon}$ and $g_{\alpha,\varepsilon}$ on $\mathcal{M}(\mathcal{K}) \times \mathcal{S}(\mathcal{H})$ by

Note that for every fixed σ , the functions $S_{\alpha}\left(\cdot\mid\mid\sigma+\varepsilon I\right)$ and $Q_{\alpha}\left(\cdot\mid\mid\sigma+\varepsilon I\right)$ are continuous for $\varepsilon>0$ and, by (12) and (13), are lower semicontinuous for $\varepsilon=0$. Hence, the integrals defining $f_{\alpha,\varepsilon}$ and $g_{\alpha,\varepsilon}$ exist for all $\varepsilon\geq0$. Furthermore, by (12), (13), and Beppo Levi's theorem,

$$f_{\alpha,0}(p,\sigma) = \lim_{\varepsilon \searrow 0} f_{\alpha,\varepsilon}(p,\sigma) = \sup_{\varepsilon > 0} f_{\alpha,\varepsilon}(p,\sigma), \qquad p \in \mathcal{M}(\mathcal{K}),$$
(24)

and the same holds if we replace $f_{\alpha,0}$ with $g_{\alpha,0}$ and $f_{\alpha,\varepsilon}$ with $g_{\alpha,\varepsilon}$.

Lemma IV.1. For every $\sigma \in \mathcal{S}(\mathcal{H})$ and $\varepsilon > 0$, $f_{\alpha,\varepsilon}(\cdot,\sigma)$ and $g_{\alpha,\varepsilon}(\cdot,\sigma)$ are affine and continuous on $\mathcal{M}(\mathcal{K})$.

Proof: The claims about the affinity are obvious, and the continuity of the functions $S_{\alpha}\left(\cdot\mid\mid\sigma+\varepsilon I\right)$ and $Q_{\alpha}\left(\cdot\mid\mid\sigma+\varepsilon I\right)$ yields, by definition, that $f_{\alpha,\varepsilon}(\cdot,\sigma)$ and $g_{\alpha,\varepsilon}(\cdot,\sigma)$ are continuous in the w*-topology.

Lemma IV.2. For every $p \in \mathcal{M}(\mathcal{K})$ and $\varepsilon > 0$, $f_{\alpha,\varepsilon}(p,\cdot)$ and $g_{\alpha,\varepsilon}(p,\cdot)$ are convex and continuous on $\mathcal{S}(\mathcal{H})$.

Proof: Convexity follows from Theorem II.1 and (15). Let $\{\sigma_k\}_{k\in\mathbb{N}}$ be a sequence in $\mathcal{S}(\mathcal{H})$, converging to some $\sigma_0\in\mathcal{S}(\mathcal{H})$. Let $f_k(\rho):=\operatorname{Tr}\rho^\alpha(\sigma_k+\varepsilon I)^{1-\alpha}$ and $f(\rho):=\operatorname{Tr}\rho^\alpha(\sigma_0+\varepsilon I)^{1-\alpha}$, $\rho\in\mathcal{K}$. Since

$$|\operatorname{Tr} \rho^{\alpha} (\sigma_{k} + \varepsilon I)^{1-\alpha} - \operatorname{Tr} \rho^{\alpha} (\sigma_{0} + \varepsilon I)^{1-\alpha}|$$

$$\leq \operatorname{Tr} \rho^{\alpha} \cdot ||(\sigma_{k} + \varepsilon I)^{1-\alpha} - (\sigma_{0} + \varepsilon I)^{1-\alpha}||_{\infty},$$

and $\operatorname{Tr} \rho^{\alpha} \leq d$ for every $\alpha \geq 0$, we see that $\lim_{k} f_{k}(\rho) = f(\rho)$ uniformly in ρ . This yields the continuity of $g_{\alpha,\varepsilon}(p,\cdot)$.

For $\alpha \in (1,2]$, $f(\rho) \geq \operatorname{Tr} \rho^{\alpha} (1+\varepsilon)^{1-\alpha} \geq (1+\varepsilon)^{1-\alpha} d^{1-\alpha}$, due to (18). For $\alpha \in (0,1)$, the operator monotonicity of the function $x \mapsto x^{1-\alpha}, x \geq 0$, yields that $f(\rho) \geq \operatorname{Tr} \rho^{\alpha} (\varepsilon I)^{1-\alpha} \geq \varepsilon^{1-\alpha}$ for all $\rho \in \mathcal{K}$. Since

$$|f_k(\rho) - f(\rho)| = f(\rho) \left| \frac{f_k(\rho)}{f(\rho)} - 1 \right| \ge \inf_{\rho \in \mathcal{K}} f(\rho) \left| \frac{f_k(\rho)}{f(\rho)} - 1 \right|,$$

we see that $f_k(\rho)/f(\rho)$ converges to 1 uniformly in ρ as $k \to \infty$, and hence

$$S_{\alpha}(\rho \| \sigma_k + \varepsilon I) - S_{\alpha}(\rho \| \sigma_0 + \varepsilon I) = \frac{1}{\alpha - 1} \log \frac{f_k(\rho)}{f(\rho)}$$

converges to 0 uniformly in ρ , due to which $\lim_{k\to\infty} f_{\alpha,\varepsilon}(p,\sigma_k) = f_{\alpha,\varepsilon}(p,\sigma_0)$.

To simplify notation, we fix an $\alpha \in (0, 2] \setminus \{1\}$ for the rest. We have the following:

Proposition IV.3. For every $\varepsilon > 0$, there exists a $\sigma_{\varepsilon} \in \mathcal{S}(\mathcal{H})$ such that

$$\max_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,\varepsilon}(p, \sigma_{\varepsilon})
= \min_{\sigma \in \mathcal{S}(\mathcal{H})} \max_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,\varepsilon}(p, \sigma) = \max_{p \in \mathcal{M}(\mathcal{K})} \min_{\sigma \in \mathcal{S}(\mathcal{H})} f_{\alpha,\varepsilon}(p, \sigma)
= \min_{\sigma \in \mathcal{S}(\mathcal{H})} \max_{\rho \in \mathcal{K}} S_{\alpha} (\rho || \sigma + \varepsilon I) = \max_{\rho \in \mathcal{K}} S_{\alpha} (\rho || \sigma_{\varepsilon} + \varepsilon I).$$
(25)

Moreover, the same relations hold if the maxima over $\mathcal{M}(\mathcal{K})$ are replaced with maxima over $\mathcal{M}_f(\mathcal{K})$.

Proof: For a fixed σ , $f_{\alpha,\varepsilon}(\cdot,\sigma)$ is continuous and, consequently, $p\mapsto \min_{\sigma\in\mathcal{S}(\mathcal{H})}f_{\alpha,\varepsilon}(p,\sigma)$ is upper semicontinuous and therefore they reach their suprema on the compact set $\mathcal{M}(\mathcal{K})$. Moreover, $f_{\alpha,\varepsilon}(p,\sigma)\leq \sup_{\rho\in\operatorname{supp} p}S_{\alpha}\left(\rho\,||\,\sigma+\varepsilon I\right),\,p\in\mathcal{M}(\mathcal{K}),\,\sigma\in\mathcal{S}(\mathcal{H}),$ yields that the maximum of $f_{\alpha,\varepsilon}(\cdot,\sigma)$ on $\mathcal{M}(\mathcal{K})$ is reached at a Dirac probability measure and hence,

$$\max_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,\varepsilon}(p,\sigma) = \max_{\rho \in \mathcal{K}} S_{\alpha} \left(\rho \mid\mid \sigma + \varepsilon I \right)$$
$$= \max_{p \in \mathcal{M}_{f}(\mathcal{K})} f_{\alpha,\varepsilon}(p,\sigma) \tag{27}$$

for every $\sigma \in \mathcal{S}(\mathcal{H})$. Continuity of $f_{\alpha,\varepsilon}(p,\cdot)$ yields that $\sigma \mapsto \max_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,\varepsilon}(p,\sigma)$ is lower semicontinuous on $\mathcal{S}(\mathcal{H})$ and hence it reaches its infimum at some point σ_{ε} , which yields $\min_{\sigma \in \mathcal{S}(\mathcal{H})} \max_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,\varepsilon}(p,\sigma) = \max_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,\varepsilon}(p,\sigma_{\varepsilon})$. The identity of the two expressions in (25) follows by Sion's minimax theorem [29], [30], due to Lemmas IV.1 and IV.2. The formulas in (26) follow from (27). The last assertion follows from (27) and the fact that $f_{\alpha,\varepsilon}|_{\mathcal{M}_f(\mathcal{K})\times\mathcal{S}(\mathcal{H})}$ also satisfies the conditions in Sion's minimax theorem.

For the rest, for every $\varepsilon > 0$ we fix a σ_{ε} as given in Proposition IV.3. Note that the compactness of $\mathcal{S}(\mathcal{H})$ yields that there exists a sequence $\{\varepsilon_k\}_{k\in\mathbb{N}}$ and a $\sigma_0 \in \mathcal{S}(\mathcal{H})$ such that $\lim_k \varepsilon_k = 0$ and $\lim_k \sigma_{\varepsilon_k} = \sigma_0$.

Proposition IV.4. Let σ_0 be a limit point as above. Then,

$$\sup_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,0}(p, \sigma_{0})$$

$$= \min_{\sigma \in \mathcal{S}(\mathcal{H})} \sup_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,0}(p, \sigma) = \sup_{p \in \mathcal{M}(\mathcal{K})} \min_{\sigma \in \mathcal{S}(\mathcal{H})} f_{\alpha,0}(p, \sigma)$$

$$= \min_{\sigma \in \mathcal{S}(\mathcal{H})} \sup_{\rho \in \mathcal{K}} S_{\alpha}(\rho || \sigma) = \sup_{\rho \in \mathcal{K}} S_{\alpha}(\rho || \sigma_{0}). \tag{29}$$

Moreover, the same relations hold if the suprema over $\mathcal{M}(\mathcal{K})$ are replaced with suprema over $\mathcal{M}_f(\mathcal{K})$.

Proof: By (24), $f_{\alpha,0}(p,\cdot)$ is lower semicontinuous on $\mathcal{S}(\mathcal{H})$ and hence so is the function $\sigma \mapsto$ $\sup_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,0}(p,\sigma), \sigma \in \mathcal{B}(\mathcal{H})_+$. Therefore, they reach their infima on $\mathcal{S}(\mathcal{H})$. For every $k \in \mathbb{N}$,

$$\max_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,0}(p, \sigma_{\varepsilon_k} + \varepsilon_k I) = \max_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,\varepsilon_k}(p, \sigma_{\varepsilon_k})$$

$$= \max_{p \in \mathcal{M}(\mathcal{K})} \min_{\sigma \in \mathcal{S}(\mathcal{H})} f_{\alpha,\varepsilon_k}(p, \sigma)$$

$$\leq \sup_{p \in \mathcal{M}(\mathcal{K})} \min_{\sigma \in \mathcal{S}(\mathcal{H})} f_{\alpha,0}(p, \sigma),$$
(30)

where the first identity is by definition, the second is due to Proposition IV.3, and the inequality follows from (24). Furthermore,

$$\sup_{p \in \mathcal{M}(\mathcal{K})} \min_{\sigma \in \mathcal{S}(\mathcal{H})} f_{\alpha,0}(p,\sigma) \leq \min_{\sigma \in \mathcal{S}(\mathcal{H})} \sup_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,0}(p,\sigma)$$

$$\leq \sup_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,0}(p,\sigma_0)$$

$$\leq \liminf_{k \to \infty} \sup_{p \in \mathcal{M}(\mathcal{K})} f_{\alpha,0}(p,\sigma_{\varepsilon_k} + \varepsilon_k I)$$

$$\leq \sup_{p \in \mathcal{M}(\mathcal{K})} \min_{\sigma \in \mathcal{S}(\mathcal{H})} f_{\alpha,0}(p,\sigma),$$

where the first two inequalities are obvious, the third one follows from the lower semicontinuity of $\sigma \mapsto$ $\sup_{p\in\mathcal{M}(\mathcal{K})} f_{\alpha,0}(p,\sigma), \ \sigma\in\mathcal{B}(\mathcal{H})_+, \ \text{and the last inequality is}$ due to (30). This gives the identities in (28), and the identities in (29) follow the same way as in Proposition IV.3. The last assertion follows by repeating the argument above with the suprema and maxima over $\mathcal{M}(\mathcal{K})$ replaced with suprema over $\mathcal{M}_f(\mathcal{K})$.

Remark IV.5. Note that the minima over $S(\mathcal{H})$ in (28) and (29) can be replaced with infima over $S(\mathcal{H})_{++}$.

Proof: The trivial inequality $(1-\varepsilon)\sigma + \varepsilon(1/d)I \ge (1-\varepsilon)\sigma$ yields

$$S_{\alpha}\left(\rho \mid\mid (1-\varepsilon)\sigma + \varepsilon(1/d)I\right) + \log(1-\varepsilon) \le S_{\alpha}\left(\rho \mid\mid \sigma\right)$$
 (31)

for every $\varepsilon \in (0,1)$, $\rho \in \mathcal{K}$ and $\sigma \in \mathcal{B}(\mathcal{H})$, and hence, for every $p \in \mathcal{M}(\mathcal{K})$,

$$f_{\alpha,0}(p,(1-\varepsilon)\sigma + \varepsilon(1/d)I) + \log(1-\varepsilon) < f_{\alpha,0}(p,\sigma).$$
 (32)

Thus,

$$\inf_{\sigma \in \mathcal{S}(\mathcal{H})_{++}} f_{\alpha,0}(p,\sigma) \ge \min_{\sigma \in \mathcal{S}(\mathcal{H})} f_{\alpha,0}(p,\sigma)$$

$$\ge \min_{\sigma \in \mathcal{S}(\mathcal{H})} f_{\alpha,0}(p,(1-\varepsilon)\sigma + \varepsilon(1/d)I)$$

$$+ \log(1-\varepsilon)$$

$$\ge \inf_{\sigma \in \mathcal{S}(\mathcal{H})_{++}} f_{\alpha,0}(p,\sigma) + \log(1-\varepsilon),$$

and by taking the supremum in ε , we get $\inf_{\sigma \in \mathcal{S}(\mathcal{H})_{++}} f_{\alpha,0}(p,\sigma) = \min_{\sigma \in \mathcal{S}(\mathcal{H})} f_{\alpha,0}(p,\sigma).$ The assertion about the other two minima can be obtained by repeating the same argument after taking the supremum over $\rho \in \mathcal{K}$ in (31) and the supremum over $p \in \mathcal{M}(\mathcal{K})$ in (32), respectively.

Remark IV.6. The first supremum in (28) and the last one in (29) can be replaced with maxima.

Proof: By Proposition IV.4,

$$\sup_{\rho \in \mathcal{K}} S_{\alpha} (\rho || \sigma_{0}) = \min_{\sigma \in \mathcal{S}(\mathcal{H})} \sup_{\rho \in \mathcal{K}} S_{\alpha} (\rho || \sigma)$$

$$\leq \sup_{\rho \in \mathcal{K}} S_{\alpha} (\rho || (1/d)I) = \sup_{\rho \in \mathcal{K}} \{\log d - S_{\alpha}(\rho)\}$$

$$< \log d.$$

Thus, $S_{\alpha}(\rho || \sigma_0)$ is finite, and therefore it is given as $S_{\alpha}(\rho || \sigma_0) = \frac{1}{\alpha - 1} \log \operatorname{Tr} \rho^{\alpha} \sigma_0^{1 - \alpha}$ for every $\rho \in \mathcal{K}$. This yields that $\rho \mapsto \overset{\alpha-1}{S_{\alpha}}(\rho || \sigma_0)$ on \mathcal{K} and $p \mapsto f_{\alpha,0}(p,\sigma_0)$ on $\mathcal{M}(\mathcal{K})$ are continuous, and hence they reach their suprema.

Since in the proofs of Propositions IV.3 and IV.4 we only used the properties of $f_{\alpha,\varepsilon}$ established in Lemmas IV.1 and IV.2, which are common with the properties of $g_{\alpha,\varepsilon}$, we have the following:

Proposition IV.7. The assertions of Propositions IV.3 and IV.4 hold true if we replace $f_{\alpha,\varepsilon}$ with $g_{\alpha,\varepsilon}$ for all $\varepsilon \geq 0$, and S_{α} with Q_{α} .

Now we are ready to prove the following:

Theorem IV.8. Let $W: \mathcal{X} \to \mathcal{S}(\mathcal{H})$ be a classical-quantum channel with compact image. Then, the capacities defined in (8)–(10) are equal to each other when $D = S_{\alpha}$ is a Rényi relative entropy with parameter $\alpha \in (0, 2]$.

Proof: The assertion is obvious for $\alpha = 1$ from the identities (2) and (3), so for the rest we assume that $\alpha \in (0, 2] \setminus \{1\}$. Let $\mathcal{K} := \operatorname{ran} W$. Proposition IV.4 yields that

$$\chi_{S_{\alpha},2}^{*}(W) = \sup_{p \in \mathcal{M}_{f}(\mathcal{X})} \inf_{\sigma \in \mathcal{S}(\mathcal{H})} \sum_{x} p(x) S_{\alpha} (W_{x} || \sigma)$$

$$= \sup_{p \in \mathcal{M}_{f}(\mathcal{K})} \inf_{\sigma \in \mathcal{S}(\mathcal{H})} \sum_{\rho \in \mathcal{K}} p(\rho) S_{\alpha} (\rho || \sigma)$$

$$= \sup_{p \in \mathcal{M}_{f}(\mathcal{K})} \min_{\sigma \in \mathcal{S}(\mathcal{H})} f_{\alpha,0}(p,\sigma)$$

$$= \min_{\sigma \in \mathcal{S}(\mathcal{H})} \sup_{\rho \in \mathcal{K}} S_{\alpha} (\rho || \sigma)$$

$$= R_{S_{\alpha}}(\operatorname{ran} W).$$

Let id be the identical channel on $\mathcal{K} = \operatorname{ran} W$, and let $f_{\alpha,0}(p,(1-\varepsilon)\sigma+\varepsilon(1/d)I)+\log(1-\varepsilon)\leq f_{\alpha,0}(p,\sigma). \ \ (32) \quad \ \text{id}: \ \rho\mapsto\delta_{\rho}\otimes\rho \ \ \text{be its lifting as in the Introduction. Using}$ Proposition IV.7, we have

$$\begin{split} \chi_{S_{\alpha},1}^{*}(W) &= \sup_{p \in \mathcal{M}_{f}(\mathcal{X})} \inf_{\sigma \in \mathcal{S}(\mathcal{H})} S_{\alpha} \left(\mathbb{E}_{p} \hat{W} \, || \, \hat{p} \otimes \sigma \right) \\ &= \sup_{p \in \mathcal{M}_{f}(\mathcal{K})} \inf_{\sigma \in \mathcal{S}(\mathcal{H})} S_{\alpha} \left(\mathbb{E}_{p} \hat{\mathrm{id}} \, || \, \hat{p} \otimes \sigma \right) \\ &= \sup_{p \in \mathcal{M}_{f}(\mathcal{K})} \inf_{\sigma \in \mathcal{S}(\mathcal{H})} \frac{1}{\alpha - 1} \log \operatorname{sign}(\alpha - 1) g_{\alpha,0}(p, \sigma) \\ &= \frac{1}{\alpha - 1} \log \operatorname{sign}(\alpha - 1) \sup_{p \in \mathcal{M}_{f}(\mathcal{K})} \min_{\sigma \in \mathcal{S}(\mathcal{H})} g_{\alpha,0}(p, \sigma) \\ &= \frac{1}{\alpha - 1} \log \operatorname{sign}(\alpha - 1) \min_{\sigma \in \mathcal{S}(\mathcal{H})} \sup_{\rho \in \mathcal{K}} Q_{\alpha} \left(\rho \, || \, \sigma \right) \\ &= \min_{\sigma \in \mathcal{S}(\mathcal{H})} \sup_{\rho \in \mathcal{K}} \frac{1}{\alpha - 1} \log \operatorname{sign}(\alpha - 1) Q_{\alpha} \left(\rho \, || \, \sigma \right) \\ &= R_{S_{\alpha}}(\operatorname{ran} W). \end{split}$$

V. THE ONE-SHOT CLASSICAL CAPACITY OF QUANTUM CHANNELS

Let $W: \mathcal{X} \to \mathcal{S}(\mathcal{H})$ be a classical-quantum channel. In order to transmit (classical) information through the channel, the sender has to encode the messages into signals at the input of the channel, and the receiver has to make a measurement at the outcome to determine which message was sent. A code is a triple (M, φ, E) , where $\{1, \ldots, M\}$ labels the possible messages to transmit, $\varphi: \{1, \ldots, M\} \to \mathcal{X}$ is the encoding map, and the positive operator valued measurement $E: \{1, \ldots, M\} \to \mathcal{B}(\mathcal{H})_+, \sum_{i=1}^M E_i = I$, is the decoding. The average probability of an erroneous decoding is given by

$$P_e(M, \varphi, E) := \frac{1}{M} \sum_{i=1}^{M} (1 - \text{Tr } W_{\varphi(i)} E_i) = 1 - P_s(M, \varphi, E),$$

where $P_s(M, \varphi, E)$ is the success probability. The one-shot ε -capacity of the channel is defined as the logarithm of the maximal number of messages that can be transmitted through the channel with error not exceeding ε :

$$C_{\varepsilon}(W) := \max\{\log M \mid \exists (M, \varphi, E) \text{ such that } P_{e}(M, \varphi, E) \leq \varepsilon\}.$$

Let $\chi_{H_r,0}^*(W)$ and $\chi_{S_\alpha,0}^*(W)$ denote the generalizations of the Holevo capacity of W as defined in (7), for a Hoeffding distance with parameter r and for a Rényi relative entropy with parameter α , respectively. For any $\varepsilon>0$ and any c>0, the one-shot ε -capacity can be lower bounded as

$$C_{\varepsilon}(W) \ge \chi_{H_{\log((1+c)/\varepsilon)},0}^*(W) - \log\left(\frac{2+c+1/c}{\varepsilon}\right)$$

$$= \sup_{0 \le \alpha < 1} \left\{ \frac{-\alpha \log\left(\frac{1+c}{\varepsilon}\right)}{1-\alpha} + \chi_{S_{\alpha},0}^*(W) \right\}$$

$$- \log\left(\frac{2+c+1/c}{\varepsilon}\right),$$

where the inequality was shown in [14], and the identity is obvious from the definition (19) of the Hoeffding distances.

While this bound might be rather loose for one single use of the channel, it is asymptotically optimal in the sense that it yields the Holevo capacity as a lower bound on the optimal asymptotic transmission rate of the channel [14].

In order to give an upper bound on the capacity, one has to find an upper bound on the success probability for any code (M,φ,E) in terms of M. Such a bound was given in [31], that we briefly outline below. Note that the function $x\mapsto x^{\frac{1}{\alpha}}$ is operator monotonic increasing for $\alpha\in[1,+\infty)$ and thus $W_{\varphi(k)}=(W_{\varphi(k)}^{\alpha})^{\frac{1}{\alpha}}\leq \left(\sum_{m=1}^{M}W_{\varphi(m)}^{\alpha}\right)^{\frac{1}{\alpha}}$. Hence, the average success probability is upper bounded as

$$P_{s}(M,\varphi,E) \leq \frac{1}{M} \sum_{k=1}^{M} \operatorname{Tr} E_{k} \left(\sum_{m=1}^{M} W_{\varphi(m)}^{\alpha} \right)^{\frac{1}{\alpha}}$$

$$= \frac{1}{M} \operatorname{Tr} \left(\sum_{m=1}^{M} W_{\varphi(m)}^{\alpha} \right)^{\frac{1}{\alpha}}$$

$$= M^{\frac{1-\alpha}{\alpha}} \operatorname{Tr} \left(\sum_{m=1}^{M} \frac{1}{M} W_{\varphi(m)}^{\alpha} \right)^{\frac{1}{\alpha}}$$

$$\leq M^{\frac{1-\alpha}{\alpha}} \sup_{p \in \mathcal{M}_{f}(\mathcal{X})} 2^{\frac{\alpha-1}{\alpha} \chi_{\alpha}(p)}, \tag{33}$$

where

$$\chi_{\alpha}(p) := \frac{\alpha}{\alpha - 1} \log \operatorname{Tr} \omega(p), \qquad \omega(p) := \left(\sum_{x \in \mathcal{X}} p(x) W_x^{\alpha} \right)^{\frac{1}{\alpha}}.$$

As it was pointed out in [13], [32], for any $\sigma \in \mathcal{S}(\mathcal{H})$ and $p \in \mathcal{M}_f(\mathcal{X})$ we have

$$S_{\alpha}\left(\mathbb{E}_{p}\hat{W}\mid\mid\hat{p}\otimes\sigma\right)$$

$$=S_{\alpha}\left(\mathbb{E}_{p}\hat{W}\mid\mid\hat{p}\otimes\frac{\omega(p)}{\operatorname{Tr}\omega(p)}\right)+S_{\alpha}\left(\frac{\omega(p)}{\operatorname{Tr}\omega(p)}\mid\mid\sigma\right)$$

$$=\chi_{\alpha}(p)+S_{\alpha}\left(\frac{\omega(p)}{\operatorname{Tr}\omega(p)}\mid\mid\sigma\right),$$
(34)

and hence

$$\chi_{\alpha}(p) = \inf_{\sigma \in \mathcal{S}(\mathcal{H})} S_{\alpha} \left(\mathbb{E}_{p} \hat{W} || \hat{p} \otimes \sigma \right), \tag{35}$$

which in turn yields

$$\sup_{p \in \mathcal{M}_f(\mathcal{X})} \chi_{\alpha}(p) = \chi_{S_{\alpha}, 1}^*(W). \tag{36}$$

The above observations lead to the following:

Theorem V.1. For any $\varepsilon > 0$, we have

$$C_{\varepsilon}(W) \le \inf_{\alpha > 1} \left\{ \chi_{S_{\alpha}, 1}^*(W) + \frac{\alpha}{\alpha - 1} \log \frac{1}{1 - \varepsilon} \right\}.$$

Proof: Assume that for a code (M, φ, E) we have $P_e(M, \varphi, E) \leq \varepsilon$. Then, by the above,

$$\log(1 - \varepsilon) \le \log P_s(M, \varphi, E) \le \frac{\alpha - 1}{\alpha} \left(\chi_{S_\alpha, 1}^*(W) - \log M \right)$$

for every $\alpha > 1$, from which the assertion follows immediately.

For each $n \in \mathbb{N}$, consider the *n*th i.i.d. extension of W, defined as $W^{(n)}: \mathcal{X}^n \to \mathcal{S}(\mathcal{H}^{\otimes n})$,

$$W^{(n)}(x_1,\ldots,x_n) := W(x_1) \otimes \ldots \otimes W(x_n).$$

The rate $R(\mathcal{C})$ of a sequence of codes $\mathcal{C}=\{C^{(n)}=(M^{(n)},\varphi^{(n)},E^{(n)})\}_{n\in\mathbb{N}}$ is $R(\mathcal{C}):=\liminf_{n\to\infty}\frac{1}{n}\log M^{(n)}$, and the asymptotic ε -capacity of W (with product encoding) is defined as

$$\overline{C}_{\varepsilon}(W) := \sup \left\{ R(\mathcal{C}) \mid \limsup_{n \to \infty} P_{e}(C^{(n)}) \le \varepsilon \right\},\,$$

where the supremum is taken over sequences of codes satisfying the indicated criterion. One can easily see that

$$\liminf_{n \to \infty} \frac{1}{n} C_{\varepsilon}(W^{(n)}) \leq \overline{C}_{\varepsilon}(W) \leq \overline{C}_{\varepsilon'}(W)
\leq \liminf_{n \to \infty} \frac{1}{n} C_{\varepsilon''}(W^{(n)})$$

for any $0 \le \varepsilon \le \varepsilon' < \varepsilon''$. The upper bound in Theorem V.1 is asymptotically sharp in the sense that it yields the Holevo capacity as an upper bound on the optimal information carrying capacity in the asymptotic limit. The details of the proof of the following Theorem are supplied in Appendix B.

Theorem V.2. Assume that ran W is compact. Then, for any $\varepsilon \in [0, 1)$,

$$\overline{C}_{\varepsilon}(W) \leq \chi_S^*(W).$$

Proof: By Theorem V.1 and Proposition B.2,

$$\overline{C}_{\varepsilon}(W) \leq \liminf_{n \to \infty} \frac{1}{n} C_{\varepsilon'}(W^{(n)})
\leq \liminf_{n \to \infty} \left\{ \frac{1}{n} \chi_{S_{\alpha}, 1}^{*}(W^{(n)}) + \frac{1}{n} \frac{\alpha}{\alpha - 1} \log \frac{1}{1 - \varepsilon'} \right\}
= \chi_{S_{\alpha}, 1}^{*}(W)$$

for any $0 < \varepsilon < \varepsilon' < 1$ and $\alpha > 1$. By Proposition B.5, the assertion follows for every $\varepsilon > 0$, and the case $\varepsilon = 0$ is immediate from $\overline{C}_0(W) \le \overline{C}_{\varepsilon}(W)$, $\varepsilon > 0$.

Remark V.3. Cutoff rates were also defined in [8] for channel coding in the following way: for $\kappa < 0$, the κ -cutoff rate $C_{\kappa}(W)$ is the largest R_0 for which

$$\limsup_{n \to \infty} \frac{1}{n} \log P_e(C^{(n)}) \le \kappa (R_0 - R)$$

for any sequence of codes with rate R, while for $\kappa > 0$, the κ -cutoff rate $C_{\kappa}(W)$ is the largest R_0 for which

$$\limsup_{n \to \infty} \frac{1}{n} \log P_s(C^{(n)}) \le \kappa(R_0 - R)$$

for any sequence of codes with rate R.

Inequality (33) and identity (36), together with the observations of Appendix B, yield that, for $\alpha > 1$,

$$\limsup_{n \to \infty} \frac{1}{n} \log P_s(C^{(n)}) \le \frac{\alpha - 1}{\alpha} (\chi_{S_\alpha, 1}^*(W) - R)$$

for any sequence of codes with rate R and hence,

$$C_{\kappa}(W) \ge \chi_{S_{\frac{1}{1}},1}^*(W), \qquad 0 < \kappa < 1.$$

The above inequality was shown to hold as an equality for classical channels in [8].

VI. REMARKS ON THE DIVERGENCE RADIUS

Let Σ be a subset of the state space $\mathcal{S}(\mathcal{H})$, and let $R_D(\Sigma)$ denote its D-radius as given in (6). A state σ^* which reaches the infimum in (6) is called a D-centre for Σ . As we have seen in the previous section, the S_α -radii of the range of a channel are related to the direct part of channel coding for $\alpha \in [0,1)$ and to the converse part for $\alpha \in (1,+\infty]$. In both cases, the asymptotically relevant quantities are the divergence radii with α close to 1. On the other hand, for state discrimination the relevant quantity turns out to be the ∞ -radius. More precisely, if $\rho_1,\ldots,\rho_r\in\mathcal{S}(\mathcal{H})$ then the optimal success probability of discriminating them by POVM measurements is given by $P_s=(1/r)\exp{(R_{S_{\max}}\{\rho_k\})}$ [33], where S_{\max} is the maxrelative entropy [24].

Related to state discrimination is the following geometrical problem: given $\rho_1, \ldots, \rho_r \in \mathcal{S}(\mathcal{H})$, find the largest q such that there exist states τ_1, \ldots, τ_r such that $q\rho_i + (1-q)\tau_i$ is independent of i. Such a family of states τ_1, \ldots, τ_r is called an optimal Helström family with parameter q in [34]. As one can easily see, the largest such q is given by $\exp(-R_{S_{\max}}\{\rho_k\})$, and $q\rho_i + (1-q)\tau_i$ is an S_{max} -centre for $\{\rho_k\}_{k=1}^r$. When r=2, the results of Holevo [35] and Helström [36] yield that the optimal success probability is given by $P_s = (1 + D)/2$, where $D := (1/2) \|\rho_1 - \rho_2\|_1$, and hence, $R_{S_{\max}}(\{\rho_1, \rho_2\}) =$ $\log(1+D)$. Moreover, an S_{max} -centre is given by $\sigma^* = (\rho_1 +$ $(2X_{+})/(1+D) = (\rho_2 + 2X_{-})/(1+D)$, where X_{+} and X_{-} are the positive and the negative parts of $\rho_1 - \rho_2$, respectively. In [38] and [37], a suboptimal Helström family was used for two states ρ_1 and ρ_2 to show Fannes type inequalities. Using instead the above optimal Helström family in the proof of [37, Proposition 1], one obtains the following:

Proposition VI.1. *Let* \mathcal{H} *be a Hilbert space and* $f : \mathcal{S}(\mathcal{H}) \to \mathbb{C}$ *be a bounded function that satisfies*

$$|f((1-\varepsilon)\rho_1 + \varepsilon \rho_2) - (1-\varepsilon)f(\rho_1) - \varepsilon f(\rho_2)| \le h_2(\varepsilon)$$
(37)

for any two states ρ_1, ρ_2 and any $\varepsilon \in [0,1]$, where $h_2(x) := -x \log x - (1-x) \log(1-x)$ is the binary entropy function. Then, for any two states ρ_1, ρ_2 on \mathcal{H} , we have

$$|f(\rho_1) - f(\rho_2)| \le 2h_2(\varepsilon) + 4\varepsilon M, \tag{38}$$

where
$$\varepsilon := \frac{\|\rho_1 - \rho_2\|_1}{2 + \|\rho_1 - \rho_2\|_1}$$
 and $M := \sup_{\rho \in \mathcal{S}(\mathcal{H})} |f(\rho)|$.

Proof: Let au_1, au_2 be the above optimal Helström family and $\sigma^*=(1-arepsilon)
ho_i+arepsilon au_i$ be the S_{\max} -centre of $\{
ho_1,
ho_2\}$. Then,

$$|f(\rho_{1}) - f(\rho_{2})|$$

$$\leq |f(\rho_{1}) - f(\sigma^{*})| + |f(\sigma^{*}) - f(\rho_{2})|$$

$$\leq \sum_{i=1}^{2} |f(\sigma^{*}) - (1 - \varepsilon)f(\rho_{i}) - \varepsilon f(\tau_{i})| + \varepsilon |f(\rho_{i})| + \varepsilon |f(\tau_{i})|$$

$$\leq 2h_{2}(\varepsilon) + 4\varepsilon M.$$

The von Neumann entropy is known to satisfy (37), which in turn yields by a simple computation that the conditional entropy and the relative entropy distance from a convex set containing a faithful state satisfy (37), too. Note that for the latter two quantities (38) yields a slight improvement of the result of [38] and of [37, Lemma 1], respectively, where the same bound was obtained with $\varepsilon = \|\rho_1 - \rho_2\|_1$.

For the case where D is the relative entropy S, it was shown in [12] that for any subset Σ of states, the S-centre is unique and is inside the closed convex hull $\overline{\operatorname{co}}\Sigma$ of Σ . This is no longer true for other Rényi relative entropies in general. For instance, for the classical probability distributions $\rho_1:=(1/2,1/4,1/4), \, \rho_2:=(1/2,1/6,1/3), \, \text{an}\,\, S_\infty$ -centre is given by $\sigma^*=(6/13,3/13,4/13), \, \text{and}\,\, \text{one}\,\, \text{can}\,\, \text{easily}\,\, \text{verify}\,\, \text{that}\,\, \text{no}\,\, S_\infty$ -centre can be found on the line segment connecting ρ_1 and ρ_2 . It is of some mathematical interest to find conditions on D ensuring the existence of a unique D-centre of Σ in $\overline{\operatorname{co}}\Sigma$ for any subset of states Σ .

VII. CONCLUDING REMARKS

The idea of representing the Rényi relative entropies as cutoff rates is from Csiszár [8], and we essentially followed his approach here. Note, however, that the analysis of the error exponents h_r , h_r , h_r in the classical case, on which the proof of [8] relies, is based on the Hellinger arc and a representation of the Hoeffding distances that have no equivalents in the quantum setting [2]. Instead, our analysis is based on an equivalent definition of the Hoeffding distances that can be defined also for quantum states, given in (19). That this definition of the Hoeffding distances have the right operational meaning was proven recently under the name of the quantum Hoeffding bound [3]–[6]. Note that this representation of the Hoeffding distances allows for a somewhat simplified proof even in the classical case. Moreover, this proof works also for the more general setting of correlated states considered in Theorem III.3.

The way to prove the identity of the different definitions of the Rényi capacities using minimax results is also from [8]. For this, the convexity of $\sigma \mapsto Q_{\alpha}(\rho || \sigma)$ and $\sigma \mapsto S_{\alpha}(\rho || \sigma)$ for every fixed ρ are essential. These are obvious in the classical case for Q_{α} , and for S_{α} when $\alpha \in (0,1)$, and were proven for S_{α} and $\alpha > 1$ in [8]. That proof, however, cannot be extended to the quantum case and, as far as we are aware, our Theorem II.1 is a new result. Note that in the quantum case the fact that $x \mapsto x^{1-\alpha}$ is not operator convex for $\alpha > 2$ yields a strong limitation, and no convexity properties of the α -relative entropies are expected to hold for parameters $\alpha > 2$. This limitation was overcome in [13], where a completely different approach was used to prove that $\chi_{S_{\alpha},1}^* = R_{S_{\alpha}}(\operatorname{ran} W)$ for all $\alpha > 1$. Another subtle technical difference between the proofs for the classical (more precisely, finite \mathcal{X}) and the general cases comes from the fact that in minimax theorems one of the sets has to be compact and convex, which in the first case can be chosen to be $\mathcal{M}_f(\mathcal{X})$, and the other space has to be convex, which is chosen to be $S(\mathcal{H})_{++}$. In the general case \mathcal{X} is usually the state space of a quantum system, which is of infinite cardinality and hence $\mathcal{M}_f(\mathcal{X})$ is convex but not compact, whereas replacing $\mathcal{M}_f(\mathcal{X})$ with $\mathcal{M}_m(\operatorname{ran} W)$ as in Appendix B yields a space that is compact but not convex. Hence we switched the role of the two spaces and chose $\mathcal{S}(\mathcal{H})$ to be the compact convex set. However, the (dis)continuity

properties of the Rényi relative entropies then wouldn't make it possible to satisfy the continuity requirements of minimax theorems, and that's why we had to use ε -perturbations in Section IV.

It is worth noting that Rényi relative entropies and the corresponding channel capacities are related to different regimes of information-theoretic tasks for the parameter values $\alpha \in (0,1)$ and for $\alpha \in (1,+\infty)$. Indeed, the first interval is related to the so-called direct part of problems, i.e., where a relevant error probability decays exponentially for rates below the optimal one, while the second interval is related to the (strong) converse regions, where a relevant success probability goes to zero (exponentially) for rates above the optimal rate. Cutoff rates are also defined in an asymmetric way, separately for the direct region ($\kappa < 0$) and for the strong converse region ($\kappa > 0$); see Remark V.3 and [8] for more details.

In the case of hypothesis testing between ρ and σ , for rates $r < S(\sigma || \rho)$, the optimal exponential decay rates of the error probabilities of the second kind are given explicitly by the Hoeffding distances $H_r(\rho || \sigma)$, which are defined through the Rényi relative entropies $S_{\alpha}(\rho || \sigma)$, $\alpha \in (0,1)$. For rates $r > S(\sigma || \rho)$, the success probabilities decay exponentially, and the optimal decay rates are known in the classical case to be given by the Han-Kobayashi bounds [2], [39], [40], defined through $S_{\alpha}(\rho || \sigma)$, $\alpha \in (1, +\infty)$. In the quantum case, however, the exact error exponents for the converse part are not known and hence it is not possible to extend the results of [8] on the cutoff rates for $\kappa > 0$ at the moment, though the results of [2], [40] give inequalities between the cutoff rates and the Rényi relative entropies that are expected to hold as equalities. For channel coding, the exact error exponents are not known for every rate value even in the classical case, but we see the same picture, i.e., the exponential decay of error probabilities for rates below the Shannon capacity can be expressed in terms of, or upper bounded by, the Rényi capacities $\chi_{S_{\alpha}}^*$ with $\alpha \in (0,1)$, while for rates above the Shannon capacity, the exponential decay rate of success probabilities can be expressed in terms of the Rényi capacities $\chi_{S_{\alpha}}^*$ with $\alpha \in (1, +\infty)$ [8].

Due to finite-size effects, the one-shot capacities are discontinuous functions of the error bar ε , and they depend on the parameters of the channel in a more intricate way than their asymptotic counterparts. As a result, it doesn't seem to be likely that they could be expressed in a similarly compact form as the asymptotic capacities, and if one is looking for some universal statement on them, applicable to all channels and all possible error bars, then probably the best one can hope for are lower and upper estimates on their values. In view of the above noted difference between the role of the intervals $\alpha \in (0,1)$ and $\alpha \in (1,+\infty)$, it seems rather natural to expect lower bounds in terms of the capacities $\chi_{S_a}^*$ with $\alpha \in (0,1)$ and upper bounds in terms of the capacities χ_S^* with $\alpha \in (1, +\infty)$. While we left the question of optimality open for the bounds provided in Section V (in fact, even to formulate what optimality might mean in this setting is a nontrivial question), it is somewhat reassuring that the optimal asymptotic capacity can be recovered by applying our bounds to several copies of the channel and letting the number of

copies go to infinity.

APPENDIX A A MINIMAX THEOREM

Let X and Y be non-empty sets and $f: X \times Y \to \overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, +\infty\}$ be a function. Obviously, for any $x_0 \in X$ and $y_0 \in Y$ we have $\inf_{x \in X} f(x, y_0) \leq \sup_{y \in Y} f(x_0, y)$ and hence,

$$\sup_{y \in Y} \inf_{x \in X} f(x, y) \le \inf_{x \in X} \sup_{y \in Y} f(x, y). \tag{39}$$

Minimax theorems give sufficient conditions on when the above inequality holds with equality. The following Lemma A.1 is a step in the proof of Sion's minimax theorem in [30], the proof of which we include for the readers' convenience. We will use the notation $[f(.\,,y)\leq c]$ to denote the level set $\{x\in X: f(x,y)\leq c\}$ for some number $c\in\mathbb{R}$, and other level sets are denoted similarly.

Lemma A.1. Assume that X is a compact topological space and f(., y) is lower semicontinuous for every $y \in Y$. Assume, moreover, that for any finite subset $Y' \subset Y$ we have

$$\inf_{x \in X} \max_{y \in Y'} f(x, y) \le \sup_{y \in Y} \inf_{x \in X} f(x, y). \tag{40}$$

Then the infima in (39) can be replaced with minima, and

$$\sup_{y \in Y} \min_{x \in X} f(x, y) = \min_{x \in X} \sup_{y \in Y} f(x, y).$$

Proof: The lower semi-continuity of f(.,y), $y \in Y$ implies the lower semi-continuity of $\sup_y f(.,y)$ and, since X is compact, all the functions f(.,y), $y \in Y$, and $\sup_y f(.,y)$ reach their infima on X. Hence, we can replace the infima with minima.

To prove the main assertion, we have to show that

$$\min_{x \in X} \sup_{y \in Y} f(x, y) \le \sup_{y \in Y} \min_{x \in X} f(x, y).$$

Let $c < \min_{x \in X} \sup_{y \in Y} f(x,y)$ or equivalently, let c be such that $\bigcap_{y \in Y} [f(.,y) \le c] = \emptyset$. Lower semicontinuity of f(.,y) yields that $[f(.,y) \le c]$ is closed (and hence compact) for every $y \in Y$ and hence, there exist finitely many y_1, \ldots, y_r such that $\bigcap_{i=1}^r [f(.,y_i) \le c] = \emptyset$ or equivalently, $c < \min_{x \in X} \max_{1 \le i \le r} f(x,y_i)$. By the assumption (40), we obtain $c < \sup_{y \in Y} \min_{x \in X} f(x,y)$. Since this holds for any $c < \min_{x \in X} \sup_{y \in Y} f(x,y)$, the assertion follows.

Corollary A.2. Let X be a compact topological space, Y be a subset of the real line and let $f: X \times Y \to \overline{\mathbb{R}}$ be a function. Assume that

- (i) f(., y) is lower semicontinuous for every $y \in Y$ and
- (ii) f(x,.) is monotonic increasing for every $x \in X$, or f(x,.) is monotonic decreasing for every $x \in X$.

Then the infima in (39) can be replaced with minima, and

$$\sup_{y \in Y} \min_{x \in X} f(x, y) = \min_{x \in X} \sup_{y \in Y} f(x, y).$$

Proof: By the monotonicity assumption, for any finite subset $Y' = \{y_1, \ldots, y_r\} \subset Y$, there exists a $y^* \in \{y_1, \ldots, y_r\}$ such that

$$\max_{1 \le i \le r} f(x, y_i) = f(x, y^*)$$

for all $x \in X$. Hence,

$$\min_{x \in X} \max_{1 \le i \le r} f(x, y_i) = \min_{x \in X} f(x, y^*) \le \sup_{y \in Y} \min_{x \in X} f(x, y).$$

Thus, all the conditions of Lemma A.1 are satisfied, from which the assertion follows.

APPENDIX B

The limit of the α -capacities

In this Appendix we collect some properties of the quantities χ_{α} and χ_{α}^* that are needed for the proof of Theorem V.2. To simplify notation, we introduce

$$\chi_{S_{\alpha}}^* := \chi_{S_{\alpha},1}^* = \chi_{S_{\alpha},2}^* = R_{S_{\alpha}}(\operatorname{ran} W),$$

where $W: \mathcal{X} \to \mathcal{S}(\mathcal{H})$ is a fixed classical-quantum channel. We start with the following:

Lemma B.1. Assume that $\alpha > 1$. Then, for any $p_1, p_2 \in \mathcal{M}_f(\mathcal{X})$, $\eta \in (0,1)$ and $\sigma \in \mathcal{S}(\mathcal{H})$,

$$S_{\alpha}\left(\mathbb{E}_{(1-\eta)p_1+\eta p_2}\hat{W} \mid\mid ((1-\eta)\hat{p}_1+\eta\hat{p}_2)\otimes\sigma\right) \tag{41}$$

$$\geq (1 - \eta) S_{\alpha} \left(\mathbb{E}_{p_{1}} \hat{W} || \hat{p}_{1} \otimes \sigma \right) + \eta S_{\alpha} \left(\mathbb{E}_{p_{2}} \hat{W} || \hat{p}_{2} \otimes \sigma \right)$$

$$(42)$$

$$\geq (1 - \eta)\chi_{\alpha}(p_1) + \eta\chi_{\alpha}(p_2). \tag{43}$$

In particular, the function $p \mapsto \chi_{\alpha}(p)$ is concave on $\mathcal{M}_f(\mathcal{X})$.

Proof: The inequality in (43) is obvious from (34). One can easily verify that the expression in (41) is equal to $+\infty$ if and only if the expression in (42) is equal to $+\infty$, and otherwise the inequality between the two follows by a straightforward computation from the concavity of the function $\frac{1}{\alpha-1}\log$. The last assertion follows by taking the infimum in σ in the inequality between (41) and (43).

The following statement is essentially Lemma 2 from [31]:

Proposition B.2. Assume that ran W is compact and $\alpha > 1$. Then

$$\chi_{S_{\alpha}}^*(W^{(n)}) = n\chi_{S_{\alpha}}^*(W), \qquad n \in \mathbb{N}.$$

Proof: Using the concavity established in Lemma B.1, one can follow the proof of Lemma 2 in [31] to obtain the assertion. (Note that in [31], \mathcal{X} was assumed to be finite, but that doesn't make a difference in the proof.)

Let
$$m := (\dim \mathcal{H})^2 + 1$$
, and let

$$\mathcal{M}_m(\operatorname{ran} W) := \{ p \in \mathcal{M}_f(\operatorname{ran} W) : |\operatorname{supp} p| \le m \}$$

denote the set of probability measures supported on not more than m points in $\operatorname{ran} W$. By Carathéodory's theorem [41, Theorem (2.3)], for every $p \in \mathcal{M}_f(\mathcal{X})$, there exists a $\tilde{p} \in \mathcal{M}_m(\operatorname{ran} W)$ such that

$$\chi_{\alpha}(p) = \tilde{\chi}_{\alpha}(\tilde{p}) := \frac{\alpha}{\alpha - 1} \log \operatorname{Tr} \left(\sum_{\omega \in \operatorname{ran} W} \tilde{p}(\omega) \omega^{\alpha} \right)^{\frac{1}{\alpha}}.$$

Note that $\tilde{\chi}$ can also be defined by replacing \mathcal{X} with ran W and W with the identity map id on ran W in (35), i.e., for each $p \in \mathcal{M}_f(\operatorname{ran} W)$,

$$\tilde{\chi}_{\alpha}(p) = \inf_{\sigma \in \mathcal{S}(\mathcal{H})} S_{\alpha} \left(\mathbb{E}_{p} \hat{\mathrm{id}} \mid\mid \hat{p} \otimes \sigma \right). \tag{44}$$

The functions χ_1 and $\tilde{\chi}_1$ are defined simply by replacing α with 1 in (35) and in (44), respectively. Note that

$$\chi_{S_{\alpha}}^{*}(W) = \sup_{p \in \mathcal{M}_{f}(\mathcal{X})} \chi_{\alpha}(p) = \sup_{p \in \mathcal{M}_{m}(\operatorname{ran} W)} \tilde{\chi}_{\alpha}(p)$$

for every $\alpha \in [0, +\infty)$.

Lemma B.3. The functions $\alpha \mapsto \chi_{\alpha}(p)$ and $\alpha \mapsto \tilde{\chi}_{\alpha}(p)$ are montonically increasing on $[0, +\infty)$ for all $p \in \mathcal{M}_f(\mathcal{X})$ and $p \in \mathcal{M}_f(\operatorname{ran} W)$, respectively, and

$$\lim_{\alpha \to 1} \chi_{\alpha}(p) = \chi_{1}(p), \qquad \lim_{\alpha \to 1} \tilde{\chi}_{\alpha}(p) = \tilde{\chi}_{1}(p).$$

Proof: The assertion on the monotonicity follows immediately from the monotonicity of the Rényi relative entropies in the parameter α . We prove the assertion on the limit separately for $\alpha \nearrow 1$ and for $\alpha \searrow 1$. In the second case, we have

$$\lim_{\alpha \searrow 1} \chi_{\alpha}(p) = \inf_{\alpha > 1} \chi_{\alpha}(p) = \inf_{\alpha > 1} \inf_{\sigma \in \mathcal{S}(\mathcal{H})} S_{\alpha} \left(\mathbb{E}_{p} \hat{W} \mid\mid \hat{p} \otimes \sigma \right)$$

$$= \inf_{\sigma \in \mathcal{S}(\mathcal{H})} \inf_{\alpha > 1} S_{\alpha} \left(\mathbb{E}_{p} \hat{W} \mid\mid \hat{p} \otimes \sigma \right)$$

$$= \inf_{\sigma \in \mathcal{S}(\mathcal{H})} S \left(\mathbb{E}_{p} \hat{W} \mid\mid \hat{p} \otimes \sigma \right) = \chi_{1}(p).$$

For fixed $p \in \mathcal{M}_f(\mathcal{X})$ and $\alpha \in [0,+\infty)$, the map $\sigma \mapsto S_\alpha\left(\mathbb{E}_p\hat{W} \mid\mid \hat{p}\otimes\sigma+\varepsilon I\right)$ is continuous on the compact set $\mathcal{S}(\mathcal{H})$ and hence the map $\sigma \mapsto S_\alpha\left(\mathbb{E}_p\hat{W} \mid\mid \hat{p}\otimes\sigma\right)$ is lower semicontinuous, due to (13). On the other hand, for fixed $\sigma \in \mathcal{S}(\mathcal{H})$, the map $\alpha \mapsto S_\alpha\left(\mathbb{E}_p\hat{W} \mid\mid \hat{p}\otimes\sigma\right)$ is monotonic increasing in α and hence, by Corollary A.2, we have

$$\begin{split} \lim_{\alpha \nearrow 1} \chi_{\alpha}(p) &= \sup_{\alpha < 1} \inf_{\sigma \in \mathcal{S}(\mathcal{H})} S_{\alpha} \left(\mathbb{E}_{p} \hat{W} \mid\mid \hat{p} \otimes \sigma \right) \\ &= \inf_{\sigma \in \mathcal{S}(\mathcal{H})} \sup_{\alpha < 1} S_{\alpha} \left(\mathbb{E}_{p} \hat{W} \mid\mid \hat{p} \otimes \sigma \right) \\ &= \inf_{\sigma \in \mathcal{S}(\mathcal{H})} S \left(\mathbb{E}_{p} \hat{W} \mid\mid \hat{p} \otimes \sigma \right) = \chi_{1}(p). \end{split}$$

The proof for $\lim_{\alpha \to 1} \tilde{\chi}_{\alpha}(p)$ goes exactly the same way.

The following Lemma was shown in [42]. For readers' conveniance, we include a proof here.

Lemma B.4. If ran W is compact then $\mathcal{M}_m(\operatorname{ran} W)$ can be equipped with a topology τ with respect to which $\mathcal{M}_m(\operatorname{ran} W)$ is compact and $\tilde{\chi}_{\alpha}$ is continuous.

Proof: Let $S_m:=\{(\lambda_1,\dots,\lambda_m):\lambda_1,\dots,\lambda_m\geq 0,\sum_{i=1}^m\lambda_i=1\}$ denote the m-dimensional probability simplex, and define $\Omega_m(W):=S_m\times(\operatorname{ran} W)^m=\{(\underline{\lambda},\underline{\omega}):\underline{\lambda}\in S_m,\,\omega_1,\dots,\omega_m\in\operatorname{ran} W\}$. Compactness of $\operatorname{ran} W$ yields that $\Omega_m(W)$ is compact with respect to its natural topology. Let $\pi_m:\Omega_m(W)\to\mathcal{M}_m(\operatorname{ran} W),\,\pi_m(\underline{\lambda},\underline{\omega}):=\sum_{i=1}^m\lambda_i\delta_{\omega_i},$ where δ_{ω_i} denotes the Dirac measure concentrated at ω_i . We define the topology τ on $\mathcal{M}_m(\operatorname{ran} W)$ to be the factor topology, i.e., the finest topology with respect to which π_m is continuous. Being the continuous image of a compact set, $\mathcal{M}_m(\operatorname{ran} W)$ is also compact. One can easily see that $\tilde{\chi}_\alpha\circ\pi_m$ is continuous on $\Omega_m(W)$, which in turn is equivalent to the continuity of $\tilde{\chi}_\alpha$ with respect to τ .

The following statement was shown in Lemma 3 of [31] for the case where \mathcal{X} is finite. Here we give an alternative proof, using the minimax theorem established in Appendix A, that covers the general case.

Proposition B.5.

$$\lim_{\alpha \to 1} \chi_{S_{\alpha}}^*(W) = \chi_S^*(W).$$

Proof: We prove separately the cases $\alpha \nearrow 1$ and $\alpha \searrow 1$. In the first case, the assertion follows immediately from Lemma B.3, as

$$\lim_{\alpha \nearrow 1} \chi_{S_{\alpha}}^{*}(W) = \sup_{\alpha \in [0,1)} \sup_{p \in \mathcal{M}_{f}(\mathcal{X})} \chi_{\alpha}(p)$$

$$= \sup_{p \in \mathcal{M}_{f}(\mathcal{X})} \sup_{\alpha \in [0,1)} \chi_{\alpha}(p)$$

$$= \sup_{p \in \mathcal{M}_{f}(\mathcal{X})} \chi_{1}(p) = \chi_{S}^{*}(W).$$

Note that the function $f(p,\alpha):=-\tilde{\chi}_{\alpha}(p)$ is monotonic decreasing in its second variable on $Y:=(1,+\infty)$ and continuous in its first variable on the compact space $X:=\mathcal{M}_m(\operatorname{ran} W)$, due to Lemma B.4. Hence, we can apply the minimiax theorem of Corollary A.2 to obtain

$$\lim_{\alpha \searrow 1} \chi_{S_{\alpha}}^{*}(W) = \inf_{\alpha > 1} \max_{p \in \mathcal{M}_{m}(\operatorname{ran} W)} \tilde{\chi}_{\alpha}(p)$$

$$= \max_{p \in \mathcal{M}_{m}(\operatorname{ran} W)} \inf_{\alpha > 1} \tilde{\chi}_{\alpha}(p)$$

$$= \max_{p \in \mathcal{M}_{m}(\operatorname{ran} W)} \tilde{\chi}_{1}(p) = \chi_{S}^{*}(W).$$

ACKNOWLEDGMENTS

M. Mosonyi is grateful to Tomohiro Ogawa and Andreas Winter for helpful discussions.

REFERENCES

- F. Hiai, D. Petz: "The proper formula for relative entropy and its asymptotics in quantum probability"; Commun. Math. Phys. 143, 99–114, (1991)
- [2] T. Ogawa, H. Nagaoka: "Strong converse and Stein's lemma in quantum hypothesis testing"; *IEEE Trans. Inform. Theory* 47, 2428–2433, (2000)
- [3] K.M.R. Audenaert, M. Nussbaum, A. Szkoła, F. Verstraete: "Asymptotic error rates in quantum hypothesis testing"; *Comm. Math. Phys.* 279, 251– 283, (2008).
- [4] M. Hayashi: "Error exponent in asymmetric quantum hypothesis testing and its application to classical-quantum channel coding"; *Phys. Rev. A* 76, 062301, (2007)
- [5] H. Nagaoka: "The converse part of the theorem for quantum Hoeffding bound"; quant-ph/0611289
- [6] F. Hiai, M. Mosonyi, T. Ogawa: "Error exponents in hypothesis testing for correlated states on a spin chain"; J. Math. Phys. 49, 032112, (2008)
- [7] A. Rényi: "On measures of entropy and information"; Proc. 4th Berkeley Symp. on Math. Statist. Probability 1, 547–561, Berkeley, CA (1961)
- [8] I. Csiszár: "Generalized cutoff rates and Rényi's information measures"; IEEE Trans. Inf. Theory 41, 26–34, (1995)
- [9] V. Vedral, M.B. Plenio, M.A. Rippin, P.L. Knight: "Quantifying entanglement"; *Phys. Rev. Lett.* 78, 2275–2279, (1997)
- [10] A.S. Holevo: "The capacity of the quantum communication channel with general signal states"; *IEEE Trans. Inform. Theory* 44, 269–273, (1998)
- [11] B. Schumacher, M.D. Westmoreland: "Sending classical information via noisy quantum channels"; *Phys. Rev. A* 56, 131–138, (1997)
- [12] M. Ohya, D. Petz, N. Watanabe: "On capacities of quantum channels"; Prob. Mat. Stat. 17, 179–196, (1997)
- [13] R. Koenig, S. Wehner: "A strong converse for classical channel coding using entangled inputs"; Phys. Rev. Lett. 103, 070504, (2009)

- [14] M. Mosonyi, N. Datta: "Generalized relative entropies and the capacity of classical-quantum channels"; J. Math. Phys. 50, 072104 (2009)
- [15] D. Petz: "Quasi-entropies for finite quantum systems"; Rep. Math. Phys. 23, 57–65, (1986)
- [16] F. Hiai, M. Mosonyi, M. Hayashi: "Quantum hypothesis testing with group symmetry"; J. Math. Phys. 50 103304 (2009)
- [17] M. Tomamichel, R. Colbeck, R. Renner: "A fully quantum asymptotic equipartition property"; *IEEE Trans. Inform. Theory* 55, 5840–5847, (2009)
- [18] T. Ando: "Concavity of certain maps and positive definite matrices and applications to Hadamard products"; *Linear Algebra Appl.* 26, 203–241, (1979)
- [19] E.H. Lieb: "Convex trace functions and the Wigner-Yanase-Dyson conjecture"; Adv. Math. 11, 267–288 (1973)
- [20] M. Ben-Bassat, J. Raviv: "Rényi's entropy and the probability of error"; IEEE Trans. Inform. Theory IT-24, 324–331, (1978)
- [21] T. Ando, F. Hiai: "Operator log-convex functions and operator means"; arXiv:0911.5267
- [22] F. Hiai, M. Ohya, M. Tsukada: "Sufficiency, KMS condition and relative entropy in von Neumann algebras"; Pacific J. Math. 96, 99–109, (1981)
- [23] I. Ekeland, R. Temam: "Convex Analysis and Variational Problems"; North-Holland Publishing Company and American Elsevier Company (1976)
- [24] N. Datta: "Min- and max- relative entropies and a new entanglement measure" *IEEE Trans. Inform. Theory* 55, 2816–2826, (2009)
- [25] F. Buscemi, N. Datta: "The quantum capacity of channels with arbitrarily correlated noise"; *IEEE Trans. Inform. Theory* 56, 1447–1460 (2010)
- [26] R. Colbeck, private communication.
- [27] M. Mosonyi, F. Hiai, T. Ogawa, M. Fannes: "Asymptotic distinguishability measures for shift-invariant quasi-free states of fermionic lattice systems"; J. Math. Phys. 49, 072104 (2008).
- [28] M. Mosonyi: "Hypothesis testing for Gaussian states on bosonic lattices"; J. Math. Phys. 50, 032104 (2009).
- [29] M. Sion: "On general minimax theorems"; Pacific J. Math. 8, 171-176, (1958)
- [30] H. Komiya: "Elementary proof for Sion's minimax theorem"; Kodai Math. J. 11, 5–7, (1988)
- [31] T. Ogawa, H. Nagaoka: "Strong converse to the quantum channel coding theorem"; *IEEE Trans. Inform. Theory* 45, 2428–2433, (1999)
- [32] R. Sibson: "Information radius"; Z. Wahrscheinlichkeitsth. Verw. Gebiete 14, 149–161, (1969)
- [33] R. Koenig, R. Renner, C. Schaffner: "The operational meaning of minand max-entropy"; IEEE Trans. Inform. Theory 55, 4337–4347, (2009)
- [34] G. Kimura, T. Miyadera, H. Imai: "Optimal State Discrimination in General Probabilistic Theories"; *Phys. Rev. A* **79**, 062306 (2009).
- [35] A.S. Holevo: "Investigations in the general theory of statistical decisions"; Proc. Steklov Inst. Math. 124, (1976) (AMS Translation: 1978 Issue 3)
- [36] C.W. Helström: "Quantum Detection and Estimation Theory"; Academic Press, (1976)
- [37] M. Horodecki, B. Synak-Radtke: "On asymptotic continuity of functions of quantum states"; J. Phys. A 39, L423–L437, (2006)
- [38] R. Alicki, M. Fannes: "Continuity of quantum conditional information"; J. Phys. A 37, L55–L57, (2004)
- [39] T.S. Han, K. Kobayashi: "The strong converse theorem for hypothesis testing"; *IEEE Trans. Inform. Theory* **35**, 178–180, (1989)
- [40] M. Hayashi: "Quantum Information: An Introduction"; Springer, (2006)
- [41] A. Barvinok: "A Course in Convexity"; American Mathematical Society, Graduate Studies in Mathematics, vol. 54, (2002)
- [42] A. Fujiwara, H. Nagaoka: "Operational capacity and pseudoclassicality of a quantum channel"; *IEEE Trans. Inform. Theory* 44, 1071–1086, (1998)