Star-forming regions in the intragroup medium of compact groups of galaxies

S. Torres-Flores^{1,2}, C. Mendes de Oliveira¹, D. F. de Mello^{3,4}, P. Amram², H. Plana^{2,6}, B. Epinat⁵ and J. Iglesias-Páramo⁷

¹Universidade de São Paulo, Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Departamento de Astronomia, São Paulo, Brazil email: storres@astro.iag.usp.br

²Laboratoire d'Astrophysique de Marseille, OAMP, Université de Provence & CNRS, 38 rue F.

Joliot–Curie, 13388 Marseille, Cedex 13, France

 $^3 \rm Observational$ Cosmology Laboratory, Code 665, Goddard Space Flight Center, Greenbelt, MD 20771, USA

⁴Catholic University of America, Washington, DC 20064, USA

⁵Laboratoire d'Astrophysique de Toulouse-Tarbes, Université de Toulouse, CNRS, 14 Avenue Edouard Belin, 31400 Toulouse, France

⁶Laboratorio de Astrofisica Teorica e Observacional, Universidade Estadual de Santa Cruz, Brazil

⁷Instituto de Astrofisica de Andalucia (CSIC), Camino Bajo de Huetor 50, 18008 Granada, Spain

Abstract.

We present the results of a multiwavelength campaign searching for young objects in the intragroup medium of seven compact groups of galaxies: HCG 2, 7, 22, 23, 92, 100 and NGC 92. We used Fabry-Perot velocity fields and rotation curves together with GALEX NUV and FUV images, optical R-band and HI maps to evaluate the stage of interaction of each group. We conclude that groups (i) HCG 7 and HCG 23 are in an early stage of interaction, (ii) HCG 2 and HCG 22 are mildly interacting, and (iii) HCG 92, HCG 100 and NGC 92 are in a late stage of evolution. Evolved groups have a population of young objects in their intragroup medium while no such population is found within the less evolved groups. We also report the discovery of a tidal dwarf galaxy candidate in the tail of NGC 92. These three groups, besides containing galaxies which have peculiar velocity fields, also show extended HI tails. Our results indicate that the advanced stage of evolution of a group together with the presence of intragroup HI clouds may lead to star formation in the intragroup medium.

Keywords. galaxies: interactions, (galaxies:) intergalactic medium, galaxies: kinematics and dynamics.

1. Introduction

Compact groups of galaxies are associations of three to seven galaxies, where the projected distances between them is of the order of their diameters, and where the group shows a low velocity dispersion, making compact groups an ideal place to study galaxy interaction and intergalactic star formation (e.g. Torres-Flores et al. 2010, de Mello et al. 2008, de Mello, Torres-Flores & Mendes de Oliveira 2008, Mendes de Oliveira et al. 2004). The main goal of this work is to search for a link between the evolutionary stage of a group and the presence of young intergalactic objects which may have formed during galaxy interactions. For this, we analyze a subsample of seven compact groups (HCG 2, 7, 22, 23, 92, 100 and NGC 92) which span a wide range of evolutionary stages, from HI

rich groups to strongly interacting groups, where the galaxies show tidal tail features and a deficiency in neutral HI gas. In order to analyze the evolutionary stage of each group, we used new Fabry-Perot velocity maps, GALEX/UV data and optical R-band images. The velocity fields and rotation curves help constraining the evolutionary stage of each compact group while ultraviolet light contains important information regarding the age of the young stellar population that may be present in the intragroup medium.

2. UV analysis

We searched for ultraviolet emitting regions in the vicinity of all seven targets, using the SExtractor software (SE, Bertin & Arnouts 1996) in the FUV, NUV and R skysubtracted images of our compact group sample. We compare the field density of regions detected in the compact group with a control sample outside the group. HCG 92 and HCG 22 have the highest field density in this study. No excess was found in HCG 2, HCG 7, HCG 23, HCG 100 and NGC 92 (Torres-Flores et al. 2009).

3. Fabry-Perot analysis

In order to constrain the evolutionary stage of each compact group, we inspected the velocity field and rotation curve of each galaxy to search for interaction indicators, in a similar way to that done by Plana et al. (2003) and Amram et al. (2003). In the case of NGC 92, it shows a prominent tidal tail in its velocity field. At the tip of this tail, there is a tidal dwarf galaxy candidate, having an age of about 40 Myrs (Torres-Flores et al. 2009).

4. Conclusions

We used multiwavelength data to study the evolutionary stages of the compact groups of galaxies HCG 2, 7, 22, 23, 92, 100 and NGC 92. New Fabry-Perot velocity fields, rotation curves and GALEX NUV/FUV images were analyzed for four and seven of these groups respectively. Groups HCG 7 and 23 are in an early stage of interaction whereas HCG 2 and 22 show limited interaction features and HCG 92, 100 and NGC 92 are in a late stage of evolution, having HI gas in the intragroup medium, galaxies with peculiar velocity fields and several young star-forming regions in the intergalactic medium.

Acknowledgements

S. T–F. acknowledges the financial support of FAPESP through the Doctoral position, under contract 2007/07973-3 and Eiffel scholarship.

References

Amram, P., Plana, H., Mendes de Oliveira, C. et al. 2003, A & A, 402, 865 Bertin, E., & Arnouts, S. 1996, A & A & S, 117, 393 de Mello, D.F., Torres–Flores, S. & Mendes de Oliveira, C. 2008, AJ, 135, 319 de Mello, D. F., Smith, L. J., Sabbi, E. et al. 2008, AJ, 135, 548 Mendes de Oliveira, C., Cypriano, E.S., & Sodré,L. Jr, Balkowski, C. 2004, ApJ, L17 Plana, H., Amram, P., Mendes de Oliveira, C. et al. 2003, AJ, 125, 1736 Torres-Flores, S., Mendes de Oliveira, C., de Mello, D. F., Amram, P., Plana, H., Epinat, B. & Iglesias-Páramo, J. 2009, A & A, 507, 723

Torres-Flores, S., Mendes de Oliveira, C., de Mello, D. F. et al. 2010, in preparation.