

Electronic zero modes of vortices in Hall states of gapped graphene

Gordon W. Semenoff

*Department of Physics and Astronomy, University of British Columbia
6224 Agricultural Road, Vancouver, British Columbia V6T 1Z1*

Recent observation of a metal-insulator phase transition in the $\nu = 0$ Hall state of graphene has inspired the idea that charge carriers in the metallic state could be fractionally charged vortices. We examine the question of whether vortices in particular gapped states of graphene and subject to external magnetic and pseudo-magnetic fields could have the mid-gap zero mode electron states which would allow them to be charged.

1. INTRODUCTION

It is now established that, at sufficiently low temperature and strong magnetic field, the four-fold degenerate Landau level which resides at the apex of the Dirac cone in graphene is split into four sublevels [1]-[5]. At least some of this splitting is attributed to spontaneous breaking of the sublattice symmetry and generation of a mass gap in the electron spectrum. The further recent observation of Kosterlitz-Thouless scaling of the resistivity near the metal-insulator transition which occurs as the strength of the field is varied [5]-[8] has inspired a number of possible explanations, perhaps the most interesting of which has fractionally charged vortices as the carriers of electric charge in the metallic phase [9][10][11]. This scenario would give another concrete physical example beyond polyacetylene [12] of topological solitons with fractional quantum numbers.

The vortices are topological defects in the condensate which creates the mass gap. In a system with particle-hole symmetry, electron energy states above and below the middle of the mass gap are paired and fractional charge can only arise from unpaired mid-gap states [13]. The possibility that the interaction of vortices with relativistic fermions can result in mid-gap states has been known for a long time [14]-[17]. The novel feature of the Hall system is the presence of a macroscopic magnetic field where the electron spectrum would normally be concentrated in Landau levels. In the absence of the mass condensate, or other symmetry breaking, a four-fold degenerate (two spins and two valleys) Landau level resides precisely at the Dirac point [18]. With a parity and time-reversal invariant mass gap $2m$, this Landau level is split into two levels which are displaced to the positive and negative mass thresholds, $E = \pm m$ [17] and, without defects, the mass gap contains no electronic states at all. It is then reasonable to ask whether a vortex defect in the condensate which creates the mass gap can bind a mid-gap electronic state. This question has been studied recently by Hou et.al. [10] and Herbut [11] who found, surprisingly, that the answer is yes. In fact, the existence of mid-gap states attached to vortices is relatively insensitive to the presence of a magnetic field. In the following, we shall examine this phenomenon in more detail. We will show that the fermion-vortex system in simultaneous magnetic and pseudo-magnetic fields gen-

erally has $|n|$ zero modes, where n is the vorticity, even when the magnetic field is a constant so that its flux diverges with the volume of the system. The exception is where the pseudo-magnetic flux inside a disc of large radius $r \rightarrow \infty$ grows faster than r and, in addition, when it is larger than the magnetic flux at large r . In that case, the number of zero modes no longer depends on the vorticity, instead there are two infinitely degenerate Landau levels at zero energy.

We shall also show that, when the pseudo-magnetic field falls off at least as fast as $1/r^2$ at $r \rightarrow \infty$, an external magnetic field has no effect on the η -invariant of the graphene Dirac operator perturbed by a constant mass term which upsets the particle-hole symmetry. η is a topological invariant and it is of physical interest because it is proportional to the electric charge of a system of electrons which is governed by the single-particle Hamiltonian in question and in a state where all of the negative energy levels are filled and all of the positive energy levels are empty. This result indicates that the fractional charge of a vortex is indeed a topological invariant and that it is independent of the external magnetic field.

Finally, we shall find interesting correlations between the sublattice components of the electron mid-gap state wavefunctions and its internal quantum numbers, such as spin or valley assignment. These correlations are analogous to the Dirac point Landau level in ungapped graphene, where, for one valley the Landau level zero mode wavefunctions all live on one of the two sublattices and the other valley zero mode wavefunctions live on the other sublattice, which sublattice depends on the sign of the external magnetic field. Zero modes in a vortex background also follow a pattern that is similar to this. We will elaborate in later Sections.

A number of scenarios for mass gap generation have been proposed [19]-[32]. To support topological vortices, the mass condensate should have an $SO(2)$ symmetry. Two proposals have appeared in recent literature. One is antiferromagnetic order with an easy plane [33]. Formation of antiferromagnetic order would be a result of the short-ranged on-site Coulomb repulsion of electrons whose role is enhanced in a strong magnetic field [27]. The Zeeman coupling of electron spins to the magnetic field would result in an “easy plane” where the antiferromagnetic order parameter lies in the plane that is perpendicular to the applied field. This is easy to demonstrate

using second order degenerate perturbation theory with the Zeeman coupling as a perturbation and the possible orientations of the antiferromagnetic condensate as the set of degenerate ground states. The other scenario with $SO(2)$ symmetry is a bond density wave called the Kekule distortion which is a result of the electron-phonon coupling[26],[9][34]-[36]. It was shown to have vortex defects which, when coupled to the graphene electron, bind an unpaired mid-gap state.

2. GRAPHENE HAMILTONIAN WITH A MASS GAP

We shall begin by considering graphene at energies near the Dirac points, and with a mass generated by antiferromagnetic order. The bipartite honeycomb lattice of graphene is a superposition of two triangular sublattices. Antiferromagnetic order gives each of the two spin states of the electron a charge density wave – spin up electrons have higher density on one of the two sublattices and spin down electrons on the other sublattice. This breaks the symmetry under interchanging the two sublattices and the charge density wave generates a parity-invariant mass term in graphene. [18]. The electrons are described by the Hamiltonian

$$H = \begin{bmatrix} i\hbar v_F \vec{\sigma} \cdot \vec{D} + \sigma^3 \vec{m} \cdot \vec{\tau} & 0 \\ 0 & -i\hbar v_F \vec{\sigma}^* \cdot \vec{D} + \sigma^3 \vec{m} \cdot \vec{\tau} \end{bmatrix} \quad (1)$$

This Hamiltonian is an 8×8 matrix. The 2×2 structure which is displayed explicitly refers to the two graphene valleys. The four additional dimensions which are suppressed are a direct product of two dimensional matrices: one on which is described by the pseudo-spin Pauli matrices σ^a act and one by the spin Pauli matrices τ^a . An alternative way of presenting this Hamiltonian is by writing the 8×8 matrices as ordered direct products of the three 2×2 matrices with unit and Pauli matrices for pseudo-spin (\mathcal{I}, σ^a), spin (\mathcal{I}, τ^b) and valleys (\mathcal{I}, η^c), in that order. In terms of these matrices, (1) has the form

$$H = i\sigma^1 \otimes \mathcal{I} \otimes \eta^3 D_1 + i\sigma^2 \otimes \mathcal{I} \otimes \mathcal{I} D_2 + \sigma^3 \otimes \tau^a \otimes \mathcal{I} m^a \quad (2)$$

Here, the covariant derivative is $\vec{D} = \vec{\nabla} - i\frac{e}{c}\vec{A}$ with \vec{A} the electromagnetic vector potential and $v_F \approx \frac{c}{300}$ is the graphene Fermi velocity. The magnetic field is $B_A \equiv \vec{\nabla} \times \vec{A}$.

It is useful to remind the reader that, the upper component of the spinors on which the pseudo-spin matrices (\mathcal{I}, σ^a) act has support on one of the graphene sublattices, the lower component on the other.

In (1) and (2), the masses of each spin polarization are $\pm|m|$, the eigenvalues of $m_a \tau^a$. If the antiferromagnetism has an “easy plane”, only two of the the three spin matrices can be used to describe possible orientations of the condensate, for example, $\vec{m} \cdot \vec{\tau} = m_1 \tau^1 + m_2 \tau^2$. We will

be interested in the circumstance where m_a can depend on spatial position and can have a vortex profile, with asymptotic limit

$$\lim_{r \rightarrow \infty} [m_1(r, \theta) + im_2(r, \theta)] = \hat{m} e^{in\theta} + \mathcal{O}\left(\frac{1}{r}\right) \quad (3)$$

with (r, θ) the polar coordinates of the plane and \hat{m} a constant.

If the mass term in (1) were absent, the existence of two valleys and two spin states give the Hamiltonian an $SU(4)$ symmetry. The mass term reduces this to an $SU(2)$ symmetry which mixes the valleys.[37] It is generated by

$$\mathcal{I} \otimes \mathcal{I} \otimes \eta^3, \sigma^2 \otimes \tau^3 \otimes \eta^1, \sigma^2 \otimes \tau^3 \otimes \eta^2 \quad (4)$$

Before we proceed, we observe that, we can redefine the matrix components of the Hamiltonian (2) by conjugating it with the matrix

$$\mathcal{I} \otimes \mathcal{I} \otimes \frac{1}{2}(\mathcal{I} + \eta^3) + \sigma^2 \otimes \tau^3 \otimes \frac{1}{2}(\mathcal{I} - \eta^3)$$

then interchanging the spin and valley labels, $\tau^a \leftrightarrow \eta^a$ and then conjugating with

$$\mathcal{I} \otimes \mathcal{I} \otimes \frac{1}{2}(\mathcal{I} + \eta^3) + \sigma^3 \otimes \mathcal{I} \otimes \frac{1}{2}(\mathcal{I} - \eta^3)$$

the resulting Hamiltonian is identical to the one with a mass term arising from the Kekule distortion that was discussed in [26] and [34]. This replacement amounts to an $SU(4)$ rotation and other orientations of the $SU(4)$ breaking terms in the Hamiltonian are possible as well. We also recall that in Ref.[34], when the mass is oriented as coming from a Kekule distortion, they add a pseudo-magnetic gauge field which couples to the Kekule condensate. In the present case where the condensate is antiferromagnetic order, the analogous gauge field would modify the covariant derivative as $\vec{\nabla} - i\frac{e}{c}\vec{A} \rightarrow \vec{\nabla} - i\frac{e}{c}\vec{A} - i\vec{V}\tau^3$ and it would add a term

$$\sigma^1 \otimes \tau^3 \otimes \eta^3 V_1 + \sigma^2 \otimes \tau^3 \otimes \mathcal{I} V_2$$

in (2). In the following, we shall include this gauge field in our analysis. In the case where the condensate is antiferromagnetic, it amounts to gauging the symmetry that rotates the antiferromagnetic order parameter, as might be done to describe a spin liquid.

3. ZERO MODES

Since the valley degrees of freedom in (1) are decoupled, we can perform our analysis for each valley separately. Consider the Hamiltonian describing electrons in one of the valleys (for simplicity, in units where $\hbar = v_f = \frac{e}{c} = 1$),

$$h = i\sigma^i \otimes \mathcal{I} \nabla_i + \sigma^i \otimes \mathcal{I} A_i + \sigma^i \otimes \tau^3 V_i + \sigma^3 \otimes \tau^a m_a \quad (5)$$

As we have suppressed the valley degree of freedom (η^a are missing), this is a 4×4 matrix differential operator.

We will search for solutions of $h\psi_0 = 0$. We note that the Hamiltonian (5) anticommutes with the matrix $\sigma^3 \otimes \tau^3$,

$$\sigma^3 \otimes \tau^3 h + h \sigma^3 \otimes \tau^3 = 0 \quad (6)$$

Zero modes of h can also be chosen to be eigenvectors of $\sigma^3 \otimes \tau^3$ and we shall show below that all of the zero modes are indeed eigenvectors with eigenvalues $\sigma^3 \otimes \tau^3 = -\text{sign}(m)$. This implies that the sublattice position and the spin orientation of the zero mode are correlated. One sublattice has $\sigma^3 = 1$ and on that sublattice the spin is polarized as $\tau^3 = \text{sign}(n)$. The other sublattice has $\sigma^3 = -1$ and the opposite spin polarization $\tau^3 = -\text{sign}(n)$. Furthermore, we shall find that, close to the vortex core, the zero modes have support on both sublattices. On the other hand, far from the vortex core, the zero modes have support on only one of the sublattices and therefore have one particular orientation of the spin. Which of the two possible sublattices is chosen depends on the orientation of the external magnetic field. Also, we note that the spin polarizations that we are discussing here are always orthogonal to the spin polarization of the condensate which is in the τ^1 - τ^2 -plane.

Also note that, if we include the second valley, its Hamiltonian differs from (5) by conjugation with the matrix $\mathcal{I} \otimes \sigma^2 \otimes \tau^3$ which interchanges the sublattices. Thus, for the other valley, there are also $|n|$ zero modes. They also have a peculiar spin polarization to sublattice relationship which is precisely the opposite of the one in the first valley.

We will begin with the special case of a rotationally covariant vortex with vorticity n : $(m_1, m_2) = m(r)(\cos n\theta, \sin n\theta)$ where $m(r) \rightarrow \hat{m}$ at $r \rightarrow \infty$ as in (3) and $m(0) = 0$. We shall also assume that the magnetic fields B_A and $B_V = \vec{\nabla} \times \vec{V}$ depend only on r . We will use the gauge $A_r = 0$ and $V_r = 0$ where $A_\theta(r) = \int_0^r r' dr' B_A(r')$ and $V_\theta(r) = \int_0^r r' dr' B_V(r')$. For example, if the magnetic fields are constants, the vector potentials would be $A_\theta(r) = \frac{B_A}{2} r^2$ and $V_\theta(r) = \frac{B_V}{2} r^2$. When written in polar coordinates, the Dirac equation for a zero mode is

$$m(r)e^{in\tau^3\theta}\tau^1 u + ie^{-i\theta} \left(\partial_r - \frac{i}{r}\partial_\theta - \frac{A_\theta + \tau^3 V_\theta}{r} \right) v = 0 \quad (7)$$

$$ie^{i\theta} \left(\partial_r + \frac{i}{r}\partial_\theta + \frac{A_\theta + V_\theta\tau^3}{r} \right) u - m(r)e^{in\tau^3\theta}\tau^1 v = 0 \quad (8)$$

The spinor components u and v are eigenstates of pseudo-spin $\sigma^3 \otimes \mathcal{I}$ with eigenvalues $+1$ and -1 , respectively. Formally, they can be defined as

$$u = \frac{1}{2} (\mathcal{I} + \sigma^3) \otimes \mathcal{I} \psi_0, \quad v = \frac{1}{2} (\mathcal{I} - \sigma^3) \otimes \mathcal{I} \psi_0$$

These project two two-component spinors, u and v , from the four-component spinor ψ_0 . u and v each have two

spin components on which the Pauli matrices τ^a act. We make the Ansätze

$$u(r, \theta) = e^{i\ell\theta} \tau^1 \tilde{u}(r) \quad (9)$$

$$v(r, \theta) = e^{-ik\theta} \tilde{v}(r) \quad (10)$$

Plugging this into (7) and (8) yields

$$m(r)e^{i(\ell+k+n\tau^3+1)\theta}\tilde{u} + i \left(\partial_r - \frac{k + A_\theta + \tau^3 V_\theta}{r} \right) \tilde{v} = 0$$

$$ie^{i(\ell+k+n\tau^3+1)\theta} \left(\partial_r - \frac{\ell - A_\theta + V_\theta\tau^3}{r} \right) u - m(r)\tilde{v} = 0$$

Now, we shall have to choose the spinors \tilde{u} and \tilde{v} to be eigenvectors of τ^3 , both with the same eigenvalue $-\text{sign}(n)$,

$$\tau^3 \tilde{u} = -\text{sign}(n)\tilde{u}, \quad \tau^3 \tilde{v} = -\text{sign}(n)\tilde{v}$$

(The reason for this choice of sign will become clear when we find the possible values of k and ℓ which both will have to be positive and have to obey (11) below.) \tilde{u} and \tilde{v} now each have only one non-zero component.

Then, the θ -dependent phases cancel if

$$\ell + k = |n| - 1 \quad (11)$$

which we shall assume holds. Consistency of (11) is a result of the correct choice of sign of the eigenvalues of τ^3 .

What remains is

$$m(r)\tilde{u} + i \left(\partial_r - \frac{k + A_\theta - \text{sign}(m)V_\theta}{r} \right) \tilde{v} = 0 \quad (12)$$

$$i \left(\partial_r - \frac{\ell - A_\theta - \text{sign}(m)V_\theta}{r} \right) \tilde{u} - m(r)\tilde{v} = 0 \quad (13)$$

Assuming that A_θ , V_θ and $m(r)$ go to zero at small r , by examining (12) and (13) at $r \sim 0$, we see that the small r behavior of the wave-functions must be

$$\tilde{u}(r) \sim r^k \tilde{u}_0 \quad (14)$$

$$\tilde{v}(r) \sim r^\ell \tilde{v}_0 \quad (15)$$

We shall need both of these solutions in order to match the solution at $r \rightarrow \infty$ where there will generally only be one solution. Both solutions are normalizable at $r = 0$ if $\ell = 0, 1, 2, \dots$ and if $k = 0, 1, 2, \dots$. However, remembering that k and ℓ are related by (11), we see that the allowed solutions are

$$(\ell, k) = (0, |n| - 1), (1, |n| - 2), \dots, (|n| - 1, 0), \quad |n| \geq 1 \quad (16)$$

We conclude that there are exactly $|n|$ possible solutions where both behaviors are normalizable at the origin.

The spinor components \tilde{u} and \tilde{v} are then both eigenvectors of spin $\mathcal{I} \otimes \tau^3$ with the same eigenvalue: $\mathcal{I} \otimes \tau^3 \tilde{u} = -\text{sign}(n)\tilde{u}$ and $\mathcal{I} \otimes \tau^3 \tilde{v} = -\text{sign}(n)\tilde{v}$. Recalling that u and \tilde{u} differ by a factor of $\mathcal{I} \otimes \tau^1$ which flips the sign of

the eigenvalue of $\mathcal{I} \otimes \tau^3$, we conclude that u and v are eigenvectors of spin with

$$\mathcal{I} \otimes \tau^3 u = +\text{sign}(n)u \quad (17)$$

$$\mathcal{I} \otimes \tau^3 v = -\text{sign}(n)v \quad (18)$$

Also, we recall that u and v are eigenvectors of pseudo-spin $\sigma^3 \otimes \mathcal{I}$:

$$\sigma^3 \otimes \mathcal{I} u = + u \quad (19)$$

$$\sigma^3 \otimes \mathcal{I} v = - v \quad (20)$$

Combining the eigenvalues tells us that u and v are eigenvectors of $\sigma^3 \otimes \tau^3$ with the same eigenvalue, $\text{sign}(n)$:

$$\sigma^3 \otimes \tau^3 u = \text{sign}(m)u \quad (21)$$

$$\sigma^3 \otimes \tau^3 v = \text{sign}(m)v \quad (22)$$

or, more succinctly

$$\sigma^3 \otimes \tau^3 \psi_0 = \text{sign}(m)\psi_0 \quad (23)$$

Remember that $\sigma^3 \otimes \tau^3$ anticommutes with the Hamiltonian, so we expect that the zero mode wave-functions ψ_0 are also eigenvectors of this matrix. Here, we see explicitly that is the case and the eigenvalue is determined by the sign of the vorticity. This establishes the basis for the discussion after equation (6) above.

Now we shall consider the large r limit of (12) and (12). If A_θ and V_θ grow slower than r at $r \rightarrow \infty$ (meaning that B_A and B_V fall off faster than $1/r$), the solution at large r is identical to the one found by Jackiw and Rossi [14] and their conclusion that there are exactly $|n|$ zero modes applies here. To generalize this, we will assume that at least one of A_θ and V_θ grow faster than r as $r \rightarrow \infty$. (This means that $B_A(r)$ or $B_V(r)$ fall off slower than $1/r$ and one of both of the fluxes $\int d^2x B_A$ or $\int d^2x B_V$ diverge.) Then (12) and (13) are solved at large r by

$$\tilde{u}(r) \sim e^{\int_0^r \frac{dr'}{r'} (A_\theta(r') - \text{sign}(m)V_\theta(r'))} \tilde{u}_0 \quad (24)$$

$$\tilde{v}(r) \sim e^{\int_0^r \frac{dr'}{r'} (-A_\theta(r') - \text{sign}(m)V_\theta(r'))} \tilde{v}_0 \quad (25)$$

We must now choose \tilde{u}_0 and \tilde{v}_0 so that the wavefunctions are normalizable.

First consider the case where $|A_\theta| > |V_\theta|$. Then, if $A_\theta > 0$ at large r , we set $\tilde{u}_0 = 0$ and (25) is normalizable at $r \rightarrow \infty$. On the other hand, if $A_\theta < 0$, we set $\tilde{v}_0 = 0$ and (24) is normalizable. Thus we see that, in the case where $|A_\theta| > |V_\theta|$, there is always one normalizable solution of the Dirac equation at large r . Once it is given the appropriate angle dependence, $e^{i\ell\theta} \tau^1 \tilde{u}(r)$ and $e^{-ik\theta} \tilde{v}(r)$ with one of the $|n|$ allowed values of k and ℓ , and continued to small r , it will become a linear combination of the two normalizable solutions that we found at $r \sim 0$. We conclude that, when $|A_\theta| > |V_\theta|$, there are exactly $|n|$ zero modes, independent of the specific asymptotic behavior of A_θ . This is in line with the conclusions in Refs. [10] and [11].

In addition, in the large r regime, the solutions reside on either one or the other sublattice (either u or v is non-zero), depending on the sign of A_θ (and therefore the magnetic flux $\phi_A = \frac{1}{2\pi} \int d^2x B_A$). Since the spin and the pseudo-spin are correlated, all zero modes in the asymptotic region have one particular spin polarization – the eigenvalue of τ^3 is $-\text{sign}(\phi_A)\text{sign}(n)$ – the zero mode in the asymptotic region is polarized in one particular direction, orthogonal to the easy plane of the condensate. The direction depends on the sign of the total magnetic flux and the sign of the vorticity.

Now, consider the case where $|A_\theta| < |V_\theta|$. To understand this case more clearly, we observe that, if $\vec{V}(x)$ is in the Coulomb gauge, $\vec{\nabla} \cdot \vec{V} = 0$, we can rewrite h as

$$h = e^{\sigma^3 \otimes \tau^3 \chi_V(x)} \left\{ i\sigma^i \otimes \mathcal{I} \nabla_i + \sigma^i \otimes \mathcal{I} A_i + \sigma^3 \otimes \tau^a m_a \right\} e^{\sigma^3 \otimes \tau^3 \chi_V(x)} \quad (26)$$

where $\chi_V(x) = \frac{1}{2\pi} \int d^2x' \ln |\vec{x} - \vec{x}'| B_V(x')$. The zero mode problem for h can be solved by finding potential zero modes when $\vec{V} = 0$ and testing their normalizability at $r = \infty$ when they are multiplied by the additional factor $e^{-\sigma^3 \otimes \tau^3 \chi_V(x)}$. In the case that we are interested in, when $e^{\pm \chi_V}$ dominates the asymptotics, there are always two normalizable solutions with correlated pseudo-spin and spin so that $\sigma^3 \otimes \tau^3 \chi_V(\infty) > 0$. When we begin at small r with one of the spinors with behavior in (14) or (15), $\begin{bmatrix} e^{i\ell\theta} \tau^1 \tilde{u}(r) \\ 0 \end{bmatrix}$ or $\begin{bmatrix} 0 \\ e^{-ik\theta} \tilde{v}(r) \end{bmatrix}$ and extrapolate to large r , they should become linear combinations of the two normalizable solutions there. In this case, there are an infinite number of zero modes for $\ell = 0, 1, 2, \dots$ or $k = 0, 1, 2, \dots$ which comprise a two-fold degenerate Landau level at zero energy. We conclude that, when $|A_\theta| < |V_\theta|$, there is a Landau level at the Dirac point. Here, by ‘‘Landau level’’ we mean an infinite set of degenerate states and they exist whenever $B_V(r)$ falls off slower than $1/r$ at $r \rightarrow \infty$.

4. INDEX AND η -INVARIANT

The index of h is the number of zero modes which have eigenvalue of $\sigma^3 \otimes \tau^3 = +1$ minus the number with eigenvalue $\sigma^3 \otimes \tau^3 = -1$. For generic smooth profiles of \vec{A} , \vec{V} and m , the index is a topological invariant that depends only on the asymptotic values of these functions. To see this we need only remember that the positive and negative energy states of h are paired: If

$$h\psi_E = E\psi_E$$

then

$$h(\sigma^3 \otimes \tau^3 \psi_E) = -E(\sigma^3 \otimes \tau^3 \psi_E)$$

and

$$\psi_{-E} = \sigma^3 \otimes \tau^3 \psi_E$$

If we deform the background fields in h smoothly, generally the eigenvalues of h move and some modes that are at $E = 0$ could move away from $E = 0$ and vice versa. However, since the non-zero modes come in positive and negative pairs, when modes move to or away from $E = 0$, they must do so in pairs. Furthermore, these pairs contain exactly one positive and one negative eigenstate of $\sigma^3 \otimes \tau^3$:

$$\sigma^3 \otimes \tau^3 (\psi_E + \psi_{-E}) = +(\psi_E + \psi_{-E})$$

and

$$\sigma^3 \otimes \tau^3 (\psi_E - \psi_{-E}) = -(\psi_E - \psi_{-E})$$

Thus the difference between the number of zero modes with $\sigma^3 \otimes \tau^3 = +1$ and $\sigma^3 \otimes \tau^3 = -1$, i.e. the index, is unchanged by a smooth deformation of the background fields.

In the discussion above, we found that when either the mass term or the magnetic field B_A dominate the asymptotics, the index is

$$\text{Index}(h) = n \quad (27)$$

On the other hand, when the pseudo-magnetic field B_V is dominant at $r \rightarrow \infty$, the index diverges, as it should equal the total number of states in two infinitely degenerate Landau levels,

$$\text{Index}(h) \sim \frac{1}{\pi} \int d^2x B_V(x) \quad (28)$$

(As we have discussed above, in order for V_θ to be more important than the mass term at $r \rightarrow \infty$, its fall off must be slow enough that the flux diverges at least logarithmically in the volume.) This quantity diverges at least logarithmically in the linear size of the 2-dimensional plane for the case in which it is valid and it would require more care to define it precisely.

Another interesting topological quantity is the η -invariant,

$$\eta(h_\epsilon) = \text{Tr sign}(h_\epsilon) \quad (29)$$

where

$$h_\epsilon = h + \sigma^3 \otimes \tau^3 \epsilon$$

It is of interest because the state of a system of electrons which is governed by the single-particle Hamiltonian h_ϵ and which has all negative energy levels occupied by electrons and all positive energy levels empty has electric charge given by the formula [17]

$$\langle Q \rangle = \frac{e}{2} \eta(h_\epsilon) \quad (30)$$

η is formally the difference of two infinite quantities, the number of states with positive energy and the number of states with negative energy. In interesting cases, these

partially cancel to leave a finite result. To make η unambiguous, we have added a term to the Hamiltonian, $h \rightarrow h_\epsilon = h + \sigma^3 \otimes \tau^3 \epsilon$. The term $\sigma^3 \otimes \tau^3 \epsilon$ is physically equivalent to an out-of-plane component of the antiferromagnetic order parameter which could be realistic. Since

$$\{h, \sigma^3 \otimes \tau^3\} = 0$$

the square of h_ϵ obeys

$$h_\epsilon^2 = h^2 + \epsilon^2 > 0$$

and h_ϵ has no zero modes. All of its eigenvalues are either positive or negative. Furthermore, since no matrix anti-commutes with h_ϵ , the spectrum of h_ϵ does not have particle-hole symmetry. Unpaired zero modes of h are also eigenfunctions of h_ϵ . They obey $h\psi_0 = 0$ and are eigenvectors of $\sigma^3 \otimes \tau^3$ with eigenvalues $+1$ or -1 . They are therefore eigenfunctions of h_ϵ with energies ϵ or $-\epsilon$. We can recover h by taking ϵ to zero. Then, the spectrum becomes entirely symmetric except for unpaired zero modes, which were assigned to the positive or negative energy states of $h + \sigma^3 \otimes \tau^3 \epsilon$.

η is a topological invariant which is entirely determined by the asymptotic behavior of $\vec{A}(x)$, $\vec{V}(x)$ and $m(x)$. We can easily derive a formula for η when we assume that the total flux $\phi_V = \frac{1}{2\pi} \int d^2x B_V(x)$ is finite, but the total electromagnetic flux $\phi_A = \frac{1}{2\pi} \int d^2x B_A(x)$ need not be. Computation of η is a straightforward generalization of that outlined in [36] and also [16],[17] and will be reviewed in the Appendix. The result is

$$\eta(h_\epsilon) = \left[\text{sign}(\epsilon) - \frac{\epsilon}{\sqrt{\epsilon^2 + \hat{m}^2}} \right] n + \frac{2\epsilon}{\sqrt{\epsilon^2 + \hat{m}^2}} \phi_V \quad (31)$$

To derive this formula, we have had to assume that $\vec{V}(x)$ is such that the covariant derivative of the mass term, $\vec{\nabla}m(x) - i\tau^3 \vec{V}(x)m(x)$ decays at least as fast as $1/r$ at large r . (For a vortex solution of Ginzburg-Landau action, this quantity generally decays exponentially.)

Like the index, η in (31) does not depend on the electromagnetic gauge field \vec{A} at all. There are a few interesting limits of (31). If we put $\epsilon \rightarrow 0^+$, we recover the index of h (27). If we put B_V to zero, we see that η has an irrational part, $-\frac{\epsilon}{\sqrt{\epsilon^2 + \hat{m}^2}} n$ which can be attributed to the polarization of continuum states by the vortex. Note that this polarization vanishes when $\epsilon \rightarrow 0$ and the continuum spectrum becomes symmetric. The irrational part of the fractional charge of a vortex in the absence of \vec{V} is identical to the one noted in Refs. [26],[35]-[36],[10]. Finally, if we decouple the vortex by setting the mass gap $\hat{m} \rightarrow 0$ we recover (28).

5. CONCLUSION

We have discussed scenarios where vortices can have mid-gap electronic bound states. The main conclusion

is that such bound states survive, practically unmodified in an external magnetic field. This in turn validates the consideration of vortices with mid-gap states as charge carriers in a Hall state.

We comment that, like polyacetylene, where spin degeneracy obscures fractional charge, so that polyacetylene actually has integer charged solitons, the system that we have discussed here has a valley degeneracy and the same phenomenon will occur. A vortex with a zero mode, actually has a pair of zero modes, one for each valley. Therefore, instead of being a two level system (zero mode occupied or unoccupied) where the states have charges $+\frac{1}{2}$ and $-\frac{1}{2}$, it is a four-level system with charges $-1, 0, 0, 1$. In polyacetylene, this leads to exotic spin-charge assignments, the charge ± 1 states of the soliton are spinless, whereas the two charge 0 states have spins $\pm\frac{1}{2}$, a topological version of spin-charge separation. A similar phenomenon would occur in graphene. When the mass condensate is a Kekule distortion, the vortex states would have spin degeneracy and the same exotic spin-charge assignments as occur in polyacetylene would now appear in two dimensions, If the condensate is an antiferromagnet, the spin is replaced by the residual SU(2) valley quantum number, and could be more difficult to see experimentally, as the symmetry is emergent and the conservation law resulting from it is not exact.

As for conductivity, as in polyacetylene, having vortices with low energy charged states makes the number of charged vortices available for conduction of electric current extremely sensitive to any displacement of the chemical potential from the charge neutral point. A signature of charged vortices of this kind should be, like in polyacetylene, an enormous variation of carrier concentration with doping.

Acknowledgments

The author acknowledges financial support from NSERC of Canada and the hospitality of the Aspen Center for Physics where part of this work was done. He also acknowledges Roman Jackiw for a discussion and some comments on the manuscript.

Appendix: Computation of the η -invariant

In this appendix, we outline the computation of η , the result of which was quoted in Eq. (31). We regulate the trace in (29) and represent the step function $\text{sign}(h)$ by an integral

$$\eta = \frac{2}{\pi} \lim_{\beta \rightarrow 0} \int_0^\infty d\omega \text{Tr} \left[\frac{h}{h^2 + \omega^2} e^{-\beta h^2} \right] \quad (\text{A.1})$$

which, using the fact that $\{\sigma^3 \tau^3, h\} = 0$, we replace by

$$\eta = \frac{2\epsilon}{\pi} \lim_{\beta \rightarrow 0} \int_0^\infty d\omega \text{Tr} \left[\sigma^3 \tau^3 \frac{e^{-\beta h^2}}{h^2 + \epsilon^2 + \omega^2} \right] \quad (\text{A.2})$$

We define the current

$$\vec{J}(x) = \text{tr} (x | i\vec{\sigma} \sigma^3 \tau^3 \frac{h_0 e^{-\beta h_0^2}}{h_0^2 + \epsilon^2 + \omega^2} | x) \quad (\text{A.3})$$

where tr indicates a trace over the Dirac and spin indices only. (Our previous trace, denoted Tr was a trace over Dirac and spin indices and position space as well $\text{Tr} \mathcal{O} = \int dx \text{tr} (x | \mathcal{O} | x)$.) Taking a divergence of the current yields the identity

$$\begin{aligned} \vec{\nabla} \cdot \vec{J} &= \vec{\nabla} \cdot \text{tr} (x | i\vec{\sigma} \sigma^3 \tau^3 \frac{h_0 e^{-\beta h_0^2}}{h_0^2 + \epsilon^2 + \omega^2} | x) \\ &= \text{tr} (x | h_0 \sigma^3 \tau^3 \frac{h_0 e^{-\beta h_0^2}}{h_0^2 + \epsilon^2 + \omega^2} - \sigma^3 \tau^3 \frac{h_0 e^{-\beta h_0^2} h_0}{h_0^2 + \epsilon^2 + \omega^2} | x) \\ &= -2 \text{tr} (x | \sigma^3 \tau^3 e^{-\beta h_0^2} | x) \\ &\quad + 2(\epsilon^2 + \omega^2) \text{tr} (x | \sigma^3 \tau^3 \frac{e^{-\beta h_0^2}}{h_0^2 + \epsilon^2 + \omega^2} | x) \end{aligned} \quad (\text{A.4})$$

Using the identity in (A.4), we rewrite (A.2) as

$$\begin{aligned} \eta &= \lim_{\beta \rightarrow 0} \int_0^\infty \frac{d\omega}{\pi} \frac{2\epsilon}{\epsilon^2 + \omega^2} \left\{ \text{Tr} \left[\sigma^3 \tau^3 e^{-\beta h_0^2} \right] \right. \\ &\quad \left. + \frac{1}{2} \int d^2 x \vec{\nabla} \cdot \text{tr} (x | i\vec{\sigma} \sigma^3 \tau^3 \frac{h_0 e^{-\beta h_0^2}}{h_0^2 + \epsilon^2 + \omega^2} | x) \right\} \quad (\text{A.5}) \end{aligned}$$

The first term in (A.5) is evaluated using an asymptotic expansion of the heat kernel,

$$\begin{aligned} (x | e^{-\beta h_0^2} | x) &= \frac{1}{4\pi\beta} + \frac{1}{4\pi} \left[\sigma^3 (B_A + \tau^3 B_V) \right. \\ &\quad \left. + \vec{\sigma} \times \vec{D}m - m^2 \right] + \mathcal{O}(\beta) \end{aligned} \quad (\text{A.6})$$

where $\vec{D}m = (\vec{\nabla} - 2i\tau^3 \vec{V})m$. When we multiply by $\sigma^3 \tau^3$ and take the trace, we see that the first term in (A.5) is

$$\lim_{\beta \rightarrow 0} \int_0^\infty \frac{d\omega}{\pi} \frac{2\epsilon}{\epsilon^2 + \omega^2} \text{Tr} \left[\sigma^3 \tau^3 e^{-\beta h_0^2} \right] = \text{sign}(\epsilon) \frac{1}{\pi} \int d^2 x B_V \quad (\text{A.7})$$

Now, we consider the second term on the right-hand-side of (A.5) which, using Gauss' theorem, can be written as a surface integral on the circle at $r = \infty$. To evaluate it, we must make some assumptions about the asymptotic behavior of the fields. We assume that the gauge field \vec{A} is such that B_A grows no faster than a constant at large r . This allows the case of a constant magnetic field. In this case asymptotically, in a radially symmetric gauge $A_\theta = \frac{B_A}{2} r^2$. We shall need to assume that the gauge field $\vec{V}(r)$ decays at least as fast as $1/r$ for large r so that $\vec{D}m \sim 1/r$

and $B_V \sim \frac{1}{r^2}$. We then can use a functional Taylor expansion of the integrand in powers of $\vec{D}m$, where it turns out that only the first (linear) order survives the traces and the asymptotic and $\beta \rightarrow 0$ limits. The trace of the zeroth order vanishes. The first and higher order terms in the expansion are finite in the limit $\beta \rightarrow 0$ which can now safely be taken. The first order in $\vec{D}m$ is

$$\lim_{\beta \rightarrow 0} \text{tr}(x | i\vec{\sigma}\sigma^3\tau^3 \frac{h_0 e^{-\beta h_0^2}}{h_0^2 + \epsilon^2 + \omega^2} | x) = \text{tr} i\vec{\sigma}\sigma^3\tau^3 \int d^2y \sigma^3 m(x) g(x, y) \vec{\sigma} \times \vec{D}m(y) g(y, x) \quad (\text{A.8})$$

where

$$g(x, y) = \langle x | \frac{1}{-D^2 - \sigma^3 B_A - \sigma^3 \tau^3 B_V + \hat{m}^2 + \epsilon^2 + \omega^2} | y \rangle \quad (\text{A.9})$$

Since the Green functions in (A.8) are short ranged and $|\vec{x}| \rightarrow \infty$, they will have support in the region where $|\vec{y}| \rightarrow \infty$ and the background fields in (A.9) can be re-

placed by their asymptotic values. Then, the Green function can be found explicitly,

$$g(x, y) = \frac{B}{4\pi} \int_0^\infty d\lambda \frac{e^{-\chi(x, y, \lambda)}}{\sinh B_A \lambda}$$

with $\chi(x, y, \lambda) = \frac{B_A}{4}(\vec{x} - \vec{y})^2 \coth B_A \lambda + \lambda(\sigma^3 B_A + \hat{m}^2 + \omega^2 + \epsilon^2) + i\frac{B_A}{2}\vec{x} \times \vec{y}$. We can now easily show that, because of the large r limit, the green functions $\dots g(x, y) \dots g(y, x) \dots$ in (A.8) can be replaced by $\frac{1}{4\pi} \frac{1}{\hat{m}^2 + \omega^2 + \epsilon^2} \delta^2(x - y)$. This can be seen by defining $\vec{x} = r\hat{x}$, rescaling $\vec{y} \rightarrow r\vec{y}$ and taking the large r limit. Then, we have

$$\lim_{\beta \rightarrow 0} \text{tr}(x | i\vec{r} \cdot \vec{\sigma}\sigma^3\tau^3 \frac{h_0 e^{-\beta h_0^2}}{h_0^2 + \epsilon^2 + \omega^2} | x) = -\frac{1}{\pi} \frac{\epsilon^{ab} m^a \partial_\theta m^b + 2m^2 V_\theta}{\hat{m}^2 + \omega^2 + \epsilon^2} \quad (\text{A.10})$$

Doing the remaining integrals, we obtain (31).

-
- [1] Y. Zhang, Z. Jiang, J.P. Small, M.S. Purewal, Y.-W. Tan, M. Fazlollahi, J.D. Chudow, J.A. Jaszczak, H.L. Störmer, and P. Kim, Phys. Rev. Lett. **96**, 136806 (2006).
- [2] D.A. Abanin, K.S. Novoselov, U. Zeitler, P.A. Lee, A.K. Geim, and L.S. Levitov, Phys. Rev. Lett. **98**, 196806 (2007).
- [3] Z. Jiang, Y. Zhang, H.L. Störmer, and P. Kim, Phys. Rev. Lett. **99**, 106802 (2007).
- [4] A.J.M. Giesbers, U. Zeitler, M.I. Katsnelson, L.A. Ponomarenko, T.M. Mohiuddin, and J.C. Maan, Phys. Rev. Lett. **99**, 206803 (2007).
- [5] J. G. Checkelsky, L. Li, and N. P. Ong, Phys. Rev. Lett. **100**, 206801 (2008).
- [6] J. G. Checkelsky, L. Li, and N. P. Ong, Phys. Rev. B **79**, 115434 (2009).
- [7] A. J. M. Giesbers, L. A. Ponomarenko, K. S. Novoselov, A. K. Geim, M. I. Katsnelson, J. C. Maan and U. Zeitler, Phys. Rev. B **80**, 201403(R) (2009).
- [8] L. Zhang, Y. Zhang, M. Khodas, T. Valla, I.A. Zaliznyak, arXiv:1003.2738.
- [9] Kentaro Nomura, Shinsei Ryu, and Dung-Hai Lee, Phys. Rev. Lett. **103**, 216801 (2009).
- [10] Chang-Yu Hou, Claudio Chamon, and Christopher Mudry, Phys. Rev. B **81**, 075427 (2010) arXiv:0909.2984v2 [cond-mat.mes-hall].
- [11] I. F. Herbut, arXiv:0910.4906 [cond-mat.mes-hall].
- [12] W. P. Su, J. R. Schrieffer and A. J. Heeger, Phys. Rev. Lett. **42**, 1698 (1979); R. Jackiw and J. R. Schrieffer, Nucl. Phys. B **190**, 253 (1981).
- [13] R. Jackiw and C. Rebbi, Phys. Rev. D **13**, 3398 (1976).
- [14] R. Jackiw and P. Rossi, Nucl. Phys. B **190**, 681 (1981).
- [15] E. J. Weinberg, Phys. Rev. D **24**, 2669 (1981).
- [16] A. J. Niemi and G. W. Semenoff, Phys. Rev. D **30**, 809 (1984).
- [17] A. J. Niemi and G. W. Semenoff, Phys. Rept. **135**, 99 (1986).
- [18] G. W. Semenoff, Phys. Rev. Lett. **53**, 2449 (1984).
- [19] K. Nomura, A.H. MacDonald, Phys. Rev. Lett. **96**, 256602 (2006).
- [20] J. Alicea and M.P.A. Fisher, Phys. Rev. B **74**, 075422 (2006).
- [21] M. O. Goerbig, R. Moessner, and B. Douçot, Phys. Rev. B **74**, 161407(R) (2006).
- [22] L. Sheng, D.N. Sheng, F.D.M. Haldane, and L. Balents, Phys. Rev. Lett. **99**, 196802 (2007).
- [23] D.A. Abanin, P.A. Lee, and L.S. Levitov, Phys. Rev. Lett. **96**, 176803 (2006).
- [24] D.A. Abanin, et al., Phys. Rev. Lett. **98**, 196806 (2007).
- [25] D.V. Khveshchenko, Phys. Rev. Lett. **87**, 206401 (2001).
- [26] C. Y. Hou, C. Chamon and C. Mudry, Phys. Rev. Lett. **98**, 186809 (2007) [arXiv:cond-mat/0609740].
- [27] V.P. Gusynin, V.A. Miransky, and I.A. Shovkovy, Phys. Rev. Lett. **73**, 3499 (1994); Phys. Rev. D **52**, 4718 (1995); Nucl. Phys. B **462**, 249 (1996).
- [28] V.P. Gusynin, V.A. Miransky, S.G. Sharapov, and I.A. Shovkovy, Phys. Rev. B **74**, 195429 (2006).
- [29] I. F. Herbut, Phys. Rev. B **76**, 085432 (2007); *ibid.* **75**, 165411 (2007).
- [30] J. N. Fuchs and P. Lederer, Phys. Rev. Lett. **98**, 016803 (2007).
- [31] N. A. Viet, H. Ajiki, T. Ando, J. Phys. Soc. Jpn. **63**, 3036 (1994), H. Ajiki, T. Ando, *ibid.* **64**, 260 (1995), *ibid.* **65**, 2976 (1996).
- [32] Y. Hatsugai, T. Fukui, H. Aoki, Physica E **40**, 1530 (2008).
- [33] I. F. Herbut, Phys. Rev. Lett. **99**, 206404 (2007) [arXiv:0704.2234 [cond-mat.str-el]].
- [34] R. Jackiw and S. Y. Pi, Phys. Rev. Lett. **100**, 110405 (2008); [arXiv:cond-mat/0701760].
- [35] C. Chamon, C. Y. Hou, R. Jackiw, C. Mudry, S. Y. Pi and A. P. Schnyder, Phys. Rev. Lett. **100**, 110405 (2008) [arXiv:0707.0293 [cond-mat.str-el]].

- [36] C. Chamon, C. Y. Hou, R. Jackiw, C. Mudry, S. Y. Pi and G. Semenoff, Phys. Rev. B **77**, 235431 (2008) [arXiv:0712.2439 [hep-th]].
- [37] If the condensate $m_a(x)$ were a constant, the residual symmetry would be $SU(2) \times SU(2)$.