## A monopole solution in a Lorentz-violating field theory

Michael D. Seifert

Dept. of Physics, Indiana University, 727 E. 3<sup>rd</sup> St., Bloomington, IN, 47405<sup>\*</sup>

I present a topological defect solution that arises in a theory where Lorentz symmetry is spontaneously broken by a rank-two antisymmetric tensor field, and I discuss its observational signatures.

PACS numbers: 11.27.+d,11.30.Cp,11.30.Qc,14.80.-j

Lorentz symmetry has been one of the cornerstones of physics for over a century. The possibility that this symmetry is approximate rather than exact is an exciting prospect for the detection of physics beyond the Standard Model, and has been the focus of many experimental searches (see Ref. [\[1\]](#page-3-1) and references therein.) Such symmetry breaking also arises in many theoretical models of current interest, including string theory [\[2\]](#page-3-2) and non-commutative geometry [\[3\]](#page-3-3).

A large class of theoretical models in which Lorentz symmetry is broken involve a field which spontaneously breaks this symmetry [\[2,](#page-3-2) [4–](#page-3-4)[6\]](#page-3-5). In such models, a vector or tensor field is postulated to have a potential energy which is minimized (and vanishes) at some field value other than zero. This behaviour is closely analogous to that of the Higgs field in the Standard Model. However, unlike the (scalar) Higgs field, a tensor field with a non-zero expectation value will provide a preferred geometric structure to Minkowski spacetime, and the vacuum solution will not have the full symmetry of the Lorentz group. Direct couplings between such a "Lorentz-violating" field and conventional matter fields will then affect the dynamics of conventional matter, just as direct Yukawa couplings between the Higgs field and fermion fields in the Standard Model give mass to the fermion fields [\[7\]](#page-3-6).

An important class of solutions that arise in theories with a spontaneously broken symmetry are topological defects. These solutions arise when the vacuum manifold of a symmetry-breaking field (i.e., the set of field values which minimize the symmetry-breaking potential) has certain non-trivial topological properties. There can then arise "domain wall," "cosmic string," or "monopole" solutions (respectively) to the equations of motion; on large scales, these solutions effectively have two, one, and zero spatial dimensions. In theories in which topological defect solutions are allowed, they will generically arise during phase transitions in the early Universe, via the Kibble mechanism [\[8\]](#page-3-7). Since isolated defects are stable, it is possible that such relics still exist in the Universe today. The presence of topological defects can also have important implications for cosmology, including inflation [\[9\]](#page-3-8) and structure formation [\[10\]](#page-3-9).

The purpose of this letter is to show that topological defect solutions can arise in theories with a tensor field that breaks Lorentz symmetry. Such solutions have not previously been described in theories where the symmetry broken is Lorentz symmetry. We will derive an explicit example of such a solution, and discuss some of its phenomenological properties.

Our theory contains an antisymmetric two-tensor field (also known as a notoph field [\[11\]](#page-3-10) or Kalb-Ramond field [\[12\]](#page-3-11)) which takes on a background expectation value [\[6\]](#page-3-5). Its action is of the form

<span id="page-0-1"></span>
$$
S_B = \int d^4x \left( -\frac{1}{6} F^{abc} F_{abc} - \frac{\lambda}{2} (B^{ab} B_{ab} - b^2)^2 \right), \quad (1)
$$

where  $B_{ab}$  is an antisymmetric tensor field and  $F_{abc}$  =  $3\partial_{[a}B_{bc]}$  is its associated field strength [\[22\]](#page-3-12). The equations of motion derived from this action are

<span id="page-0-0"></span>
$$
\partial^c F_{cab} - 2\lambda (B^{cd} B_{cd} - b^2) B_{ab} = 0.
$$
 (2)

Our first task is to show that the vacuum manifold of this theory has the correct topology. For a localized topological defect solution to exist in three spatial dimensions, the vacuum manifold must either be disconnected, contain a non-contractible loop, or contain a non-contractible two-sphere; mathematically, it is necessary that one of the vacuum manifold's homotopy groups  $\pi_0(M_{\text{vac}}), \pi_1(M_{\text{vac}}), \text{ or } \pi_2(M_{\text{vac}})$  be non-trivial. From  $(2)$ , we can see that any constant field  $B_{ab}$  such that  $B_{ab}B^{ab} = b^2$  is a solution of the equation of motion. In terms of Cartesian coordinate components, this condition is

$$
-2\sum_{i}(B_{0i})^{2} + 2\sum_{i\n(3)
$$

where the summations run over spatial indices (i.e.,  $\{i, j\} = 1, 2, 3$ .) This condition defines a five-dimensional submanifold of the six-dimensional field space of  $B_{ab}$ . This submanifold is homeomorphic to  $S^2 \times \mathbb{R}^3$ : for any choice of the three components  $B_{0i}$ , the three components  $B_{ij}$  are constrained to lie on a sphere whose radius squared is  $\frac{1}{2}b^2 + B_{0i}B_0^i$ . Thus, the topology of the vacuum manifold allows for a monopole solution, since it contains a non-contractible two-sphere (i.e.,  $\pi_2(M_{\text{vac}})$  = Z.)

These topological conditions on the vacuum manifold are necessary but not sufficient for the existence of a monopole solution; we must still find a solution of the equations of motion valid throughout space. If such solutions exist in this theory, we expect the simplest ones to be static and spherically symmetric. In standard spherical coordinates, the most general antisymmetric twotensor with this property can be written in terms of two functions of r:

$$
B_{tr} = -B_{rt} = f(r), \quad B_{\theta\phi} = -B_{\phi\theta} = g(r)r^2\sin\theta.
$$
 (4)

In terms of this ansatz, the equation of motion [\(2\)](#page-0-0) has two non-vanishing components, telling us that

<span id="page-1-0"></span>
$$
-2\lambda(-2f^2 + 2g^2 - b^2)f = 0
$$
 (5a)

and

<span id="page-1-1"></span>
$$
\frac{\partial}{\partial r}\left(\frac{\partial g}{\partial r} + \frac{2}{r}g\right) - 2\lambda(-2f^2 + 2g^2 - b^2)g = 0.
$$
 (5b)

From the first of these equations [\(5a\)](#page-1-0), we see that either f must vanish or  $B_{ab}$  must be in its vacuum manifold throughout spacetime. Since we only want our field to approach the vacuum manifold asymptotically, we set  $f = 0$ . Defining  $\tilde{r}$  and  $\tilde{g}$  such that  $g = b\tilde{g}/\sqrt{2}$  and  $r = \tilde{r}/(\sqrt{2\lambda b})$ , the second equation [\(5b\)](#page-1-1) becomes

$$
\frac{\partial}{\partial \tilde{r}} \left( \frac{\partial \tilde{g}}{\partial \tilde{r}} + \frac{2}{\tilde{r}} \tilde{g} \right) - (\tilde{g}^2 - 1)\tilde{g} = 0.
$$
 (6)

This equation is identical (up to rescaling) to that obeyed by topological defects arising in a theory containing a triplet of Lorentz scalars with a spontaneously broken global  $O(3)$  symmetry [\[13\]](#page-3-13), for which a solution broken global  $O(3)$  symmetry [13], for which a solution<br>with  $g(0) = 0$  and  $g(r) \to b/\sqrt{2}$  as  $r \to \infty$  is known to exist. No closed-form expression for this solution is known, although series techniques [\[14\]](#page-3-14) or numerical integration [\[15\]](#page-3-15) can be used to approximate it. Asymptotically, the solution can be shown to obey

$$
\tilde{g}(\tilde{r}) = 1 - \frac{1}{2\tilde{r}^2} - \frac{3}{2\tilde{r}^4} + \dots
$$
 (7)

In the absence of any direct coupling between  $B_{ab}$  and conventional matter fields, the principal method of detection of these monopoles will be gravitational, via the coupling of their stress-energy to the metric. The stressenergy of the field  $B_{ab}$  in a flat background is

$$
T_{ab} = F_{acd}F_b{}^{cd} - \frac{1}{6}\eta_{ab}F_{cde}F^{cde}
$$

$$
-\frac{\lambda}{2}\eta_{ab}(B^2 - b^2)^2 + 4\lambda(B^2 - b^2)B_{ac}B_b{}^c, \quad (8)
$$

where  $B^2 \equiv B_{ab}B^{ab}$ .

While the field profile of our tensor monopole is identical to that of the  $O(3)$  scalar monopole (as noted above), their respective stress-energies differ in two important respects. First, since the kinetic term for  $B_{ab}$  in the action [\(1\)](#page-0-1) is not simply of a "gradient-squared" form, the kinetic terms in the stress-energy tensor will differ from the analogous terms in the scalar monopole stress-energy.



<span id="page-1-3"></span>FIG. 1: Densities and pressures of the antisymmetric tensor monopole and the  $O(3)$  scalar monopole. Overall scaling for the  $O(3)$  stress-energy is chosen so that the asymptotic fall-off of  $\rho$  is the same in both solutions.

Second, the last term in Equation [\(8\)](#page-1-2) has no analogue in the scalar case. It arises from the differentiation of the potential  $V(B^2)$  with respect to the metric, since the "square" of a tensor field (unlike that of a scalar) depends on the metric.

The components of the stress-energy for our tensor monopole are compared to those of the  $O(3)$  scalar monopole in Figure [1.](#page-1-3) Notably, while the radial and tangential pressures of the  $O(3)$  monopole are negative, those of our tensor monopole are positive; this difference is primarily due to the differences between the kinetic terms of the two theories. In flat spacetime, the leadingorder asymptotic behaviour of the energy density  $\rho$ , the radial pressure  $P_r$ , and the tangential pressure  $P_\theta$  can be shown to be

$$
\rho \approx P_r \approx \frac{4\lambda b^4}{\tilde{r}^2}, \qquad \qquad P_\theta \approx \frac{\lambda b^4}{\tilde{r}^4}.
$$
 (9)

To see the gravitational effects of our solution, we promote the metric  $g_{ab}$  to a dynamical field. Assuming spherical symmetry and staticity, we can write our line element as

<span id="page-1-2"></span>
$$
ds^{2} = -M^{2}(r)dt^{2} + N^{2}(r)dr^{2} + r^{2}(d\theta^{2} + \sin^{2}\theta d\phi^{2}).
$$
 (10)

The independent components of the Einstein equation are then equivalent to

<span id="page-1-4"></span>
$$
\frac{2 N'}{r N} + \frac{1}{r^2} (N^2 - 1)
$$
  
=  $\frac{\epsilon}{2b^2} \left( \left( g' + \frac{2}{r} g \right)^2 + \frac{\lambda}{2} N^2 (2g^2 - b^2)^2 \right)$  (11a)

and

$$
\frac{2}{r}\frac{M'}{M} - \frac{1}{r^2}(N^2 - 1) \n= \frac{\epsilon}{2b^2} \left( \left( g' + \frac{2}{r}g \right)^2 - \frac{\lambda}{2} N^2 (2g^2 - b^2)^2 \right), \quad (11b)
$$

where primes denote differentiation with respect to  $r$  and  $\epsilon \equiv 16\pi G b^2$ . Meanwhile, the field equation of motion becomes

$$
\frac{N}{M}\frac{\partial}{\partial r}\left(\frac{M}{N}\left(g' + \frac{2}{r}g\right)\right) - 2\lambda N^2(2g^2 - b^2)g = 0 \quad (12)
$$

While this system of equations is non-linear and rather intractable, in the context of gravitational fields we are most interested in the far-field effects. A useful approximation for investigating this regime in monopole solutions (though one that must be used with some care in this case [\[16\]](#page-3-16)) is the BPS limit [\[17,](#page-3-17) [18\]](#page-3-18). In this limit, we take  $\lambda \to 0$  and look for solutions of the Einstein equation [\(11\)](#page-1-4) where the field takes on its asymptotic value tion [\(11\)](#page-1-4) where the field takes on its asymptotic value  $g = b/\sqrt{2}$  everywhere. In this limit, the equations (11) have the exact solution

$$
M^{2}(r) = C_{2} \frac{\tilde{r}^{1+\epsilon} + C_{1}}{\tilde{r}^{1-\epsilon}} \tag{13}
$$

$$
N^{2}(r) = (1+\epsilon)\frac{\tilde{r}^{1+\epsilon}}{\tilde{r}^{1+\epsilon} + C_{1}},\tag{14}
$$

where  $C_1$  and  $C_2$  are constants of integration, the latter of which can be set to unity via rescaling of t.

As noted above, the field profiles of our antisymmetric tensor monopoles and those of global  $O(3)$  monopoles are quite similar; however, their gravitational fields have important qualitative differences. Both cases share the same asymptotic spatial geometry: for both, the slices of constant t are simply flat space with a spherical deficit angle. However, unlike the scalar monopole case, the component  $g_{tt}$  in our solution grows without bound as  $r \to \infty$ : from [\(13\)](#page-2-0), we see that  $M(r) \propto r^{\epsilon}$ . This exceedingly slow divergence is due to the slow fall-off of the stress-energy components, specifically the combination  $\rho + P_r + 2P_\theta$  [\[23\]](#page-3-19). Such a power-law divergence might indicate that the solution for an isolated monopole is fundamentally non-static; this would be analogous to antide Sitter space written in spherical coordinates, where  $g_{tt} \propto r^2$  asymptotically. In a more physically realistic situation, we only expect this solution to be valid out to some finite radius, where the effects of larger structure (on galactic or cosmic scales) take over. Since we expect to have  $\epsilon \ll 1$ , the far-field geometry should be sufficiently flat that we can "patch" our solution into one describing the appropriate larger-scale structure.

The effects of this geometry on test particles, specifically test photons, would be the primary method by which these monopoles could be directly detected. Two main effects on photons can be envisioned: gravitational redshift and the bending of light rays. The first of these can be shown to be minimal: if our mass scale  $b$  is well below the Planck scale, the fractional gravitational redshift experienced by a photon in this background will be within no more than two orders of magnitude of  $\epsilon$  [\[16\]](#page-3-16). Such effects will thus be negligible.

A more interesting effect arises in the deflection of light rays in this background. Using standard techniques, we can calculate that to leading order a light ray propagating in this background will be deflected by an angle [\[16\]](#page-3-16)

$$
\delta\phi \approx \frac{3\pi}{2}\epsilon. \tag{15}
$$

Note that at this order, the deflection angle is independent of the "apparent impact parameter." Rather, with respect to light bending, the spacetime behaves as though it has a spherical deficit angle of  $\frac{3}{2}\epsilon$ . If a monopole were perfectly aligned between an observer and a point source, the observer would see this point source as a ring with angular diameter  $\delta \phi \times l/(d+l)$ , where d is the distance from the observer to the monopole and  $l$  the distance from the monopole to the source. A monopole slightly off of the line of sight (but still sufficiently close to it) would give rise to two images on the sky at this same angular separation.

<span id="page-2-0"></span>From this point of view, the signature of an antisymmetric tensor monopole is effectively the same as that of a global  $O(3)$  monopole: both give rise to apparent spherical deficit angles, albeit with differing dependence on their respective mass scales. Any lensing-derived bounds placed on the current cosmological abundance of global  $O(3)$  monopoles will thus have a bearing on the abundance of our tensor monopoles as well, and vice versa.

I am unaware of any direct observational searches specifically targeted at this type of gravitational lensing. However, a search for angular deficits arising from cosmic strings, using images obtained as part of the Great Observatories Origins Deep Survey, was recently performed [\[19\]](#page-3-20). This search looked for morphologically similar galaxies correlated on opposite sides of a line on the sky. Similar techniques could be used to search for global monopoles; in this case the doubled images would be correlated inside and outside of a circle on the sky rather than on opposite sides of a line. While such structures are expected to be rare (see below), a search for double images of distant galaxies would yield important information; even if no monopoles were detected, one could in principle use a null result from such a search to bound the mass scale b associated with our tensor field.

Other bounds on the abundance of  $O(3)$  monopoles have also been derived [\[20,](#page-3-21) [21\]](#page-3-22). While such bounds are likely adaptable to the present case, it is important to emphasize that they are not immediately applicable in the same way that gravitational lensing bounds would be. The main reason for this difference is that the stress-energy (and thus the far-field metric) of our tensor monopole has important qualitative differences from that of the  $O(3)$  monopole; in particular, the combination  $\rho + P_r + 2P_\theta$  falls off as  $r^{-2}$  for our tensor monopole, but as  $r^{-4}$  (due to cancellation of the  $r^{-2}$  dependence) for the  $O(3)$  scalar monopole.

A final question is what the expected monopole density in the current Universe should be. While the Kibble mechanism predicts that one monopole per Hubble volume (to within an order of magnitude) should form during a phase transition in the early Universe, the subsequent field dynamics might cause monopoles and antimonopoles to recombine in the subsequent cosmological evolution. Many of the arguments and counterarguments concerning the recombination of global  $O(3)$ scalar monopoles [\[13\]](#page-3-13) would seem to apply here. As in the scalar monopole case, a definitive answer to this question is likely to require computational simulation.

Such simulations of scalar monopoles [\[21\]](#page-3-22) have found that the density of such structures remains at approximately four monopoles per Hubble volume throughout both the radiation- and matter-dominated eras. However, the above noted differences between the gravitational fields of scalar monopoles and tensor monopoles might cause recombination to proceed quite differently in the tensor case. While it still seems plausible that the results from scalar monopole simulations would hold for our model as well (up to an order of magnitude), such a statement should be regarded only as a conjecture.

If such a monopole were to be detected, it would obviously be of great import to physics. Such an observation would represent both the first field observed (other than the metric) that is not described by the Standard Model, and the first field observed to break Lorentz symmetry. As such, observation of a topological defect of this type would provide invaluable insight into the roles played by fundamental symmetries in physics.

Helpful discussions with B. Altschul, V. A. Kostelecký, and M. Uhlmann concerning this work are gratefully acknowledged. This research was supported in part by the United States Department of Energy under Grant No. DE-FG02-91ER40661.

- <span id="page-3-0"></span><sup>∗</sup> Electronic address: [mdseifer@indiana.edu](mailto:mdseifer@indiana.edu)
- <span id="page-3-1"></span>[1] V. A. Kostelecký and N. Russell, arXiv  $0801.0287v3$  $(2010)$ .
- <span id="page-3-2"></span>[2] V. A. Kostelecký and S. Samuel, Phys. Rev. D 40, 1886 (1989).
- <span id="page-3-3"></span>[3] S. M. Carroll, J. A. Harvey, V. A. Kostelecký, C. D. Lane, and T. Okamoto, Phys. Rev. Lett. 87, 141601 (2001).
- <span id="page-3-4"></span>[4] V. A. Kostelecký, Phys. Rev. D 69, 105009 (2004).
- [5] R. Bluhm and V. A. Kostelecký, Phys. Rev. D **71**, 065008 (2005).
- <span id="page-3-5"></span>[6] B. Altschul, Q. G. Bailey, and V. A. Kostelecký, Phys. Rev. D 81, 065028 (2010).
- <span id="page-3-6"></span>[7] D. Colladay and V. A. Kostelecký, Phys. Rev. D 58, 116002 (1998).
- <span id="page-3-7"></span>[8] T. W. B. Kibble, J. Phys. A 9, 1387 (1976).
- <span id="page-3-8"></span>[9] A. H. Guth, Phys. Rev. D 23, 347 (1981).
- <span id="page-3-9"></span>[10] R. Durrer, M. Kunz, and A. Melchiorri, Phys. Rep. 364, 1 (2002).
- <span id="page-3-10"></span>[11] V. I. Ogievetskii and I. V. Polubarinov, Sov. J. Nuc. Phys. 4, 156 (1967).
- <span id="page-3-11"></span>[12] M. Kalb and P. Ramond, Phys. Rev. D 9, 2273 (1974).
- <span id="page-3-13"></span>[13] M. Barriola and A. Vilenkin, Phys. Rev. Lett. **63**, 341 (1989).
- <span id="page-3-14"></span>[14] X. Shi and X. Li, Class. Quant. Grav. 8, 761 (1991).
- <span id="page-3-15"></span>[15] D. Harari and C. Loustó, Phys. Rev. D 42, 2626 (1990).
- <span id="page-3-16"></span>[16] M. D. Seifert, in preparation.
- <span id="page-3-17"></span>[17] E. B. Bogomol'nyi, Sov. J. Nuc. Phys. 24, 449 (1976).
- <span id="page-3-18"></span>[18] M. Prasad and C. Sommerfield, Phys. Rev. Lett. **35**, 760 (1975).
- <span id="page-3-20"></span>[19] J. Christiansen, E. Albin, K. James, J. Goldman, D. Maruyama, and G. Smoot, Phys. Rev. D 77, 123509 (2008).
- <span id="page-3-21"></span>[20] W. Hiscock, Phys. Rev. Lett. **64**, 344 (1990).
- <span id="page-3-22"></span>[21] D. P. Bennett and S. H. Rhie, Phys. Rev. Lett. 65, 1709 (1990).
- <span id="page-3-12"></span>[22] We will use a metric signature of  $(-, +, +, +)$ , and units where  $c = h = 1$ .
- <span id="page-3-19"></span>[23] This combination is proportional to the tt-component of the trace-reversed stress-energy tensor, which couples to the tt-component of the metric in the linearized approximation.