

Solar and Atmospheric Neutrinos: Limitations for Direct Dark Matter Searches

A. Gütlein, W. Potzel, C. Ciemniak, F. von Feilitzsch,
N. Haag, M. Hofmann, C. Isaila, T. Lachenmaier,
J.-C. Lanfranchi, L. Oberauer, S. Pfister, S. Roth,
M. von Sivers, R. Strauß, and A. Zöller

Physik-Department E15,
Technische Universität München, D-85748 Garching,
Germany

September 10, 2021

Abstract

In experiments for direct dark matter searches, neutrinos coherently scattering off nuclei can produce similar events as Weakly Interacting Massive Particles (WIMPs). To reach sensitivities better than $\sim 10^{-10}$ pb for the elastic WIMP nucleon spin-independent cross section in the zero-background limit, energy thresholds for nuclear recoils should be $\gtrsim 2.05$ keV for CaWO_4 , $\gtrsim 4.91$ keV for Ge, $\gtrsim 2.89$ keV for Xe, $\gtrsim 8.62$ keV for Ar and $\gtrsim 15.93$ keV for Ne as target material. Atmospheric neutrinos limit the achievable sensitivity for the background-free direct dark matter search to $\gtrsim 10^{-12}$ pb.

Coherent Neutrino Nucleus Scattering (CNNS) is a neutral-current interaction (exchange of a virtual Z^0 boson) and thus independent of the neutrino flavour. In a CNNS event, momentum and recoil energy are transferred to the nucleus. For low transferred momenta where the Z^0 -wavelength and the radius of the nucleus are about equal, the neutrino scatters off all nucleons coherently leading to an enhanced cross section [1].

The count rate R_{th} in a detector due to CNNS above a threshold E_{th} of

the nuclear recoil energy is given by [2]

$$R_{th} = \int_{E_{th}}^{\infty} dE_{rec} \frac{dR(E_{rec})}{dE_{rec}} = N_t \int_{E_{th}}^{\infty} dE_{rec} \int_{\sqrt{\frac{E_{rec}M}{2}}}^{\infty} dE_{\nu} \Phi(E_{\nu}) \frac{d\sigma(E_{\nu}, E_{rec})}{dE_{rec}}, \quad (1)$$

where M is the mass of the target nucleus, E_{rec} its recoil energy; N_t is the total number of target nuclei, $\Phi(E_{\nu})$ the neutrino flux at a neutrino energy E_{ν} ; $\frac{dR(E_{rec})}{dE_{rec}}$ is the recoil spectrum and $\frac{d\sigma(E_{\nu}, E_{rec})}{dE_{rec}}$ is the differential scattering cross section which includes the Helm form factor [3], [2], see below.

Assuming an isothermal WIMP halo, the recoil spectra (see figure 1)




Figure 1: Recoil spectra of WIMPs for different target materials assuming a WIMP mass of 60 GeV and a WIMP-nucleon cross section of 10^{-44} cm².

of WIMPs scattering coherently off various target materials were calculated according to [4]

$$\frac{dR(E_{rec})}{dE_{rec}} = \frac{c_1 N_A \rho_D}{2\sqrt{\pi} \mu_1^2} \sigma_{WN} |F(q)|^2 v_0 \frac{A^2}{E_0} e^{-\frac{c_2 E_{rec}}{E_0 r}} \quad (2)$$

where c_1 , c_2 are constants describing the annular modulation of the WIMP flux, N_A is Avogadro's number, ρ_D the local WIMP density, μ_1 the reduced mass for $A=1$, σ_{WN} the WIMP-nucleon cross section, $F(q)$ is the Helm form factor [3], v_0 the velocity of the earth relative to the galaxy, A the mass number, E_0 the kinetic energy of the WIMPs; $r = 4 \frac{M_D M_T}{(M_D + M_T)^2}$ a kinematic factor, M_D the WIMP mass and M_T the mass of the target nucleus.

Since recoil energies due to solar neutrinos are small, such events can be rejected by a proper choice of the energy threshold. If one allows only 0.1 counts per ton-year, the energy regions listed in the second column of table 1 are obtained. As an upper limit, 100 keV was assumed. This value is

Material	energy region solar neutrinos	max. exposure	energy region atmosph. neutrinos
Ne	15.93 - 100 keV	35.9 ton-years	15.93 - 100 keV
Ar	8.62 - 100 keV	9.2 ton-years	8.62 - 100 keV
Ge	4.91 - 100 keV	4.3 ton-years	4.91 - 100 keV
Xe	2.89 - 100 keV	3.0 ton-years	2.89 - 50 keV
NaI	11.48 - 100 keV	6.5 ton-years	11.48 - 50 keV
W in CaWO ₄	2.05 - 100 keV	3.9 ton-years	2.05 - 40 keV

Table 1: The second column denotes the optimal energy regions for the WIMP search taking solar neutrinos into account. In the third column the exposures leading to 0.1 atmospheric-neutrino events are given if an optimal energy region according to the last column is selected for the WIMP search.

uncritical since the number of WIMP events drops exponentially with energy. The corresponding exclusion plots (see figure 2) for various target materials




Figure 2: Exclusion plots for different target materials (Exposure 1 ton-year, 90% confidence level).

were obtained for an exposure of 1 ton-year, 90% confidence level assuming zero events in the optimal energy region of table 1 (second column).

For WIMP-nucleon cross sections below 10^{-46} cm^2 (10^{-10} pb) solar neutrinos have to be taken into account. For WIMP masses $> 10 \text{ GeV}$, Xe and W in CaWO_4 are the most promising target materials.

For sensitivities below 10^{-48} cm^2 (10^{-12} pb) also atmospheric neutrinos are becoming an important background source [2]. The third column of table 1 shows the maximal exposure if only 0.1 atmospheric-neutrino events within the optimal energy window given by the last column are allowed. Again the upper limit of this energy window is found to be uncritical. For example, if for Xe the upper threshold is increased from 50 to 100 keV, the maximum exposure is reduced to 2.9 ton-years.

This work was supported by funds of the Deutsche Forschungsgemeinschaft DFG (Transregio 27: Neutrinos and Beyond), the Munich Cluster of Excellence (Origin and Structure of the Universe), and the Maier-Leibnitz-Laboratorium (Garching).

References

- [1] A. Drukier and L. Stodolsky, *Phys. Rev. D* **30** (1984) 2295
- [2] A. Gütlein et al., *Astropart. Phys.* **34** (2010) 90
- [3] R.H. Helm, *Phys. Rev.* **104** (1956) 1466
- [4] J.D. Lewin and P.F. Smith, *Astropart. Phys.* **6** (1996) 87