

Experimental realities refuting existence of $p=0$ condensate
in a system of interacting bosons : I. Electron bubble

Yatendra S. Jain

Department of Physics
North-Eastern Hill University, Shillong - 793022, India

Abstract

Physical reality of the existence of electron bubble in liquid ${}^4\text{He}$ (or ${}^3\text{He}$) renders a *clear experimental evidence* for a quantum particle (in an interacting environment as seen by electron in liquid helium) to occupy exclusively a space of size $\lambda/2$ that, obviously, depends on its energy/momentum. This unequivocally proves that *no particle* in a system of interacting bosons such as liquid ${}^4\text{He}$ has momentum $p = 0$; in stead, *all particles* in the ground state of such a system are in the single quantum state of energy $\varepsilon_o = h^2/8md^2$ or momentum $p = h/2d$.

PACS : 67.25.D, 67.10.-j

Key words : BEC, bosons, electron-bubble, He-4

© by authors

Superfluidity and related properties of the low temperature phase of a *system of interacting bosons* (SIB) such as liquid ${}^4\text{He}$ [1] and trapped dilute gases [2] are, conventionally, attributed to an assumed existence of $p = 0$ condensate, -a macroscopic fraction [$n_{p=0} = N_{p=0}/N$] of its total number of particles, N , occupying zero momentum ($p = 0$) state of a single particle confined to its volume, V ; remaining particles, $N - N_{p=0}$, believed to occupy different states of non-zero momenta, k_1, k_2, k_3, \dots etc., (in wave number unit) form non-condensate component. This follows from the statistical analysis of non-interacting bosons [3] and Bogoliubov's field theoretical study of a system of weakly interacting bosons [4]. Analyzing this distribution (believed to be the origin of superfluidity and related aspects of a SIB for about seven decades now), we recently discovered [5] that it does not correspond to lowest possible energy and, as such, it is unacceptable for the ground state (G-state) of any physical system. Our analysis [5] also reveals that in the state of least possible energy, needed for the G-state, of a SIB all particles have identically equal energy $\varepsilon_o = h^2/8md^2$ (equivalent momentum $q_o = \pi/d$) which represents the G-state energy (momentum) of a particle trapped in a cavity formed by its nearest neighbors and the state is shown to have least possible energy. Consequently not even a single particle is expected to have $p = 0$. In other words, our analysis reveals that the laws of nature, which define the G-state of a physical system to have minimum possible energy, forbid the existence of $p = 0$ condensate and the said non-condensate in the G-state of a SIB. In this paper, we use the physical reality of the existence of an *electron bubble* (EB) [6,7] in liquid helium to place these inferences on unshakable foundation.

An excess electron in liquid helium exclusively occupies a self created spherical cavity (known as electron bubble) of certain radius when it assumes its *lowest possible energy* in the cavity; to create the said cavity, electron exerts its zero-point force on helium atoms in its surroundings and works against the forces originating from inter-atomic interactions and external pressure on the liquid [6,7]. It is evident that the bubble formation is a consequence of the facts that: (i) the electron experiences a strong short range repulsion with helium atoms which forbids its binding with them and (ii) an electron is a quantum particle which behaves as a wave packet whose size, $\lambda/2 = h/p = h/\sqrt{(2mE)}$, increases with the decrease in its energy E (or equivalent momentum p). This implies that any quantum particle that experiences similar repulsion with helium atoms should have similar state in liquid helium and this is found to be true with positron [8] and other particles (ions, atoms, molecules, etc. [9]).

Since ${}^4\text{He}$ atom is also a quantum particle and it also has a short range strong repulsion with other ${}^4\text{He}$ atoms, each ${}^4\text{He}$ atom in its lowest possible energy in liquid ${}^4\text{He}$ should exclusively occupy a spherical cavity with an energy $h^2/8mv^{2/3}$ where v is the volume of the cavity. Assuming that i - atom occupies v_i volume we can express the total sum of the G-state energy of N particles in the liquid as

$$E = \sum_i^N \frac{h^2}{8mv_i^{2/3}} \quad (1)$$

with

$$V = \sum_i^N v_i \quad (2)$$

A simple algebra reveals that E has its minimum value only and only when all v_i are identically equal to $V/N = d^3$. This concludes that the right value of the G-state energy of a SIB like

liquid ${}^4\text{He}$ and trapped dilute gases should be

$$E_o = N \frac{h^2}{8md^2} = N\varepsilon_o \quad (3)$$

which clearly proves that no particle in the G-state of a SIB has zero energy/momentum. Evidently, the laws of nature, responsible for the physical existence of EB, forbid the existence of even a single particle to have $p < h/2d$; the question of a particle having $p = 0$ or a macroscopic fraction of particles in a SIB having $p = 0$ does not arise.

The reasons, for which an electron in liquid ${}^4\text{He}$ or ${}^3\text{He}$ exclusively occupies a large size spherical cavity (*electron bubble*) are, obviously, applicable to each ${}^4\text{He}$ atom in liquid ${}^4\text{He}$ which should occupy exclusively a spherical cavity with certain energy $\varepsilon_o = h^2/8md^2$ or equivalent momentum $h/2d$. Thus the physical reality of the existence of an EB in liquid ${}^4\text{He}$ and ${}^3\text{He}$ unequivocally establishes that no ${}^4\text{He}$ atom in the G-state of liquid ${}^4\text{He}$ has $p = 0$ confirming that the nature forbids $p = 0$ condensate of helium atoms in the ground state of liquid ${}^4\text{He}$. This conclusion, obviously holds true for the atoms in Bose Einstein condensate of trapped dilute gases. In fact no particle of such gases has zero momentum because the wave mechanics of a particle in harmonic trap has to have a least energy of $(3/2)h\nu$ (where ν is the fundamental frequency of oscillations of the particle in the trap). While we find that Eqns.(1-3) are consistent with excluded volume condition [10] which states that each particle such as ${}^4\text{He}$ atom occupies certain volume *exclusively* due to its short range hard core interaction with neighboring particles, it is important to note that Kleban [10], in his paper published long back in 1973, concluded that $p = 0$ condensate contradicts the excluded volume condition. However, his inference was not given due importance for so long for an undue bias in favor $p = 0$ condensate as the origin of superfluidity and related properties of a SIB.

In what follows, we compare the momentum distribution of particles in the G-state of a system of non-interacting bosons [Fig 1(A)], and the momentum distributions of bosons in the G-state of a SIB as found by Bogoliubov [4] with that concluded by our studies [5], respectively, depicted in Fig.1(B) and Fig.1(C), for their better perception. It is clear that 100% particles in a SIB at absolute zero occupy a state of non-zero momentum, $p = h/2d$ as depicted in Fig.1(C). Although, the existence of an electron bubble clearly proving that a quantum particle in its G-state exclusively occupies certain space of the size, $\lambda/2$, of its wave packet particularly when it interacts with its neighbors through a strong repulsive force, it is unfortunate that this reality was never used to determine the G-state of liquid ${}^4\text{He}$, liquid ${}^3\text{He}$ and electron fluid in conductors. Had it been so, our understanding of superfluidity and superconductivity would have been as simple as what is concluded in [11, 12] and the G-state of a SIB [as depicted by Fig.1(C)] could have been concluded long back.

References

- [1] C. Enss, S. Hunklinger, Low Temperature Physics, Springer Verlag, Berlin, 2005.
- [2] F. Dalfovo, S. Giorgini, L.P. Pitaevskii, and S. Stringari, Rev. Mod. Phys. 71, 463-512 (1999), and Arxiv/Cond-mat/9806038.

- [3] R.K. Pathria, Statistical Mechanics, Pergamon Press Oxford (1976).
- [4] N.N. Bogoliubov, J. Phys, USSR, 11, 23-32 (1947).
- [5] Y.S. Jain, *The $p=0$ condensate is a myth*, arxiv:cond-mat/1008.240v2
- [6] M. Rosenblit, J. Jortner, Phys. Rev. Lett. 75, 4079-4082 (1995); M. Farnik, U. Henne, B. Samelin, and J.P. Toennies, Phys. Rev. Lett. 81, 3892-3895 (1998)
- [7] H. Marris, and S. Balibar, Phys. To-day 53, 29-34 (2000)
- [8] P. Hautojarvi, M.T. Loponen, and K. Rytola, J. Phys. Atom. Mole. Phys. 9, 411-422 (1976)
- [9] J.P. Toennies, and A.F. Vilesov, Ann. Rev. Phys. Chem. 49, 1-41 (1998)
- [10] P. Kleban, Phys. Lett. A 49, 19-20 (1973).
- [11] (a) Y. S. Jain, Ind. J. Phys. 79, 1009-14 (2005); the factor $|\sin(\mathbf{k}\cdot\mathbf{r}/2)|$ in Eqn.5 for Ψ^+ in this paper should be read as $\sin|(\mathbf{k}\cdot\mathbf{r}/2)|$ and $E_g(T)/Nk_B T$ in Eqn.25 should be read as $E_g(T)/Nk_B$
 (b) Y.S. Jain, Macro-orbitals and microscopic theory of a system of interacting bosons, arXiv:cond-mat/0606571
- [12] Y.S. Jain, Basic foundations of the microscopic theory of super conductivity, arXiv:cond-mat/0603784, (2006).






Fig.1 : Schematic of distribution of N bosons in their ground state. (A) All the N particles occupy $p = 0$ state in a system of non-interacting bosons, (B) depletion of $p = 0$ condensate (*i.e.* only a fraction $n_{p=0} = N_{p=0}/N$ of N bosons occupy $p = 0$ state) in weakly interacting boson system as predicted by Bogoliubov model [4], and (C) all the N particles occupy a state of $p = p_o = \hbar q_o = h/2d$ and $\hbar K = 0$ as concluded by this study and our recent analysis [5].