Comment on: "Hawking Radiation from Ultrashort Laser Pulse Filaments"

In a recent paper [1] Belgiorno *et al* claimed to have observed the analog of the Hawking effect created by light pulses in silica glass because of the detection of radiation in a frequency range in which what they called "phase horizons" existed. Unfortunately, while the observations are very interesting, the cause of the radiation is not understood, and we feel it is not justified to call this a detection of the Hawking effect in an analog system.

The Hawking effect is the observation of thermal radiation (with the temperature being determined by the geometry) from a time independent system, in which the radiation is caused by the (quasi) exponential tearing apart of the waves in the vicinity of the horizon. (This applies to black hole horizons, for white holes it would be the time-reversed process, i.e., squeezing of waves.) Since the part of the (torn apart) wave-packet beyond the horizon can have negative energy, the other part may acquire positive energy and constitutes the Hawking radiation. Out of these five conditions, only the last one (i.e., the negative energy beyond the "phase horizon" which is related to the Landau criterion, see below) applies to the above experiment. Thus, it is an important step towards the observation of Hawking radiation, but not more.

Even in the frame of the pulse, the system is not time independent – the pulse itself lasts only a brief time (shorter than the time scale set by the surface gravity, i.e., Hawking temperature, see below) and has spacetime dependent sub-structure (due to the difference between the phase velocity of the radiation creating the pulse and the speed of the pulse itself). In the frame of the pulse, the photons co-moving with the pulse satisfying Eq. (1) in [1] approach zero frequency due to $\omega_{\rm frame}^{\rm pulse} \approx \omega_{\rm frame}^{\rm lab} - \boldsymbol{v}_{\rm pulse} \cdot \boldsymbol{k}$, making the creation of particles by that time dependence easy from an energetic point of view. (Momentum conservation is another matter.) In fact, Eq. (1) in [1] is closely related to the Landau criterion for particle creation $\omega_{\rm frame}^{\rm pulse} = 0$.

Second, there is no exponential tearing (or compaction) by the "horizon". Since the group velocity of the photons under consideration is always smaller than the speed of the pulse [1], it just passes through them (i.e., there is no "group velocity horizon"). As an instructive example, consider the dispersion relation of a massive particle $\omega^2 = c^2 k^2 + m^2 c^4 / \hbar^2$. As in the experiment [1], the phase velocity $\omega/k > c$ is larger than the group velocity $d\omega/dk < c$ and a perturbation with v > cwould have phase horizons but no group horizons. In a suitable Lorentz frame, this corresponds to an instantaneous, time-dependent perturbation with no horizon.

Third, the condition $\omega_{\text{frame}}^{\text{lab}} - \boldsymbol{v}_{\text{pulse}} \cdot \boldsymbol{k} = 0$ applies only to waves (in the frequency range of interest) which are moving in the same direction as the pulse. Thus Hawking radiation would also occur in this direction only. How-

ever, the authors of [1] observed photons at 90 degrees to the propagation direction of the pulse. The unpolarized nature of the observed radiation seems to rule out scattering of co-moving radiation as a source.

As a fourth an final point, we note that the interpretation of this emission as Hawking radiation yields Hawking temperatures which are far too low to explain the observations. Even if the spectrum was deformed by dispersion and no longer Planckian, the following estimates for the energy and number of emitted photons would still yield the correct orders of magnitude. In the frame of the pulse, the Hawking temperature would be given by the gradient of the speed of light cin the medium $T_{\text{Hawking}} = \hbar |\partial c / \partial x| / (2\pi k_{\text{B}})$ which yields $\hbar |\partial \delta n / \partial x| / (2\pi k_{\rm B} n^2)$, where n = 1/c is the refractive index and δn its change due to the Kerr effect. Since the non-linearity δn is quite small $\delta n \approx 10^{-3}$, this temperature could only create the observed photons at around 800 nm if one assumed that δn changes on a sub-nm length scale, which seems unrealistic given that the wavelength of the photons generating the pulse is around 1000 nm. Note that the transformation to the lab frame increases the frequency (though not the number) of the photons emitted in forward direction, but this Doppler shift does not apply to any photons emitted to the side.

In addition, the Hawking effect would correspond to thermal radiation where the number of particles created goes as $\sigma AT^3_{\text{Hawking}}(\Delta \vartheta)^2$. Here $A \sim L^2$ is the area of the horizon and $\Delta \vartheta$ is the solid angle into which the radiation is created. Both are very small: due to the "phase horizon" condition, one has $\Delta \vartheta = \mathcal{O}(10^{-2} \text{ rad})$ and the core size L of a Bessel pulse with 7° is a few μ m. Together with $T_{\text{Hawking}} \propto \delta n \approx 10^{-3}$, one obtains estimates for the number of created particles which are several orders of magnitude too small to explain the data. An even simpler perturbation theory estimate for the maximum number of photons emitted per unit time scales with $(\delta n)^2$, showing serious difficulties with the interpretation as a stationary quantum process. Nevertheless, we admire the experimental technique of [1] and we think that such a set-up may well provide the first observation of spontaneous Hawking emission in an analogue system.

Ralf Schützhold^{1,*} and William G. Unruh^{2,3,+}

¹Fakultät für Physik, Universität Duisburg-Essen,

D-47048 Duisburg, Germany

²Department of Physics and Astronomy, University of British Columbia, Vancouver B.C., V6T 1Z1 Canada ³Institute for Theoretical Physics, Utrecht University, NL-3584 CE Utrecht, The Netherlands

*ralf.schuetzhold@uni-due.de,

⁺unruh@phas.ubc.ca

^[1] F. Belgiorno et al, Phys. Rev. Lett. 105, 203901 (2010).