The integrability of the 2-Toda lattice on a simple Lie algebra

Khaoula Ben Abdeljelil

Abstract

We define the 2-Toda lattice on every simple Lie algebra \mathfrak{g} , and we show its Liouville integrability. We show that this lattice is given by a pair of Hamiltonian vector fields, associated with a Poisson bracket which results from an \mathcal{R} -matrix of the underlying Lie algebra. We construct a big family of constants of motion which we use to prove the Liouville integrability of the system. We achieve the proof of their integrability by using several results on simple Lie algebras, \mathcal{R} -matrices, invariant functions and root systems.

Contents

1	Intr	roduction	2
2	\mathcal{R} -n	natrices, Poisson structures and functions in involution on the square	
	of a	a Lie algebra	4
	2.1	Construction of \mathcal{R} -matrices on \mathfrak{g}^2	5
	2.2	Linear Poisson structures on $\mathfrak{g}^* \times \mathfrak{g}^*$	7
	2.3	Casimirs and functions in involution on $\mathfrak{g}^* \times \mathfrak{g}^*$	7
3	The	e Liouville integrability of the 2-Toda lattice	12
	3.1	Definition of the 2-Toda lattice	12
	3.2	The 2-Toda lattice is a Hamiltonian system	12
	3.3	The integrability of the 2-Toda lattice	15
		3.3.1 The restriction of \mathcal{F} to \mathcal{T}^2 is involutive for $\{\cdot,\cdot\}_{\mathcal{R}}$	16
		3.3.2 The restriction of \mathcal{F} to \mathcal{T}^2 is independent	17
		3.3.3 The exact number of functions	20
	3.4	The integrability for the quadratic Poisson \mathcal{R} -bracket	22
		3.4.1 The Liouville integrability of the 2-Toda lattice on $\mathfrak{gl}_n(\mathbf{C})$ for the	
		linear Poisson \mathcal{R} -bracket	22
		3.4.2 The integrability of the 2-Toda lattice on $\mathfrak{gl}_n(\mathbf{C})$ for the quadratic	
		Poisson \mathcal{R} -bracket	23

3.5	The re	elation between the 2-Toda lattice and the Toda lattice	27
	3.5.1	The Toda lattice	27
	3.5.2	Restriction of the 2-Toda lattice and construction of the Toda lattice	28

1 Introduction

The 2-Toda lattice associated with $\mathfrak{sl}_n(\mathbf{C})$ is the pair of differential equations given by the Lax equations

$$\begin{cases}
\frac{\partial(L,M)}{\partial t} = [(L_u,L_u),(L,M)], \\
\frac{\partial(L,M)}{\partial s} = [(M_\ell,M_\ell),(L,M)],
\end{cases} (1)$$

where (L, M) are traceless matrices of the form

$$(L,M) = \left(\begin{pmatrix} a_{11} & 1 & & & 0 \\ a_{21} & a_{22} & \ddots & & \\ \vdots & & \ddots & 1 \\ a_{n1} & \cdots & a_{n,n-1} & a_{nn} \end{pmatrix}, \begin{pmatrix} b_{11} & \cdots & \cdots & b_{1n} \\ b_{21} & \ddots & & \vdots \\ & \ddots & \ddots & \vdots \\ 0 & & b_{n,n-1} & b_{nn} \end{pmatrix} \right), \tag{2}$$

and where L_u is the upper triangular part of L and M_ℓ is the strictly lower triangular part of M.

The 2-Toda lattice was first introduced in the context of infinite-dimensional matrices [1], [3], [13]. This system is Hamiltonian with respect to a Poisson structure, associated with an \mathcal{R} -matrix with $\frac{1}{2}$ Trace L^2 and $\frac{1}{2}$ Trace M^2 as Hamiltonians. These Hamiltonians admit the Ad-invariant functions on $\mathfrak{gl}((\infty)) \times \mathfrak{gl}((\infty))$ as Poisson commuting constants of motion. In the finite-dimensional setting, i.e., on $\mathfrak{sl}_n(\mathbf{C})$, this system is still Hamiltonian with respect to a Poisson structure associated with an \mathcal{R} -matrix similar to that of the infinite-dimensional setting. The family of Ad-invariant functions are again in involution with respect to the Poisson structure associated with the \mathcal{R} -matrix but their number of independent Ad-invariant functions, which is 2n-2, is much too small compared with the dimension of the phase space of the 2-Toda lattice, which is $n^2 + 2n - 3$.

The main purpose of the present article is to introduce a family of functions which contains the Ad-invariant functions and is large enough to give the Liouville integrability of the 2-Toda lattice. This family is $\mathcal{F} := (F_{j,i}, 1 \leq i \leq n-1 \text{ and } 0 \leq j \leq i+1)$, where $F_{j,i}$ is the function on $\mathfrak{sl}_n(\mathbf{C})$ constructed from the the relation

Trace
$$(\lambda L - M)^{i+1} = \sum_{j=0}^{i+1} (-1)^{m_i+1-j} \lambda^j F_{j,i}(L, M).$$

The Toda lattice can be defined for every simple Lie algebra [7]. In the same spirit we introduce the 2-Toda system for every simple Lie algebra and we show its Liouville integrability. We recall that, according to [2, Definition 4.13], in the general case a system $(M, \{\cdot, \cdot\}, \mathcal{F})$ is Liouville integrable if $(M, \{\cdot, \cdot\})$ is a Poisson manifold of rank 2r and $\mathcal{F} = (F_1, \ldots, F_s)$ is involutive and independent, with $s = \dim M - r$.

The main result of this paper is given in Section 3: we show the Liouville integrability of the 2-Toda lattice, not only on $\mathfrak{sl}_n(\mathbf{C})$, but also on an arbitrary simple Lie algebra \mathfrak{g} . We define this lattice in this general context. We choose a simple Lie algebra \mathfrak{g} ; then we denote by \mathfrak{h} a Cartan subalgebra, by $\alpha_1, \ldots, \alpha_\ell$ the simple roots of \mathfrak{g} with respect to \mathfrak{h} , by e_1, \ldots, e_ℓ the corresponding eigenvectors, and by \mathfrak{g}_i the subspace spanned by the eigenvectors associated with roots of length $i \in \mathbb{Z}$. The 2-Toda lattice, associated with \mathfrak{g} , is the pair of differential equations on \mathcal{T}^2 given by the following Lax pair equations

$$\begin{cases}
\frac{\partial(L,M)}{\partial t} = [(L_+,L_+),(L,M)], \\
\frac{\partial(L,M)}{\partial s} = [(M_-,M_-),(L,M)],
\end{cases}$$
(3)

where (L, M) belongs to the phase space $\mathcal{T}^2 := \sum_{i \leq 0} \mathfrak{g}_i \times \sum_{i \geq -1} \mathfrak{g}_i + (\sum_{i=1}^{\ell} e_i, 0)$, and here, L_+ (resp. M_-) stands for the projection of $L \in \mathfrak{g}$ (resp. $M \in \mathfrak{g}$) on $\sum_{i \geq 0} \mathfrak{g}_i$ (resp. $\sum_{i \leq 0} \mathfrak{g}_i$).

We construct a linear Poisson structure with respect to which the 2-Toda lattice is Hamiltonian. This linear Poisson structure comes from an \mathcal{R} -matrix of the type studied in Section 2, namely from the endomorphism of \mathfrak{g}^2 given by $\mathcal{R}(x,y) = (R(x-y)+y,R(x-y)+x)$, for all $x,y\in\mathfrak{g}$, where $R:=P_+-P_-$ is the classical R-matrix on a simple Lie algebra, associated with the Lie algebra splitting $\mathfrak{g}=\sum_{i\geqslant 0}\mathfrak{g}_i\oplus\sum_{i<0}\mathfrak{g}_i$. Using the Killing form of \mathfrak{g} , the product Lie algebra \mathfrak{g}^2 is equipped with a bilinear, symmetric, Ad-invariant, non-degenerate form, which allows us to identify \mathfrak{g}^2 with its dual and hence to equip \mathfrak{g}^2 with a linear Poisson structure that we denote by $\{\cdot,\cdot\}_{\mathcal{R}}$. We then establish that the phase space \mathcal{T}^2 of the 2-Toda lattice is a Poisson submanifold of $(\mathfrak{g}^2,\{\cdot,\cdot\}_{\mathcal{R}})$ (see Propositions 8) and the Hamiltonian vector fields of functions $H(x,y):=\frac{1}{2}\langle x|x\rangle$ and $\tilde{H}(x,y):=\frac{1}{2}\langle y|y\rangle$ are tangent to \mathcal{T}^2 and describe on \mathcal{T}^2 the equations of motion (3) for the 2-Toda lattice (see Proposition 10).

We construct the integrable system. Let (P_1, \ldots, P_ℓ) be a generating family of the algebra of Ad-invariant functions, chosen to be of respective degrees $m_1 + 1, \ldots, m_\ell + 1$. For all $1 \leq i \leq \ell$ and $0 \leq j \leq m_i + 1$, we define $F_{j,i} \in \mathcal{F}(\mathfrak{g}^2)$ to be the coefficient in λ^j of the polynomial $(x, y) \mapsto P_i(\lambda x - y)$, and we consider the following family of functions on \mathfrak{g}^2 :

$$\mathcal{F} = (F_{j,i}, 1 \leqslant i \leqslant \ell \text{ and } 0 \leqslant j \leqslant m_i + 1).$$

Our main result is the following theorem:

Theorem The triplet $(\mathcal{T}^2, \mathcal{F}_{|\mathcal{T}^2}, \{\cdot, \cdot\}_{\mathcal{R}})$ is an integrable system.

To prove this result, we proceed as follows:

- We show in Proposition 14 that \mathcal{F} is involutive for $\{\cdot,\cdot\}_{\mathcal{R}}$: this result is a particular case of the second point of Theorem 4 that we prove in Section 2.
- We show in Proposition 17 the independence of \mathcal{F} restricted to \mathcal{T}^2 by using a theorem of Raïs [10].
- We compute the rank of the restriction of the Poisson \mathcal{R} -bracket to \mathcal{T}^2 . This is done by establishing a Poisson isomorphism between $(\mathcal{T}^2, \{\cdot, \cdot\}_{\mathcal{R}})$ and the product Poisson manifold $(\mathfrak{g}^*, \{\cdot, \cdot\}) \times (\mathfrak{g}_0 \oplus \mathfrak{g}_1, \{\cdot, \cdot\}_R)$, where $\{\cdot, \cdot\}$ is the Lie-Poisson bracket on \mathfrak{g}^* , and $\{\cdot, \cdot\}_R$ is the above R-bracket (restricted to $\mathfrak{g}_0 \oplus \mathfrak{g}_1$; see Proposition 18).
- We check that $\operatorname{card} \mathcal{F}_{|\mathcal{T}^2} = \dim \mathcal{T}^2 \frac{1}{2}\operatorname{Rk}(\mathcal{T}^2, \{\cdot\,,\cdot\}_{\mathcal{R}}).$

We also define the 2-Toda from on $\mathfrak{gl}_n(\mathbf{C})$ and we show its Liouville integrability with respect to the linear and quadratic Poisson \mathcal{R} -bracket. This result is mainly based on the fact that the Hamiltonian vector fields of the linear and quadratic Poisson structure are essentially identical.

In the last subsection, by restricting the 2-Toda lattice on \mathfrak{g} to a well chosen affine subspace, we find again the usual Toda lattice.

2 R-matrices, Poisson structures and functions in involution on the square of a Lie algebra

In this section we fix $(\mathfrak{g}, [\cdot, \cdot])$ a Lie algebra over \mathbf{F} , with $\mathbf{F} = \mathbf{R}$ or $\mathbf{F} = \mathbf{C}$. The vector space $\mathfrak{g}^2 := \mathfrak{g} \times \mathfrak{g}$, endowed with the Lie bracket

$$[(x,y),(z,s)] := ([x,z],[y,s]), \qquad \forall x,y,z,s \in \mathfrak{g},$$
 (4)

is a Lie algebra.

We construct an \mathcal{R} -matrix of \mathfrak{g}^2 with the help of an endomorphism R of \mathfrak{g} satisfying some conditions. With this \mathcal{R} -matrix, when \mathfrak{g} is finite dimensional, we construct a linear Poisson bracket on the dual \mathfrak{g}^{2^*} of \mathfrak{g}^2 , we explain the construction of a large family of functions on \mathfrak{g}^{2^*} which commute for this linear Poisson structure and we spell out the expressions of their Hamiltonian vector fields. We also give some Casimir functions.

2.1 Construction of R-matrices on \mathfrak{g}^2

We begin by recalling some properties and definitions of R-matrices (see [2, Section 4.4]). Let \mathfrak{g} be a Lie algebra. A (vector space) endomorphism R of \mathfrak{g} is called an R-matrix of \mathfrak{g} if the bilinear map $[\cdot,\cdot]_R:\mathfrak{g}\times\mathfrak{g}\to\mathfrak{g}$, defined for all $x,y\in\mathfrak{g}$ by

$$[x,y]_R := \frac{1}{2}([Rx,y] + [x,Ry]), \tag{5}$$

defines a (second) Lie bracket on \mathfrak{g} , which is then called a Lie *R*-bracket. Let $B_R : \mathfrak{g} \times \mathfrak{g} \to \mathfrak{g}$ be defined, for all $x, y \in \mathfrak{g}$, by

$$B_R(x,y) := [Rx, Ry] - R([Rx, y] + [x, Ry]). \tag{6}$$

The bracket $[\cdot,\cdot]_R$ satisfies the Jacobi identity if and only if

$$[B_R(x,y),z] + [B_R(y,z),x] + [B_R(z,x),y] = 0, \quad \forall x,y,z \in \mathfrak{g}.$$
 (7)

A sufficient condition for (7) to be satisfied is that there exists a constant $c \in \mathbf{F}$ such that

$$B_R(x,y) = -c^2[x,y], \qquad \forall x,y \in \mathfrak{g}.$$
 (8)

We call (8) the modified classical Yang-Baxter equation (mCYBE) of \mathfrak{g} of constant c.

We construct in the following proposition an \mathcal{R} -matrix of \mathfrak{g}^2 by using an endomorphism of \mathfrak{g} which satisfies some condition which generalizes the modified classical Yang-Baxter equation of \mathfrak{g}^2 .

Proposition 1 Let \mathfrak{g} be a Lie algebra, $\mathfrak{z}(\mathfrak{g})$ its center and $c \in \mathbf{F}$ a constant. Let R be an endomorphism of \mathfrak{g} and \mathcal{R} the endomorphism of \mathfrak{g}^2 defined, for every $(x,y) \in \mathfrak{g}^2$, by

$$\mathcal{R}(x,y) := (R(x-y) + cy, R(x-y) + cx). \tag{9}$$

The endomorphism \mathcal{R} is an \mathcal{R} -matrix of \mathfrak{g}^2 if and only if, for every $x, y \in \mathfrak{g}$,

$$B_R(x,y) + c^2[x,y] \in \mathfrak{z}(\mathfrak{g}).$$
 (10)

In the particular case where \mathfrak{g} is a complex semi-simple Lie algebra, this is in turn equivalent to R being a solution of (mCYBE) of constant c.

Proof. Using Definition (6) of $B_{\mathcal{R}}$ and replacing \mathcal{R} by its Expression (9), we obtain

$$B_{\mathcal{R}}((x,y),(z,s)) = (B_{\mathcal{R}}(x-y,z-s) + c^2([x-y,z-s] - [x,z]), B_{\mathcal{R}}(x-y,z-s) + c^2([x-y,z-s] - [y,s]).$$
(11)

According to Formula (7), \mathcal{R} is an \mathcal{R} -matrix of \mathfrak{g}^2 if and only if, for every (x, y), (x', y') and (x'', y'') in \mathfrak{g}^2 ,

$$[B_{\mathcal{R}}((x,y),(x',y')),(x'',y'')] + \circlearrowleft = (0,0), \tag{12}$$

where $\circlearrowleft = \operatorname{cycl}((x, y), (x', y'), (x'', y''))$. By Formula (11), the left hand side of Equation (12) is equal to

$$([B_R(x-y,x'-y')+c^2[x-y,x'-y'],x''],[B_R(x-y,x'-y')+c^2[x-y,x'-y'],y''])+\circlearrowleft,$$
(13)

where we used the Jacobi identities $[[x, x'], x''] + \circlearrowleft = 0$ and $[[y, y'], y''] + \circlearrowleft = 0$. Now, if \mathcal{R} is an \mathcal{R} -matrix of \mathfrak{g}^2 , then (13) is equal to zero (even without circular permutation), so Formula (10) holds. If Formula (10) holds then the first component of the left hand side of (13) is equal to zero:

$$[B_R(x-y,x'-y')+c^2[x-y,x'-y'],x''] + \operatorname{cycl}((x,y),(x',y'),(x'',y'')) = 0, \tag{14}$$

for every $x, y, x', y', x'', y'' \in \mathfrak{g}$; in particular, for x' = x'' = 0, we obtain

$$[B_R(y',y'') + c^2[y',y''], x] = 0, \quad \forall y', y'', x \in \mathfrak{g},$$

hence the endomorphism \mathcal{R} is an \mathcal{R} -matrix of \mathfrak{g}^2 holds.

For future use, we also state the following corollary, which is a consequence of Formula (11) and already appeared in [11] (with c = 1).

Corollary 2 Let \mathfrak{g} be a Lie algebra and let R be an endomorphism of \mathfrak{g} . If R is a solution of (mCYBE) of \mathfrak{g} of constant c, then the endomorphism \mathcal{R} of \mathfrak{g}^2 , defined, for all $(x,y) \in \mathfrak{g}^2$, by

$$\mathcal{R}(x,y) := (R(x-y) + cy, R(x-y) + cx),$$

is a solution of (mCYBE) of \mathfrak{g}^2 of constant c.

Example 3 Let $\mathfrak{g} := \mathfrak{g}_+ \oplus \mathfrak{g}_-$ be a Lie algebra splitting (i.e., \mathfrak{g} is, as a vector space, the direct sum of Lie subalgebras \mathfrak{g}_+ and \mathfrak{g}_-). Let P_\pm be the projections of \mathfrak{g} on \mathfrak{g}_\pm and let $R := P_+ - P_-$. Since R is a solution of (mCYBE) of \mathfrak{g} of c = 1, Corollary 2 implies that the endomorphism \mathcal{R} , defined for every $x, y \in \mathfrak{g}$ by $\mathcal{R}(x, y) = (R(x - y) + y, R(x - y) + x)$ is a solution of (mCYBE) of \mathfrak{g}^2 of c = 1. Replacing R by its expression $P_+ - P_-$, we verify that

$$\mathcal{R}(x,y) = (x_{+} - x_{-} + 2y_{-}, y_{-} - y_{+} + 2x_{+})$$

$$= (x_{+} + y_{-}, x_{+} + y_{-}) - (x_{-} - y_{-}, y_{+} - x_{+}), \tag{15}$$

where $x = x_+ + x_-$ and $y = y_+ + y_-$ are the decompositions of x and y with respect to the splitting $\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$. Since the Lie subalgebras

$$\mathfrak{g}_+^2:=\{(x,x) \ \mid \ x\in\mathfrak{g}\} \quad and \quad \mathfrak{g}_-^2:=\{(x,y) \ \mid \ x\in\mathfrak{g}_- \ and \ y\in\mathfrak{g}_+\}$$

have a trivial intersection, (15) implies that $\mathfrak{g}^2 = \mathfrak{g}_+^2 \oplus \mathfrak{g}_-^2$ is a Lie algebra splitting and that \mathcal{R} is the difference of the projections on \mathfrak{g}_+^2 and \mathfrak{g}_-^2 .

Note that independently of Proposition 1 or Corollary 2 the above example shows that from an R-matrix of \mathfrak{g} , which is the difference of the projections on two supplementary Lie subalgebras of \mathfrak{g} , the endomorphism \mathcal{R} given by the Formula (9) is automatically an \mathcal{R} -matrix of \mathfrak{g}^2 , which is the difference of projections on two supplementary Lie subalgebras of \mathfrak{g}^2 .

2.2 Linear Poisson structures on $\mathfrak{g}^* \times \mathfrak{g}^*$

From now and until the end of the article, we assume that \mathfrak{g} is a finite-dimensional Lie algebra and we denote by $\mathcal{F}(\mathfrak{g})$ the algebra of smooth functions on \mathfrak{g} (if \mathfrak{g} is a Lie algebra over \mathbf{R}) or holomorphic functions on \mathfrak{g} (if \mathfrak{g} is a Lie algebra over \mathbf{C}). Let \mathcal{R} be an \mathcal{R} -matrix of \mathfrak{g}^2 . The dual $(\mathfrak{g}^2)^*$ of the Lie algebra $(\mathfrak{g}^2, [\cdot, \cdot]_{\mathcal{R}})$ admits a Lie-Poisson structure, defined for every $F, G \in \mathcal{F}((\mathfrak{g}^2)^*)$ at $\varphi \in (\mathfrak{g}^2)^*$ by

$$\begin{aligned}
\{F,G\}_{\mathcal{R}}(\varphi) : &= \langle \varphi, [\mathsf{d}_{\varphi}F, \mathsf{d}_{\varphi}G]_{\mathcal{R}} \rangle \\
&= \frac{1}{2} \langle \varphi, [\mathcal{R}\mathsf{d}_{\varphi}F, \mathsf{d}_{\varphi}G] + [\mathsf{d}_{\varphi}F, \mathcal{R}\mathsf{d}_{\varphi}G] \rangle,
\end{aligned} (16)$$

where $d_{\varphi}F \in ((\mathfrak{g}^2)^*)^* \simeq \mathfrak{g}^2$ is the differential of F at the point $\varphi \in (\mathfrak{g}^2)^*$. We identify $(\mathfrak{g}^2)^*$ with $\mathfrak{g}^* \times \mathfrak{g}^*$ as follows:

$$\Phi : \mathfrak{g}^* \times \mathfrak{g}^* \to (\mathfrak{g}^2)^*$$
$$(\xi, \eta) \mapsto ((x, y) \mapsto \langle (\xi, \eta), (x, y) \rangle) := \langle \xi, x \rangle - \langle \eta, y \rangle.$$

By means of this identification, the Poisson structure (16) can be transported to the vector space $\mathfrak{g}^* \times \mathfrak{g}^*$. This transported Poisson structure is given, for all functions $F, G \in \mathcal{F}(\mathfrak{g}^* \times \mathfrak{g}^*)$ at $(\xi, \eta) \in \mathfrak{g}^* \times \mathfrak{g}^*$, by

$$\{F,G\}_{\mathcal{R}}(\xi,\eta) := \left\langle (\xi,\eta), \left[\mathsf{d}_{(\xi,\eta)}F, \mathsf{d}_{(\xi,\eta)}G\right]_{\mathcal{R}} \right\rangle,\tag{17}$$

where $d_{(\xi,\eta)}F \in (\mathfrak{g}^* \times \mathfrak{g}^*)^* \simeq \mathfrak{g} \times \mathfrak{g}$. We call this Poisson structure the Poisson \mathcal{R} -bracket on $\mathfrak{g}^* \times \mathfrak{g}^*$.

2.3 Casimirs and functions in involution on $\mathfrak{g}^* \times \mathfrak{g}^*$

Let \mathfrak{g} be a finite-dimensional Lie algebra. We denote by \mathbf{G} a connected Lie group whose Lie algebra is \mathfrak{g} . We denote by $\mathcal{F}(\mathfrak{g}^*)^{\mathbf{G}}$ the algebra of Ad*-invariant functions on \mathfrak{g}^* .

According to the Adler-Kostant-Symes theorem [2, Theorem 4.37], the Ad*-invariant functions on $\mathfrak{g}^* \times \mathfrak{g}^*$ (i.e., functions in $\mathcal{F}(\mathfrak{g}^* \times \mathfrak{g}^*)^{\mathbf{G} \times \mathbf{G}}$) are in involution for the Poisson structure associated with an \mathcal{R} -matrix. But, when \mathfrak{g} is a semi-simple Lie algebra, this family of functions is too small to insure the Liouville integrability. We construct a larger family of functions in involution.

¹Which contains the Ad*-invariant functions on $\mathfrak{g}^* \times \mathfrak{g}^*$.

Theorem 4 Let $\lambda \in \mathbf{F}$ be a constant and let ψ_{λ} be the map

$$\psi_{\lambda} : \mathfrak{g}^* \times \mathfrak{g}^* \to \mathfrak{g}^* (\xi, \eta) \mapsto \lambda \xi - \eta.$$
 (18)

Let R be an endomorphism² of \mathfrak{g} and $c \in \mathbf{F}$ a constant. Assume that the endomorphism \mathcal{R} of \mathfrak{g}^2 defined for every $(x,y) \in \mathfrak{g}^2$ by

$$\mathcal{R}(x,y) := (R(x-y) + cy, R(x-y) + cx) \tag{19}$$

is an \mathcal{R} -matrix of \mathfrak{g}^2 , and denote by $\{\cdot,\cdot\}_{\mathcal{R}}$ the Poisson \mathcal{R} -bracket on $\mathfrak{g}^* \times \mathfrak{g}^*$. Then:

- (1) For every $F \in \mathcal{F}(\mathfrak{g}^*)^{\mathbf{G}}$, the function $F \circ \psi_1$ is a Casimir for $\{\cdot,\cdot\}_{\mathcal{R}}$.
- (2) For every $F, G \in \mathcal{F}(\mathfrak{g}^*)^{\mathbf{G}}$ and every $\lambda, \gamma \in \mathbf{F}$, the functions $F \circ \psi_{\lambda}$ and $G \circ \psi_{\gamma}$ are in involution for $\{\cdot, \cdot\}_{\mathcal{R}}$. In particular, if F and G are polynomials of degree respectively l and k, then the functions $F_0, \ldots, F_l, G_0, \ldots, G_k$, defined, for every $(\xi, \eta) \in \mathfrak{g}^* \times \mathfrak{g}^*$, by

$$F(\psi_{\lambda}(\xi,\eta)) = \sum_{i=0}^{l} \lambda^{i} F_{i}(\xi,\eta)$$
 and $G(\psi_{\gamma}(\xi,\eta)) = \sum_{j=0}^{k} \gamma^{j} G_{j}(\xi,\eta),$

are in involution for $\{\cdot,\cdot\}_{\mathcal{R}}$.

- (3) If c = 1, the map $\psi_1 : (\mathfrak{g}^* \times \mathfrak{g}^*, \{\cdot, \cdot\}_{\mathcal{R}}) \to (\mathfrak{g}^*, \{\cdot, \cdot\})$ is a Poisson morphism.
- (4) For every $H \in \mathcal{F}(\mathfrak{g}^*)^{\mathbf{G}}$, the Hamiltonian vector field $\mathcal{X}_{H \circ \psi_{\lambda}} := \{\cdot, H \circ \psi_{\lambda}\}_{\mathcal{R}}$ is given at $(\xi, \mu) \in \mathfrak{g}^* \times \mathfrak{g}^*$ by

$$\mathcal{X}_{H \circ \psi_{\lambda}}(\xi, \eta) = \frac{1}{2} (1 - \lambda) \operatorname{ad}_{((R - cI)\mathsf{d}_{\lambda\xi - \eta}H, (R + cI)\mathsf{d}_{\lambda\xi - \eta}H)}^{*}(\xi, \eta). \tag{20}$$

In particular, if $\mathfrak{g} := \mathfrak{g}_+ \oplus \mathfrak{g}_-$ is a Lie algebra splitting and $R := P_+ - P_-$ is the difference of the projections on \mathfrak{g}_+ and \mathfrak{g}_- , we have

$$\mathcal{X}_{H \circ \phi_{\lambda}}(\xi, \eta) = (1 - \lambda) \operatorname{ad}^{*}_{((-\mathsf{d}_{\lambda \xi - \eta} H)_{-}, (\mathsf{d}_{\lambda \xi - \eta} H)_{+})}(\xi, \eta). \tag{21}$$

To prove this theorem, we need the following lemmas, the proof of the first of which is left to the reader.

Lemma 5 Let $F \in \mathcal{F}(\mathfrak{g}^*)$, $(\xi, \eta) \in \mathfrak{g}^* \times \mathfrak{g}^*$ and $\lambda \in \mathbf{F}$ be a constant.

(1) The differential of $F \circ \psi_{\lambda}$ at $(\xi, \eta) \in \mathfrak{g}^* \times \mathfrak{g}^*$ is given by

$$\mathsf{d}_{(\xi,\eta)}(F \circ \psi_{\lambda}) = (\lambda \mathsf{d}_{\lambda\xi-\eta} F, \mathsf{d}_{\lambda\xi-\eta} F). \tag{22}$$

 $^{^{2}}R$ is not necessarily an R-matrix.

(2) Let R be an endomorphism of \mathfrak{g} and let \mathcal{R} be the endomorphism of \mathfrak{g}^2 defined in (19). Then

$$\mathcal{R} \,\mathsf{d}_{(\xi,\eta)}(F \circ \psi_{\lambda}) = (\lambda - 1)(R \,\mathsf{d}_{\lambda\xi - \eta}F, R \,\mathsf{d}_{\lambda\xi - \eta}F) + c(\mathsf{d}_{\lambda\xi - \eta}F, \lambda \mathsf{d}_{\lambda\xi - \eta}F). \tag{23}$$

Lemma 6 Let $F, G \in \mathcal{F}(\mathfrak{g}^*)^{\mathbf{G}}$.

(1) The differentials of F and G commute at every point of \mathfrak{g}^* , i.e.,

$$[\mathsf{d}_{\xi}F, \mathsf{d}_{\xi}G] = 0, \qquad \forall \xi \in \mathfrak{g}^*. \tag{24}$$

(2) For every $\xi, \eta \in \mathfrak{g}^*$ and every $a, b \in \mathbf{F}$,

$$\langle a\xi + b\eta, [\mathsf{d}_{\xi}F, \mathsf{d}_{\eta}G] \rangle = 0. \tag{25}$$

Proof. Using the Ad^* -invariance of F and G, we obtain

$$\operatorname{ad}_{\mathsf{d}_{\varepsilon}F}^* \xi = 0 \qquad \text{and} \qquad \operatorname{ad}_{\mathsf{d}_{\varepsilon}G}^* \xi = 0. \tag{26}$$

Stated differently, $d_{\xi}F \in \mathfrak{g}^{\xi}$ and $d_{\xi}G \in \mathfrak{g}^{\xi}$ where \mathfrak{g}^{ξ} is the centralizer of ξ . For ξ regular, according to [12, Proposition 19.7.5], the centralizer \mathfrak{g}^{ξ} is abelian, so Formula (24) holds at least for every regular point ξ of \mathfrak{g}^* . Since the set of regular elements of \mathfrak{g}^* is dense in \mathfrak{g}^* (see [12, Proposition 19.7.5]), the continuous map $\xi \to [d_{\xi}F, d_{\xi}G]$ is zero at all points. This establishes (24). The second point of the lemma follows directly from (26).

Let us now prove Theorem 4.

Proof. (1) Let $K \in \mathcal{F}(\mathfrak{g}^* \times \mathfrak{g}^*)$ and let $(\xi, \eta) \in \mathfrak{g}^* \times \mathfrak{g}^*$. The Poisson \mathcal{R} -bracket between $\psi_1^* F = F \circ \psi_1$ and K at (ξ, η) is given by

$$\begin{split} \{\psi_1^*F,K\}_{\mathcal{R}}(\xi,\eta) &= &\frac{1}{2}\left\langle (\xi,\eta), [\mathcal{R}\,\mathsf{d}_{(\xi,\eta)}(F\circ\psi_1),\mathsf{d}_{(\xi,\eta)}K]\right\rangle \\ &+ &\frac{1}{2}\left\langle (\xi,\eta), [\mathsf{d}_{(\xi,\eta)}(F\circ\psi_1),\mathcal{R}\,\mathsf{d}_{(\xi,\eta)}K]\right\rangle. \end{split}$$

According to (22) and (23), we obtain $\mathsf{d}_{(\xi,\eta)}(F \circ \psi_1) = (\mathsf{d}_{\xi-\eta}F, \mathsf{d}_{\xi-\eta}F)$ and $\mathcal{R} \, \mathsf{d}_{(\xi,\eta)}(F \circ \psi_1) = c(\mathsf{d}_{\xi-\eta}F, \mathsf{d}_{\xi-\eta}F)$, so

$$\{\psi_1^*F, K\}_{\mathcal{R}}(\xi, \eta) = \frac{1}{2} \left\langle (\xi, \eta), [(\mathsf{d}_{\xi - \eta}F, \mathsf{d}_{\xi - \eta}F), \mathcal{R} \, \mathsf{d}_{(\xi, \eta)}K + c \mathsf{d}_{(\xi, \eta)}K] \right\rangle. \tag{27}$$

Using Formula (19), we have that $\mathcal{R} d_{(\xi,\eta)}K + cd_{(\xi,\eta)}K$ is of the form (x,x) for some $x \in \mathfrak{g}$, the explicit expression of which is not needed in this proof. Then (27) becomes

$$\{\psi_1^* F, K\}_{\mathcal{R}}(\xi, \eta) = \frac{1}{2} \langle (\xi, \eta), [(\mathsf{d}_{\xi - \eta} F, \mathsf{d}_{\xi - \eta} F), (x, x)] \rangle$$
$$= \frac{1}{2} \langle \xi - \eta, [\mathsf{d}_{\xi - \eta} F, x] \rangle$$
$$= 0,$$

where we used that F is an Ad*-invariant function on \mathfrak{g}^* in the last line. This shows that $\psi_1^* F$ is a Casimir on $((\mathfrak{g}^2)^*, \{\cdot, \cdot\}_{\mathcal{R}})$.

(2) Let us prove that the Poisson \mathcal{R} -bracket between $F \circ \psi_{\lambda}$ and $G \circ \psi_{\gamma}$ at an arbitrary point $(\xi, \eta) \in \mathfrak{g}^* \times \mathfrak{g}^*$ is equal to zero. According to (17),

$$\begin{aligned}
\left\{\psi_{\lambda}^{*}F, \psi_{\gamma}^{*}G\right\}_{\mathcal{R}}(\xi, \eta) &= \frac{1}{2}\left\langle (\xi, \eta), \left[\mathcal{R}\,\mathsf{d}_{(\xi, \eta)}(F \circ \psi_{\lambda}), \mathsf{d}_{(\xi, \eta)}(G \circ \psi_{\gamma})\right]\right\rangle \\
&+ \frac{1}{2}\left\langle (\xi, \eta), \left[\mathsf{d}_{(\xi, \eta)}(F \circ \psi_{\lambda}), \mathcal{R}\,\mathsf{d}_{(\xi, \eta)}(G \circ \psi_{\gamma})\right]\right\rangle.
\end{aligned} (28)$$

Using Formulae (22) and (23), the Poisson \mathcal{R} -bracket $\{\psi_{\lambda}^* F, \psi_{\gamma}^* G\}_{\mathcal{R}}(\xi, \eta)$ of (28) becomes

$$\begin{split} \left\{ \psi_{\lambda}^* F, \psi_{\gamma}^* G \right\}_{\mathcal{R}} (\xi, \eta) &= \frac{1}{2} \left\langle (\xi, \eta), [(\lambda - 1)(R \operatorname{d}_{\lambda \xi - \eta} F, R \operatorname{d}_{\lambda \xi - \eta} F), (\gamma \operatorname{d}_{\gamma \xi - \eta} G, \operatorname{d}_{\gamma \xi - \eta} G)] \right\rangle \\ &+ \frac{1}{2} \left\langle (\xi, \eta), [(c \operatorname{d}_{\lambda \xi - \eta} F, c \lambda \operatorname{d}_{\lambda \xi - \eta} F), (\gamma \operatorname{d}_{\gamma \xi - \eta} G, \operatorname{d}_{\gamma \xi - \eta} G)] \right\rangle \\ &- ((F, \lambda) \leftrightarrow (G, \gamma)) \\ &= \frac{(\lambda - 1)}{2} \left\langle \gamma \xi - \eta, [R \operatorname{d}_{\lambda \xi - \eta} F, \operatorname{d}_{\gamma \xi + \eta} G] \right\rangle \\ &+ \frac{c \gamma}{2} \left\langle \xi, [\operatorname{d}_{\lambda \xi - \eta} F, \operatorname{d}_{\gamma \xi - \eta} G] \right\rangle - \frac{c \lambda}{2} \left\langle \eta, [\operatorname{d}_{\lambda \xi - \eta} F, \operatorname{d}_{\gamma \xi - \eta} G] \right\rangle \\ &- ((F, \lambda) \leftrightarrow (G, \gamma)) \\ &= \frac{c \gamma}{2} \left\langle \xi, [\operatorname{d}_{\lambda \xi - \eta} F, \operatorname{d}_{\gamma \xi - \eta} G] \right\rangle - \frac{c \lambda}{2} \left\langle \eta, [\operatorname{d}_{\lambda \xi - \eta} F, \operatorname{d}_{\gamma \xi - \eta} G] \right\rangle \\ &- ((F, \lambda) \leftrightarrow (G, \gamma)), \end{split}$$

where we have used the Ad*-invariance of F and G on \mathfrak{g}^* to simplify the expression. According to Formula (25), all four terms of the previous expression vanish if $\lambda \neq \gamma$. If $\lambda = \gamma$, the bracket $\{\psi_{\lambda}^* F, \psi_{\gamma}^* G\}_{\mathcal{R}}$ becomes

$$\left\{\psi_{\lambda}^*F,\psi_{\lambda}^*G\right\}_{\mathcal{R}}(\xi,\eta) = c\lambda \left\langle \xi - \eta, \left[\mathsf{d}_{\lambda\xi-\eta}F,\mathsf{d}_{\lambda\xi-\eta}G\right]\right\rangle,$$

which is zero in view of Formula (24).

We now suppose that F and G are polynomials of degrees l and k, so for all $(\xi, \eta) \in \mathfrak{g} \times \mathfrak{g}^*$, $F \circ \psi_{\lambda}(\xi, \eta) = \sum_{i=0}^{l} \lambda^i F_i(\xi, \eta)$ and $G \circ \psi_{\gamma}(\xi, \eta) = \sum_{j=0}^{k} \gamma^j G_j(\xi, \eta)$; hence

$$\{F \circ \psi_{\lambda}, G \circ \psi_{\gamma}\}_{\mathcal{R}} = \sum_{i=0}^{l} \sum_{j=0}^{k} \lambda^{i} \gamma^{j} \{F_{i}, G_{j}\}_{\mathcal{R}}.$$
 (29)

As we just showed, the term on the left hand side of Equation (29) is zero, for all $\lambda, \gamma \in \mathbf{F}$. Therefore, all the coefficients of $\sum_{i=0}^{l} \sum_{j=0}^{k} \lambda^{i} \gamma^{j} \{F_{i}, G_{j}\}_{\mathcal{R}}$ are zero.

(3) We assume that c = 1. Let $F, G \in \mathcal{F}(\mathfrak{g}^*)$ and let (ξ, η) be a point in $\mathfrak{g}^* \times \mathfrak{g}^*$. From (28), (23) and (22), it follows that the Poisson \mathcal{R} -bracket between $F \circ \psi_1$ and $G \circ \psi_1$ at (ξ, η) is given by

$$\{F \circ \psi_1, G \circ \psi_1\}_{\mathcal{P}}(\xi, \eta) = \langle \xi - \eta, [\mathsf{d}_{\xi - \eta} F, \mathsf{d}_{\xi - \eta} G] \rangle = \{F, G\} (\psi_1(\xi, \eta)).$$

Hence ψ_1^* is a Poisson map.

(4) Let K be a function on $\mathfrak{g}^* \times \mathfrak{g}^*$ and (ξ, η) be an element of $\mathfrak{g}^* \times \mathfrak{g}^*$. A direct computation gives

$$\mathcal{X}_{H \circ \psi_{\lambda}}(\xi, \eta)[K] = \{K, H \circ \psi_{\lambda}\}_{\mathcal{R}}(\xi, \eta)
= \frac{1}{2} \left\langle (\xi, \eta), \left[\mathcal{R} \, \mathsf{d}_{(\xi, \eta)} K, \mathsf{d}_{(\xi, \eta)} (H \circ \psi_{\lambda}) \right] + \left[\mathsf{d}_{(\xi, \eta)} K, \mathcal{R} \, \mathsf{d}_{(\xi, \eta)} (H \circ \psi_{\lambda}) \right] \right\rangle.$$
(30)

In order to rewrite this formula, we temporarily use $(x,y) := \mathsf{d}_{(\xi,\eta)}K$. According to (19), $\mathcal{R}\,\mathsf{d}_{(\xi,\eta)}K = \mathcal{R}(x,y)$ is given by

$$\mathcal{R} \mathsf{d}_{(\xi,\eta)} K = (R(x-y), R(x-y)) + c(y,x). \tag{31}$$

Formulae (31) and (22) allow one to rewrite the first term of (30) in the following manner:

$$\begin{split} \left\langle (\xi,\eta), \left[\mathcal{R} \mathsf{d}_{(\xi,\eta)} K, \mathsf{d}_{(\xi,\eta)} (H \circ \psi_{\lambda}) \right] \right\rangle \\ &= \left\langle (\xi,\eta), \left[(R(x-y), R(x-y)), (\lambda \mathsf{d}_{\lambda\xi-\eta} H, \mathsf{d}_{\lambda\xi-\eta} H) \right] \right\rangle \\ &+ c \left\langle (\xi,\eta), \left[(y,x), (\lambda \mathsf{d}_{\lambda\xi-\eta} H, \mathsf{d}_{\lambda\xi-\eta} H) \right] \right\rangle \\ &= \left\langle \lambda\xi - \eta, \left[R(x-y), \mathsf{d}_{\lambda\xi-\eta} H \right] \right\rangle + c \left\langle \lambda\xi, \left[y, \mathsf{d}_{\lambda\xi-\eta} H \right] \right\rangle - c \left\langle \eta, \left[x, \mathsf{d}_{\lambda\xi-\eta} H \right] \right\rangle \\ &= c \left\langle \eta, \left[y, \mathsf{d}_{\lambda\xi-\eta} H \right] \right\rangle - c \left\langle \lambda\xi, \left[x, \mathsf{d}_{\lambda\xi-\eta} H \right] \right\rangle, \end{split} \tag{32}$$

where, in the second line, we used, three times, the Ad^* -invariance of H. Thus

$$\left\langle (\xi, \eta), \left[\mathcal{R} \, \mathsf{d}_{(\xi, \eta)} K, \mathsf{d}_{(\xi, \eta)} (H \circ \psi_{\lambda}) \right] \right\rangle = \left\langle (\xi, \eta), \left[(c \lambda \mathsf{d}_{\lambda \xi - \eta} H, c \mathsf{d}_{\lambda \xi - \eta} H), (x, y) \right] \right\rangle. \tag{33}$$

By replacing the first term of (30) by its expression given in (33) and by using Formula (23) to express the second term of (30), we rewrite $\mathcal{X}_{H\circ\psi_{\lambda}}(\xi,\eta)[F]$ as follows:

$$\begin{split} \mathcal{X}_{H\circ\psi_{\lambda}}(\xi,\eta)[K] &= \frac{1}{2} \left\langle (\xi,\eta), [(c\lambda\mathsf{d}_{\lambda\xi-\eta}H,c\mathsf{d}_{\lambda\xi-\eta}H),(x,y)] \right\rangle \\ &- \frac{(\lambda-1)}{2} \left\langle (\xi,\eta), [(R\,\mathsf{d}_{\lambda\xi-\eta}H,R\,\mathsf{d}_{\lambda\xi-\eta}H),(x,y)] \right\rangle \\ &- \frac{1}{2} \left\langle (\xi,\eta), [(c\mathsf{d}_{\lambda\xi-\eta}H,c\lambda\mathsf{d}_{\lambda\xi-\eta}H),(x,y)] \right\rangle \\ &= \frac{(\lambda-1)}{2} \left\langle (\xi,\eta), [((cI-R)\mathsf{d}_{\lambda\xi-\eta}H,-(R+cI)\mathsf{d}_{\lambda\xi-\eta}H),\mathsf{d}_{(\xi,\eta)}K] \right\rangle. \end{split}$$

We then deduce Formula (20). When \mathfrak{g} is a Lie algebra splitting and $R = P_+ - P_-$, we have c = 1, $R - cI = -2P_-$ and $R + cI = 2P_+$. This implies that (20) gives (21).

3 The Liouville integrability of the 2-Toda lattice

In this section we define the 2-Toda lattice for every complex simple Lie algebra and we prove its Liouville integrability.

In order to define the 2-Toda lattice for every simple Lie algebra, we need some notation. Let \mathfrak{g} be a simple Lie algebra of rank ℓ , with Killing form $\langle \cdot | \cdot \rangle$. We choose \mathfrak{h} , a Cartan subalgebra with roots system Φ , and $\Pi = (\alpha_1, \ldots, \alpha_\ell)$, a system of simple roots with respect to \mathfrak{h} . For every α in $\Phi \setminus \{-\Pi, \Pi\}$, we denote by e_{α} a non-zero eigenvector associated with eigenvalue α , and, for every $1 \leq i \leq \ell$, we denote by e_i and e_{-i} non-zero eigenvectors associated respectively with α_i and $-\alpha_i$. The Lie algebra $\mathfrak{g} = \sum_{k \in \mathbb{Z}} \mathfrak{g}_k$ is endowed with the natural grading (i.e., for every $k, l \in \mathbb{Z}$, $[\mathfrak{g}_k, \mathfrak{g}_l] \subset \mathfrak{g}_{k+l}$) defined by $\mathfrak{g}_0 := \mathfrak{h}$ and, for every $k \in \mathbb{Z}$, $\mathfrak{g}_k := \langle e_{\alpha} \mid \alpha \in \Phi, |\alpha| = k \rangle$, for $|\alpha|$ the length of the root α , i.e., $|\alpha|$ is $\sum_{i=1}^{\ell} a_i$ for $\alpha = \sum_{i=1}^{\ell} a_i \alpha_i$. In the sequel, we shall use the following property: $\langle \mathfrak{g}_k \mid \mathfrak{g}_l \rangle = 0$ if $k + l \neq 0$. We introduce the following notation

$$\begin{split} \mathfrak{g}_{< k} &:= \sum_{i < k} \mathfrak{g}_i, & \mathfrak{g}_{\leqslant k} &:= \sum_{i \leqslant k} \mathfrak{g}_i, \\ \mathfrak{g}_{> k} &:= \sum_{i > k} \mathfrak{g}_i, & \mathfrak{g}_{\geqslant k} &:= \sum_{i \geqslant k} \mathfrak{g}_i. \end{split}$$

Also, $\mathfrak{g}_+ := \mathfrak{g}_{\geqslant 0}$ and $\mathfrak{g}_- := \mathfrak{g}_{< 0}$.

3.1 Definition of the 2-Toda lattice

The next definition gives again the definition given in (1) when specialized to the case $\mathfrak{g} = \mathfrak{sl}_n(\mathbf{C})$, taking for \mathfrak{h} the Lie subalgebra of diagonal matrices.

Definition 7 The 2-Toda lattice associated with a simple Lie algebra \mathfrak{g} is the system of differential equations given by the following Lax equations:

$$\frac{\partial(L,M)}{\partial t} = [(L_+,L_+),(L,M)], \tag{34}$$

$$\frac{\partial(L,M)}{\partial s} = [(M_-,M_-),(L,M)], \tag{35}$$

where (L,M) is an element of the phase space of the 2-Toda lattice $\mathcal{T}^2 := \mathfrak{g}_{\leq 0} \times \mathfrak{g}_{\geqslant -1} + (\sum_{i=1}^{\ell} e_i, 0)$, where $L_+ := P_+(L)$, $M_- := P_-(M)$, and where P_{\pm} is the projection of \mathfrak{g} on \mathfrak{g}_+ .

3.2 The 2-Toda lattice is a Hamiltonian system

We recall from Example 3 that when $\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$ is a Lie algebra splitting then $\mathfrak{g}^2 = \mathfrak{g}_+^2 \oplus \mathfrak{g}_-^2$ is also a Lie algebra splitting, where

$$g_{+}^{2} := \{(x, x) \mid x \in \mathfrak{g}\} \text{ et } \mathfrak{g}_{-}^{2} := \{(x, y) \mid x \in \mathfrak{g}_{-} \text{ et } y \in \mathfrak{g}_{+}\}.$$
 (36)

Also, for every $(x, y) \in \mathfrak{g}^2$ we have $(x, y) = (x, y)_+ + (x, y)_-$, where

$$\begin{cases}
(x,y)_{+} = (x_{+} + y_{-}, x_{+} + y_{-}) \in \mathfrak{g}_{+}^{2}, \\
(x,y)_{-} = (x_{-} - y_{-}, y_{+} - x_{+}) \in \mathfrak{g}_{-}^{2}.
\end{cases}$$
(37)

Let \mathcal{R} be the difference of the projections of \mathfrak{g}^2 on the Lie subalgebras \mathfrak{g}_+^2 and \mathfrak{g}_-^2 . According to Example 3, we have

$$\mathcal{R}(x,y) = (x_{+} - x_{-} + 2y_{-}, y_{-} - y_{+} + 2x_{+}) \tag{38}$$

$$= (R(x-y) + y, R(x-y) + x), (39)$$

where R is the difference of the projections of \mathfrak{g} on \mathfrak{g}_+ and on \mathfrak{g}_- . We provide \mathfrak{g}^2 with the following Ad-invariant, non-degenerate symmetric bilinear form:

$$\langle \cdot | \cdot \rangle_2 : \mathfrak{g}^2 \times \mathfrak{g}^2 \to \mathbf{C}$$

$$((x_1, y_1), (x_2, y_2)) \mapsto \langle x_1 | x_2 \rangle - \langle y_1 | y_2 \rangle.$$

$$(40)$$

We use it to identify \mathfrak{g}^2 with its dual and we obtain according to (16) a linear Poisson structure on \mathfrak{g}^2 , defined for every $F, G \in \mathcal{F}(\mathfrak{g}^2)$ at $(x, y) \in \mathfrak{g}^2$ by

$$\{F,G\}_{\mathcal{R}}(x,y) = \frac{1}{2} \langle (x,y) | [\mathcal{R}\nabla_{(x,y)}F, \nabla_{(x,y)}G] + [\nabla_{(x,y)}F, \mathcal{R}\nabla_{(x,y)}G] \rangle_2, \tag{41}$$

where $\nabla_{(x,y)}F$ is the gradient of F at $(x,y) \in \mathfrak{g}^2$ (with respect to $\langle \cdot | \cdot \rangle_2$), i.e.,

$$\left\langle \nabla_{(x,y)}F\left|\left(z,s\right)\right\rangle _{2}=\left\langle \mathsf{d}_{(x,y)}F,(z,s)\right\rangle ,\qquad\forall(z,s)\in\mathfrak{g}^{2}.$$

We show that the phase space \mathcal{T}^2 is equipped with a Poisson structure and the equations of motion of the 2-Toda lattice are Hamiltonian.

Proposition 8 \mathcal{T}^2 is a Poisson submanifold of $(\mathfrak{g}^2, \{\cdot, \cdot\}_{\mathcal{R}})$.

We use the following lemma to show the above proposition.

Lemma 9 Let \mathfrak{g} be a Lie algebra equipped with a non-degenerate, symmetric, bilinear form $\langle \cdot | \cdot \rangle$. Let $a \in \mathfrak{g}$ and let E be a subspace of \mathfrak{g} . We suppose that:

- (1) The orthogonal³ E^{\perp} of E is a Lie ideal of $(\mathfrak{g}, [\cdot, \cdot])$.
- (2) For every $x, y \in \mathfrak{g}$, we have $\langle a \mid [x, y] \rangle = 0$.

Then a+E and E are Poisson submanifolds of $(\mathfrak{g}, \{\cdot,\cdot\})$, equipped with its linear Poisson structure $\{\cdot,\cdot\}$. The map $x \to a+x$ is a Poisson isomorphism between them. Moreover E is Poisson isomorphic to the Lie-Poisson manifold $(\mathfrak{g}/E^{\perp})^*$, also equipped with its linear Poisson structure.

³Here the orthogonality is with respect to the form $\langle \cdot | \cdot \rangle$.

Proof. We start to show that E is a Poisson submanifold of $(\mathfrak{g}, \{\cdot, \cdot\})$. Let $\mathcal{J} := \langle F \in \mathcal{F}(\mathfrak{g}) \mid F \equiv 0$ on $E \rangle$. To show that E is a Poisson submanifold is equivalent to show that \mathcal{J} is Poisson ideal. Let $F \in \mathcal{J}$ and let $x \in E$, we notice that $\nabla_x F \in E^{\perp}$. According to the first condition of the proposition, for every $G \in \mathcal{F}(\mathfrak{g})$, we have $[\nabla_x f, \nabla_x G] \in E^{\perp}$. This implies that $\{F, G\}(x) = \langle x \mid [\nabla_x f, \nabla_x G] \rangle = 0$. Then \mathcal{J} is a Poisson ideal.

We now show that E + a is a Poisson submanifold. Let $\mathcal{I} := \langle F \in \mathcal{F}(\mathfrak{g}^2) \mid F \equiv 0$ on $E + a \rangle$, $F \in \mathcal{I}$ and let $x + a \in E + a$. Notice that $\nabla_{x+a}F$ is an element of E^{\perp} , which is according to first condition of these proposition is a Lie ideal of $(\mathfrak{g}, [\cdot, \cdot])$. Then, for every $G \in \mathcal{F}(\mathfrak{g})$, $[\nabla_{x+a}F, \nabla_{x+a}G] \in E^{\perp}$. Let now compute the Poisson bracket between F and $G \in \mathcal{F}(\mathfrak{g})$ at a point x + a.

$$\{F,G\} (x+a) = \langle x+a | [\nabla_{x+a}F, \nabla_{x+a}G] \rangle$$
$$= \langle x | [\nabla_{x+a}F, \nabla_{x+a}G] \rangle$$
$$= 0,$$

where we have used the second condition of proposition to justify the transition from first to second line and we have zero in the last line because $x \in E$ and $[\nabla_{x+a}F, \nabla_{x+a}G] \in E^{\perp}$.

Let now show the translation by a is a Poisson isomorphism between \mathfrak{g} and \mathfrak{g} . We denote by T_a the translation by a, defined for every $x \in \mathfrak{g}$, by $T_a(x) = x + a$. For every $F, G \in \mathcal{F}(\mathfrak{g})$ and every $x \in \mathfrak{g}$, we have

$$\begin{aligned} \left\{ F \circ T_{a}, G \circ T_{a} \right\} (x) &= \left\langle x \, | \, \left[\nabla_{x} (F \circ T_{a}), \nabla_{x} (G \circ T_{a}) \right] \right\rangle \\ &= \left\langle x \, | \, \left[\nabla_{T_{a}(x)} F, \nabla_{T_{a}(x)} G \right] \right\rangle \\ &= \left\langle x + a \, | \, \left[\nabla_{T_{a}(x)} F, \nabla_{T_{a}(x)} G \right] \right\rangle \\ &= \left\langle F, G \right\} (T_{a}(x)), \end{aligned}$$

where we have used Condition (2) to justify the transition from second to third line. Since E is a Poisson submanifold of $(\mathfrak{g}, \{\cdot, \cdot\})$ the restriction of the Poisson isomorphism T_a to E is also a Poisson isomorphism on its image.

We now prove Proposition 8.

Proof. Its easy to verify that \mathcal{T}^2 admits the following description as an affine subspace of \mathfrak{g}^2 :

$$\mathcal{T}^2 = (e, e) + \Delta(\mathfrak{g}_0 \oplus \mathfrak{g}_{-1}) \oplus \mathfrak{g}_{-}^2, \tag{42}$$

where $\Delta(\mathfrak{g}_0 \oplus \mathfrak{g}_{-1}) := \{(x,x) \mid x \in \mathfrak{g}_0 \oplus \mathfrak{g}_{-1}\}$ and $e = \sum_{i=1}^{\ell} e_i$. We check that the two assumptions of Lemma 9 are satisfied, with a = (e,e), $E = \Delta(\mathfrak{g}_0 \oplus \mathfrak{g}_{-1}) \oplus \mathfrak{g}_{-}^2$, $\mathfrak{g} = \mathfrak{g}^2$ and $[\cdot,\cdot] = [\cdot,\cdot]_{\mathcal{R}}$.

(1) It is clear that $E = \mathfrak{g}_{\leq 0} \times \mathfrak{g}_{\geq -1}$. The orthogonal of E is $E^{\perp} = \mathfrak{g}_{<0} \times \mathfrak{g}_{>1}$, which is a subspace of \mathfrak{g}_{-}^2 and it is an ideal of $(\mathfrak{g}^2, [\cdot, \cdot]_{\mathcal{R}})$, because, for every $(x, y) \in \mathfrak{g}_{<0} \times \mathfrak{g}_{>1}$ and $(z, s) \in \mathfrak{g}^2$,

$$[(x,y),(z,s)]_{\mathcal{R}} \in [\mathfrak{g}_{<0} \times \mathfrak{g}_{>1},\mathfrak{g}_{<0} \times \mathfrak{g}_{\geqslant 0}] \subset \mathfrak{g}_{<-1} \times \mathfrak{g}_{>1} \subset \mathfrak{g}_{<0} \times \mathfrak{g}_{>1}.$$

(2) For every $(x, y), (x', y') \in \mathfrak{g}^2$, we have

$$\left\langle (e,e) \,|\, [(x,y),(x',y')]_{\mathcal{R}} \right\rangle = \left\langle (e,e) \,|\, [(x,y)_+,(x',y')_+] - [(x,y)_-,(x',y')_-] \right\rangle_2. \tag{43}$$

Since $[(x,y)_+,(x',y')_+]=(u,u)$ for some $u \in \mathfrak{g}$ it follows that the first term on the right side of Equation (43) is zero. Also $[(x,y)_-,(x',y')_-] \in \mathfrak{g}_{<-1} \times \mathfrak{g}_{>0}$, which is orthogonal to $(e,e) \in \mathfrak{g}_1 \times \mathfrak{g}_1$; hence the second term on the right hand side of Equation (43) is zero.

Proposition 10 Let $H, \tilde{H} \in \mathcal{F}(\mathfrak{g}^2)$ be defined at every point (x, y) of \mathfrak{g}^2 by

$$H(x,y) := \frac{1}{2} \langle x \, | \, x \rangle \quad \text{ and } \quad \tilde{H}(x,y) := \frac{1}{2} \langle y \, | \, y \rangle \,. \tag{44}$$

The Hamiltonian vector field $\mathcal{X}_H := \{\cdot, H\}_{\mathcal{R}}$ (resp. $\mathcal{X}_{\tilde{H}} := \{\cdot, \tilde{H}\}_{\mathcal{R}}$) is tangent to \mathcal{T}^2 and describes on \mathcal{T}^2 the equation of motion (34) (resp. (35)) for the 2-Toda lattice.

Proof. Let $F \in \mathcal{F}(\mathfrak{g}^2)$ and let $(L, M) \in \mathcal{T}^2$. We have

$$\mathcal{X}_{H}[F](L,M) = \{F,H\}_{\mathcal{R}}(L,M)
= \frac{1}{2} \langle (L,M) | \left[\mathcal{R} \nabla_{(L,M)} F, \nabla_{(L,M)} H \right] \rangle_{2}
+ \frac{1}{2} \langle (L,M) | \left[\nabla_{(L,M)} F, \mathcal{R} \nabla_{(L,M)} H \right] \rangle_{2}
= -\frac{1}{2} \langle \left[(L,M), \mathcal{R} \nabla_{(L,M)} H \right] | \nabla_{(L,M)} F \rangle_{2},$$
(45)

where we have used the Ad-invariance on \mathfrak{g}^2 of H to justify the transition from second to third equality. We deduce from Equation (45) that $\mathcal{X}_H(L,M) = \frac{1}{2} \left[\mathcal{R} \nabla_{(L,M)} H, (L,M) \right]$. Hence according to Formula (37), we have

$$\mathcal{X}_H(L,M) = [(\nabla_{(L,M)}H)_+, (L,M)] = [(L,0)_+, (L,M)] = [(L_+,L_+), (L,M)].$$

To show that $\mathcal{X}_{\tilde{H}}$ describes Equation (35) it suffices to repeat the same reasoning of \mathcal{X}_H . Since \mathcal{T}^2 is a Poisson submanifold of $(\mathfrak{g}^2, \{\cdot, \cdot\})$, the Hamiltonian vector fields \mathcal{X}_H and $\mathcal{X}_{\tilde{H}}$ are tangent to \mathcal{T}^2 .

3.3 The integrability of the 2-Toda lattice

According to [5, Theorem 7.3.8], for every simple Lie algebra \mathfrak{g} of rank ℓ , there exist ℓ homogeneous, independent, Ad-invariant polynomials P_1, \ldots, P_ℓ which generate the algebra of Ad-invariant polynomial functions on \mathfrak{g} and which are of degree, respectively, $m_1 + 1, \ldots, m_\ell + 1$, where m_1, \ldots, m_ℓ are the exponents of \mathfrak{g} (we note that $m_1 \leq \ldots \leq m_\ell$). Each P_i induces $m_i + 2$ functions $F_{j,i} \in \mathcal{F}(\mathfrak{g}^2)$, as follows:

$$P_i(\lambda x - y) = \sum_{0 \le j \le m_i + 1} (-1)^{m_i + 1 - j} \lambda^j F_{j,i}(x, y).$$
 (46)

Every function $F_{j,i}$ for $1 \le i \le \ell$ and $0 \le j \le m_i + 1$ is homogeneous of degree j with respect to its first variable and of degree $m_i + 1 - j$ with respect to its second variable.

Notation 11 We denote by \mathcal{F} the family of functions on \mathfrak{g}^2 given by

$$\mathcal{F} := (F_{j,i}, 1 \leqslant i \leqslant \ell \text{ and } 0 \leqslant j \leqslant m_i + 1). \tag{47}$$

Remark 12 (1) The functions $F_{0,1}$ and $F_{2,1}$ are the Hamiltonians of the 2-Toda lattice introduced in Proposition 10.

(2) The functions $F_{0,i}$ and $F_{m_i+1,i}$ for $1 \le i \le \ell$ are Ad-invariant functions on \mathfrak{g}^2 . According to the Adler-Kostant-Symes theorem [2, Theorem 4.37] they are in involution with respect to the bracket $\{\cdot,\cdot\}_{\mathcal{R}}$. Also they are independent on \mathfrak{g}^2 . They are therefore a good candidates for giving the Liouville integrability of the 2-Toda lattice. However their cardinal 2ℓ is very small compared to dim $\mathcal{T}^2 - \frac{1}{2} \operatorname{Rk}(\mathcal{T}^2, \{\cdot,\cdot\}_{\mathcal{R}})$.

Since, as we noticed in Remark 12, the Hamiltonians of the 2-Toda lattice appear among the functions composing \mathcal{F} , the next theorem gives the Liouville integrability of the 2-Toda lattice.

Theorem 13 The triplet $(\mathcal{T}^2, \mathcal{F}_{|\mathcal{T}^2}, \{\cdot, \cdot\}_{\mathcal{R}})$ is an integrable system.

Proof. According to the definition of integrability in the sense of Liouville (see [2, Definition 4.13]), to prove Theorem 4 we must show that

- (1) $\mathcal{F}_{|\mathcal{T}^2}$ is involutive for the Poisson \mathcal{R} -bracket $\{\cdot\,,\cdot\}_{\mathcal{R}}$.
- (2) $\mathcal{F}_{|\mathcal{T}^2}$ is independent.
- (3) The cardinal of the restriction of \mathcal{F} to \mathcal{T}^2 satisfies

$$\operatorname{card} \mathcal{F}_{|\mathcal{T}^2} = \dim \mathcal{T}^2 - \frac{1}{2} \operatorname{Rk}(\mathcal{T}^2, \{\cdot, \cdot\}_{\mathcal{R}}). \tag{48}$$

The proofs of these three points are given in respectively Proposition 14, Proposition 17 and Proposition 19, which are given in the next three subsections. \Box

3.3.1 The restriction of \mathcal{F} to \mathcal{T}^2 is involutive for $\{\cdot\,,\cdot\}_{\mathcal{R}}$

Unlike in the Toda lattice case, it is not the Adler-Kostant-Symes theorem that gives us the commutativity of the considered family of functions \mathcal{F} , which is not formed only by the Ad-invariant functions. This commutativity arises from the fact our phase space is a Poisson submanifold of $(\mathfrak{g}^2, \{\cdot, \cdot\}_{\mathcal{R}})$, and from the second item of Theorem 4.

Proposition 14 The family of functions $\mathcal{F}_{|\mathcal{T}^2}$ is involutive for the Poisson structure $\{\cdot,\cdot\}_{\mathcal{R}}$.

Proof. Since the polynomials P_1, \ldots, P_ℓ are Ad-invariant on \mathfrak{g} (and hence Ad*-invariant upon identifying \mathfrak{g} with \mathfrak{g}^*), according to the second point of Theorem 4 the family \mathcal{F} is involutive on $(\mathfrak{g}^2, \{\cdot, \cdot\}_{\mathcal{R}})$. Furthermore, \mathcal{T}^2 being a Poisson submanifold of $(\mathfrak{g}^2, \{\cdot, \cdot\}_{\mathcal{R}})$ (see Proposition 8), the restriction of \mathcal{F} to \mathcal{T}^2 is involutive.

3.3.2 The restriction of \mathcal{F} to \mathcal{T}^2 is independent

We use an unpublished result of Raïs [10], which establishes the independence of a large family of functions on \mathfrak{g}^2 . We state this result below and the proof is in [4, Section 1].

Theorem 15 Let P_1, \ldots, P_ℓ be a generating family of homogeneous polynomials of the algebra of Ad-invariant polynomial functions on \mathfrak{g} . Let e and h be two elements of \mathfrak{g} such that e is regular and [h, e] = 2e.

For every $F \in \mathcal{F}(\mathfrak{g})$, and every $y \in \mathfrak{g}$, we denote by $\mathsf{d}_y^k F$ the differential of order k of F at y. Denote by $V_{k,i}$, for every $1 \leqslant i \leqslant \ell$ and $0 \leqslant k \leqslant m_i$, the element of \mathfrak{g} defined by

$$\langle V_{k,i} | z \rangle = \left\langle \mathsf{d}_h^{k+1} P_i, (e^k, z) \right\rangle, \qquad \forall z \in \mathfrak{g},$$
 (49)

where, for every $x \in \mathfrak{g}$ and $k \in \mathbb{N}$, x^k is shorthand for (x, \ldots, x) (k times).

- (1) The family $\mathcal{F}_1 := (V_{k,i}, 1 \leq i \leq \ell \text{ and } 0 \leq k \leq m_i)$ is linearly independent.
- (2) The subspace generated by \mathcal{F}_1 is the Lie subalgebra formed by the sum of the all eigenspaces of ad_h associated with positive or zero eigenvalues.

We first prove, using the first point of Theorem 15, the independence of \mathcal{F} (which is a family $\mathcal{F}(\mathfrak{g}^2)$) at a well chosen point $(e,h) \in \mathcal{T}^2 \cap \mathfrak{g}^2$. Afterwards, using the second point of Theorem 15, we show that the restriction of \mathcal{F} to the phase space \mathcal{T}^2 of the 2-Toda lattice is also an independent family of functions.

Proposition 16 Let $h \in \mathfrak{h}$ be such that [h, e] = 2e.

- (1) The polynomial functions $F_{0,1}(x,y), \ldots, F_{0,\ell}(x,y)$ only depend on the second variable y. Their differentials at the point (e,h) are independent;
- (2) We denote by $\frac{\partial}{\partial x}$ the partial differential with respect to the variable x. The $\frac{1}{2}(\dim \mathfrak{g} + \ell)$ partial derivatives $\frac{\partial F_{j,i}}{\partial x}$, $1 \leq i \leq \ell$ and $1 \leq j \leq m_i + 1$ are independent at the point (e,h);
- (3) The family $(d_{(e,h)}F_{i,i}, 1 \le i \le \ell, 0 \le j \le m_i + 1)$ is independent.

Proof. (1) For every $1 \leq i \leq \ell$ and every $(x,y) \in \mathfrak{g}^2$, $F_{0,i}(x,y)$ is the term of degree 0 in λ of $P_i(\lambda x - y)$, that is $P_i(y)$. Hence, the function $F_{0,i}$, for $1 \leq i \leq \ell$, is a homogeneous polynomial of degree $m_i + 1$ which only depends on the second variable y. Moreover, according to two theorems of Kostant [6, Theorem 9] and [7, Theorem 5.2], the differentials of the polynomials P_1, \ldots, P_ℓ are independent at every regular point of \mathfrak{g} . In particular,

they are independent at the regular⁴ point h. In conclusion the differentials of the functions $F_{0,1}, \ldots, F_{0,\ell}$ are independent at (e, h).

(2) For $1 \leq i \leq \ell$, the Taylor Formula applied to the polynomial P_i , $1 \leq i \leq \ell$, at $\lambda x - y$, yields

$$P_{i}(\lambda x - y) = \sum_{k=0}^{m_{i}+1} (-1)^{m_{i}+1-k} \frac{\lambda^{k}}{k!} \left\langle \mathsf{d}_{y}^{k} P_{i}, x^{k} \right\rangle. \tag{50}$$

By identifying the coefficients of Equations (50) and (46), we obtain, for every $1 \le i \le \ell$ and $1 \le k \le m_i + 1$, the equality

$$F_{k,i}(x,y) = \frac{1}{k!} \left\langle \mathsf{d}_y^k P_i, x^k \right\rangle. \tag{51}$$

By differentiating $F_{k+1,i}$ with respect to the variable x, we obtain

$$\left\langle \frac{\partial F_{k+1,i}}{\partial x}(x,y), z \right\rangle = \frac{1}{k!} \left\langle \mathsf{d}_y^{k+1} P_i, (x^k, z) \right\rangle, \qquad \forall k = 0, \dots, m_i, \text{ and } \forall z \in \mathfrak{g}.$$
 (52)

In particular, according to Theorem 15, when (x, y) = (e, h), Equation (52) becomes

$$\left\langle \frac{\partial F_{k+1,i}}{\partial x}(e,h), z \right\rangle = \frac{1}{k!} \left\langle \mathsf{d}_{h}^{k+1} P_{i}, (e^{k}, z) \right\rangle = \frac{1}{k!} \left\langle V_{k,i} \, | \, z \right\rangle, \qquad \forall z \in \mathfrak{g}, \tag{53}$$

for every $1 \leqslant i \leqslant \ell$ and $1 \leqslant k \leqslant m_i + 1$. According to Theorem 15, the family $(V_{k,i}, 1 \leqslant i \leqslant \ell \text{ and } 0 \leqslant k \leqslant \ell)$ is independent. This implies the independence of the family $(\frac{\partial F_{k,i}}{\partial x}(e,h), 1 \leqslant i \leqslant \ell \text{ and } 1 \leqslant k \leqslant m_i + 1)$.

(3) Denote by M the matrix

$$M = (\mathsf{d}_{(e,h)}F_{k,i}, 0 \leqslant k \leqslant m_i + 1 \text{ and } 1 \leqslant i \leqslant \ell)$$

$$= \begin{pmatrix} \frac{\partial F_{0,1}}{\partial x}(e,h) & \frac{\partial F_{0,1}}{\partial y}(e,h) \\ \vdots & \vdots \\ \frac{\partial F_{0,\ell}}{\partial x}(e,h) & \frac{\partial F_{0,\ell}}{\partial y}(e,h) \\ \frac{\partial F_{1,1}}{\partial x}(e,h) & \frac{\partial F_{1,1}}{\partial y}(e,h) \\ \vdots & \vdots \\ \frac{\partial F_{m_\ell+1,\ell}}{\partial x}(e,h) & \frac{\partial F_{m_\ell+1,\ell}}{\partial y}(e,h) \end{pmatrix}.$$

According to item (1), the form of the matrix M is

$$M = \left(\begin{array}{cc} 0 & A \\ B & * \end{array}\right),$$

The element h of \mathfrak{g} is regular. Indeed, since it verifies [h, e] = 2e with e regular, it belongs to a principal $\mathfrak{sl}(2)$ triple (see [12, Theorem 32.1.5]).

where $A = (\frac{\partial F_{0,i}}{\partial y}(e,h), 1 \leqslant i \leqslant \ell)$ and $B = (\frac{\partial F_{k,i}}{\partial x}(e,h), 1 \leqslant k \leqslant m_i + 1 \text{ and } 1 \leqslant i \leqslant \ell)$. According to (1) and (2), the ℓ rows of the matrix A and the $\frac{1}{2}(\dim \mathfrak{g} + \ell)$ rows of the matrix B are independent. The independence of the rows of M yields the independence of the differentials of the family of functions \mathcal{F} at $(e,h) \in \mathcal{T}^2$.

Proposition 17 The restriction of the family of functions \mathcal{F} to the phase space \mathcal{T}^2 of the 2-Toda lattice is an independent family.

Proof. It suffices to show that the restrictions on the tangent space $\mathbf{T}_{(e,h)}\mathcal{T}^2$ of the differentials

$$(\mathsf{d}_{(e,h)}F_{j,i}, \ 1 \leqslant i \leqslant \ell \text{ et } 0 \leqslant j \leqslant m_i + 1)$$

are independent. Since the tangent space of \mathcal{T}^2 is $\mathfrak{g}_{\leq 0} \times \mathfrak{g}_{\geqslant -1}$ and since $(\mathsf{d}_{(e,h)}F_{0,i}, 1 \leq i \leq \ell)$ vanish on $\mathfrak{g}_{\leq 0} \times \{0\}$, it suffices to show that:

(a) The restriction to the space $\{0\} \times \mathfrak{h}$ of the linear forms

$$(\mathsf{d}_{(e,h)}F_{0,i}, \ 1 \leqslant i \leqslant \ell)$$

is an independent family.

(b) The restriction to the space $\mathfrak{g}_{\leq 0} \times \{0\}$ of the linear independent forms

$$(\mathsf{d}_{(e,h)}F_{j,i}, \ 1 \leqslant i \leqslant \ell \text{ and } 1 \leqslant j \leqslant m_i + 1)$$

is an independent family.

According to Definition (46), $F_{0,i}(x,y) = P_i(y)$ for every $1 \le i \le \ell$ and every $(x,y) \in \mathfrak{g}^2$. Then $\frac{\partial F_{0,i}}{\partial x}(e,h) = 0$ and

$$\left\langle \frac{\partial F_{0,i}}{\partial y}(e,h), z \right\rangle = \left\langle \nabla_h P_i \,|\, z \right\rangle, \qquad \forall z \in \mathfrak{g}.$$
 (54)

Since $\nabla_h P_i \in \mathfrak{h}$ (because $[h, \nabla_h P_i] = 0$ and $h \in \mathfrak{h}$ is regular), the restriction to \mathfrak{h} of the family $(\frac{\partial F_{0,1}}{\partial y}(e,h),\ldots,\frac{\partial F_{0,\ell}}{\partial y}(e,h))$ is an independent family of linear forms. This implies the independence of the restriction to $\{0\} \times \mathfrak{h}$ of $\mathsf{d}_{(e,h)}F_{0,1},\ldots,\mathsf{d}_{(e,h)}F_{0,\ell}$.

According to equation (53), for every $1 \leq i \leq \ell$ and $0 \leq k \leq \ell$,

$$\langle V_{k,i} | z \rangle = k! \left\langle \frac{\partial F_{k+1,i}}{\partial x} (e,h), z \right\rangle.$$
 (55)

The second point of Theorem 15 shows that the subspace generated by the family $(V_{k,i}, 1 \le i \le \ell \text{ and } 0 \le k \le m_i)$ is contained in the Lie subalgebra obtained by summing the eigenspaces of ad_h , associated with the nonnegative eigenvalues, which is in our case $\mathfrak{g}_{\ge 0}$. This proves, using Equation (55), that the restriction to $\mathfrak{g}_{\le 0}$ of the family of linear forms $(\frac{\partial F_{k,i}}{\partial x}(e,h), 1 \le i \le \ell \text{ and } 1 \le k \le m_i + 1)$ is an independent family. Therefore the restriction to $\mathfrak{g}_{\le 0} \times \{0\}$ of $(\mathsf{d}_{(e,h)}F_{k,i}, 1 \le i \le \ell \text{ and } 1 \le i \le m_i + 1)$ is an independent family.

3.3.3 The exact number of functions

According to Equation (47), the cardinal of \mathcal{F} is related to the exponents m_i , $1 \leq i \leq \ell$, as follows

card
$$\mathcal{F} = \sum_{i=1}^{\ell} (m_i + 2) = \sum_{i=1}^{\ell} m_i + 2\ell.$$

Since $\sum_{i=1}^{\ell} m_i = \frac{1}{2} (\dim \mathfrak{g} - \ell)$ (see [5, Theorem 7.3.8]) and $\mathcal{F}_{|\mathcal{T}^2}$ is an independent family, we have card $\mathcal{F}_{|\mathcal{T}^2} = \frac{1}{2} (\dim \mathfrak{g} + 3\ell)$. The dimension of \mathcal{T}^2 is equal to $\dim \mathfrak{g} + 2\ell$. In conclusion, the relation below is satisfied:

$$\operatorname{card} \mathcal{F}_{|\mathcal{T}^2} = \dim \mathcal{T}^2 - \frac{1}{2}\operatorname{Rk}(\mathcal{T}^2, \{\cdot\,,\cdot\}_{\mathcal{R}})$$

if and only if $Rk(\mathcal{T}^2, \{\cdot, \cdot\}_{\mathcal{R}}) = \dim \mathfrak{g} + \ell$. We need therefore to prove this last result, which will be done in Proposition 19 below.

The rank of the restriction of $\{\cdot,\cdot\}_{\mathcal{R}}$ to \mathcal{T}^2 : The purpose of this part is to compute the rank of the Poisson manifold $(\mathcal{T}^2,\{\cdot,\cdot\}_{\mathcal{R}})$, i.e., the maximum of the rank at x of the Poisson \mathcal{R} -bracket for every $x \in \mathcal{T}^2$. We begin by establishing an isomorphism between the Poisson manifold $(\mathcal{T}^2,\{\cdot,\cdot\}_{\mathcal{R}})$ and a product Poisson manifold.

Before doing this, notice that it follows from Lemma 9 applied to a = 0 and $E = \mathfrak{g}_0 \oplus \mathfrak{g}_1$, $\mathfrak{g} = \mathfrak{g}$ and $[\cdot, \cdot] = [\cdot, \cdot]_R$ that (the subspace) $\mathfrak{g}_0 \oplus \mathfrak{g}_1$ is a Poisson submanifold of $(\mathfrak{g}, \{\cdot, \cdot\}_R)$, since the orthogonal of $\mathfrak{g}_0 \oplus \mathfrak{g}_1$ is a Lie ideal of $(\mathfrak{g}, [\cdot, \cdot]_R)$.

Proposition 18 The Poisson manifold $(\mathcal{T}^2, \{\cdot, \cdot\}_R)$ is isomorphic to the product Poisson manifold $(\mathfrak{g}^*, \{\cdot, \cdot\}) \times (\mathfrak{g}_0 \oplus \mathfrak{g}_1, \{\cdot, \cdot\}_R)$, where $\{\cdot, \cdot\}$ is the Lie-Poisson bracket on \mathfrak{g}^* , R is the difference of projections on \mathfrak{g}_+ and \mathfrak{g}_- , and $\{\cdot, \cdot\}_R$ is the Poisson R-bracket on \mathfrak{g} (restricted to $\mathfrak{g}_0 \oplus \mathfrak{g}_1$).

Proof. As we saw in the proof of Proposition 8, a := (e, e) and $E := \mathfrak{g}_{-}^2 \oplus \Delta(\mathfrak{g}_0 \oplus \mathfrak{g}_{-1})$ satisfy Conditions (1) and (2) of Lemma 9; hence $\mathcal{T}^2 = (e, e) + E$ and E are isomorphic Poisson submanifolds of $(\mathfrak{g}^2, \{\cdot, \cdot\}_{\mathcal{R}})$. According to Lemma 9 again, E is Poisson isomorphic to $(\mathfrak{g}^2/E^{\perp})^*$, where \mathfrak{g}^2/E^{\perp} is equipped with the quotient of the Lie bracket $[\cdot, \cdot]_{\mathcal{R}}$ with respect to the Lie ideal E^{\perp} . In conclusion, $(\mathcal{T}^2, \{\cdot, \cdot\}_{\mathcal{R}})$ is Poisson isomorphic to $((\mathfrak{g}^2/E^{\perp})^*, \{\cdot, \cdot\}')$, where $\{\cdot, \cdot\}'$ is the Lie Poisson bracket of $(\mathfrak{g}^2/E^{\perp})^*$. Let us determine the quotient Lie algebra \mathfrak{g}^2/E^{\perp} . The Lie algebra \mathfrak{g}^2 (endowed with the bracket $[\cdot, \cdot]_{\mathcal{R}}$) is isomorphic to the direct sum of the Lie algebras \mathfrak{g} , $\mathfrak{g}_{\leqslant -1}$ and $\mathfrak{g}_{\geqslant 0}$, the isomorphism being given, for every $x \in \mathfrak{g}, y_- \in \mathfrak{g}_-, y_+ \in \mathfrak{g}_+$, by

$$\mathfrak{g} \times \mathfrak{g}_{-} \times \mathfrak{g}_{+} \simeq \mathfrak{g}^{2}
(x, y_{-}, y_{+}) \rightarrow (x, x) + (y_{-}, y_{+}).$$
(56)

The Lie algebra isomorphism (56) identifies the subspace $E^{\perp} = \mathfrak{g}_{-} \times \mathfrak{g}_{>1} \subset \mathfrak{g}^{2}$ with $(0,\mathfrak{g}_{-},\mathfrak{g}_{\geq 2})$. Hence the quotient $\mathfrak{g}^{2}/E^{\perp}$ is the direct sum of the Lie algebra \mathfrak{g} (endowed with the usual bracket) and the quotient Lie algebra $\mathfrak{g}_{+}/\mathfrak{g}_{\geq 2}$.

We conclude that \mathcal{T}^2 is Poisson isomorphic to the dual of the direct sum Lie algebra $\mathfrak{g} \oplus (\mathfrak{g}_+/\mathfrak{g}_{\geqslant 2})$, endowed with the Lie-Poisson structure, and then is isomorphic to the product of the Poisson manifold \mathfrak{g}^* (endowed with the Lie-Poisson structure) with $(\mathfrak{g}_+/\mathfrak{g}_{\geqslant 2})^*$ (endowed with the Lie-Poisson structure). To complete the proof, it suffices to recall (see Lemma 9) that $\mathfrak{g}_0 \oplus \mathfrak{g}_1$ is a Poisson submanifold of $(\mathfrak{g}, \{\cdot, \cdot\}_R)$, isomorphic to the Lie-Poisson structure on the dual of the quotient Lie algebra $\mathfrak{g}/(\mathfrak{g}_0 \oplus \mathfrak{g}_1)^{\perp} = \mathfrak{g}_+/\mathfrak{g}_{\geqslant 2}$.

Proposition 19 The rank of the restriction of the Poisson \mathcal{R} -bracket $\{\cdot,\cdot\}_{\mathcal{R}}$ to the manifold \mathcal{T}^2 is dim $\mathfrak{g} + \ell$. As a consequence, the following relation is satisfied:

$$\operatorname{card} \mathcal{F}_{|\mathcal{T}^2} = \dim \mathcal{T}^2 - \frac{1}{2}\operatorname{Rk}(\mathcal{T}^2, \{\cdot\,,\cdot\}_{\mathcal{R}}).$$

Proof. According to Proposition 18 the Poisson submanifold $(\mathcal{T}^2, \{\cdot, \cdot\}_{\mathcal{R}})$ is isomorphic to the product manifold $(\mathfrak{g}^*, \{\cdot, \cdot\}) \times (\mathfrak{g}_0 \oplus \mathfrak{g}_1, \{\cdot, \cdot\}_R)$. This result proves that the restriction of the rank of Poisson \mathcal{R} -bracket $\{\cdot, \cdot\}_{\mathcal{R}}$ to the manifold \mathcal{T}^2 is the sum of the rank of the Lie-Poisson structure on \mathfrak{g}^* , which is dim $\mathfrak{g} - \ell$ (see [12, Proposition 29.3.2]), and the rank of the Poisson \mathcal{R} -bracket on $\mathfrak{g}_0 \oplus \mathfrak{g}_1$.

We calculate the latter rank and show that it is 2ℓ , which finishes the proof. Let $(z_1, \ldots, z_{2\ell})$ be the coordinate system on $\mathfrak{g}_0 \oplus \mathfrak{g}_1$, defined by

$$\begin{cases} z_i(x) = \langle h_i | x \rangle, \\ z_{i+\ell}(x) = \langle e_{-i} | x \rangle, \end{cases} \quad \forall i \in \{1 \dots, \ell\},$$

where as before, for $1 \leq i \leq \ell$, the element e_{-i} is a non-zero eigenvector associated with the root $-\alpha_i$. The Lie-Poisson brackets between these coordinate functions are given by the following formulae:

$$\begin{cases}
\{z_i, z_j\}_R = \{z_{i+\ell}, z_{j+\ell}\}_R = 0, \\
\{z_{i+\ell}, z_j\}_R = c_{ij} z_{i+\ell},
\end{cases}$$

$$\forall i, j \in \{1, \dots, \ell\},$$

where $C := (c_{ij})_{1 \leqslant i,j \leqslant \ell}$ is the Cartan matrix of \mathfrak{g} . Then the Poisson matrix $M = (\{z_i, z_j\}_R)_{1 \leqslant i,j \leqslant 2\ell}$ is equal to

$$M = \left(\begin{array}{cc} 0 & -A^T \\ A & 0 \end{array}\right),$$

where A := vC and $v := diag(z_{1+\ell}, \dots, z_{2\ell})$. Since C is invertible, the rank of A at any point for which $(z_{1+\ell}, \dots, z_{2\ell}) = (1, \dots, 1)$ is ℓ ; therefore the rank of M is 2ℓ , which implies that the rank of the Poisson R-structure on $\mathfrak{g}_0 \oplus \mathfrak{g}_1$ is 2ℓ .

3.4 The integrability for the quadratic Poisson \mathcal{R} -bracket

In this subsection we study the integrability of the 2-Toda lattice on some Lie algebra \mathfrak{g} with respect to a quadratic Poisson \mathcal{R} -bracket. To construct a quadratic Poisson \mathcal{R} -bracket on \mathfrak{g}^2 it is necessary that \mathfrak{g} is an associative algebra of finite dimensional and \mathcal{R} and its antisymmetric part \mathcal{R}_- are solutions of (mCYBE). For this it suffices to choose $\mathfrak{g} = \mathfrak{gl}_n(\mathbf{C})$ and $\mathcal{R}(x,y) = (R(x-y)+y,R(x-y)+x)$, where where $R = P_+ - P_-$, P_+ (resp. P_-) being the projection of $\mathfrak{gl}_n(\mathbf{C})$ on the subalgebra of upper (resp. strictly lower) triangular matrices (we will show again later that \mathcal{R}_- is a solution of (mCYBE) of \mathfrak{g}^2).

In this subsection we show the integrability of a system of equations which is exactly the system (1), up to the fact that we do no longer assume the matrices L, M to be traceless. We denote by $\mathcal{T}^{2'}$ the phase space of the 2-Toda lattice, i.e.,

$$(L,M) = \begin{pmatrix} a_{11} & 1 & 0 \\ a_{21} & a_{22} & \ddots \\ \vdots & & \ddots & 1 \\ a_{n1} & \cdots & a_{n,n-1} & a_{nn} \end{pmatrix}, \begin{pmatrix} b_{11} & \cdots & \cdots & b_{1n} \\ b_{21} & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & & b_{n,n-1} & b_{nn} \end{pmatrix} \}$$

and we name as the 2-Toda lattice on $\mathfrak{gl}_n(\mathbf{C})$ the system of differential equations (1) with the constraint $(L, M) \in \mathcal{T}^{2'}$.

Before studying the integrability of the 2-Toda lattice with respect to the quadratic Poisson \mathcal{R} -bracket we will study the integrability of the latter system with respect to the linear Poisson \mathcal{R} -bracket.

3.4.1 The Liouville integrability of the 2-Toda lattice on $\mathfrak{gl}_n(\mathbf{C})$ for the linear Poisson $\mathcal{R}\text{-bracket}$

We equip $\mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C})$ with the Ad-invariant, non-degenerate, symmetric, bilinear form $\langle \cdot | \cdot \rangle_2$, defined for every $(x, y), (x', y') \in \mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C})$ by

$$\langle (x,y) | (x',y') \rangle_2 := \langle x | x' \rangle - \langle y | y' \rangle = \operatorname{Trace}(xx') - \operatorname{Trace}(yy').$$
 (57)

As in Subsection 3.2 we consider $\mathcal{R}(x,y) = (R(x-y) + y, R(x-y) + x)$ and we consider the linear Poisson \mathcal{R} -bracket⁵, defined for every $F, G \in \mathcal{F}(\mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C}))$ at $(x,y) \in \mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C})$, by

$$\{F,G\}_{\mathcal{R}}(x,y) = \frac{1}{2}(\left\langle (x,y) \,|\, [\mathcal{R}\nabla_{(x,y)}F,\nabla_{(x,y)}G]\right\rangle_2 + \left\langle (x,y) \,|\, [\nabla_{(x,y)}F,\mathcal{R}\nabla_{(x,y)}G]\right\rangle_2).$$

⁵According to Corollary 2, the endomorphism \mathcal{R} is an \mathcal{R} -matrix on $\mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C})$.

By using the proofs of Propositions 8 and 10 we show that the phase space $\mathcal{T}^{2'}$ is a Poisson submanifold of $(\mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C}), \{\cdot, \cdot\}_{\mathcal{R}})$ and $\mathcal{X}_H := \{\cdot, H\}_{\mathcal{R}}$ and $\mathcal{X}_{H'} := \{\cdot, H'\}_{\mathcal{R}}$ where $H(x, y) = \frac{1}{2} \operatorname{Trace} x^2$ and $H'(x, y) = \frac{1}{2} \operatorname{Trace} y^2$ describes on $\mathcal{T}^{2'}$ the equations of motion of the 2-Toda lattice.

Let us now study the integrability of the 2-Toda lattice on $\mathfrak{gl}_n(\mathbf{C})$. Let P_i , for every $i \in \mathbf{N}$ be the Ad-invariant function of $\mathfrak{gl}_n(\mathbf{C})$ defined for all $x \in \mathfrak{gl}_n(\mathbf{C})$ by $P_i(x) = \frac{1}{i+1}\operatorname{Trace}(x^{i+1})$. We define $F_{j,i} \in \mathcal{F}(\mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C}))$ by $P_i(\lambda x - y) = \sum_{j=0}^{i+1} (-1)^{m_i+1-j} \lambda^j F_{j,i}(x,y)$ and we define

$$\mathcal{F}' = (F_{j,i}, 0 \leqslant i \leqslant n-1 \text{ and } 0 \leqslant j \leqslant i+1).$$

Note that the functions that make up \mathcal{F}' are the functions that make up \mathcal{F} together with the functions $F_{1,0}(x,y) := \operatorname{Trace}(x)$ and $F_{0,0}(x,y) := \operatorname{Trace}(y)$. We have the following proposition.

Proposition 20 The triplet $(\mathcal{T}^{2'}, \mathcal{F}'_{|_{\mathcal{T}^{2'}}}, \{\cdot, \cdot\}_{\mathcal{R}})$ is an integrable system.

Proof. According to the item (2) of Theorem 4 the family \mathcal{F}' is involutive for the Poisson bracket $\{\cdot,\cdot\}_{\mathcal{R}}$. Since \mathcal{F}' is an independent family on $\mathcal{T}^{2'}$ (this follows from the independence of the differentials $dF_{0,0}$ and $dF_{1,0}$ at all points of $\mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C})$ and the independence of \mathcal{F} on the submanifold \mathcal{T}^2 , which is the submanifold of $\mathcal{T}^{2'}$ defined by $F_{1,0} = F_{0,0} = 0$), and since the cardinal of \mathcal{F}' is $\frac{n(n+3)}{2}$ we have, according to [2, Proposition 4.12], the inequality

$$Rk(\mathcal{T}^{2'}, \{\cdot, \cdot\}_{\mathcal{R}}) \leqslant 2(\dim(\mathcal{T}^{2'}) - Card(\mathcal{F}')) = n^2 + n - 2.$$
(58)

Furthermore $\operatorname{Rk}(\mathcal{T}^{2'}, \{\cdot\,,\cdot\}_{\mathcal{R}}) \geqslant \operatorname{Rk}(\mathcal{T}^2, \{\cdot\,,\cdot\}_{\mathcal{R}}) = n^2 + n - 2$. This implies that the rank of the restriction to $\mathcal{T}^{2'}$ of $\{\cdot\,,\cdot\}_{\mathcal{R}}^Q$ is exactly $n^2 + n - 2$. Then $\operatorname{card} \mathcal{F'}_{|\mathcal{T}^{2'}} = \dim \mathcal{T}^{2'} - \frac{1}{2}\operatorname{Rk}(\mathcal{T}^{2'}, \{\cdot\,,\cdot\}_{\mathcal{R}}^Q)$. This completes the proof.

3.4.2 The integrability of the 2-Toda lattice on $\mathfrak{gl}_n(\mathbf{C})$ for the quadratic Poisson $\mathcal{R}\text{-bracket}$

Let the quadratic \mathcal{R} -bracket be defined for every $F, G \in \mathcal{F}(\mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C}))$ at $(x, y) \in \mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C})$, by

$$\{F,G\}_{\mathcal{R}}^{Q}(x,y) := \frac{1}{2} \langle [(x,y), \nabla_{(x,y)}F] | \mathcal{R}((x,y)\nabla_{(x,y)}G + \nabla_{(x,y)}G(x,y)) \rangle_{2}$$

$$-(F \leftrightarrow G).$$

$$(59)$$

Since $\mathcal{R}_{-}(x,y) = \frac{1}{2}(\mathcal{R} - \mathcal{R}^*)(x,y) = (R_{-}(x-y) + y, R_{-}(x-y) + x)$ and since both the endomorphism $R = P_{+} - P_{-}$ and its antisymmetric part $R_{-} = \frac{1}{2}(R - R^*)$ are solutions of (mCYBE) of $\mathfrak{gl}_{n}(\mathbf{C})$ of constant c = 1, it follows from Corollary 2 that \mathcal{R} and \mathcal{R}_{-} are solutions of (mCYBE) of $\mathfrak{gl}_{n}(\mathbf{C}) \times \mathfrak{gl}_{n}(\mathbf{C})$ of constant c = 1. According to [8, Section 4] the

 \mathcal{R} -bracket (59) is indeed a Poisson bracket, that we call the quadratic Poisson \mathcal{R} -bracket (on $\mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C})$).

By a direct computation in coordinates we can prove the following proposition.

Proposition 21 The phase space $\mathcal{T}^{2'}$ of the 2-Toda lattice on $\mathfrak{gl}_n(\mathbf{C})$ is a Poisson submanifold of $(\mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C}), \{\cdot, \cdot\}_{\mathcal{R}}^Q)$.

We show the following proposition.

Proposition 22 Let P_i , for every $i \in \mathbf{N}$ be the Ad-invariant functions of $\mathfrak{gl}_n(\mathbf{C})$ defined for all $x \in \mathfrak{gl}_n(\mathbf{C})$ by $P_i(x) = \frac{1}{i+1}\operatorname{Trace}(x^{i+1})$ and let $\phi_{\lambda} : \mathfrak{gl}_n(\mathbf{C}) \times \mathfrak{gl}_n(\mathbf{C}) \to \mathfrak{gl}_n(\mathbf{C})$, for every $\lambda \in \mathbf{F}$, be defined by $\phi_{\lambda} : (x,y) \to \lambda x - y$.

- (1) For every $i, j \in \mathbf{N}$ and every $\lambda, \gamma \in \mathbf{C}$, the functions $P_i \circ \phi_{\lambda}$ and $P_j \circ \phi_{\gamma}$ are in involution for $\{\cdot, \cdot\}_{\mathcal{R}}^Q$.
- (2) The Hamiltonian vector field $\mathcal{X}_{P_i \circ \phi_{\lambda}}^Q := \{\cdot, P_i \circ \phi_{\lambda}\}_{\mathcal{R}}^Q$ is given by

$$\mathcal{X}_{P_i \circ \phi_{\lambda}}^{Q} = -[(x, y), ((R - I)(\lambda x - y)^{i+1}, (R + I)(\lambda x - y)^{i+1})]. \tag{60}$$

Proof. (1) According to (59) and (22), for every $x, y \in \mathfrak{g}$,

$$\begin{aligned}
&\{P_i \circ \phi_{\lambda}, P_j \circ \phi_{\mu}\}_{\mathcal{R}}^{Q}(x, y) \\
&= \frac{1}{2} \langle [(x, y), (\lambda(\lambda x - y)^i, (\lambda x - y)^i)] | \mathcal{R}(\mu x (\mu x - y)^j, y (\mu x - y)^j) \rangle_2 \\
&+ \frac{1}{2} \langle [(x, y), (\lambda(\lambda x - y)^i, (\lambda x - y)^i)] | \mathcal{R}(\mu (\mu x - y)^j x, (\mu x - y)^j y) \rangle_2 \\
&- (j, \mu) \longleftrightarrow (i, \lambda).
\end{aligned}$$

By replacing \mathcal{R} by its expression $\mathcal{R}(x,y) = (R(x-y) + y, R(x-y) + x)$, we obtain

$$\begin{aligned} &\{P_{i} \circ \phi_{\lambda}, P_{j} \circ \phi_{\mu}\}_{\mathcal{R}}^{Q}(x, y) \\ &= \left\langle \left[(x, y), (\lambda(\lambda x - y)^{i}, (\lambda x - y)^{i}) \right] \middle| \left(R((\mu x - y)^{j+1}), R((\mu x - y)^{j+1}) \right) \right\rangle_{2} \\ &+ \frac{1}{2} \left\langle \left[(x, y), (\lambda(\lambda x - y)^{i}, (\lambda x - \mu)^{i}) \right] \middle| \left(y(\mu x - y)^{j}, \mu x(\mu x - y)^{j} \right) \right\rangle_{2} \\ &+ \frac{1}{2} \left\langle \left[(x, y), (\lambda(\lambda x - y)^{i}, (\lambda x - \mu)^{i}) \right] \middle| \left((\mu x - y)^{j} y, \mu(\mu x - y)^{j} x \right) \right\rangle_{2} \\ &- (j, \mu) \longleftrightarrow (i, \lambda). \end{aligned}$$

By using Formula (57) we obtain

$$\begin{aligned} \{P_i \circ \phi_{\lambda}, P_j \circ \phi_{\mu}\}_{\mathcal{R}}^Q(x, y) &= \left\langle \left[\lambda x - y, (\lambda x - y)^i\right] \mid R((\mu x - y)^{j+1}) \right\rangle \\ &+ \frac{1}{2} \left\langle \left[\lambda x, (\lambda x - y)^i\right] \mid y(\mu x - y)^j + (\mu x - y)^j y \right\rangle \\ &- \frac{1}{2} \left\langle \left[y, (\lambda x - y)^i\right] \mid \mu x(\mu x - y)^j + \mu(\mu x - y)^j x \right) \right\rangle \\ &- (j, \mu) \longleftrightarrow (i, \lambda). \end{aligned}$$

Since $[\lambda x, (\lambda x - y)^i] = [y, (\lambda x - y)^i]$

$$\begin{aligned}
\{P_i \circ \phi_{\lambda}, P_j \circ \phi_{\mu}\}_{\mathcal{R}}^{Q}(x, y) &= -\frac{1}{2} \left\langle [y, (\lambda x - y)^i] \,|\, 2(\mu x - y)^{j+1} \right\rangle - (j, \mu) \longleftrightarrow (i, \lambda) \\
&= \left\langle y \,|\, [(\lambda x - y)^i, (\mu x - y)^{j+1}] \right\rangle - (j, \mu) \longleftrightarrow (i, \lambda) \\
&= 0,
\end{aligned}$$

where we used Lemma 6 to provide the last line.

(2) Let K be a function of \mathfrak{g}^2 ; we denote by $(a,b) = \nabla_{(x,y)}K$, according to (59) and (22),

$$\mathcal{X}_{P_{i}\circ\phi_{\lambda}}^{Q}(x,y)[K] = \frac{1}{2} \left\langle \left[(x,y), (a,b) \right] \middle| \mathcal{R}(\lambda x (\lambda x - y)^{i} + \lambda (\lambda x - y)^{i} x, y (\lambda x - y)^{i} + (\lambda x - y)^{i} y) \right\rangle \\
- \frac{1}{2} \left\langle \left[(x,y), (\lambda (\lambda x - y)^{i}, (\lambda x - y)^{i}) \right] \middle| \mathcal{R}(xa + ax, yb + by) \right\rangle.$$
(61)

We introduce the shorthand B = xa + ax - yb - by. By using the expression for \mathcal{R} the Hamiltonian vector field (61) becomes

$$\begin{split} \mathcal{X}^Q_{P_i \circ \phi_{\lambda}}(x,y)[K] &= \left\langle \left[(x,y), (a,b) \right] | \left(R((\lambda x-y)^{i+1}), R((\lambda x-y)^{i+1}) \right) \right\rangle_2 \\ &+ \frac{1}{2} \left\langle \left[(x,y), (a,b) \right] | \left(y(\lambda x-y)^i + (\lambda x-y)^i y, \lambda x(\lambda x-y)^i + \lambda (\lambda x-y)^i x \right) \right\rangle_2 \\ &- \frac{1}{2} \left\langle \left[(x,y), (\lambda (\lambda x-y)^i, (\lambda x-y)^i) \right] | \left(R(B), R(B) \right) \right\rangle_2 \\ &- \frac{1}{2} \left\langle \left[(x,y), (\lambda ((x-y)^i, (\lambda x-y)^i) \right] | \left(yb + by, xa + ax \right) \right\rangle_2 \\ &= &- \left\langle \left[(x,y), (R((\lambda x-y)^{i+1}), R((\lambda x-y)^{i+1})) \right] | \left(a,b \right) \right\rangle_2 \\ &- \frac{1}{2} \left\langle \left[(x,y), (y(\lambda x-y)^i + (\lambda x-y)^i y, \lambda x(\lambda x-y)^i + \lambda (\lambda x-y)^i x \right) \right] | \left(a,b \right) \right\rangle_2 \\ &- \frac{1}{2} \left\langle \left[(x,y), (\lambda x-y)^i \right] | R(B) \right\rangle \\ &- \frac{1}{2} \left\langle \left[(x,x), (\lambda x-y)^i \right] | yb + by \right\rangle + \frac{1}{2} \left\langle \left[(x,x), (\lambda x-y)^i \right] | xa + ax \right\rangle, \end{split}$$

Moreover according to the Ad-invariance of P_i and the property $\langle x \,|\, yz \rangle = \langle xy \,|\, z \rangle$, $\forall x,y,z \in \mathfrak{g}$, we obtain

$$\begin{split} \mathcal{X}^Q_{P_i \circ \phi_\lambda}(x,y)[K] &= - \big\langle [(x,y), (R((\lambda x-y)^{i+1}), R((\lambda x-y)^{i+1}))] \, | \, (a,b) \big\rangle_2 \\ &- \frac{1}{2} \big\langle [(x,y), (y(\lambda x-y)^i + (\lambda x-y)^i y, \lambda x (\lambda x-y)^i + \lambda (\lambda x-y)^i x)] \, | \, (a,b) \big\rangle_2 \\ &- \frac{1}{2} \left\langle y[x, \lambda (\lambda x-y)^i] + [x, \lambda (\lambda x-y)^i] y \, | \, b \right\rangle \\ &+ \frac{1}{2} \left\langle x[y, (\lambda x-y)^i] + [y, (\lambda x-y)^i] x \, | \, a \right\rangle. \end{split}$$

We then deduce that

$$\mathcal{X}_{P_{i} \circ \phi_{\lambda}}^{Q}(x,y) = -[(x,y), (R((\lambda x - y)^{i+1}), R((\lambda x - y)^{i+1}))]$$

$$-\frac{1}{2}[(x,y), (y(\lambda x - y)^{i} + (\lambda x - y)^{i}y, \lambda x(\lambda x - y)^{i} + \lambda(\lambda x - y)^{i}x)]$$

$$+\frac{1}{2}(x[y, (\lambda x - y)^{i}] + [y, (\lambda x - y)^{i}]x, y[x, \lambda(\lambda x - y)^{i}] + [x, \lambda(\lambda x - y)^{i}]y).$$

Let us compute separately the two last lines:

$$\begin{split} &-\frac{1}{2}[(x,y),(y(\lambda x-y)^{i}+(\lambda x-y)^{i}y,\lambda x(\lambda x-y)^{i}+\lambda(\lambda x-y)^{i}x)]\\ &+\frac{1}{2}(x[y,(\lambda x-y)^{i}]+[y,(\lambda x-y)^{i}]x,y[x,\lambda(\lambda x-y)^{i}]+[x,\lambda(\lambda x-y)^{i}]y)\\ &=-(x(\lambda x-y)^{i}y-y(\lambda x-y)^{i}x,\lambda y(\lambda x-y)^{i}x-\lambda x(\lambda x-y)^{i}y)\\ &=([x,(\lambda x-y)^{i+1}],-\lambda[x,(\lambda x-y)^{i+1}])\\ &=[(x,y),((\lambda x-y)^{i+1},-(\lambda x-y)^{i+1})]. \end{split}$$

Then we have Formula (60).

Let us compare the Hamiltonian vector fields for the quadratic \mathcal{R} -Poisson bracket and the linear \mathcal{R} -Poisson bracket. Formula (20) gives, in our case, the following expression for the Hamiltonian vector field of $P_{i+1} \circ \phi_{\lambda}$ with respect to the linear \mathcal{R} -Poisson structure:

$$\mathcal{X}_{P_{i+1} \circ \phi_{\lambda}} = \frac{1}{2} (\lambda - 1) [(x, y), ((R - I)(\lambda x - y)^{i+1}, (R + I)(\lambda x - y)^{i+1})]. \tag{62}$$

Comparing Formulae (62) and (60), we obtain, for $\lambda \neq 1$,

$$\mathcal{X}_{P_i \circ \phi_{\lambda}}^Q(x, y) = \frac{2}{1 - \lambda} \mathcal{X}_{P_{i+1} \circ \phi_{\lambda}}(x, y). \tag{63}$$

The relation (63) implies that for every $0 \le i \le n-1$, we have

$$\begin{cases}
\mathcal{X}_{F_{0,i}}^{Q} = 2\mathcal{X}_{F_{0,i+1}}, \\
\mathcal{X}_{F_{j,i}}^{Q} - \mathcal{X}_{F_{j-1,i}}^{Q} = 2\mathcal{X}_{F_{j,i+1}}, & 1 \leq j \leq i+1, \\
\mathcal{X}_{F_{i+1,i}}^{Q} = -2\mathcal{X}_{F_{i+2,i+1}}.
\end{cases} (64)$$

Notice also that the Hamiltonian vector fields $\mathcal{X}_{F_{0,0}}^{Q}$ and $\mathcal{X}_{F_{1,0}}$ are precisely the equations of the 2-Toda lattice on $\mathfrak{gl}_{n}(\mathbf{C})$ as can be shown by specializing (64) to i=0 and the first item of Remark 12.

Theorem 23 The triplet $(\mathcal{T}^{2'}, \mathcal{F}'_{|_{\mathcal{T}^{2'}}}, \{\cdot, \cdot\}_{\mathcal{R}}^{Q})$ is an integrable system.

Proof. The involutivity of the family \mathcal{F}' on $(\mathfrak{g}^2, \{\cdot, \cdot\}_{\mathcal{R}})$ follows from item (1) in Proposition 22. Since \mathcal{F}' is an independent family on $\mathcal{T}^{2'}$ (see the proof of Proposition 20), and since the cardinal of \mathcal{F}' is $\frac{n(n+3)}{2}$, we have, according to [2, Proposition 4.12], the inequality

$$Rk(\mathcal{T}^{2'}, \{\cdot, \cdot\}_{\mathcal{R}}^{Q}) \leqslant 2(\dim(\mathcal{T}^{2'}) - Card(\mathcal{F}')) = n^{2} + n - 2.$$

$$(65)$$

Moreover, according to Formula (64), the family of vector fields $\mathcal{X}_{\mathcal{F}'} := (\mathcal{X}_{F_{j,i}}^Q, 0 \leqslant i \leqslant n-1 \text{ and } 0 \leqslant j \leqslant i)$ and the family of vector fields $\mathcal{X}_{\mathcal{F}} := (\mathcal{X}_{F_{j,i}}, 1 \leqslant i \leqslant n-1 \text{ and } 1 \leqslant j \leqslant i+1)$ have the same rank at all points. By choosing a point in \mathcal{T}^2 , we can deduce from the fact that \mathcal{F} is an integrable system on \mathcal{T}^2 (Theorem 13) that this rank is at least the cardinal of $\mathcal{F} := \frac{n(n+3)}{2} - 2$) minus the number of independent Casimir functions on \mathcal{T} for the linear bracket $\{\cdot,\cdot\}_{\mathcal{R}} := (n-1)$. Since \mathcal{F}' is involutive, we have therefore the inequality

$$\operatorname{Rk}\left\{\cdot,\cdot\right\}_{\mathcal{R}}^{Q} \geqslant 2(\operatorname{Rk}(\mathcal{X}_{\mathcal{F}'})) = 2(\operatorname{Rk}(\mathcal{X}_{\mathcal{F}})) \geqslant 2(\frac{n(n+3)}{2} - 2 - n + 1) = n^2 + n - 2.$$

Together with (65), this implies that the rank of the restriction to $\mathcal{T}^{2'}$ of $\{\cdot,\cdot\}_{\mathcal{R}}^{Q}$ is exactly n^2+n-2 . The identity card $\mathcal{F'}_{|\mathcal{T}^{2'}}=\dim \mathcal{T}^{2'}-\frac{1}{2}\operatorname{Rk}(\mathcal{T}^{2'},\{\cdot,\cdot\}_{\mathcal{R}}^{Q})$ follows and completes the proof.

3.5 The relation between the 2-Toda lattice and the Toda lattice

In this section we show that the Toda lattice is a restriction of the 2-Toda lattice. We begin by recalling the Liouville integrable system of the Toda lattice.

3.5.1 The Toda lattice

We give some definitions and properties of the *Toda lattice*, which will be useful afterwards in this section.

Definition 24 (1) The phase space \mathcal{T} of the Toda lattice is the affine subspace of \mathfrak{g} given by

$$\mathcal{T} := \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 + e, \tag{66}$$

where $e = \sum_{i=1}^{\ell} e_i$.

(2) The Toda lattice is the system of differential equations on \mathcal{T} given by the Lax equation

$$\dot{A} = [A_+, A], \tag{67}$$

where A_+ is the projection of A on \mathfrak{g}_+ .

We consider the endomorphism $R := P_+ - P_-$ of \mathfrak{g} , the difference of the projections on \mathfrak{g}_+ and \mathfrak{g}_- . The Poisson R-bracket on \mathfrak{g} is defined, for every $F, G \in \mathcal{F}(\mathfrak{g})$ and every $x \in \mathfrak{g}$, by

$$\{F,G\}_R := \frac{1}{2} \langle x \mid [R\nabla_x F, \nabla_x G] + [\nabla_x F, R\nabla_x G] \rangle. \tag{68}$$

Theorem 25 [9, Section 4.1] (1) The affine subspace \mathcal{T} of \mathfrak{g} is a Poisson submanifold of $(\mathfrak{g}, \{\cdot, \cdot\}_R)$.

- (2) Let $H \in \mathcal{F}(\mathfrak{g})$, defined for every $x \in \mathfrak{g}$ by $H(x) = \frac{1}{2} \langle x | x \rangle$. The equation of the Hamiltonian field $\mathcal{X}_H := \{\cdot, H\}_R$ is the equation of motion (67) of the Toda lattice.
- (3) Let P_1, \ldots, P_ℓ be a generating family of homogeneous polynomials of the algebra of Ad-invariant functions on \mathfrak{g} of degree respectively, $m_1 + 1, \ldots, m_\ell + 1$. We define $\mathcal{F}_0 := (P_1, \ldots, P_\ell)$. The triplet $(\mathcal{F}_0, \{\cdot, \cdot\}_R, \mathcal{T})$ is Liouville integrable system and the equation of motion of the Toda lattice is

$$\dot{A} := \{\cdot, P_1\}_R(A) = [(\nabla_A P_1)_+, A]. \tag{69}$$

3.5.2 Restriction of the 2-Toda lattice and construction of the Toda lattice

The phase space \mathcal{T}^2 of the 2-Toda lattice decomposes as

$$\mathcal{T}^2 := \mathfrak{g}_{\leqslant -1} \times \mathfrak{g}_{\geqslant 0} \oplus \Delta(\mathfrak{g}_{-1} \oplus \mathfrak{g}_0) + (e, e), \tag{70}$$

where $\Delta(\mathfrak{g}_{-1} \oplus \mathfrak{g}_0) := \{(x,x) \mid x \in \mathfrak{g}_{-1} \oplus \mathfrak{g}_0\}.$

Theorem 26 Let \mathfrak{g}^2 be the Lie algebra $\mathfrak{g} \times \mathfrak{g}$ with $\{\cdot,\cdot\}_{\mathcal{R}}$ the \mathcal{R} -Poisson bracket, \mathcal{T}^2 the phase space of the 2-Toda lattice and $\mathcal{T}' := \Delta(\mathfrak{g}_{-1} \oplus \mathfrak{g}_0) + (e,e)$.

- (1) The submanifold \mathcal{T}' is a Poisson submanifold of $(\mathfrak{g}^2, \{\cdot\,,\cdot\}_{\mathcal{R}})$.
- (2) Let $(\mathcal{T}, \{\cdot, \cdot\}_R)$ be the phase space of the Toda lattice, equipped with the Poisson R-bracket given as in (68). The map

$$\varphi : (\mathcal{T}, \{\cdot, \cdot\}_R) \to (\mathcal{T}', \{\cdot, \cdot\}_R)
 x \mapsto (x, x)$$
(71)

is an isomorphism of integrable systems.

Proof. (1) It is clear that $\mathcal{T}' = \Delta(\mathfrak{g}_{-1} \oplus \mathfrak{g}_0 + e) = \mathcal{T}^2 \cap \mathfrak{g}_+^2$. According to Proposition 8, \mathcal{T}^2 is a Poisson submanifold of $(\mathfrak{g}^2, \{\cdot, \cdot\}_{\mathcal{R}})$ while \mathfrak{g}_+^2 is a Poisson submanifold of $(\mathfrak{g}^2, \{\cdot, \cdot\}_{\mathcal{R}})$ because the orthogonal of \mathfrak{g}^2 , which is \mathfrak{g}_+^2 , is a Lie ideal for the bracket $[\cdot, \cdot]_{\mathcal{R}}$; their intersection \mathcal{T}' is then a Poison submanifold of $(\mathfrak{g}^2, \{\cdot, \cdot\}_{\mathcal{R}})$.

(2) By using the coordinate functions on \mathcal{T} and on \mathcal{T}' we show that φ is a Poisson isomorphism. Furthermore, the functions of the integrable system $\mathcal{F} = (F_{k,i}, 1 \leq i \leq \ell \text{ and } 0 \leq \ell$

 $k \leq m_i + 1$), restricted to \mathcal{T}' , and pulled back on \mathcal{T} by φ , give again the functions of the family \mathcal{F}_0 of the Toda lattice. Specifically, for every i in $1, \ldots, \ell$ and every $0 \leq k \leq m_i + 1$, the functions $F_{k,i}$ are all equal (up to multiplicative constants) to the function P_i . In fact, for every $x \in \mathfrak{g}_0 \oplus \mathfrak{g}_{-1}$,

$$P_i \circ \phi_{\lambda}(\varphi(x)) = P_i(\phi_{\lambda}(x, -x))$$

$$= P_i(\lambda x - x)$$

$$= (\lambda - 1)^{m_i + 1} P_i(x),$$

and so

$$\sum_{k=0}^{m_i+1} (-1)^{m_i+1-k} \lambda^k F_{k,i}(\varphi(x)) = \sum_{k=0}^{m_i+1} (-1)^{m_i+1-k} \lambda^k C_{m_i+1}^k P_i(x)$$

and $F_{k,i}(\varphi(x)) = C_{m_i+1}^k P_i(x)$. In conclusion, $\varphi : \mathcal{T} \to \mathcal{T}'$ is Poisson isomorphic and $\varphi^* \mathcal{F} = \mathcal{F}_0$ (see item (3) of Theorem 25 for the definition of \mathcal{F}_0). This proves the claim. \square

References

- [1] Mark Adler and Pierre van Moerbeke. String-orthogonal polynomials, string equations, and 2-Toda symmetries. *Comm. Pure Appl. Math.*, 50(3):241–290, 1997.
- [2] Mark Adler, Pierre van Moerbeke, and Pol Vanhaecke. Algebraic integrability, Painlevé geometry and Lie algebras, volume 47 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, 2004.
- [3] Guido Carlet. The Hamiltonian structures of the two-dimensional Toda lattice and R-matrices. Lett. Math. Phys., 71(3):209–226, 2005.
- [4] Dennis DeTurck, Hubert Goldschmidt, and Janet Talvacchia. Connections with prescribed curvature and Yang-Mills currents: the semi-simple case. *Ann. Sci. École Norm. Sup.* (4), 24(1):57–112, 1991.
- [5] Jacques Dixmier. Enveloping algebras, volume 11 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 1996. Revised reprint of the 1977 translation.
- [6] Bertram Kostant. Lie group representations on polynomial rings. Bull. Amer. Math. Soc., 69:518–526, 1963.
- [7] Bertram Kostant. The solution to a generalized Toda lattice and representation theory. *Adv. in Math.*, 34(3):195–338, 1979.

- [8] Luen Chau Li and Serge Parmentier. Nonlinear Poisson structures and r-matrices. Comm. Math. Phys., 125(4):545–563, 1989.
- [9] A. M. Perelomov. *Integrable systems of classical mechanics and Lie algebras. Vol. I.* Birkhäuser Verlag, Basel, 1990. Translated from the Russian by A. G. Reyman.
- [10] Mustapha Rais. Sur les données des polynômes invariants sur une algèbre de Lie semi-simple complexe (lettre à Hubert Goldschmidt,1988).
- [11] Michael A. Semenov-Tian-Shansky. Dressing transformations and Poisson group actions. *Publ. Res. Inst. Math. Sci.*, 21(6):1237–1260, 1985.
- [12] Patrice Tauvel and Rupert W. T. Yu. *Lie algebras and algebraic groups*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2005.
- [13] Kimio Ueno and Kanehisa Takasaki. Toda lattice hierarchy. II. *Proc. Japan Acad. Ser. A Math. Sci.*, 59(6):215–218, 1983.