WEAK μ -EQUALITY IS DECIDABLE

WIL DEKKERS, RADBOUD UNIVERSITY, NIJMEGEN, THE NETHERLANDS

Abstract

In this paper we consider the set \mathbb{T}_{μ} of μ -types, an extension of the set \mathbb{T} of simple types freely generated from a set A of atomic types and the type constructor \rightarrow , by a new operator μ , to explicitly denote solutions of recursive equations like $A =_{\mu} A \rightarrow \beta$. We show that this so-called weak μ -equality for μ -types is decidable by defining a derivation system for weak μ -equality based on standard reduction for μ -types such that the number of nodes in a derivation tree for $A =_{\mu} B$ is bounded as a function of A, B. We give two proofs. One for decidability of $=$ _μ for α -equivalence classes of μ -types and one for decidability of $=_{\mu/\alpha}$ for μ -types themselves. Both proofs are straightfroward and elementary.

Introduction

In Cardone and Coppo[1] a proof method is given to show decidability of weak μ -equality, using standard reduction and a special purpose proof system. It turned out recently that their treatment of α -conversion was not completely correct. Nethertheless their result is correct and in Endrullis, Grabmayer, Klop and van Oostrom[2] three proofs for the decidability are given. Two of the given proofs are inspired by the Cardone-Coppo proof strategy. The third proof uses the theory of regular languages and is totally different. This paper is based on the first two proofs of Endrullis et al. These proofs are ingenuous but they are also technical and intricate. Our aim is to give simpler and more straightforward proofs.

In their first proof Endrullis et al. work with μ -types themselves, not with α -equivalence classes of μ -types. The proof constructs a decision procedure for equality of types based on standard reduction for types. In that proof they need the theory of α -avoiding developed by van Oostrom^[3]. In the second proof they work with α -equivalence classes and the corresponding reduction relation \rightarrow_{μ} for classes and they show the decidability of $=_{\mu}$. Our proof is more or less the other way round. Based on the first proof of Endrullis et al. of decidabiliy of equality for types we give a proof of decidability of $=$ for classes of types and then from that proof we derive a first order proof of decidability of equality for types where the theory of α -avoiding is not needed. Both proofs are straightforward and elementary.

So in our first proof \mathbb{T}_{μ} is he set of α -equivalence classes of types and A, B stand for classes of types. We start by defining a set $\mathbb{T}_{\mu'}$, containing \mathbb{T}_{μ} , of so-called annotated (classes of) μ -types and a reduction $\rightarrow_{\mu'} : \mathbb{T}_{\mu'} \rightarrow \mathbb{T}_{\mu'}$ that is closely related with standard reduction on \mathbb{T}_{μ} . Then we define a derivation

system $\vdash_{\mu'}$ for equality of elements of $\mathbb{T}_{\mu'}$ such that for $A, B \in \mathbb{T}_{\mu}$ one has

 $\vdash_{\mu'} A = B \Leftrightarrow A$ and B have a common standard reduct.

We show for $a \in \mathbb{T}_{\mu'}$ that $SC(a) \equiv \{b \in \mathbb{T}_{\mu'} | a \to_{\mu'}^* b\}$ is bounded as a function in a and also that for $a = b$ occurring in a derivation tree for $\vdash_{\mu'} A = B$ we have $a \in SC(A)$ and $b \in SC(B)$. So the number of nodes occurring in a derivation tree for $\vdash_{\mu'} A = B$ is bounded as a function in A, B. So $\vdash_{\mu'} A = B$ is decidable and hence $A =_{\mu} B$ is decidable, because of the fact that

 $A =_{\mu} B \Leftrightarrow A$ and B have a common standard reduct.

Then in part 2 from this result we derive easily that $=_{\mu/\alpha}$ on types themselves can be decided by first order means via a decision procedure for $=_{\mu/\alpha}$ on types.

1. A decision procedure for weak μ -equality via α -equivalence classes of types

1.1. DEFINITION (Of \mathbb{T}_{μ} and \rightarrow_{μ}). (i) Let $\mathbb{A} = \mathbb{A}_{\infty}$ be an infinite set of type atoms $\alpha, \beta, \gamma \cdots$ considered as type variables for the purpose of binding and substitution. The set of μ -types, notation \mathbb{T}_{μ} , is defined as follows. $\mathbb{T}_{\mu} = \alpha | A_1 \rightarrow A_2 | \mu \alpha A_1$.

(ii) On \mathbb{T}_{μ} we define the notion of α -reduction and α -conversion via the contraction rule

$$
\mu\alpha.A \mapsto_{\alpha} \mu\alpha'.A[\alpha := \alpha'],
$$
 provided $\alpha' \notin \text{FV}(A).$

The relation \rightarrow_{α} is the least compatible relation containing \rightarrow_{α} and $=\alpha$ is the least congruence containing \mapsto_{α} .

(iii) Define on \mathbb{T}_{μ} a notion of μ -reduction via the contraction rule \mapsto_{μ}

$$
\mu\beta.A_1 \mapsto_{\mu} A_1[\beta := \mu\beta.A_1]
$$

The relation $\rightarrow_{\mu} \subseteq \mathbb{T}_{\mu} \times \mathbb{T}_{\mu}$ is the compatible closure of \rightarrow_{μ} . That is

$$
A_1 \rightarrow_{\mu} B_1 \Rightarrow \mu \beta. A_1 \rightarrow_{\mu} \mu \beta. B_1;
$$

\n
$$
A_1 \rightarrow_{\mu} A'_1 \Rightarrow (A_1 \rightarrow A_2) \rightarrow_{\mu} (A'_1 \rightarrow A_2);
$$

\n
$$
A_2 \rightarrow_{\mu} A'_2 \Rightarrow (A_1 \rightarrow A_2) \rightarrow_{\mu} (A_1 \rightarrow A'_2).
$$

(iv) \rightarrow^*_{μ} denotes zero or more \rightarrow_{μ} steps and $=_{\mu}$ is the least congruence containing \mapsto_{μ} .

(v) In this first part we work with α -equivalence classes of types and we asume that all bound and free variables are different. Especially in $\mu\alpha_1...\mu\alpha_n$. A all α_i are different. 1.2. DEFINITION. Let [A] denote the α -equivalence class of A and let $\mathbb{T}_{[\mu]}$ denote the set of α -equivalence classes. \rightarrow_{μ} and $=_{\mu}$ can easily be lifted to relations $\rightarrow_{[\mu]}$ and $=_{[\mu]}$ on $\mathbb{T}_{[\mu]}$. In part 2 of this paper we will consider \mathbb{T}_{μ} , \rightarrow_{μ} and \models_{μ} but in this part 1 we work with $\mathbb{T}_{[\mu]}, \rightarrow_{[\mu]}$ and $=_{[\mu]}$ and we simply write $\mathbb{T}_{\mu}, \rightarrow_{\mu}$ and $=_{\mu}$. Also we write A for $|A|$.

1.3. DEFINITION. [Of standard reduction \rightarrow_{st}^*]

(a) $A \rightarrow_{st}^* A$

(b) If $A_1[\beta] := \mu \beta A_1 \rightarrow_{st}^* B$, then the composition $\mu \beta A_1 \rightarrow A_1[\beta] := \mu \beta A_1 \rightarrow_{st}^* B$ is a standard reduction.

(c) If $A_1 \rightarrow_{st}^* B_1$, then the induced reduction $\mu \beta \cdot A_1 \rightarrow_{st}^* \mu \beta \cdot B_1$ is a standard reduction. (d) If $A_1 \rightarrow_{st}^* A'_1$ and $A_2 \rightarrow_{st}^* A'_2$, then each reduction $(A_1 \rightarrow A_2) \rightarrow_{\mu}^* (A'_1 \rightarrow A'_2)$ generated by interleaving is a standard reduction.

1.4. EXAMPLE. (i) Let $A \equiv \mu \alpha \mu \beta \alpha$. Then $A \rightarrow_{\mu} \mu \beta_1 A \rightarrow_{\mu} \mu \beta_1 \mu \beta_2 A \rightarrow_{\mu} \cdots$ is a standard reduction.

(ii) Let $B \equiv \mu \alpha \cdot \alpha \rightarrow \alpha$. Then $B \rightarrow_{\mu} (B \rightarrow B) \rightarrow_{\mu} (B \rightarrow B) \rightarrow B \rightarrow_{\mu} \cdots$ is a standard reduction.

(iii) Let $C \equiv \mu \alpha \mu \beta \mu \gamma \alpha$. Then $C \rightarrow_{\mu} \mu \alpha \mu \gamma \alpha \rightarrow_{\mu} \mu \gamma_1 (\mu \alpha \mu \gamma \alpha)$ is not a standard reduction.

1.5. REMARK. We have the well known facts.

- (i) The relation \rightarrow_{μ} is Church Rosser (CR);
- (ii) If $A \to_{\mu}^* B$ then there is a standard reduction $A \to_{st}^* B$.

We illustrate (ii) in the above remark by an example.

1.6. EXAMPLE. Let $A = \mu \alpha \mu \beta \cdot C[\alpha, \beta]$. Here $C[.,.]$ is a so-called context. The variables α and β can occur several times in it. Now let $B = \mu\alpha$. C[α , $\mu\beta$. C]. Then

$$
A \to_{\mu} B = \mu \alpha. C[\alpha, \mu \beta. C[\alpha, \beta]] \to_{\mu} C[B, \mu \beta. C[B, \beta]].
$$

We can make this two-step reduction into a standard reduction as follows.

$$
A \to_{\mu} \mu \beta.C[A,\beta] \to_{\mu} C[A,\mu\beta.C[A,\beta]]
$$

and now reduce each occurrence of A to B , from left to right.

Let for the moment $A \downarrow B$ denote that A and B have a common standard reduct. Our aim is to show that $A =_{\mu} B$ is decidable. By (ii) of the above remark we are ready if we show that $A \downarrow B$ is decidable. However a direct proof of that is not so easy because in general a type A may have infinitely many standard reducts as is shown by (i) and (ii) of Example [1.4.](#page-9-0)

We remedy the two problems given by the example by changing the notion of standard reduction (and also the set of types) roughly as follows. In clause (c) of the notion of standard reduction we add the condition $\beta \in FV(A_1)$. Then the number of $\mu \beta_i$ as in (i) of the example is bounded. Moreover clause (d) is changed into $(A_1 \rightarrow A_2) \rightarrow A_i$ $i = 1, 2$ to remedy the problem shown in (ii).

This is formalised as follows.

1.7. DEFINITION (Of the set of annotated types $\mathbb{T}_{\mu'}$). (i) The set of annotated types $\mathbb{T}_{\mu'}$ is defined by $\mathbb{T}_{\mu'} \equiv \{(\mu \alpha_1) \dots (\mu \alpha_n) A \mid A \in \mathbb{T}_{\mu}, \alpha_i \in FV(A), \alpha_i \neq \alpha_j \text{ if } i \neq j \},\$

(ii) We consider the annotated types modulo α -equivalence as follows $(\mu \alpha_1)$ $(\mu \alpha_n)A \equiv (\mu \alpha'_1)$ $(\mu \alpha'_n)A[\alpha_i := \alpha'_i]$ $'_{i}$] $(\alpha'_{i} \notin \text{FV}(A) \cup \text{BV}(A)).$ And implicitly α'_i $\gamma_i \neq \alpha'_j$ $'_{j}$ if $i \neq j$,

1.8. REMARK. (i) In Endrullis et al. [2] a set $AnnTer(\mu)$ is defined by $AnnTer(\mu)$ = $\{(\mu\alpha_1)....(\mu\alpha_n)A \mid A \in \mathbb{T}_\mu\}$. So they don't have the conditions $\alpha_i \in FV(A)$ and $\alpha_i \neq \alpha_j$ if $i \neq j$. Moreover they don't work modulo α -conversion.

(ii) The intuition for $(\mu \alpha_1)$ $(\mu \alpha_n)A$ is that it stands for $\mu \alpha_1$ $\mu \alpha_n.A \in \mathbb{T}_\mu$ where the $\mu \alpha_i$ at the root are frozen. This will become clear in Definition [1.10.](#page-3-0)

(iii) Note that $\mathbb{T}_{\mu} \subset \mathbb{T}_{\mu'}$.

1.9. NOTATION. (i) We denote $(\mu \alpha_1)$ $(\mu \alpha_n)A$ simply as $(\mu \alpha_1 \dots \alpha_n)A$ and mostly we write $(\mu \vec{\alpha})A$.

(ii) a, b, c....denote elements of $\mathbb{T}_{\mu'}$ and as before A, B, C.... stand for elements of \mathbb{T}_{μ} .

(iii) If $\vec{\alpha}$ is a sequence of (different) type variables then $\vec{\alpha}$ |a denotes the sequence that arises from $\vec{\alpha}$ by omitting the variables that do not occur freely in a. Similarly for $\vec{\alpha} |a \cap b$. $(\mu \vec{\alpha}) \Box A$ stands for $(\mu (\vec{\alpha} \Box A)) A$.

If $S \subseteq \mathbb{T}_{\mu'}$ then $(\mu \vec{\alpha}) \square S = \{ (\mu \vec{\alpha}) \square a \mid a \in S \}.$

(iv) If $S \subset \mathbb{T}_{\mu'}$ then |S| stands for the number of elements of S.

(v) $\{\vec{\alpha}\}\$ stands for the set of variables occurring in $\vec{\alpha}$. When we write $\{\alpha_1, ..., \alpha_n\}$ we always assume $\alpha_i \neq \alpha_j$ for $i \neq j$.

1.10. DEFINITION. $(0f \rightarrow_{\mu'}: \mathbb{T}_{\mu'} \rightarrow \mathbb{T}_{\mu'}$ and $SC(a)$)

(i) The reduction relation $\rightarrow_{\mu'}: \mathbb{T}_{\mu'} \rightarrow \mathbb{T}_{\mu'}$ is defined as follows.

$$
(\mu \vec{\alpha})\mu \beta \cdot A_1 \rightarrow_{\mu'} (\mu \vec{\alpha}\beta)A_1 \text{ if } \beta \in FV((A_1);
$$

$$
(\mu \vec{\alpha})\mu \beta \cdot A_1 \rightarrow_{\mu'} (\mu \vec{\alpha})A_1[\beta := \mu \beta \cdot A_1];
$$

$$
(\mu \vec{\alpha})(A_1 \rightarrow A_2) \rightarrow_{\mu'} (\mu \vec{\alpha}) \square A_i \quad i = 1, 2;
$$

(ii)
$$
SC(a) = \{b \in \mathbb{T}_{\mu'} \mid a \to_{\mu'}^* b\}.
$$

1.11. REMARK. (i) Note that $\rightarrow_{\mu'}$ is not compatible with μ or \rightarrow . So for example we do not have $\mu \alpha \mu \beta \cdot A_1 \rightarrow_{\mu'} \mu \alpha \cdot A_1[\beta] := \mu \beta \cdot A_1].$

(ii) Now the outline of the decidability proof is as follows.

–We define a derivation system $\vdash_{\mu'} a = b$ for types $a, b \in \mathbb{T}_{\mu'}$, closely connected with the reduction relation $\rightarrow_{\mu'}$ and we show for $A, B \in \mathbb{T}_{\mu}$:

 $\vdash_{\mu'} A = B \Leftrightarrow A$ and B have a common standard reduct.

Hence we have $A =_{\mu} B \Leftrightarrow \vdash_{\mu'} A = B$.

– We note that for all nodes $c = d$ occurring in a derivation tree for $\vdash_{\mu'} a = b$ we have $c \in SC(a)$ and $d \in SC(b)$.

– We show that $|SC(a)|$ is bounded as a function in a. Hence the number of different nodes in a derivation tree for $\vdash_{\mu'} a = b$ is bounded as a function in a, b.

– We conclude that $\vdash_{\mu'} a = b$ is decidable and hence so is $A =_{\mu} B$.

We start by defining the derivation system $\vdash_{\mu'} a = b$ for $a, b \in \mathbb{T}_{\mu'}$. We give the definition via representatives, types instead of α -equivalence classes of types, and we show that it is invariant under α -conversion. Hence this defines in fact a relation on the set of α -equivalence classes.

1.12. DEFINITION. (Of $\vdash_{\mu'} a = b$ for $a, b \in \mathbb{T}_{\mu'}$)

Note that in the above definition A, B stand for types, not α -equivalence classes of types and $=_{\alpha}$ on $\mathbb{T}_{\mu'}$ is given by

$$
A =_{\alpha} A' \Rightarrow (\mu \alpha_1) \dots (\mu \alpha_n) A' =_{\alpha} (\mu \alpha'_1) \dots (\mu \alpha'_n) A'[\alpha_i := \alpha'_i] \ (\alpha'_i \notin \text{FV}(A)).
$$

Note also that in the rule (μ -freezing) implicitly $\gamma \in FV(A_1)$, $\delta \in FV(B_1)$ and in rule (decomposition) $\{\vec{\alpha_i}\}\subseteq \text{FV}(A_i), \{\vec{\beta_i}\}\subseteq \text{FV}(B_i)$.

In a decision procedure of $\vdash_{\mu'} a = b$ in fact these rules are used upside down. That explains the names of the rules.

1.13. LEMMA. Let $a =_{\alpha} a', b =_{\alpha} b'$. Then

$$
\vdash_{\mu'} a = b \Rightarrow \vdash_{\mu'} a' = b'.
$$

PROOF. By an easy induction on the definition of $\vdash_{\mu'} a = b$.

So Definition [1.12](#page-3-1) defines in fact a relation on the set of α -equivalence classes. We denote that relation also by $\vdash_{\mu'}$.

1.14. LEMMA. (i)
$$
\text{FV}(\mu\beta.A_1) = \text{FV}(A_1[\beta := \mu\beta.A_1]) = \text{FV}((\mu\beta)A_1).
$$

\n(ii) $A \rightarrow_{\mu}^* B \Rightarrow \text{FV}(A) = \text{FV}(B).$
\n(iii) $a \rightarrow_{\mu'}^* b \Rightarrow \text{FV}(b) \subseteq \text{FV}(a).$
\n(iv) $\vdash_{\mu'} (\mu\vec{\alpha})A = (\mu\vec{\beta})B \Rightarrow |\{\vec{\alpha}\}| = |\{\vec{\beta}\}|, \text{FV}((\mu\vec{\alpha})A) = \text{FV}((\mu\vec{\beta})B)$

For easier use in the following we rewrite Definition [1.12](#page-3-1) for α -equivalence classes as follows.

Г

1.15. DEFINITION. (Of $\vdash_{\mu'} a = b$ for α -equivalence classes a, b .)

(axiom)	$a = a$
$(\text{left } \mu\text{-step})$	$(\mu \vec{\alpha})A_1[\beta := \mu \beta.A_1] = (\mu \vec{\alpha})B$ $(\mu \vec{\alpha}) \mu \beta \cdot A_1 = (\mu \vec{\alpha}) B$
$(\text{right }\mu\text{-step})$	$(\mu \vec{\alpha})A = (\mu \vec{\alpha})B_1[\beta := \mu \beta.B_1]$ $(\mu \vec{\alpha})A = (\mu \vec{\alpha})\mu\beta.B_1$
$(\mu$ -freezing)	$(\mu \vec{\alpha} \beta) A_1 = (\mu \vec{\alpha} \beta) B_1$ $(\mu \vec{\alpha}) \mu \beta \cdot A_1 = (\mu \vec{\alpha}) \mu \beta \cdot B_1$
(decomposition)	$\{\vec{\alpha}\}\subseteq \mathrm{FV}(A_1\rightarrow A_2)$ $\vec{\alpha} \restriction A_1 = \vec{\alpha_1}, \vec{\alpha} \restriction A_2 = \vec{\alpha_2}$ $(\mu \vec{\alpha_1})A_1 = (\mu \vec{\alpha_1})B_1$ $(\mu \vec{\alpha_2})A_2 = (\mu \vec{\alpha_2})B_2$ $(\mu \vec{\alpha})A_1 \rightarrow A_2 = (\mu \vec{\alpha})B_1 \rightarrow B_2$

The following is an example of a derivation of $\vdash_{\mu'} a = b$. We reason backwards, so we write down $a = b$ and using the rules upside down we hope to finish with only axioms. 1.16. Example.

$$
\mu\alpha\beta\gamma\cdot((\mu\delta.\alpha)\to\gamma) = \mu\alpha\beta\cdot(\alpha\to(\mu\gamma.\alpha\to\gamma))
$$

\n
$$
\downarrow \quad \mu-\text{freezing}
$$

\n
$$
(\mu\alpha)\mu\beta\gamma\cdot((\mu\delta.\alpha)\to\gamma) = (\mu\alpha)\mu\beta\cdot(\alpha\to(\mu\gamma.\alpha\to\gamma))
$$

\n
$$
\downarrow \quad \text{left and right } \mu-\text{step}
$$

\n
$$
(\mu\alpha)\mu\gamma\cdot((\mu\delta.\alpha)\to\gamma) = (\mu\alpha)(\alpha\to(\mu\gamma.\alpha\to\gamma))
$$

\n
$$
\downarrow \quad \text{left } \mu-\text{step}
$$

\n
$$
(\mu\alpha)((\mu\delta.\alpha)\to\mu\gamma\cdot((\mu\delta.\alpha)\to\gamma)) = (\mu\alpha)(\alpha\to(\mu\gamma.\alpha\to\gamma))
$$

\n
$$
\downarrow \quad \text{decomposition}
$$

\n
$$
(\mu\alpha)(\mu\delta.\alpha) = (\mu\alpha)\alpha \qquad (\mu\alpha)\mu\gamma\cdot((\mu\delta.\alpha)\to\gamma) = (\mu\alpha)(\mu\gamma.\alpha\to\gamma)
$$

\nleft
$$
\downarrow \mu-\text{freezing}
$$

\n
$$
(\mu\alpha)\alpha = (\mu\alpha)\alpha \qquad (\mu\alpha\gamma)((\mu\delta.\alpha)\to\gamma) = (\mu\alpha\gamma)(\alpha\to\gamma)
$$

\n
$$
\downarrow \text{decomposition}
$$

\n
$$
(\mu\alpha)\mu\delta.\alpha = (\mu\alpha)\alpha \qquad (\mu\gamma)\gamma = (\mu\gamma)\gamma
$$

\n
$$
\downarrow \text{left } \mu-\text{step}
$$

\n
$$
(\mu\alpha)\alpha = (\mu\alpha)\alpha
$$

1.17. REMARK. Let $A = \mu \alpha \beta \gamma \cdot ((\mu \delta \alpha) \rightarrow \gamma)$ and $B = \mu \alpha \beta \cdot (\alpha \rightarrow (\mu \gamma \cdot \alpha \rightarrow \gamma))$. In the example above we gave a proof of $\vdash_{\mu'} A = B$. In fact one could try other proofs. For example one could start with a left μ -step and a right μ -step. Then one encounters at some stage again a node $A = B$. So one could go on for ever and the result would be an infinite tree. But of course one should stop developing the sub branch at that node. In general if in a derivation of $\vdash_{\mu'} a = b$ we arrive at a node $c = d$ that we encountered already higher up in the branch from the root to that node then we stop at that node. Further on we will show that the number of nodes in the resulting tree is bounded as a function of a, b.

But first we will show in the next proposition that $\vdash_{\mu'} A = B$ iff A and B have a common standard reduct.

1.18. PROPOSITION. The following two assertions are equivalent

(i)
$$
\vdash_{\mu'}
$$
 $(\mu \vec{\alpha})A = (\mu \vec{\alpha})B;$

(ii) A and B have a common standard reduct.

PROOF. (i)⇒(ii). By induction on the derivation of $\vdash_{\mu'} (\mu \vec{\alpha})A = (\mu \vec{\alpha})B$.

(axiom) $(\mu \vec{\alpha})A = (\mu \vec{\alpha})A$. This is clearly ok.

(left μ -step) $\frac{(\mu \vec{\alpha})A_1[\beta := \mu \beta.A_1] = (\mu \vec{\alpha})B}{(\vec{\alpha})A_1[\beta + \mu \beta + \beta]}$ $(\mu \vec{\alpha})\mu \beta.A_1 = (\mu \vec{\alpha})B$

By the induction hypothsis $A_1[\beta := \mu \beta A_1]$ and B have a common standard reduct, hence the same holds for $\mu\beta.A_1$ and B by (b) of Definition [1.3.](#page-9-1)

(right μ -step) Similarly.

$$
(\mu\text{-freezing}) \quad \frac{(\mu\vec{\alpha}\beta)A_1 = (\mu\vec{\alpha}\beta)B_1}{(\mu\vec{\alpha})\mu\beta.A_1 = (\mu\vec{\alpha})\mu\beta.B_1}
$$

By the induction hypothesis A_1 and B_1 have a common standard reduct, hence the same holds for $\mu\beta.A_1$ and $\mu\beta.B_1$.

$$
\{\vec{\alpha}\} \subseteq \text{FV}(A_1 \rightarrow A_2) \quad \vec{\alpha} | A_1 = \vec{\alpha_1}, \quad \vec{\alpha} | A_2 = \vec{\alpha_2}
$$
\n
$$
\text{(decomposition)} \quad \frac{(\mu \vec{\alpha_1}) A_1 = (\mu \vec{\alpha_1}) B_1 \quad (\mu \vec{\alpha_2}) A_2 = (\mu \vec{\alpha_2}) B_2}{(\mu \vec{\alpha}) A_1 \rightarrow A_2 = (\mu \vec{\alpha}) B_1 \rightarrow B_2}
$$

By the induction hypothesis A_1 and B_1 have a common standard reduct and the same holds for A_2 and B_2 . So it also holds for $A_1 \rightarrow A_2$ and $B_1 \rightarrow B_2$.

(ii) \Rightarrow (i). By induction on the definition of $A \rightarrow_{st}^* C$ and $B \rightarrow_{st}^* C$ in Definition [1.3.](#page-9-1)

Case (a) for both A and B. Then $B = A$ and we have $\vdash_{\mu'} (\mu \vec{\alpha}) A = (\mu \vec{\alpha}) A$ by axiom.

Case (b) for A (and similarly for B). We have $\vdash_{\mu'} (\mu \vec{\alpha})A_1[\beta := \mu \beta.A_1] = (\mu \vec{\alpha})B$ and we get $\vdash_{\mu'} (\mu \vec{\alpha})\mu\beta.A_1 = (\mu \vec{\alpha})B$ by (left μ -step).

Case (c) for A an for B. Now $A = \mu \beta A_1$ and $B = \mu \beta B_1$, $A_1 \rightarrow_{st}^* C_1 \leftarrow_{st}^* B_1$ and therefore $\mu\beta.A_1 \rightarrow_{st}^* \mu\beta.C_1 \leftarrow_{st}^* \mu\beta.B_1.$

Note that $FV(A_1) = FV(B_1)$ by Lemma [1.14\(](#page-4-0)ii) We distinguish two subcases

Subcase (c_1) $\beta \in FV(A_1) \cap FV(B_1)$. By the induction hypothesis we have $\vdash_{\mu'} (\mu \vec{\alpha} \beta) A_1 = (\mu \vec{\alpha} \beta) B_1$ and hence by (μ -freezing) $\vdash_{\mu'} (\mu \vec{\alpha}) \mu \beta . A_1 = (\mu \vec{\alpha}) \mu \beta . B_1.$

Subcase (c_2) $\beta \notin FV(A_1), \beta \notin FV(B_1).$ Now we have by the induction hypothesis $\vdash_{\mu'} (\mu \vec{\alpha})A_1 = (\mu \vec{\alpha}).B_1$, hence we get by (left μ -step) and (right μ -step) $\vdash_{\mu'} (\mu \vec{\alpha}) \mu \beta A_1 = (\mu \vec{\alpha}) \mu \beta B_1$.

Case (d) for A and for B. Now we have $A_1 \rightarrow_{st}^* C_1 \leftarrow_{st}^* B_1$, $A_2 \rightarrow_{st}^* C_2 \leftarrow_{st}^* B_2$ and hence $A = A_1 \rightarrow A_2 \rightarrow_{st}^* C_1 \rightarrow C_2 \leftarrow_{st}^* B_1 \rightarrow B_2$. Note that by Lemma [1.14\(](#page-4-0)ii) we have $FV(A_i) = FV(B_i)$ and $FV(A_1 \rightarrow A_2) = FV(B_1 \rightarrow B_2)$. Let $\{\vec{\alpha}\}\subseteq FV(A_1 \rightarrow B_2)$ A_2) so that $(\mu \vec{\alpha})A_1 \rightarrow A_2, (\mu \vec{\alpha})B_1 \rightarrow B_2 \in \mathbb{T}_{\mu'}$. Define $\vec{\alpha}_i = \vec{\alpha} \upharpoonright A_i, i = 1, 2$. Then $(\mu \vec{\alpha_i}) A_i, \ (\mu \vec{\alpha_i}) B_i \in \mathbb{T}_{\mu'}$ and by the induction hypothesis, two times, we have $\vdash_{\mu'} (\mu \vec{\alpha_1}) A_1 =$ $(\mu \vec{\alpha_1})B_1, \vdash_{\mu'} (\mu \vec{\alpha_2})A_2 = (\mu \vec{\alpha_2})B_2.$

Now we have treated all cases. If we have Case (c) for A and Case (a) for B then in fact we have Case (c) for both A and B. If we have Case (d) for A and Case (a) for B then in fact we have Case (d) for both A and B. Finally Case (c) for A and Case (d) for B (or vice versa) cannot occur at the same time.

Now we will finish the proof of decidability of $=$ _µ by showing that the number of different nodes $c = d$ occurring in a derivation tree of $\vdash_{\mu'} a = b$ is bounded as a function in a, b .

Note that for all these nodes $c = d$ we have $c \in \mathcal{SC}(a)$, $d \in \mathcal{SC}(b)$. (See Definition [1.10\)](#page-3-0) We start by showing that $|SC(a)|$ is bounded as a function in a.

1.19. LEMMA. $(\mu \vec{\alpha}) \square((\mu \vec{\beta}) \square a) = (\mu \vec{\alpha} \vec{\beta}) \square a$.

PROOF. Immediate. Note that by the variable convention $\vec{\alpha}$ and $\vec{\beta}$ have no variables in common.

1.20. LEMMA. $SC(\alpha) = {\alpha};$ $SC(\mu\beta.A_1) = {\mu\beta.A_1} \cup (\mu\beta) \square SC(A_1) \cup SC(A_1[\beta := \mu\beta.A_1]);$ $SC(A_1 \rightarrow A_2) = \{A_1 \rightarrow A_2\} \cup SC(A_1) \cup SC(A_2);$ $SC((\mu\vec{\alpha})A) = (\mu\vec{\alpha})\Box SC(A).$

PROOF. By induction on the structure of $a \in \mathbb{T}_{\mu'}$, using the preceding lemma. Note that the second clause also holds for $\beta \notin FV(A_1)$ because then $(\mu \beta) \square SC(A_1) = SC(A_1)$ $\mathrm{SC}(A_1[\beta := \mu \beta.A_1])$ by Lemma [1.14\(](#page-4-0)iii).

Now $SC([A_1|\beta := \mu\beta.A_1])$ could be intricate but we will show in Lemma [1.22](#page-8-0) that we have in fact $SC(\mu\beta.A_1) \subseteq {\mu\beta.A_1} \cup (\mu\beta) \square SC(A_1) \cup SC(A_1)[\beta := \mu\beta.A_1].$

1.21. LEMMA. If $A[\beta := B] \rightarrow_{\mu'}^* b$ then we have either

- (i) This reduction is $A[\beta := B] \rightarrow_{\mu'}^* B \rightarrow_{\mu'}^* b$
- (ii) This reduction is a $[\beta := B]$ instance of $A \to_{\mu'}^* a$ (So $b = a[\beta := B]$).

PROOF. By induction on the length of the reduction $A[\beta := B] \to_{\mu'}^* b$.

If the length is zero then we have (ii). Else we have $A[\beta := B] \rightarrow_{\mu'}^* b' \rightarrow_{\mu'} b$. The induction hypothesis for $A[\beta := B] \to_{\mu'}^* b'$ gives that we have either (i) or (ii) as follows (i) $A[\beta := B] \to_{\mu'}^* B \to_{\mu'}^* b'$ Then we have also case (i) for $A[\beta := B] \to_{\mu'}^* b$.

(ii) The reduction $A[\beta := B] \to_{\mu'}^* b'$ is a $[\beta := B]$ instance of $A \to_{\mu'}^* a'.$ So $b' = a'[\beta := B]$ We distinguish subcases for a' as follows.

(iia) $a' = \beta$. Then we have Case (i) (for $A[\beta := B] \to_{\mu'}^* b$).

(iib) $a' = \alpha \neq \beta$ or $a' = (\mu \alpha) \cdot \alpha$. These cases cannot occur because then $b' = a'$ is in nf.

(iic) $a' = (\mu \vec{\alpha}) \mu \gamma . A'$ with $\beta \notin {\vec{\alpha}, \gamma}$.

In this case $(\mu \vec{\alpha})$ at the root is very innocent. It is frozen and occurs in the same way at the root in all types occurring in the proof in this part (iic). Therefore for simplicity we assume $\{\vec{\alpha}\} = \emptyset$. Now $b' = \mu \gamma A'[\beta := B]$. We distinguish two subcases.

(iic1) $\mu \gamma . A'[\beta := B] \rightarrow_{\mu'} (\mu \gamma) . (A'[\beta := B]) (*)$ Then $\gamma \in \text{FV}(A'[\beta := \widehat{B}])$, hence also $\gamma \in \text{FV}(A')$, because $\gamma \notin \text{FV}(B)$ by the variable convention. So (*) is a $\hat{\beta} := B$ instance of $\mu \gamma \hat{A}' \to_{\mu'} (\mu \gamma) \hat{A}'$. In total we have that $A[\beta := B] \to_{\mu'}^* \dot{B}' \to_{\mu'} b$ is an instance of $A \to_{\mu'}^* \mu \gamma \cdot A' \to_{\mu'} (\mu \gamma) \cdot A'$ and we are in Case (ii).

(iic2) $b' = \mu \gamma . A'[\beta := B] \rightarrow_{\mu'} A'[\beta := B][\gamma := \mu \gamma . A'[\beta := B]] = b.$ By the substitution lemma we get $b' = \mu \gamma A'[\beta := \beta] \rightarrow_{\mu'} \mathring{A}'[\gamma := \mu \gamma A'][\beta := \beta] = b.$ Now we have that $A[\beta := B] \to_{\mu'}^* a'[\beta := B] \to_{\mu'} a[\beta := B]$ is an instance of $A \to_{\mu'}^* a' =$ $\mu \gamma \cdot A' \rightarrow_{\mu'} a = A'[\gamma := \mu \gamma \cdot A']$ and we are in Case (ii).

(iid) $a' = (\mu \vec{\alpha})(A_1 \rightarrow A_2)$. Then $b' = (\mu \vec{\alpha})(A_1[\beta := B] \rightarrow A_2[\beta := B]) \rightarrow_{\mu'} (\mu \vec{\alpha}) \square (A_i[\beta := B])$ $[B]$ = b and $a' \rightarrow_{\mu'} (\mu \vec{\alpha}) \square A_i$. Now we are in case (ii) again.

1.22. LEMMA. $SC(\mu\beta.A_1) \subseteq {\mu\beta.A_1} \cup (\mu\beta) \square SC(A_1) \cup SC(A_1)[\beta := \mu\beta.A_1].$

PROOF. Let $\mu\beta \cdot A_1 \rightarrow_{\mu'}^* b$ be a reduction path of minimal length. If the length is zero then $b = \mu \beta A_1$. Now let the length be bigger then zero. We distinguish two cases.

Case (i) $\mu\beta.A_1 \rightarrow_{\mu'} (\mu\beta).A_1 \rightarrow_{\mu'}^* b$ Now $\beta \in FV(A_1)$ and $b \in SC((\mu \beta).A_1) = (\mu \beta) \Box SC(A_1)$.

Case(ii) $\mu\beta.A_1 \rightarrow_{\mu'} A_1[\beta := \mu\beta.A_1] \rightarrow_{\mu'}^* b.$ We distinguish two subcases according to Lemma [1.21](#page-7-0) (iia) $\mu \beta \cdot A_1 \rightarrow_{\mu'} A_1[\beta := \mu \beta \cdot A_1] \rightarrow_{\mu'}^* \mu \beta \cdot A_1 \rightarrow_{\mu'}^* b$. This cannot occur because it is not a reduction of minimal length. (iib) $A_1[\beta := \mu \beta A_1] \rightarrow \mu^* \phi$ is a $[\beta := \mu \beta A_1]$ instance of $A_1 \rightarrow \mu^* \phi$. Now $a \in SC(A_1)$ hence $b = a[\beta := \mu \beta.A_1] \in SC(A_1)[\beta := \mu \beta.A_1]$.

1.23. Definition.

$$
l(\alpha) = 1;
$$

\n
$$
l(A_1 \to A_2) = 1 + l(A_1) + l(A_2);
$$

\n
$$
l(\mu\beta.A_1) = 1 + l(A_1);
$$

\n
$$
l((\mu\vec{\alpha}).A) = l(A).
$$

1.24. LEMMA. $|SC((\mu \vec{\alpha})A)| = |SC(A)| \leq 3^{l(A)}$.

PROOF. By Lemmas [1.20](#page-7-1) and [1.22.](#page-8-0) \blacksquare

1.25. LEMMA. Let $n = n((\mu \vec{\alpha})A = (\mu \vec{\alpha})B)$ be the number of different nodes in a complete derivation treee for $\vdash_{\mu'} (\mu \vec{\alpha})A = (\mu \vec{\alpha})B$. Then

$$
n \leq 3^{(3^{l(A)+l(B)})}.
$$

PROOF. Easy. Note that each node in the derivation tree has at most three daughters.

1.26. COROLLARY. (i) $\vdash_{\mu'} (\mu \vec{\alpha})A = (\mu \vec{\alpha})B$ is decidable.

(ii) $\vdash_{\mu'} A = B$ is decidable.

1.27. COROLLARY. $A =_{\mu} B$ is decidable.

PROOF. By Remark [1.5](#page-10-0) and Proposition [1.18.](#page-6-0)

2. A decision procedure of weak μ -equality for types themselves

In this subsection we work with the μ -types themselves instead of their α -equivalence classes. On this set of types $\rightarrow_{\mu/\alpha}$ is the reduction relation on the set of types \mathbb{T}_{μ} generated by \rightarrow_{μ} and \rightarrow_{α} . We show that the relation $=_{\mu/\alpha}$ on \mathbb{T}_{μ} is decidable via the first order decision procedure of Definition [1.12](#page-3-1) for types.

2.1. DEFINITION. $\rightarrow_{\mu/\alpha}$ is the reduction relation on the set of types \mathbb{T}_{μ} generated by \rightarrow_{μ} and \rightarrow_{α} .

 $=\mu/\alpha$ is the transitive, symmetric, reflexive closure of $\rightarrow_{\mu/\alpha}$. $\rightarrow_{\mu/\alpha}^*$ denotes zero or more $\rightarrow_{\mu/\alpha}$ steps.

We will show

$$
A =_{\mu/\alpha} B \Leftrightarrow \vdash_{\mu'} A = B.
$$

where $\vdash_{\mu'}$ derives equations between types in $\mathbb{T}_{\mu'}$, as given in Definition [1.12](#page-3-1) and moreover that this relation is decidable.

From now on α -quivalence classes in \mathbb{T}_{μ} and $\mathbb{T}_{\mu'}$ are denoted by [A] and [a]. $=_{\lbrack \mu \rbrack}$ stands for equality on the set of α -equivalence classes [A] and $\vdash_{\lbrack \mu' \rbrack}$ denotes the derivation system of Definition [1.15](#page-4-1) for classes [a] where $a \in \mathbb{T}_{\mu'}$.

2.2. LEMMA. $\vdash_{\mu'} a = b \Leftrightarrow \vdash_{[\mu']} [a] = [b]$ and both derivations can be done by the "same" steps. Equivalently they have the same length.

PROOF. Immediate

2.3. LEMMA. $A =_{\mu/\alpha} B \Leftrightarrow [A] =_{[\mu]} [B].$

PROOF. Immediate

Combining Remark [1.5,](#page-10-0) Proposition [1.18](#page-6-0) and Lemmas [2.2](#page-9-2) and [2.3](#page-9-1) we get

2.4. COROLLARY. $A =_{\mu/\alpha} B \Leftrightarrow \vdash_{\mu'} A = B$.

Finally by Lemma [2.2](#page-9-2) we have that $\vdash_{\mu'} A = B$ is decidable. In fact there is a bijection from the derivation tree for $\vdash_{\mu'} A = B$ on the derivation tree for $\vdash_{\lbrack \mu' \rbrack} [A] = [B]$. So we have the following

2.5. THEOREM. $A =_{\mu/\alpha} B$ is decidable via the first order decision procedure $\vdash_{\mu'} A = B$. $\overline{}$

2.6. Remark. In this part 2 we do not have Lemma [1.22](#page-8-0) for types. It only holds modulo α -equivalence.

References

[1] Felice Cardone and Mario Coppo. Decidability Properties of Recursive Types. In C. Blundo and C. Laneve, editors, ICTCS 2003 volume 2841 of LNCS, pages 242-255. Springer, 2003.

[2] Jörg Endrullis, Clemens Grabmayer, Jan Willem Klop, Vincent van Oostrom. On equal μ -terms. 2011. Free university Amsterdam.

[3] V. van Oostrom. FD à la Melliès. February 1997. Free University Amsterdam.