

Exploring the origin of magnetic fields in massive stars: A survey of O-type stars in clusters and in the field^{*}

S. Hubrig¹, M. Schöller², N. V. Kharchenko^{1,3}, N. Langer⁴, W. J. de Wit⁵, I. Ilyin¹, A. F. Kholtygin⁶,
A. E. Piskunov^{1,7}, N. Przybilla⁸, and the MAGORI collaboration

¹ Astrophysikalisches Institut Potsdam, An der Sternwarte 16, 14482 Potsdam, Germany

² European Southern Observatory, Karl-Schwarzschild-Str. 2, 85748 Garching, Germany

³ Main Astronomical Observatory, 27 Academica Zabolotnogo Str., 03680 Kiev, Ukraine

⁴ Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, 53121 Bonn, Germany

⁵ European Southern Observatory, Alonso de Cordova 3107, Santiago, Chile

⁶ Astronomical Institute, Saint-Petersburg State University, Saint-Petersburg, Russia

⁷ Institute of Astronomy of the Russian Acad. Sci., 48 Pyatnitskaya Str., 109017 Moscow, Russia

⁸ Dr.-Karl-Rehms-Sternwarte Bamberg & ECAP, Universität Erlangen-Nürnberg, Sternwartstr. 7, 96049 Bamberg, Germany

Received date / Accepted date

ABSTRACT

Context. Although important effects of magnetic fields in massive stars are suggested by recent models and observations, only a small number of massive O-type stars have been investigated for magnetic fields until now. Additional observations are of utmost importance to constrain the conditions which enable the presence of magnetic fields and give first trends about their occurrence rate and field strength distribution.

Aims. To investigate statistically whether magnetic fields in massive stars are ubiquitous or appear in stars with specific spectral classification, certain ages, or in a special environment, we acquired 41 new spectropolarimetric observations for 36 stars. Among the observed sample roughly half of the stars are probable members of clusters at different ages, whereas the remaining stars are field stars not known to belong to any cluster or association.

Methods. Spectropolarimetric observations were obtained during three different nights using the low-resolution spectropolarimetric mode of FORS2 (FOcal Reducer low dispersion Spectrograph) mounted on the 8-m Antu telescope of the VLT. To assess the membership in open clusters and associations, we used astrometric catalogues with the best currently available kinematic and photometric data.

Results. A field at a significance level of 3σ was detected in ten O-type stars. Importantly, the largest longitudinal magnetic fields were measured in two Of?p stars: $\langle B_z \rangle = -381 \pm 122$ G for CPD-28 2561 and $\langle B_z \rangle = -297 \pm 62$ G for HD 148937, previously detected by us as magnetic. The obtained observations of HD 148937 on three different nights indicate that the magnetic field is slightly variable. Our new measurements support our previous conclusion that large-scale organized magnetic fields with polar field strengths in excess of 1 kG are not widespread among O-type stars. Among the stars with a detected magnetic field, only one star, HD 156154, belongs to an open cluster at high membership probability. According to previous kinematic studies, four magnetic O-type stars in the sample are well-known candidate runaway stars.

Key words. polarization - stars: early-type - stars: magnetic field - stars: kinematics and dynamics — open clusters and associations: general

1. Introduction

Magnetic fields play an important role in astrophysical phenomena of the universe at various scales. In galaxies, dynamo models associated with various MHD instabilities occurring in the interstellar medium (ISM) are used to explain the formation of the galactic structure (e.g., Gomez & Cox 2004; Bonanno & Urpin 2008). Magnetic fields play a role in the evolution of interstellar molecular clouds and in the star formation process, where the cloud collapse is probably taking place along the magnetic field lines (e.g. Alves et al. 2008). They are also present at all stages of stellar evolution, from young T Tauri stars and Ap/Bp stars to the end products: white dwarfs, neutron stars, and magnetars.

On the other hand, role of magnetic fields in massive O-type stars and Wolf-Rayet (WR) stars remains unknown. No definitive magnetic field was ever detected in WR stars and presently only less than a dozen O stars have published magnetic fields. Also, the theories on the origin of magnetic fields in O-type stars are still poorly developed, mostly due to the fact that the distribution of magnetic field strengths in massive stars from the ZAMS to more evolved stages has not yet been studied.

In our study we focus on magnetic fields of massive stars observed in different environments: in open clusters at different ages and in the field. The results of our recent kinematic analysis of known magnetic O-type stars using the best available astrometric, spectroscopic, and photometric data indicates that the presence of a magnetic field is more frequently detected in candidate runaway stars than

^{*} Based on observations obtained at the European Southern Observatory, Paranal, Chile (ESO programme 085.D-0667(A)).

in stars belonging to clusters or associations (Hubrig et al. 2011a). As the sample of stars with magnetic field detections is still very small, a study of a larger sample is urgently needed to confirm the detected trend by dedicated magnetic field surveys of O stars in clusters/associations and in the field. We were granted four nights in 2010 May with FORS 2 at the VLT to survey magnetic fields in massive stars, but due to deteriorated weather conditions, only half of the granted time could be used for observations. Notwithstanding, the obtained results allow us to preliminarily constrain the conditions which enable the presence of magnetic fields and give first trends about their occurrence rate and field strength distribution. This information is critical for answering the principal question of the possible origin of magnetic fields in massive stars.

In the following, we present 41 new measurements of magnetic fields in 36 massive stars using FORS 2 at the VLT in spectropolarimetric mode. Our observations and the obtained results are described in Sect. 2, and their discussion is presented in Sect. 3.

2. Observations and results

Spectropolarimetric observations were carried out in 2010 May 20–23 in visitor mode at the European Southern Observatory with FORS2 mounted on the 8-m Antu telescope of the VLT. This multi-mode instrument is equipped with polarization analyzing optics, comprising super-achromatic half-wave and quarter-wave phase retarder plates, and a Wollaston prism with a beam divergence of $22''$ in standard resolution mode¹. Polarimetric spectra were obtained with the GRISM 600B and the narrowest slit width of $0''.4$ to achieve a spectral resolving power of $R \sim 2000$. The use of the mosaic detector made of blue optimized E2V chips and a pixel size of $15 \mu\text{m}$ allowed us to cover a large spectral range, from 3250 to 6215 \AA , which includes all hydrogen Balmer lines from $H\beta$ to the Balmer jump. The spectral types and the visual magnitudes of the studied stars are listed in Table 1.

A detailed description of the assessment of the longitudinal magnetic-field measurements using FORS 2 is presented in our previous papers (e.g., Hubrig et al. 2004a, 2004b, and references therein). The mean longitudinal magnetic field, $\langle B_z \rangle$, was derived using

$$\frac{V}{I} = -\frac{g_{\text{eff}} e \lambda^2}{4\pi m_e c^2} \frac{1}{I} \frac{dI}{d\lambda} \langle B_z \rangle, \quad (1)$$

where V is the Stokes parameter that measures the circular polarisation, I is the intensity in the unpolarised spectrum, g_{eff} is the effective Landé factor, e is the electron charge, λ is the wavelength, m_e the electron mass, c the speed of light, $dI/d\lambda$ is the derivative of Stokes I , and $\langle B_z \rangle$ is the mean longitudinal magnetic field.

Longitudinal magnetic fields were measured in two ways: using only the absorption hydrogen Balmer lines or using the entire spectrum including all available absorption lines. The lines that show evidence for emission were not used in the determination of the magnetic field strength. The feasibility of longitudinal magnetic field

¹ The spectropolarimetric capabilities of FORS 1 were moved to FORS 2 in 2009.

Table 1. List of O-type stars observed with FORS 2. Spectral classifications are listed according to the Galactic O Star Catalogue (Maíz-Apellániz et al. 2004).

Name	Other Identifier	V	Spectral Type
CPD–28 2561	SAO 174826	10.09	O6.5 fp
CPD–47 2963	SAO 220701	8.45	O4 III (f)
CPD–58 2620	CD–58 3529	9.27	O6.5 V ((f))
HDE 303311	CPD–58 2652	8.98	O5 V
HD 93129B	CPD–58 2618B	7.16	O3.5 V ((f+))
HD 93204	CPD–59 2584	8.48	O5 V ((f))
HDE 303308	CPD–59 2623	8.14	O4 V ((f+))
HD 93403	CPD–58 2680	7.30	O5 III (f) var sec
HD 93843	CPD–59 2732	7.33	O5 III(f) var
HD 105056	GS Mus	7.34	ON9.7 Ia e
HD 115455	CPD–61 3575	7.97	O7.5 III ((f))
HD 120521	CPD–57 6339	8.56	O8 Ib (f)
HD 123590	CPD–61 4382	7.62	O7/8
HD 125206	CPD–60 5298	7.94	O9.5 IV: (n)
HD 125241	CPD–60 5300	8.31	O8.5 Ib (f)
HD 130298	CPD–55 6191	9.29	O6.5 III (n)(f)
HD 148937	CPD–47 7765	6.77	O6.5 f?p
HD 328856	CPD–46 8218	8.50	O9.5III*
HD 152233	CPD–41 7718	6.59	O6 III: (f)p
HD 152247	CPD–41 7732	7.20	O9.5 II-III
HD 152249	HR 6263	6.47	OC9.5 Iab
HD 153426	CPD–38 6624	7.49	O9 II-III
HD 153919	V884 Sco	6.53	O6.5 Ia f+
HD 154368	V1074 Sco	6.18	O9.5 Iab
HD 154643	CPD–34 6733	7.15	O9.5 III
HD 154811	CPD–46 8416	6.93	OC9.7 Iab
HD 156154	CPD–35 6916	8.04	O8 Iab (f)
HD 156212	CPD–27 5605	7.95	O9.7 Iab
HD 165319	BD–14 4880	7.93	O9.5Iab*
HD 315033	CPD–24 6163	8.90	B3*
HD 168075	BD–13 4925	8.73	O6 V ((f))
HD 168112	BD–12 4988	8.55	O5 III (f)
HD 169582	BD–09 4729	8.70	O6 I f
HD 171589	BD–14 5131	8.21	O7 II (f)
HD 175754	BD–19 5242	7.02	O8 II ((f))
HD 187474	V3961 Sgr	5.32	Ap*

Notes: * indicates Spectral Types taken from SIMBAD.

measurements in massive stars using low-resolution spectropolarimetric observations was demonstrated by previous studies of O and B-type stars (e.g., Hubrig et al. 2006, 2008, 2009a, 2011b). To check that the instrument was functioning properly, we observed the magnetic Ap star HD 187474, which has a well studied longitudinal magnetic field, during the night of May 23 at rotation phase 0.66. HD 187474 has a rotation period of 6.4 yr and a longitudinal magnetic field ranging roughly from -2 kG to 2 kG . The measured value of the magnetic field, $\langle B_z \rangle_{\text{all}} = -1249 \pm 47 \text{ G}$, fits very well to the observations at the same phase presented by Landstreet & Mathys (2000).

Although we were granted four nights for our survey, due to unfavorable weather conditions (snow and high humidity) only four stars could be observed during the first night, 14 stars during the second night, none during the third night, and 23 during the last night. Most of the targets were observed only once. The exceptions were the stars HD 328856, HD 153426, and HD 168112, which we were able to observe twice. HD 148937 was observed three times to assess the magnetic field variability over the rotation cycle.

Table 2. Longitudinal magnetic fields measured with FORS2 in the studied sample. All quoted errors are 1σ uncertainties.

Name	MJD	$\langle B_z \rangle_{\text{all}}$ [G]	$\langle B_z \rangle_{\text{hydr}}$ [G]	Comment
CPD–28 2561	55338.969	-381 ± 122	-534 ± 167	ND
CPD–47 2963	55337.094	-190 ± 62	-154 ± 96	ND
CPD–58 2620	55339.020	-53 ± 71	-66 ± 88	
HDE 303311	55337.131	-56 ± 40	-19 ± 61	
HD 93129B	55337.179	-49 ± 44	-79 ± 78	
HD 93204	55339.053	22 ± 46	16 ± 66	
HDE 303308	55339.078	122 ± 54	137 ± 96	
HD 93403	55339.116	39 ± 41	-15 ± 88	
HD 93843	55339.099	-157 ± 42	-173 ± 56	ND
HD 105056	55337.208	-93 ± 48	-156 ± 63	
HD 115455	55337.230	4 ± 47	13 ± 64	
HD 120521	55337.257	25 ± 44	-3 ± 62	
HD 123590	55339.133	19 ± 42	56 ± 70	
HD 125206	55337.307	12 ± 64	8 ± 91	
HD 125241	55339.188	24 ± 39	16 ± 72	
HD 130298	55339.159	113 ± 38	193 ± 62	ND
HD 148937	55336.307	-297 ± 62	-293 ± 85	CD
	55337.285	-204 ± 71	-225 ± 103	
	55339.206	-290 ± 85	-389 ± 129	
HD 328856	55336.370	-173 ± 53	-155 ± 65	ND
	55339.223	-149 ± 48	-75 ± 72	
HD 152233	55339.289	-74 ± 52	-76 ± 74	
HD 152247	55339.261	34 ± 52	86 ± 64	
HD 152249	55339.275	-22 ± 51	-15 ± 72	
HD 153426	55336.338	-27 ± 53	-10 ± 62	
	55339.246	-171 ± 55	-275 ± 70	ND
HD 153919	55337.341	-213 ± 68	-119 ± 95	ND
HD 154368	55339.324	-74 ± 38	-77 ± 63	
HD 154643	55339.340	110 ± 34	121 ± 52	ND
HD 154811	55337.327	91 ± 39	59 ± 64	
HD 156154	55337.358	-118 ± 38	-167 ± 54	ND
HD 156212	55337.377	-104 ± 42	-51 ± 63	
HD 165319	55339.306	-44 ± 48	-38 ± 74	
HD 315033	55337.402	-41 ± 41	-36 ± 52	
HD 168075	55339.389	17 ± 42	3 ± 65	
HD 168112	55336.418	-74 ± 53	-66 ± 74	
	55339.362	112 ± 62	40 ± 96	
HD 169582	55339.415	-87 ± 58	-124 ± 88	
HD 171589	55339.445	-62 ± 140	-119 ± 180	
HD 175754	55337.426	82 ± 67	66 ± 77	
HD 187474	55339.433	-1249 ± 22	-1253 ± 32	Ap star

Apart from this star, which has a rotation period of seven days (Nazé et al. 2008), no exact rotation periods are known for the other stars in our sample.

The results of our magnetic field measurements are presented in Table 2. In the first two columns, we provide the star names and the modified Julian dates at the middle of the exposures. In Cols. 3 and 4 we present the longitudinal magnetic field $\langle B_z \rangle_{\text{all}}$ using the whole spectrum and the longitudinal magnetic field $\langle B_z \rangle_{\text{hyd}}$ using only the hydrogen lines. All quoted errors are 1σ uncertainties. In Col. 5, we identify new detections by ND and in the case of HD 148937 the confirmed detection is marked by CD.

Ten stars of our sample, CPD–28 2561, CPD–47 2963, HD 93843, HD 130298, HD 148937, HD 328856, HD 153426, HD 153919, HD 154643, and HD 156154, show evidence for the presence of a magnetic field.

Importantly, the strongest magnetic fields are detected in both Of?p stars CPD–28 2561 and HD 148937. Walborn (1973) introduced the Of?p category for massive O stars displaying recurrent spectral variations in certain spectral

lines, sharp emission or P Cygni profiles in He I and the Balmer lines, and strong C III emission lines around 4650 Å. Only five Galactic Of?p stars are presently known: HD 108, NGC 1624-2, CPD–28 2561, HD 148937, and HD 191612 (Walborn et al. 2010). Our observations of CPD–28 2561 reveal a magnetic field at 3.1σ level using the whole spectrum and at 3.2σ level using Balmer lines. The study of radial velocity variation of Levato et al. (1988) indicated the presence of variability of a few emission lines with a probable period of 17 days. Walborn et al. (2010) report that CPD–28 2561 undergoes extreme spectral transformations very similar to those of HD 191612, on a timescale of weeks, exhibiting variable emission intensity of the C III $\lambda\lambda$ 4647-4650-4652 triplet. The detection of a mean longitudinal magnetic field $\langle B_z \rangle = -254 \pm 81$ G in the Of?p star HD 148937 using FORS1 at the VLT was previously reported by Hubrig et al. (2008). An extensive multiwavelength study of HD 148937 was carried out by Nazé et al. (2008), who detected small-scale variations of He II 4686 and the Balmer lines with a period of 7 days and an over-

abundance of nitrogen by a factor of 4 compared to the Sun. The periodicity of spectral variations in hydrogen and He lines was re-confirmed using additional higher resolution spectroscopic material indicating similarity to the other Of?p stars HD 108 and HD 191612 (Naze et al. 2010). Our spectropolarimetric observations of this star indicate that the magnetic field is variable, but due to the low number of measurements it is not possible to verify the period deduced from spectroscopic observations. The magnetic field of this star was observed at 4.8σ , 2.9σ , and 3.4σ levels on three different nights using all absorption lines.

The remaining three Of?p stars are located in the Northern hemisphere and cannot be reached with FORS 2 at the VLT. To study magnetic fields in HD 108 and HD 191612 we recently used polarimetric spectra obtained with the SOFIN spectrograph installed at the Nordic Optical Telescope (Hubrig et al. 2010). As a result, we detected a longitudinal magnetic field $\langle B_z \rangle = -168 \pm 35$ G in the Of?p star HD 108, which is in agreement with the longitudinal magnetic field measurement of the order of -150 G recently reported by Martins et al. (2010). For the star HD 191612 with a rotation period of 537.6 d (Howarth et al. 2007) we measured a longitudinal magnetic field $\langle B_z \rangle = 450 \pm 153$ G at rotation phase 0.43 (Hubrig et al. 2010). The only previously published magnetic field measurement for this star showed a negative longitudinal magnetic field $\langle B_z \rangle = -220 \pm 38$ G at rotation phase 0.24 (Donati et al. 2006), indicating a change of polarity over ~ 100 days. No attempt has yet been made to measure the magnetic field of NGC 1624-2. Clearly the recent results of magnetic field measurements in Of?p stars imply a tight relation between the observed properties of the Of?p star group and the presence of a magnetic field.

For the star CPD-472963 we achieved a 3.1σ detection using all absorption lines. According to Walborn et al. (2010) this star belongs to the Ofc category, which consists of normal spectra with C III $\lambda\lambda$ 4647-4650-4652 emission lines of comparable intensity to those of the Of defining lines N III $\lambda\lambda$ 4634-4640-4642. The authors indicate that the Ofc phenomenon occurs primarily in certain associations and young clusters. However, the available kinematic and photometric data do not indicate cluster or association membership for CPD-472963. The origin of the magnetic field in this star is probably different compared to that of other magnetic O-type stars, as non-thermal radio emission, which is frequently observed in binary systems with colliding winds, was detected by Benaglia et al. (2001). On the other hand, the membership of CPD-472963 in a binary or multiple system has not been investigated yet. The authors suggest that the non-thermal radiation from this star possibly comes from strong shocks in the wind itself and/or in the wind colliding region if the star has a massive early-type companion. Both optical and radio observations reveal the presence of a second source separated by $5''$.

According to Walborn et al. (2010), also the star HD 93843, with a 3.7σ detection achieved using all absorption lines, belongs to the Ofc category. Prinja et al. (1998) monitored the stellar wind of this star using IUE time series. They identified systematic changes in the absorption troughs of the Si IV and N V resonance lines with a repeatability of wind structures with a period of 7.1 days. Noteworthy, the authors suggest the presence of a magnetic field as one of the possible mechanisms to explain the cyclical wind perturbation. On the other hand, three other

stars of the Ofc category included in our survey, HD 93204, HDE 303308, and HD 93403, do not show the presence of a magnetic field at a 3σ level.

The star HD 130298 with a longitudinal magnetic field observed at 3.1σ level using the Balmer lines, is known as an object with a bow shock. Noriega-Crespo et al. (1997) used ISSA/IRAS archival spectra to identify stars surrounded by extended infrared emission at $60\mu\text{m}$, which is a signature of wind bow shocks. The bow shocks are usually associated with runaway early-type stars with typical wind velocities of $500\text{--}3000\text{ km s}^{-1}$ and mass loss rates $\sim 10^{-5} - 10^{-6} M_{\odot} \text{ yr}^{-1}$ (see e.g. Puls et al. 1996).

The two stars HD 328856 and HD 153426, both with magnetic field detections, were observed on two different nights, namely the first and the fourth night of our observing run. For HD 328856 we obtained on these nights 3.3σ and 3.1σ level detections, respectively, using all absorption lines. Based on the photometric membership probability, this star is a member of the compact open cluster Hogg 22 in the Ara region at an age of 5 Myr and a distance of about 1300 pc (see Sect. 3). On the other hand, its proper motions indicate that HD 328856 is not fully co-moving with the other cluster members, deviating from the cluster mean proper motion by $\sim 2\sigma$ (for more details on membership probabilities see Kharchenko et al. 2004). The observations of the star HD 153426 revealed the presence of a mean longitudinal magnetic field at the 3.9σ level using Balmer lines on the fourth night. The non-detection of the magnetic field for HD 153426 on the first observing night can probably be explained by the strong dependence of the longitudinal magnetic field on the rotational aspect. HD 153426 is a double-lined spectroscopic binary with unknown orbit parameters and was considered by de Wit et al. (2005) as a star in a newly detected cluster. Using the best presently available kinematic data on young open clusters, Schilbach & Röser (2008) suggested that HD 153426 was ejected from the cluster Hogg 22. Their back-tracing procedure indicates that the encounter time for HD 153426, i.e. the time when the star was ejected, is about 8.1 Myr, while the age of the cluster Hogg 22 is only 5 Myr.

The star HD 153919 was observed only once, revealing the presence of a mean longitudinal magnetic field at the 3.9σ level, using all absorption lines. The study of Ankay et al. (2001) suggested that this star is a runaway X-ray binary, ejected from the OB association Sco OB1 about 2 Myr ago due to the supernova of 4U1700-37's progenitor. They considered this star as a companion to 4U1700-37, most likely a neutron star powered by wind accretion (e.g., Jones et al. 1973). Since 4U1700-37 is a candidate for a low-mass black hole (Brown et al. 1996), this system can be similar to the optical component (the O9.7 Iab supergiant) in the system Cyg X-1, for which the presence of a variable weak magnetic field was recently detected using a FORS 1 spectropolarimetric time series over the orbital period of 5.6 days (Karitskaya et al. 2010). Schilbach & Röser (2008) identified the origin of this field star in the cluster NGC 6231 (the open cluster inside Sco OB1) at an age of about 6.5 Myr and their back-tracing procedure indicates that the star was ejected from the cluster 1.1 Myr ago.

The longitudinal magnetic field for the star HD 154643 was observed at 3.2σ level using all absorption lines. De Wit et al. (2005) characterise this star as a candidate runaway star associated with the young cluster Bochum 13. However, Schilbach & Röser (2008) identified the origin of this field

Table 3. Probable members in open clusters and associations.

Object name	ASCC number	P_{kin} [%]	P_{phot} [%]	Cluster	dist [pc]	$\log t$ [yr]	$(\mu_X, \mu_Y, \sigma_\mu)^{\text{star}}$ [mas/yr]	$(\mu_X, \mu_Y, \sigma_\mu)^{\text{cl}}$ [mas/yr]
HD 47839*	1021435	62	100	NGC 2264	660	6.81	-3.84, -2.50, 0.94	-2.70, -3.50, 0.25
CPD-58 2620	2232461	44	1	Trumpler 14	2753	6.67	-3.77, -2.20, 2.37	-3.91, 3.65, 0.60
HDE 303311	2232607	91	27	Trumpler 14	2753	6.67	-2.91, 2.64, 2.40	-3.91, 3.65, 0.60
HD 93129B	2232449	0	100	Trumpler 14	2753	6.67	-9.80, 10.86, 1.96	-3.91, 3.65, 0.60
HD 93204	2232588	75	0	Trumpler 16	2842	6.90	-9.33, 3.16, 1.70	-11.10, 4.02, 0.42
HDE 303308	2232708	19	0	Trumpler 16	2842	6.90	-6.44, 2.54, 1.79	-11.10, 4.02, 0.42
HD 105056	2396832	12	100	ASCC 69	1000	7.91	-4.56, -1.81, 0.87	7.91, -7.52, 0.41
HD 115455	2338543	90	72	Stock 16	1640	6.78	-2.56, 1.08, 2.02	-3.31, -0.03, 0.66
HD 120521	2256830	0	100	Platais 10	246	8.20	-5.55, 0.40, 1.39	-29.10, -10.73, 0.40
HD 123590	2345821	83	83	ASCC 77	2200	6.99	-4.96, -2.45, 0.99	-4.59, -1.49, 0.47
HD 328856	2074567	22	100	Hogg 22	1297	6.70	0.78, -1.37, 1.26	-0.84, -4.39, 0.43
HD 135240*	2351294	61	100	ASCC 79	800	6.86	-3.00, -2.65, 0.85	-2.67, -4.10, 0.44
HD 135591*	2351425	67	100	ASCC 79	800	6.86	-3.00, -2.65, 0.85	-2.67, -4.10, 0.44
HD 152233	1974789	13	100	NGC 6231	1250	6.81	-4.11, -8.05, 2.40	-0.39, -1.99, 0.46
HD 152247	1974811	95	100	NGC 6231	1250	6.81	-0.55, -1.14, 1.91	-0.39, -1.99, 0.46
HD 152249	1974812	84	100	NGC 6231	1250	6.81	0.13, -0.81, 1.36	-0.39, -1.99, 0.46
HD 153919	1877055	0	100	NGC 6281	494	8.51	1.42, 4.84, 1.04	-2.96, -3.75, 0.23
HD 154368	1877523	88	100	ASCC 88	1900	7.17	3.80, -2.08, 1.05	2.89, -2.00, 0.32
<i>HD 155806*</i>	1783567	98	100	Sco OB4	1100	6.82	0.33, -2.02, 0.80	0.46, -2.22, 0.15
<i>HD 156154</i>	1878846	82	100	Bochum 13	1077	7.08	-0.88, -2.53, 1.64	-0.28, -1.20, 0.30
<i>HD 164794*</i>	1593528	69	100	NGC 6530	1322	6.67	1.92, -0.40, 1.09	2.01, -1.81, 0.51
HD 315033	1593655	97	100	NGC 6530	1322	6.67	2.33, -2.51, 2.09	2.01, -1.81, 0.51
HD 167263*	1595683	97	100	Sgr OB7	1860	6.64	1.56, -1.53, 1.19	1.80, -1.20, 0.26
HD 168075	1407400	73	100	NGC 6611	1719	6.72	2.81, -1.14, 1.18	1.60, -0.35, 0.41
HD 168112	1407419	73	19	NGC 6604	1696	6.64	1.67, -2.20, 1.69	-0.36, -2.68, 0.31
HD 153426	1876676	0	0	Hogg 22	1297	6.70(6.91)	-0.21, -0.09, 1.38	-0.84, -4.39, 0.43
HD 154643	1877779	0	0	ASCC 88	1900	7.17(6.15)	3.66, -1.71, 1.55	2.89, -2.00, 0.32
HD 169582	1323022	0	0	NGC 6604	1696	6.64(6.74)	-0.40, -1.63, 1.14	-0.36, -2.68, 0.31
HD 171589	1409170	0	0	NGC 6618	1814	7.78(6.52)	5.10, 3.74, 1.66	3.04, 0.33, 0.92
HD 175754	1503597	0	0	ASCC 93	2500	7.72(7.01)	1.49, 2.15, 1.01	-2.18, -1.80, 0.55

Notes. For each star we give in the first two columns the object name and the corresponding catalogue number. The kinematic and photometric probabilities for cluster membership presented in Cols. 3 und 4 were calculated according to the procedures described by Kharchenko et al. (2004). The cluster names, the distances and the ages are presented in the next three columns. The proper motions and their errors for stars and clusters are given in the last columns. All cluster data are taken from Kharchenko et al. (2005a, 2005b). The five stars at the end of the table are currently not members of the indicated clusters, and were ejected from them t_{enc} years ago (Schilbach, Roeser 2008). The encounter times t_{enc} of these stars are given in brackets. The three stars with magnetic fields that are most probable cluster members are presented in italics.

star in the cluster ASCC 88 at an age of about 14.8 Myr, ejected 1.4 Myr ago.

The star HD 156154, for which we achieved a 3.1σ detection using all absorption lines, seems to be the only star with a high cluster membership probability among the presented O-type stars with detected magnetic fields (see Sect. 3). According to kinematic and photometric criteria belongs this star to the open cluster Bochum 13 at an age of 12 Myr and a distance of about 1 kpc.

3. Discussion

A lot of effort has been put into the research of massive stars in recent years in order to properly model the effects of rotation, stellar winds, and surface chemical composition. However, possible paths for the formation of magnetic O-type stars were not analysed yet with modern theories for the evolution of single and binary stars. Clearly, the number of massive stars with detected magnetic fields is still small, and the available data are insufficient to prove statistically whether magnetic fields in massive stars are ubiquitous or appear in specific stars with certain stellar parameters and in a special environment. On the other hand, the observations of magnetic fields in massive stars accumulated over

the last few years can be used to preliminarily constrain the conditions which enable the appearance of magnetic fields and give first trends about their occurrence and field strength distribution.

Since no longitudinal magnetic fields stronger than 300 G were detected in our study (apart from the rather large field in the Of?p star CPD-28 2561), we confirm our previous conclusion (Hubrig et al. 2008) that large-scale, dipole-like, magnetic fields with polar field strengths higher than 1 kG are not widespread among O type stars. Our study presents the results of a magnetic field survey in 36 massive stars. Among them, 19 stars can be related to open clusters and associations at different age. The data on the cluster membership of these probable cluster O-type stars are presented in Table 3. To increase the significance of our statistic assessment, we present in the same table the data for an additional six probable cluster O-type stars, which have been studied during the last years by Hubrig et al. (2008, 2009b, 2011a), marked for convenience by an asterisk. As database for the compilation of Table 3 we used the All-sky Compiled Catalogue of 2.5 million stars (ASCC-2.5, 3rd version) of Kharchenko & Roeser (2009). We note that for the calculation of kinematic membership proba-




Fig. 1. The positions of the stars studied for cluster membership in the colour-magnitude diagram. Different symbols indicate stars with different membership probabilities: Squares stand for stars with cluster membership probability larger than 60%, circles for stars with a probability between 14% and 60%, and triangles for membership probability between 1% and 14%. Non-members and runaway stars are marked by dots and crosses, respectively. One runaway star, HD 171589, does not appear in this figure, as its colour and magnitude do not fit the presented parameter space (see Table 4). The three stars with magnetic fields, HD 155806, HD 156154, and HD 164794, with high cluster membership probabilities are denoted by filled squares. Isochrones for $\log t = 6.6, 6.8, 7.0,$ and 7.2 are presented by solid lines and the zero-age main sequence is shown as a dashed line.

bility only proper motions were used. According to Dias et al. (2002; Version 3.1 (24/11/2010)), one of the previously studied O-type stars, the star HD 152408 (Hubrig et al. 2008), is projected on the cluster Collinder 316, which presents a large group of bright stars superposed on Trumpler 24 at the age $\log t = 6.92$. We have not included this star in Table 3, as no membership criteria are discussed in this work. According to Mason et al. (1998) and Pourbaix et al. (2004)², among the stars presented in Table 3, six stars, HD 47839, HD 135240, HD 152233, HD 153919, HD 154368, and HD 167263, are members of spectroscopic binary systems, with orbital periods between 3.4 and 9247 days.

From the inspection of kinematic and photometric membership probabilities, and following the membership criteria described by Kharchenko et al. (2004), five stars in Table 3, CPD-58 2620, HD 93129B, HDE 303308, HD 105056, HD 152233, have a rather low cluster membership probability. Two other stars, HD 120521 and HD 153919, are not kinematic members of the oldest clusters Platais 10 and NGC 6281, respectively, and should be regarded as field stars projected against the clusters by chance. Among the remaining stars, only in three stars, HD 155806, HD 156154, and HD 164794, weak magnetic fields have been detected (more details on the kinematic study of HD 155806 and HD 164794 can be found in Hubrig et al. 2011a). In Fig. 1 we display the positions of the stars

² <http://sb9.astro.ulb.ac.be/mainform.cgi>

massive stars




Fig. 2. The age distribution of studied probable cluster members. The three stars with magnetic fields, HD 155806, HD 156154, and HD 164794, are denoted by horizontal lines.

listed in Table 3 in the colour-magnitude diagram. The absolute magnitudes M_V and intrinsic colours $(B - V)_0$ were calculated using cluster distances and $E(B - V)$ values presented by Kharchenko et al. (2005a, 2005b). The theoretical isochrones were calculated by the Padova group (Girardi et al. 2002) and the values for the zero-age main sequence were retrieved from Schmidt-Kaler (1982). The absolute magnitudes and colours are presented in Table 4. The uncertainty in the distance modulus is assumed to be 0.2. The positions of the magnetic stars with high cluster membership probabilities, HD 156154, HD 155806, and HD 164794, do not reveal any sign of a specific distribution, which would hint at the origin of their magnetic fields at a certain evolutionary age.

In Fig. 2 we present the age distribution of the most probable cluster members. While the age of HD 155806 and HD 164794 is similar to the bulk of the studied cluster O-type stars, the star HD 156154 is somewhat older at an age of ~ 12 Myr. HD 155806 is classified as an Oe star, possibly representing the higher mass analogues of classical Be stars (e.g. Walborn 1973). Only six members are suggested to belong to this group of stars (e.g. Negueruela et al. 2004). The star HD 164794 is a spectroscopic double-lined system with an orbital period of 2.4 yr, known as emitting non-thermal radio-emission, probably associated with colliding winds (Nazé et al. 2010). No specific information can be found in the literature about the luminous supergiant HD 156154.

The available observations seem to indicate that the presence of a magnetic field is more frequently detected in field stars than in stars belonging to clusters or associations. It is generally accepted that the majority of massive stars form in star clusters and associations, and studies of kinematical properties of the massive star field population indicate that a major part of these stars can be traced back to their parent open clusters or associations (e.g. Schilbach & Roeser 2008). Pflamm-Altenburg & Kroupa (2010) recently

Table 4. Absolute magnitudes and intrinsic colours of stars studied for the cluster membership.

Object name	M_V	$(B - V)_0$
HD 47839	-4.557	-0.266±0.007
CPD-58 2620	-4.239	-0.436±0.040
HDE 303311	-4.632	-0.370±0.030
HD 93129B	-6.315	-0.285±0.020
HD 93204	-5.381	-0.586±0.026
HDE 303308	-5.756	-0.433±0.027
HD 105056	-3.145	-0.143±0.011
HD 115455	-4.563	-0.329±0.011
HD 120521	-4.011	-0.277±0.019
HD 123590	-5.017	-0.187±0.011
HD 328856	-4.066	-0.282±0.038
HD 135240	-4.942	-0.253±0.012
HD 135591	-4.576	-0.261±0.004
HD 152233	-5.317	-0.320±0.017
HD 152247	-4.656	-0.276±0.018
HD 152249	-5.471	-0.259±0.019
HD 153919	-2.406	0.090±0.013
HD 154368	-6.781	-0.047±0.006
HD 155806	-5.214	-0.254±0.006
HD 156154	-4.562	-0.242±0.020
HD 164794	-5.603	-0.293±0.007
HD 315033	-2.956	-0.203±0.030
HD 167263	-6.125	-0.257±0.006
HD 168075	-4.566	-0.319±0.035
HD 168112	-5.545	-0.381±0.028
HD 153426	-5.109	-0.548±0.013
HD 154643	-5.775	-0.278±0.011
HD 169582	-5.382	-0.474±0.028
HD 171589	-8.036	-1.340±0.021
HD 175754	-5.838	-0.379±0.011

discussed in their work whether massive stars can form in isolation in the galactic field. According to de Wit et al. (2005), only a few per cent of all O-type stars can be considered as formed outside a cluster environment. Pflamm-Altenburg & Kroupa considered the two-step-ejection process, which presents the combination of the dynamical and the supernova ejection scenario with the result that massive field stars produced via this ejection process for the vast majority of cases cannot be traced back to their parent star clusters. These stars can be mistakenly considered as massive stars formed in isolation. While this can not be proven, the observed numbers of field O stars is consistent with this idea.

For the newly detected magnetic O-type stars, HD 153426, HD 153919, and HD 154643, Schilbach & Roeser (2008) suggested that the three stars were ejected from the clusters Hogg 22, NGC 6231, and ASCC 88, respectively. On the other hand, none of the four magnetic Of?p stars is known to belong to a cluster or an association. The study of the evolutionary state of HD 108 and HD 191612 indicates that both stars are significantly evolved (Martins et al. 2010). Our kinematic study of the Of?p star HD 148937 showed that it possesses a space velocity of 32 km s^{-1} with respect to the Galactic open cluster system, with the velocity component $U = -26$ directed opposite from the Galactic center and the velocity component $W = -13$ directed from the Galactic plane (Hubrig et al. 2011a). These rather large velocities indicate that this star can be considered as a candidate runaway star.

It is striking that the major part of previously detected magnetic O-type stars are candidate runaway stars (Hubrig et al. 2011a, 2011c). Also in the sample of O-stars with magnetic fields detected in this work, four other stars, HD 130298, HD 153426, HD 153919, and HD 154643, are mentioned in the literature as candidate runaway stars. In the past, two mechanisms were discussed to explain the existence of runaway stars: In one scenario, close multibody interactions in a dense cluster environment cause one or more stars to be scattered out of the region (e.g. Leonard & Duncan 1990). For this mechanism, runaways are ejected in dynamical three- or four-body interactions. An alternative mechanism involves a supernova explosion within a close binary, ejecting the secondary due to the conservation of momentum (Zwicky 1957; Blaauw 1961). However, none of these scenarios consider the possibility how a massive star can acquire a magnetic field during the ejection process. Clearly, these findings generate a strong motivation to carry out a kinematic study of all stars previously surveyed for magnetic fields to search for a correlation between the kinematic status and the presence of a magnetic field.

Based on the still very limited magnetic surveys in massive stars, we cannot yet answer the question if O-type stars are magnetic in certain evolutionary states and in a specific environment. Open star clusters and associations are very useful laboratories to test star formation and stellar evolution. The ages of our subsample of three stars with magnetic fields do not contradict the idea, that it is drawn from the general distribution of cluster ages. We have to keep in mind though that we have a very small number statistics. It appears that our observations are consistent with the assumption that the presence of a magnetic field can be expected in stars of different classification categories. Although it was possible to recognize a few hot Of?p magnetic stars as being peculiar on the basis of their spectral morphology, prior to their field detection (Walborn 2006), the presence of a magnetic field can also be expected in stars of other classifications. Future magnetic field measurements are urgently needed to constrain the conditions controlling the presence of magnetic fields in hot stars, and the implications of these fields on their physical parameters and evolution.

Acknowledgements. NVK and AEP thank for support by DFG grant RO 528/10-1. AEP acknowledges support of the RFBR grant 10-02-91338.

References

- Alves, F. O., Franco, G. A. P., & Girart, J. M. 2008, *A&A*, 486, L13
Ankay, A., Kaper, L., de Bruijne, J. H. J., et al. 2001, *A&A*, 370, 170
Benaglia, P., Cappa, C. E., & Koribalski, B. S. 2001, *A&A*, 372, 952
Blaauw, A. 1961, *Bull. Astron. Inst. Netherlands*, 15, 265
Bonanno, A., & Urpin, V. 2008, *MNRAS*, 388, 1679
Brown, G. E., Weingartner, J. C., & Wijers, R. A. M. J. 1996, *ApJ*, 463, 297
de Wit, W. J., Testi, L., Palla, F., & Zinnecker, H. 2005, *A&A*, 437, 247
Dias, W. S., Alessi, B. S., Moitinho, A., & Lépine, J. R. D. 2002, *A&A*, 389, 871
Donati, J.-F., Howarth, I. D., Bouret, J.-C., et al. 2006, *MNRAS*, 365, L6
Girardi, L., Bertelli, G., Bressan, A., et al. 2002, *A&A*, 391, 195
Gómez, G. C., & Cox, D. P. 2004, *ApJ*, 615, 744
Howarth, I. D., Walborn, N. R., Lennon, D. J., et al. 2007, *MNRAS*, 381, 433
Hubrig, S., Kurtz, D. W., Bagnulo, S., et al. 2004a, *A&A*, 415, 661
Hubrig, S., Szeifert, T., Schöller, M., et al. 2004b, *A&A*, 415, 685

- Hubrig, S., Briquet, M., Schöller, M., et al. 2006, *MNRAS*, 369, L61
- Hubrig, S., Schöller, M., Schnerr, R. S., et al. 2008, *A&A*, 490, 793
- Hubrig, S., Schöller, M., Savanov, I., et al. 2009a, *AN*, 330, 708
- Hubrig, S., Stelzer, B., Schöller, M., et al. 2009b, *A&A*, 502, 283
- Hubrig, S., Ilyin, I., & Schöller, M. 2010, *AN*, 331, 781
- Hubrig, S., Kharchenko, N. V., & Schöller, M. 2011a, *AN*, 332, 65
- Hubrig, S., Schöller, M., Ilyin, I. et al. 2011b, *ApJL*, 726, L5
- Hubrig, S., Oskinova, L.M., Schöller, M. 2011c, *AN*, 332, 147, also arXiv:1101.5500
- Jones, C., Forman, W., Tananbaum, H., et al. 1973, *ApJ*, 181, L43
- Karitskaya, E. A., Bochkarev, N. G., Hubrig, S., et al. 2010, *IBVS*, 5950, 1
- Kharchenko, N. V., Piskunov, A. E., Röser, S., et al. 2004, *AN*, 325, 740
- Kharchenko, N. V., Piskunov, A. E., Röser, S., Schilbach, E., & Scholz, R.-D. 2005a, *A&A*, 438, 1163
- Kharchenko, N. V., Piskunov, A. E., Röser, S., Schilbach, E., & Scholz, R.-D. 2005b, *A&A*, 440, 403
- Kharchenko, N. V., & Roeser, S. 2009, “All-sky Compiled Catalogue of 2.5 million stars”
- Landstreet, J. D., & Mathys, G. 2000, *A&A*, 359, 213
- Leonard, P. J. T., & Duncan, M. J. 1990, *AJ*, 99, 608
- Levato, H., Morrell, N., Garcia, B., & Malaroda, S. 1988, *ApJS*, 68, 319
- Maíz-Apellániz, J., Walborn, N. R., Galué, H. Á., & Wei, L. H. 2004, *ApJS*, 151, 103
- Martins, F., Donati, J.-F., Marcolino, W. L. F., et al. 2010, *MNRAS*, 407, 1423
- Mason, B. D., Gies, D. R., Hartkopf, W. I., et al. 1998, *AJ*, 115, 821
- Nazé, Y., Walborn, N. R., Rauw, G., et al. 2008, *AJ*, 135, 1946
- Nazé, Y., ud-Doula, A., Spano, M., et al. 2010, *A&A*, 520, A59
- Negueruela, I., Steele, I. A., & Bernabeu, G. 2004, *AN*, 325, 749
- Noriega-Crespo, A., van Buren, D., & Dgani, R. 1997, *AJ*, 113, 780
- Pflamm-Altenburg, J., & Kroupa, P. 2010, *MNRAS*, 404, 1564
- Pourbaix, D., Tokovinin, A. A., Batten, A. H., et al. 2004, *A&A*, 424, 727
- Prinja, R. K., Massa, D., Howarth, I. D., & Fullerton, A. W. 1998, *MNRAS*, 301, 926
- Puls, J., Kudritzki, R.-P., Herrero, A., et al. 1996, *A&A*, 305, 171
- Schilbach, E., & Röser, S. 2008, *A&A*, 489, 105
- Schmidt-Kaler, Th. 1982, In: *Landolt-Börnstein Numerical Data and Functional Relationships in Science and Technology, New Series, Group IV*, eds. K. Schafer & H. H. Voigt, Springer-Verlag Press: Berlin-Heidelberg, New York
- Walborn, N. R. 1973, *AJ*, 78, 1067
- Walborn, N. R. 2006, In: “The Ultraviolet Universe: Stars from Birth to Death”, 26th meeting of the IAU, Joint Discussion 4, 16-17 August 2006, Prague, Czech Republic, JD04, #19
- Walborn, N. R., Sota, A., Maíz Apellániz, J., et al. 2010, *ApJ*, 711, L143
- Zwicky, F. 1957, “Morphological astronomy”, Springer, Berlin