A DIAZ-METCALF TYPE INEQUALITY FOR POSITIVE LINEAR MAPS AND ITS APPLICATIONS

MOHAMMAD SAL MOSLEHIAN¹, RITSUO NAKAMOTO² AND YUKI SEO³

ABSTRACT. We present a Diaz-Metcalf type operator inequality as a reverse Cauchy-Schwarz inequality and then apply it to get the operator versions of Pólya-Szegö's, Greub-Rheinboldt's, Kantorovich's, Shisha-Mond's, Schweitzer's, Cassels' and Klamkin-McLenaghan's inequalities via a unified approach. We also give some operator Grüss type inequalities and an operator Ozeki-Izumino-Mori-Seo type inequality. Several applications are concluded as well.

1. Introduction

The Cauchy–Schwarz inequality plays an essential role in mathematical inequalities and its applications. In a semi-inner product space $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ the Cauchy–Schwarz inequality reads as follows

$$|\langle x, y \rangle| \le \langle x, x \rangle^{1/2} \langle y, y \rangle^{1/2} \qquad (x, y \in \mathscr{H}).$$

There are interesting generalizations of the Cauchy–Schwarz inequality in various frameworks, e.g. finite sums, integrals, isotone functionals, inner product spaces, C^* -algebras and Hilbert C^* -modules; see [5, 6, 7, 13, 17, 20, 9] and references therein. There are several reverses of the Cauchy–Schwarz inequality in the literature: Diaz–Metcalf's, Pólya–Szegö's, Greub–Rheinboldt's, Kantorovich's, Shisha–Mond's, Ozeki–Izumino–Mori–Seo's, Schweitzer's, Cassels' and Klamkin–McLenaghan's inequalities.

Inspired by the work of J.B. Diaz and F.T. Metcalf [4], we present several reverse Cauchy–Schwarz type inequalities for positive linear maps. We give a unified treatment of some reverse inequalities of the classical Cauchy–Schwarz type for positive linear maps.

Throughout the paper $\mathbb{B}(\mathcal{H})$ stands for the algebra of all bounded linear operators acting on a Hilbert space \mathcal{H} . We simply denote by α the scalar multiple αI of the identity operator $I \in \mathbb{B}(\mathcal{H})$. For self-adjoint operators A, B the partially

²⁰¹⁰ Mathematics Subject Classification. Primary 46L08; Secondary 26D15, 46L05, 47A30, 47A63

Key words and phrases. Diaz-Metcalf type inequality, reverse Cauchy-Schwarz inequality, positive map, Ozeki-Izumino-Mori-Seo inequality, operator inequality.

ordered relation $B \leq A$ means that $\langle B\xi, \xi \rangle \leq \langle A\xi, \xi \rangle$ for all $\xi \in \mathcal{H}$. In particular, if $0 \leq A$, then A is called positive. If A is a positive invertible operator, then we write 0 < A. A linear map $\Phi : \mathscr{A} \to \mathscr{B}$ between C^* -algebras is said to be positive if $\Phi(A)$ is positive whenever so is A. We say that Φ is unital if Φ preserves the identity. The reader is referred to [9] for undefined notations and terminologies.

2. OPERATOR DIAZ-METCALF TYPE INEQUALITY

We start this section with our main result. Recall that the geometric operator mean $A \sharp B$ for positive operators $A, B \in \mathbb{B}(\mathcal{H})$ is defined by

$$A \sharp B = A^{\frac{1}{2}} \left(A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \right)^{\frac{1}{2}} A^{\frac{1}{2}}$$

if 0 < A.

Theorem 2.1. Let $A, B \in \mathbb{B}(\mathcal{H})$ be positive invertible operators and $\Phi : \mathbb{B}(\mathcal{H}) \to \mathbb{B}(\mathcal{H})$ be a positive linear map.

- (i) If $m^2A \leq B \leq M^2A$ for some positive real numbers m < M, then the following inequalities hold:
 - Operator Diaz-Metcalf inequality of first type

$$Mm\Phi(A) + \Phi(B) \le (M+m)\Phi(A\sharp B);$$

• Operator Cassels inequality

$$\Phi(A)\sharp\Phi(B) \leq \frac{M+m}{2\sqrt{Mm}}\Phi(A\sharp B);$$

• Operator Klamkin–McLenaghan inequality

$$\Phi(A\sharp B)^{\frac{-1}{2}}\Phi(B)\Phi(A\sharp B)^{\frac{-1}{2}} - \Phi(A\sharp B)^{\frac{1}{2}}\Phi(A)^{-1}\Phi(A\sharp B)^{\frac{1}{2}} \le (\sqrt{M} - \sqrt{m})^2;$$

• Operator Kantorovich inequality

$$\Phi(A)\sharp\Phi(A^{-1}) \le \frac{M^2 + m^2}{2Mm}$$
.

- (ii) If $m_1^2 \leq A \leq M_1^2$ and $m_2^2 \leq B \leq M_2^2$ for some positive real numbers $m_1 < M_1$ and $m_2 < M_2$, then the following inequalities hold:
 - Operator Diaz-Metcalf inequality of second type

$$\frac{M_2 m_2}{M_1 m_1} \Phi(A) + \Phi(B) \le \left(\frac{M_2}{m_1} + \frac{m_2}{M_1}\right) \Phi(A \sharp B);$$

• Operator Pólya–Szegö inequality

$$\Phi(A)\sharp\Phi(B) \le \frac{1}{2} \left(\sqrt{\frac{M_1 M_2}{m_1 m_2}} + \sqrt{\frac{m_1 m_2}{M_1 M_2}} \right) \Phi(A\sharp B);$$

• Operator Shisha-Mond inequality

$$\Phi(A\sharp B)^{\frac{-1}{2}}\Phi(B)\Phi(A\sharp B)^{\frac{-1}{2}} - \Phi(A\sharp B)^{\frac{1}{2}}\Phi(A)^{-1}\Phi(A\sharp B)^{\frac{1}{2}} \le \left(\sqrt{\frac{M_2}{m_1}} - \sqrt{\frac{m_2}{M_1}}\right)^2;$$

• Operator Grüss type inequality

$$\Phi(A)\sharp\Phi(B) - \Phi(A\sharp B) \le \frac{\sqrt{M_1 M_2} \left(\sqrt{M_1 M_2} - \sqrt{m_1 m_2}\right)^2}{2\sqrt{m_1 m_2}} \min\left\{\frac{M_1}{m_1}, \frac{M_2}{m_2}\right\}.$$

Proof. (i) If $m^2A \leq B \leq M^2A$ for some positive real numbers m < M, then $m^2 \leq A^{\frac{-1}{2}}BA^{\frac{-1}{2}} \leq M^2$.

(ii) If $m_1^2 \leq A \leq M_1^2$ and $m_2^2 \leq B \leq M_2^2$ for some positive real numbers $m_1 < M_1$ and $m_2 < M_2$, then

$$m^2 = \frac{m_2^2}{M_1^2} \le A^{\frac{-1}{2}} B A^{\frac{-1}{2}} \le \frac{M_2^2}{m_1^2} = M^2$$
. (2.1)

In any case we then have

$$\left(M - \left(A^{\frac{-1}{2}}BA^{\frac{-1}{2}}\right)^{1/2}\right) \left(\left(A^{\frac{-1}{2}}BA^{\frac{-1}{2}}\right)^{1/2} - m\right) \ge 0,$$

whence

$$Mm + A^{\frac{-1}{2}}BA^{\frac{-1}{2}} \le (M+m)\left(A^{\frac{-1}{2}}BA^{\frac{-1}{2}}\right)^{\frac{1}{2}}.$$

Hence

$$MmA + B \le (M+m)A^{1/2} \left(A^{\frac{-1}{2}}BA^{\frac{-1}{2}}\right)^{\frac{1}{2}}A^{1/2} = (M+m)A\sharp B.$$
 (2.2)

Since Φ is a positive linear map, (2.2) yields the operator Diaz–Metcalf inequality of first type as follows:

$$Mm\Phi(A) + \Phi(B) \le (M+m)\Phi(A\sharp B). \tag{2.3}$$

In the case when (ii) holds we get the following, which is called the *operator Diaz–Metcalf inequality of second type*:

$$\frac{M_2 m_2}{M_1 m_1} \Phi(A) + \Phi(B) \le \left(\frac{M_2}{m_1} + \frac{m_2}{M_1}\right) \Phi(A \sharp B).$$

Following the strategy of [21], we apply the operator geometric–arithmetic inequality to $Mm\Phi(A)$ and $\Phi(B)$ to get:

$$\sqrt{Mm}(\Phi(A)\sharp\Phi(B)) = (Mm\Phi(A))\sharp\Phi(B) \le \frac{1}{2}(Mm\Phi(A) + \Phi(B)).$$
 (2.4)

It follows from (2.3) and (2.4) that

$$\Phi(A)\sharp\Phi(B) \le \frac{M+m}{2\sqrt{Mm}}\Phi(A\sharp B),$$

which is said to be the *operator Cassels inequality* under the assumption (i); see also [16]. Under the case (ii) we can represent it as the following inequality being called the *operator Pólya–Szegö inequality* or the *operator Greub–Rheinboldt inequality*:

$$\Phi(A)\sharp\Phi(B) \le \frac{1}{2} \left(\sqrt{\frac{M_1 M_2}{m_1 m_2}} + \sqrt{\frac{m_1 m_2}{M_1 M_2}} \right) \Phi(A\sharp B). \tag{2.5}$$

It follows from (2.5) that

$$\Phi(A)\sharp\Phi(B) - \Phi(A\sharp B) \leq \left(\frac{1}{2}\left(\sqrt{\frac{M_1M_2}{m_1m_2}} + \sqrt{\frac{m_1m_2}{M_1M_2}}\right) - 1\right)\Phi(A\sharp B)
= \frac{\left(\sqrt{M_1M_2} - \sqrt{m_1m_2}\right)^2}{2\sqrt{m_1m_2}\sqrt{M_1M_2}}\Phi(A\sharp B).$$
(2.6)

It follows from (2.1) that

$$\frac{m_2}{M_1} A \le A^{1/2} \left(A^{\frac{-1}{2}} B A^{\frac{-1}{2}} \right)^{1/2} A^{1/2} \le \frac{M_2}{m_1} A,$$

SO

$$\frac{m_1^2 m_2}{M_1} \le A \sharp B \le \frac{M_1^2 M_2}{m_1} \,. \tag{2.7}$$

Now, (2.6) and (2.7) yield that

$$\Phi(A)\sharp\Phi(B) - \Phi(A\sharp B) \le \frac{\left(\sqrt{M_1 M_2} - \sqrt{m_1 m_2}\right)^2}{2\sqrt{m_1 m_2}\sqrt{M_1 M_2}} \frac{M_1^2 M_2}{m_1}.$$

An easy symmetric argument then follows that

$$\Phi(A) \sharp \Phi(B) - \Phi(A \sharp B) \le \frac{\sqrt{M_1 M_2} \left(\sqrt{M_1 M_2} - \sqrt{m_1 m_2}\right)^2}{2\sqrt{m_1 m_2}} \min \left\{ \frac{M_1}{m_1}, \frac{M_2}{m_2} \right\} ,$$

presenting a Grüss type inequality.

If A is invertible and Φ is unital and $m_1^2 = m^2 \le A \le M^2 = M_1^2$, then by putting $m_2^2 = 1/M^2 \le B = A^{-1} \le 1/m^2 = M_2^2$ in (2.5) we get the following operator Kantorovich inequality:

$$\Phi(A)\sharp\Phi(A^{-1}) \le \frac{M^2 + m^2}{2Mm} \,. \tag{2.8}$$

It follows from (2.3) that

$$\Phi(A\sharp B)^{\frac{-1}{2}}\Phi(B)\Phi(A\sharp B)^{\frac{-1}{2}} - \Phi(A\sharp B)^{\frac{1}{2}}\Phi(A)^{-1}\Phi(A\sharp B)^{\frac{1}{2}}$$

$$\leq M + m - Mm\Phi(A\sharp B)^{\frac{-1}{2}}\Phi(A)\Phi(A\sharp B)^{\frac{-1}{2}} - \Phi(A\sharp B)^{\frac{1}{2}}\Phi(A)^{-1}\Phi(A\sharp B)^{\frac{1}{2}}$$

$$\leq M + m - 2\sqrt{Mm} - \left(\sqrt{Mm}\left(\Phi(A\sharp B)^{\frac{-1}{2}}\Phi(A)\Phi(A\sharp B)^{\frac{-1}{2}}\right)^{1/2}\right)^{1/2}$$

$$- \left(\Phi(A\sharp B)^{\frac{1}{2}}\Phi(A)^{-1}\Phi(A\sharp B)^{\frac{1}{2}}\right)^{1/2}\right)^{2}$$

$$\leq (\sqrt{M} - \sqrt{m})^{2}, \tag{2.9}$$

that is, an operator Klakmin-Mclenaghan inequality when (i) holds. Under (ii), we get the following operator Shisha-Szegö inequality from (2.9):

$$\Phi(A\sharp B)^{\frac{-1}{2}}\Phi(B)\Phi(A\sharp B)^{\frac{-1}{2}}-\Phi(A\sharp B)^{\frac{1}{2}}\Phi(A)^{-1}\Phi(A\sharp B)^{\frac{1}{2}}\leq \left(\sqrt{\frac{M_2}{m_1}}-\sqrt{\frac{m_2}{M_1}}\right)^2\;.$$

3. Applications

If (a_1, \ldots, a_n) and (b_1, \ldots, b_n) are *n*-tuples of real numbers with $0 < m_1 \le a_i \le M_1$ $(1 \le i \le n), 0 < m_2 \le b_i \le M_2$ $(1 \le i \le n)$, we can consider the positive linear map $\Phi(T) = \langle Tx, x \rangle$ on $\mathbb{B}(\mathbb{C}^n) = M_n(\mathbb{C})$ and let $A = \operatorname{diag}(a_1^2, \ldots, a_n^2)$, $B = \operatorname{diag}(b_1^2, \ldots, b_n^2)$ and $x = (1, \ldots, 1)^t$ in the operator inequalities above to get the following classical inequalities:

• Diaz–Metcalf inequality [4]

$$\sum_{k=1}^{n} b_k^2 + \frac{m_2 M_2}{m_1 M_1} \sum_{k=1}^{n} a_k^2 \le \left(\frac{M_2}{m_1} + \frac{m_2}{M_1}\right) \sum_{k=1}^{n} a_k b_k.$$

• Pólya–Szegő inequality [23]

$$\frac{\sum_{k=1}^{n} a_k^2 \sum_{k=1}^{n} b_k^2}{\left(\sum_{k=1}^{n} a_k b_k\right)^2} \le \frac{1}{4} \left(\sqrt{\frac{M_1 M_2}{m_1 m_2}} + \sqrt{\frac{m_1 m_2}{M_1 M_2}} \right)^2;$$

• Shisha–Mond inequality [24]

$$\frac{\sum_{k=1}^{n} a_k^2}{\sum_{k=1}^{n} a_k b_k} - \frac{\sum_{k=1}^{n} a_k b_k}{\sum_{k=1}^{n} b_k^2} \le \left(\sqrt{\frac{M_1}{m_2}} - \sqrt{\frac{m_1}{M_2}}\right)^2;$$

• A Grüss type inequality

$$\left(\sum_{k=1}^{n} a_k^2\right)^{1/2} \left(\sum_{k=1}^{n} b_k^2\right)^{1/2} - \sum_{k=1}^{n} a_k b_k \\
\leq \frac{\sqrt{M_1 M_2} \left(\sqrt{M_1 M_2} - \sqrt{m_1 m_2}\right)^2}{2\sqrt{m_1 m_2}} \min\left\{\frac{M_1}{m_1}, \frac{M_2}{m_2}\right\}.$$

Using the same argument with a positive *n*-tuple (a_1, \ldots, a_n) of real numbers with $0 < m \le a_i \le M$ $(1 \le i \le n)$, $x = \frac{1}{\sqrt{n}}(1, \ldots, 1)^t$, we get from Kantorovich inequality that

• Schweitzer inequality [2]

$$\left(\frac{1}{n}\sum_{i=1}^{n}a_{i}^{2}\right)\left(\frac{1}{n}\sum_{i=1}^{n}a_{i}^{-2}\right) \leq \frac{(M^{2}+m^{2})^{2}}{4M^{2}m^{2}}.$$

If (a_1, \ldots, a_n) and (b_1, \ldots, b_n) are *n*-tuples of real numbers with $0 < m \le a_i/b_i \le M$ $(1 \le i \le n)$, we can consider the positive linear map $\Phi(T) = \langle Tx, x \rangle$ on $\mathbb{B}(\mathbb{C}^n) = M_n(\mathbb{C})$ and let $A = \operatorname{diag}(a_1^2, \ldots, a_n^2)$, $B = \operatorname{diag}(b_1^2, \ldots, b_n^2)$ and $x = (\sqrt{w_1}, \ldots, \sqrt{w_n})^t$ based on the weight $\bar{\mathbf{w}} = (w_1, \ldots, w_n)$, in the operator inequalities above to get the following classical inequalities:

• Cassels inequality [25]

$$\frac{\sum_{k=1}^{n} w_k a_k^2 \sum_{k=1}^{n} w_k b_k^2}{\left(\sum_{k=1}^{n} w_k a_k b_k\right)^2} \le \frac{(M+m)^2}{4mM};$$

• Klamkin–McLenaghan inequality [14]

$$\sum_{k=1}^{n} w_k a_k^2 \sum_{k=1}^{n} w_k b_k^2 - \left(\sum_{k=1}^{n} w_k a_k b_k\right)^2 \le \left(\sqrt{M} - \sqrt{m}\right)^2 \sum_{k=1}^{n} w_k a_k b_k \sum_{k=1}^{n} w_k a_k^2.$$

Using the same argument, we obtain a weighted form of the Pólya–Szegö inequality as follows:

• Grueb–Rheinboldt inequality [10]

$$\frac{\sum_{k=1}^{n} w_k a_k^2 \sum_{k=1}^{n} w_k b_k^2}{\left(\sum_{k=1}^{n} w_k a_k b_k\right)^2} \le \frac{\left(M_1 M_2 + m_1 m_2\right)^2}{4 m_1 m_2 M_1 M_2}.$$

One can assert the integral versions of discrete results above by considering $L^2(X,\mu)$, where (X,μ) is a probability space, as a Hilbert space via $\langle h_1,h_2\rangle = \int_X h_1 \overline{h_2} d\mu$, multiplication operators $A,B \in \mathbb{B}(L^2(X,\mu))$ defined by $A(h) = f^2 h$ and $B(h) = g^2 h$ for bounded $f,g \in L^2(X,\mu)$ and a positive linear map Φ by $\Phi(T) = \int_X T(1) d\mu$ on $\mathbb{B}(L^2(X,\mu))$. For instance, let us state integral versions

of the Cassels and Klamkin–McLenaghan inequalities. These two inequalities are obtained, first for bounded positive functions $f, g \in L^2(X, \mu)$ and next for general positive functions $f, g \in L^2(X, \mu)$ as the limits of sequences of bounded positive functions.

Corollary 3.1. Let (X, μ) be a probability space and $f, g \in L^2(X, \mu)$ with $0 \le mg \le f \le Mg$ for some scalars 0 < m < M. Then

$$\int_X f^2 d\mu \int_X g^2 d\mu \le \frac{(M+m)^2}{4Mm} \left(\int_X f g d\mu \right)^2$$

and

$$\int_X f^2 d\mu \int_X g^2 d\mu - \left(\int_X fg d\mu\right)^2 \le \left(\sqrt{M} - \sqrt{m}\right)^2 \int_X fg d\mu \int_X f^2 d\mu.$$

Considering the positive linear functional $\Phi(R) = \sum_{i=1}^{n} \langle R\xi_i, \xi_i \rangle$ on $\mathbb{B}(\mathcal{H})$, where $\xi_1, \ldots, \xi_n \in \mathcal{H}$, we get the following versions of the Diaz–Metcalf and Pólya–Szegö inequalities in a Hilbert space.

Corollary 3.2. Let \mathscr{H} be a Hilbert space, let $\xi_1, \ldots, \xi_n \in \mathscr{H}$ and let $T, S \in \mathbb{B}(\mathscr{H})$ be positive operators satisfying $0 < m_1 \le T \le M_1$ and $0 < m_2 \le S \le M_2$. Then

$$\frac{M_2 m_2}{M_1 m_1} \sum_{i=1}^n \|T\xi_i\|^2 + \sum_{i=1}^n \|S\xi_i\|^2 \le \left(\frac{M_2}{m_1} + \frac{m_2}{M_1}\right) \sum_{i=1}^n \|(T^2 \sharp S^2)^{1/2} \xi_i\|^2$$

and

$$\left(\sum_{i=1}^{n} \|T\xi_i\|^2\right)^{1/2} \left(\sum_{i=1}^{n} \|S\xi_i\|^2\right)^{1/2}$$

$$\leq \frac{1}{2} \left(\sqrt{\frac{M_1 M_2}{m_1 m_2}} + \sqrt{\frac{m_1 m_2}{M_1 M_2}}\right) \sum_{i=1}^{n} \|(T^2 \sharp S^2)^{1/2} \xi_i\|^2.$$

4. A Grüss type inequality

In this section we obtain another Grüss type inequality, see also [18]. Let \mathscr{A} be a C^* -algebra and let \mathscr{B} be a C^* -subalgebra of \mathscr{A} . Following [1], a positive linear map $\Phi: \mathscr{A} \to \mathscr{B}$ is called a left multiplier if $\Phi(XY) = \Phi(X)Y$ for every $X \in \mathscr{A}, Y \in \mathscr{B}$.

The following lemma is interesting on its own right.

Lemma 4.1. Let Φ be a unital positive linear map on \mathscr{A} , $A \in \mathscr{A}$ and M, m be complex numbers such that

$$Re((M-A)^*(A-m)) \ge 0.$$
 (4.1)

Then

$$\Phi(|A|^2) - |\Phi(A)|^2 \le \frac{1}{4}|M - m|^2. \tag{4.2}$$

Proof. For any complex number $c \in \mathbb{C}$, we have

$$\Phi(|A|^2) - |\Phi(A)|^2 = \Phi(|A-c|^2) - |\Phi(A-c)|^2. \tag{4.3}$$

Since for any $T \in \mathcal{A}$ the operator equality

$$\frac{1}{4}|M-m|^2 - \left|T - \frac{M+m}{2}\right|^2 = \text{Re}\left((M-T)(T-m)^*\right)$$

holds, the condition (4.1) implies that

$$\Phi\left(\left|A - \frac{M+m}{2}\right|^2\right) \le \frac{1}{4}|M-m|^2.$$
(4.4)

Therefore, it follows from (4.3) and (4.4) that

$$\Phi(|A|^2) - |\Phi(A)|^2 \le \Phi(|A - \frac{M+m}{2}|^2)$$

$$\le \frac{1}{4}|M-m|^2.$$

Remark 4.2. If (i) Φ is a unital positive linear map and A is a normal operator or (ii) Φ is a 2-positive linear map and A is an arbitrary operator, then it follows from [3] that

$$0 \le \Phi(|A|^2) - |\Phi(A)|^2 \ . \tag{4.5}$$

Condition (4.5) is stronger than positivity and weaker than 2-positivity; see [8]. Another class of positive linear maps satisfying (4.5) are left multipliers, cf. [1, Corollary 2.4].

Lemma 4.3. Let a positive linear map $\Phi : \mathscr{A} \to \mathscr{B}$ be a unital left multiplier. Then

$$|\Phi(A^*B) - \Phi(A)^*\Phi(B)|^2 \le \|\Phi(|A|^2) - |\Phi(A)|^2 \| (\Phi(|B|^2) - |\Phi(B)|^2)$$
 (4.6)

Proof. If we put $[X,Y] := \Phi(X^*Y) - \Phi(X)^*\Phi(Y)$, then \mathscr{A} is a right pre-inner product C^* -module over \mathscr{B} , since $\Phi(X^*Y)$ is a right pre-inner product \mathscr{B} -module, see [1, Corollary 2.4]. It follows from the Cauchy–Schwarz inequality in pre-inner product C^* -modules (see [15, Proposition 1.1]) that

$$|\Phi(A^*B) - \Phi(A)^*\Phi(B)|^2 = [B, A][A, B]$$

$$\leq ||[A, A]||[B, B]|$$

$$= ||\Phi(A^*A) - \Phi(A)^*\Phi(A)|| (\Phi(B^*B) - \Phi(B)^*\Phi(B))$$

and hence (4.6) holds.

Theorem 4.4. Let a positive linear map $\Phi : \mathscr{A} \to \mathscr{B}$ be a unital left multiplier. If $M_1, m_1, M_2, m_2 \in \mathbb{C}$ and $A, B \in \mathscr{A}$ satisfy the following conditions:

$$Re(M_1 - A)^*(A - m_1) \ge 0$$
 and $Re(M_2 - B)^*(B - m_2) \ge 0$,

then

$$|\Phi(A^*B) - \Phi(A)^*\Phi(B)| \le \frac{1}{4}|M_1 - m_1||M_2 - m_2|.$$

Proof. By Löwner–Heinz theorem, we have

$$|\Phi(A^*B) - \Phi(A)^*\Phi(B)|$$

$$\leq \|\Phi(|A|^2) - |\Phi(A)|^2\|^{\frac{1}{2}} (\Phi(|B|^2) - |\Phi(B)|^2)^{\frac{1}{2}}$$
 (by Lemma 4.3)
$$\leq \frac{1}{4}|M_1 - m_1| |M_2 - m_2|$$
 (by Lemma 4.1).

5. Ozeki–Izumino–Mori–Seo type inequality

Let $a = (a_1, ..., a_n)$ and $b = (b_1, ..., b_n)$ be n-tuples of real numbers satisfying $0 \le m_1 \le a_i \le M_1$ and $0 \le m_2 \le b_i \le M_2$ (i = 1, ..., n).

Then Ozeki-Izumino-Mori-Seo inequality [12, 22] asserts that

$$\sum_{i=1}^{n} a_i^2 \sum_{i=1}^{n} b_i^2 - \left(\sum_{i=1}^{n} a_i b_i\right)^2 \le \frac{n^2}{3} \left(M_1 M_2 - m_1 m_2\right)^2. \tag{5.1}$$

In [12] they also showed the following operator version of (5.1): If A and B are positive operators in $\mathbb{B}(\mathcal{H})$ such that $0 < m_1 \le A \le M_1$ and $0 < m_2 \le B \le M_2$ for some scalars $m_1 \le M_1$ and $m_2 \le M_2$, then

$$(A^{2}x, x)(B^{2}x, x) - (A^{2} \sharp B^{2}x, x)^{2} \le \frac{1}{4\gamma^{2}} (M_{1}M_{2} - m_{1}m_{2})^{2}$$
 (5.2)

for every unit vector $x \in H$, where $\gamma = \max\{\frac{m_1}{M_1}, \frac{m_2}{M_2}\}$.

Based on the Kantorovich inequality for the difference, we present an extension of Ozeki–Izumino–Mori–Seo inequality (5.2) as follows.

Theorem 5.1. Suppose that $\Phi : \mathbb{B}(\mathcal{H}) \to \mathbb{B}(\mathcal{H})$ is a positive linear map such that $\Phi(I)$ is invertible and $\Phi(I) \leq I$. Assume that $A, B \in \mathbb{B}(\mathcal{H})$ are positive invertible operators such that $0 < m_1 \leq A \leq M_1$ and $0 < m_2 \leq B \leq M_2$ for some scalars $m_1 \leq M_1$ and $m_2 \leq M_2$. Then

$$\Phi(B^2)^{\frac{1}{2}}\Phi(A^2)\Phi(B^2)^{\frac{1}{2}} - |\Phi(B^2)^{-\frac{1}{2}}\Phi(A^2\sharp B^2)\Phi(B^2)^{\frac{1}{2}}|^2 \le \frac{(M_1M_2 - m_1m_2)^2}{4} \times \frac{M_2^2}{m_2^2} \tag{5.3}$$

and

$$\Phi(A^2)^{\frac{1}{2}}\Phi(B^2)\Phi(A^2)^{\frac{1}{2}} - |\Phi(A^2)^{-\frac{1}{2}}\Phi(A^2\sharp B^2)\Phi(A^2)^{\frac{1}{2}}|^2 \le \frac{(M_1M_2 - m_1m_2)^2}{4} \times \frac{M_1^2}{m_1^2}.$$
(5.4)

Proof. Define a normalized positive linear map Ψ by

$$\Psi(X) := \Phi(A)^{-\frac{1}{2}} \Phi(A^{\frac{1}{2}} X A^{\frac{1}{2}}) \Phi(A)^{-\frac{1}{2}}$$

By using the Kantorovich inequality for the difference, it follows that

$$\Psi(X^2) - \Psi(X)^2 \le \frac{(M-m)^2}{4} \tag{5.5}$$

for all $0 < m \le X \le M$ with some scalars $m \le M$. As a matter of fact, we have

$$\begin{split} \Psi(X^2) - \Psi(X)^2 &\leq \Psi((M+m)X - Mm) - \Psi(X)^2 \\ &= -\left(\Psi(X) - \frac{M+m}{2}\right)^2 + \frac{(M-m)^2}{4} \\ &\leq \frac{(M-m)^2}{4}. \end{split}$$

If we put $X = (A^{-\frac{1}{2}}BA^{-\frac{1}{2}})^{\frac{1}{2}}$, then due to

$$0 < (m =) \sqrt{\frac{m_2}{M_1}} \le X \le \sqrt{\frac{M_2}{m_1}} (= M)$$

we deduce from (5.5) that

$$\Phi(A)^{-\frac{1}{2}}\Phi(B)\Phi(A)^{-\frac{1}{2}} - \left(\Phi(A)^{-\frac{1}{2}}\Phi(A\sharp B)\Phi(A)^{-\frac{1}{2}}\right)^2 \le \frac{(\sqrt{M_1M_2} - \sqrt{m_1m_2})^2}{4M_1m_1}.$$

Pre- and post-multiplying both sides by $\Phi(A)$, we obtain

$$\Phi(A)^{\frac{1}{2}}\Phi(B)\Phi(A)^{\frac{1}{2}} - |\Phi(A)^{-\frac{1}{2}}\Phi(A\sharp B)\Phi(A)^{\frac{1}{2}}|^{2} \leq \frac{(\sqrt{M_{1}M_{2}} - \sqrt{m_{1}m_{2}})^{2}}{4M_{1}m_{1}}\Phi(A)^{2}
\leq \frac{(\sqrt{M_{1}M_{2}} - \sqrt{m_{1}m_{2}})^{2}}{4} \times \frac{M_{1}}{m_{1}},$$

since $0 \le \Phi(A)^2 \le M_1^2$. Replacing A and B by A^2 and B^2 respectively, we have the desired inequality (5.4). Similarly, one can obtain (5.3).

Remark 5.2. If Φ is a vector state in (5.3) and (5.4), then we get Ozeki–Izumino–Mori–Seo inequality (5.2).

References

- [1] Lj. Arambašić, D. Bakić and M.S. Moslehian, A treatment of the Cauchy–Schwarz inequality in C*-modules, J. Math. Anal. Appl. (to appear), arXiv:0905.3509v2.
- [2] P.S. Bullen, A dictionary of inequalities, Pitman Monographs and Surveys in Pure and Applied Mathematics, 97. Longman, Harlow, 1998.
- [3] M.-D. Choi, Some assorted inequalities for positive linear maps on C*-algebras, J. Operator Theory 4 (1980), no. 2, 271–285.
- [4] J.B. Diaz and F.T. Metcalf, Stronger forms of a class of inequalities of G. Pólya-G. Szegö and L.V. Kantorovich, Bull. Amer. Math. Soc. 69 (1963), 415–418.
- [5] S.S. Dragomir, A survey on Cauchy-Bunyakovsky-Schwarz type discrete inequalities, J. Inequal. Pure Appl. Math. 4 (2003), no. 3, Article 63, 142 pp.
- [6] S.S. Dragomir, A counterpart of Schwarz inequality in inner product spaces, RGMIA. Res. Rep. Coll. 6 (2003), Article 18.
- [7] S.S. Dragomir, Advances in Inequalities of the Schwarz, Triangle and Heisenberg Type in Inner Product Spaces, Nova Science Publishers, Inc., New York, 2007.
- [8] D.E. Evans and J.T. Lewis, Dilations of irreversible evolutions in algebraic quantum theory, Comm. Dublin Inst. Adv. Studies. Ser. A, no. 24, 1977.
- [9] T. Furuta, J.M. Hot, J.E. Pečarić and Y. Seo, Mond-Pečarić method in operator inequalities. Inequalities for bounded selfadjoint operators on a Hilbert space, Monographs in Inequalities 1. Zagreb: Eelement, 2005.
- [10] W. Greub and W. Rheinboldt, On a generalisation of an inequality of L.V. Kantorovich, Proc. Amer. Math. Soc. 10 (1959), 407–415.
- [11] D. Ilišević and S. Varošanec, On the Cauchy-Schwarz inequality and its reverse in semiinner product C*-modules, Banach J. Math. Anal. 1 (2007), no. 1, 78–84.
- [12] S. Izumino, H. Mori and Y. Seo, On Ozeki's inequality, J. Inequal. Appl. 2 (1998), 235–253.
- [13] M. Joiţa, On the Cauchy-Schwarz inequality in C^* -algebras, Math. Rep. (Bucur.) **3(53)** (2001), no. 3, 243–246.
- [14] M.S. Klamkin and R.G. Mclenaghan, An ellipse inequality, Math. Mag. 50 (1977), 261–263.
- [15] E.C. Lance, Hilbert C*-Modules, London Math. Soc. Lecture Note Series 210, Cambridge Univ. Press, 1995.
- [16] E.-Y. Lee, A matrix reverse Cauchy-Schwarz inequality, Linear Algebra Appl. 430 (2009), no. 2-3, 805–810.
- [17] M.S. Moslehian and L.-E. Persson, Reverse Cauchy-Schwarz inequalities for positive C*-valued sesquilinear forms, Math. Inequal. Appl. 12 (2009), no. 4, 701–709.

- [18] M.S. Moslehian and R. Rajić, *Grüss inequality for n-positive linear maps*, Linear Alg. Appl. **433** (2010), no. 8-10, 1555–1560.
- [19] G.J. Murphy, C*-algebras and Operator Theory, Academic Press, Boston, 1990.
- [20] C.P. Niculescu, Converses of the Cauchy–Schwarz inequality in the C*-framework, An. Univ. Craiova Ser. Mat. Inform. **26** (1999), 22–28.
- [21] M. Niezgoda, Accretive operators and Cassels inequality, Linear Algebra Appl. 433 (2009), no. 1, 136–142.
- [22] N. Ozeki, On the estimation of the inequalities by the maximum, or minimum values(in Japanese), J.College Arts Ci. Chiba Univ., 5 (1968), 199–203.
- [23] G. Pólya and G. Szegö, Aufgaben und Lehrsätze aus der Analysis, Vol. 1, Berlin 1925, pp. 57 and 213–214.
- [24] O. Shisha and B. Mond, *Bounds on differences of means*, Inequalities I, New York-London, 1967, 293–308.
- [25] G.S. Watson, Serial correlation in regression analysis I, Biometrika 42 (1955), 327–342.
- ¹ Department of Pure Mathematics, Centre of Excellence in Analysis on Algebraic Structures (CEAAS), Ferdowsi University of Mashhad, P.O. Box 1159, Mashhad 91775, Iran.

E-mail address: moslehian@ferdowsi.um.ac.ir and moslehian@ams.org

² FACULTY OF ENGINEERING, IBARAKI UNIVERSITY, HITACHI, IBARAKI 316-0033, JAPAN. E-mail address: r-naka@net1.jway.ne.jp

³ Faculty of Engineering, Shibaura Institute of Technology, 307 Fukasaku, Minuma-ku, Saitama-city, Saitama 337-8570, Japan.

E-mail address: yukis@sic.shibaura-it.ac.jp