

Posterior consistency in linear models under shrinkage priors

BY A. ARMAGAN

SAS Institute Inc., Cary, North Carolina 27513, USA
artin.armagan@sas.com

AND D. B. DUNSON

Department of Statistical Science, Duke University, Durham, North Carolina 27708, USA
dunson@stat.duke.edu

AND J. LEE

Department of Statistics, Seoul National University, Seoul, 151-747, Korea
leejyc@gmail.com

AND W. U. BAJWA

Department of Electrical and Computer Engineering, Rutgers University, Piscataway, New Jersey 08854, USA
waheed.bajwa@rutgers.edu

AND N. STRAWN

Department of Mathematics, Duke University, Durham, North Carolina 27708, USA
nstrawn@math.duke.edu

SUMMARY

We investigate the asymptotic behavior of posterior distributions of regression coefficients in high-dimensional linear models as the number of dimensions grows with the number of observations. We show that the posterior distribution concentrates in neighborhoods of the true parameter under simple sufficient conditions. These conditions hold under popular shrinkage priors given some sparsity assumptions.

Some key words: Bayesian Lasso; Generalized double Pareto prior; Heavy tails; High-dimensional data; Horseshoe prior; Posterior consistency; Shrinkage estimation.

1. INTRODUCTION

Consider the linear model $y_n = X_n \beta_n^0 + \varepsilon_n$, where y_n is an n -dimensional vector of responses, X_n is the $n \times p_n$ design matrix, $\varepsilon_n \sim N(0, \sigma^2 I_n)$ with known σ^2 , and some of the components of β_n^0 are zero. Let $\mathcal{A}_n = \{j : \beta_{nj}^0 \neq 0, j = 1, \dots, p_n\}$ and $|\mathcal{A}_n| = q_n$ denote the set of indices and number of nonzero elements in β_n^0 .

49 In studying the behavior of regression methods in high-dimensional settings, it is in-
 50 creasingly common to allow the number of candidate predictors p_n to grow with sample
 51 size n . This is realistic in many applications. In genomics the number of predictors tends
 52 to be larger by design for studies with more subjects. In collecting single nucleotide poly-
 53 morphisms, gene expression, proteomics and so on, one can obtain an immense number
 54 of candidate predictors. However, when n is small, attempting to measure and include
 55 all such predictors in the statistical analysis seems unreasonable, so that one tends to
 56 collect and analyze increasing subsets of an effectively unbounded number of candidate
 57 predictors as sample size increases. In such applications, we are often interested in in-
 58 ferences on the model parameters as much as building a predictive model in order to
 59 understand the associations between the response and the candidate predictors.

60 Our setup is not new, and we follow Ghosal (1999) who also focused on asymptotic
 61 properties of the posterior on the regression coefficients assuming known σ^2 and growing
 62 p_n . The increasing p_n paradigm induces some challenges relative to the traditional liter-
 63 ature on posterior consistency in that growing dimension of β_n^0 results in a changing ℓ_2
 64 neighborhood around β_n^0 . This makes it more challenging to show that the posterior as-
 65 signs all such neighborhoods probability converging to one. One way to bypass this issue
 66 is to focus on the predictive distribution of y_n given X_n as in Jiang (2007). However, this
 67 does not address the common interest in inferences on the regression coefficients. Ghosal
 68 (1999) and Bontemps (2011) provide results on asymptotic normality of the posteriors
 69 in linear models for $p_n^4 \log p_n = o(n)$ and $p_n \leq n$, respectively. As a corollary, Ghosal
 70 (1999) states posterior consistency results in linear models when $p_n^3 \log n/n \rightarrow 0$ under
 71 the usual assumptions on X_n . However, both Ghosal (1999) and Bontemps (2011) require
 72 Lipschitz conditions ensuring that the prior is sufficiently flat in a neighborhood of the
 73 true β_n^0 . Such conditions are restrictive when using shrinkage priors that are designed to
 74 concentrate on sparse β_n vectors.

75 Our main contribution is providing a simple sufficient condition on the prior con-
 76 centration to achieve the desired asymptotic posterior behavior when $p_n = o(n)$. Our
 77 particular focus is on shrinkage priors, including the Laplace, Student's t , generalized
 78 double Pareto, and horseshoe-type priors (Johnstone & Silverman, 2004; Carvalho et al.,
 79 2010; Armagan et al., 2011, 2013). There is a rich methodological and applied literature
 80 supporting such priors but a lack of theoretical results.

83 2. SUFFICIENT CONDITIONS FOR POSTERIOR CONSISTENCY

84 Our results on posterior consistency rely on the following assumptions as $n \rightarrow \infty$:

85 (A1) Let $p_n = o(n)$;

86 (A2) Let $\Lambda_{n \min}$ and $\Lambda_{n \max}$ be the smallest and the largest singular values of
 87 X_n , respectively. Then $0 < \Lambda_{\min} < \liminf_{n \rightarrow \infty} \Lambda_{n \min}/\sqrt{n} \leq \limsup_{n \rightarrow \infty} \Lambda_{n \max}/\sqrt{n} <$
 88 $\Lambda_{\max} < \infty$;

89 (A3) Let $\sup_{j=1, \dots, p_n} |\beta_{nj}^0| < \infty$;

90 (A4) Let $q_n = o\{n^{1-\rho/2}/(\sqrt{p_n \log n})\}$ for $\rho \in (0, 2)$;

91 (A5) Let $q_n = o(n/\log n)$.

92 Assumptions (A4) and (A5) will be used in different settings.

93
 94
 95 LEMMA 1. Let $\mathcal{B}_n := \{\beta_n : \|\beta_n - \beta_n^0\| > \epsilon\}$ where $\epsilon > 0$. To test $H_0 : \beta_n = \beta_n^0$ vs $H_1 :$
 96 $\beta_n \in \mathcal{B}_n$, we define a test function $\Phi_n(y_n) = I(y_n \in \mathcal{C}_n)$ where the critical region is $\mathcal{C}_n :=$

97 $\{y_n : \|\hat{\beta}_n - \beta_n^0\| > \epsilon/2\}$ and $\hat{\beta}_n = (X_n^T X_n)^{-1} X_n^T y_n$. Then, under assumptions (A1) and
 98 (A2), as $n \rightarrow \infty$,
 99

- 100 1. $E_{\beta_n^0}(\Phi_n) \leq \exp\{-\epsilon^2 n \Lambda_{\min}^2 / (16\sigma^2)\}$,
 101 2. $\sup_{\beta_n \in \mathcal{B}_n} E_{\beta_n}(1 - \Phi_n) \leq \exp\{-\epsilon^2 n \Lambda_{\min}^2 / (16\sigma^2)\}$.
 102

103 **THEOREM 1.** *Given Lemma 1, the posterior of β_n under prior $\Pi_n(\beta_n)$ is strongly con-*
 104 *sistent, that is, for any $\epsilon > 0$, $\Pi_n(\mathcal{B}_n | y_n) = \Pi_n(\beta_n : \|\beta_n - \beta_n^0\| > \epsilon | y_n) \rightarrow 0$ pr β_n^0 -almost*
 105 *surely as $n \rightarrow \infty$, if*
 106

$$107 \Pi_n \left(\beta_n : \|\beta_n - \beta_n^0\| < \frac{\Delta}{n^{\rho/2}} \right) > \exp(-dn)$$

108 for all $0 < \Delta < \epsilon^2 \Lambda_{\min}^2 / (48 \Lambda_{\max}^2)$ and $0 < d < \epsilon^2 \Lambda_{\min}^2 / (32\sigma^2) - 3\Delta \Lambda_{\max}^2 / (2\sigma^2)$ and some
 109 $\rho > 0$.
 110

111
 112 Theorem 1 provides a simple sufficient condition on the concentration of the prior
 113 around sparse β_n^0 . We use Theorem 1 to provide conditions on β_n^0 under which specific
 114 shrinkage priors achieve posterior consistency focusing on priors that assume independent
 115 and identically distributed elements of β_n .
 116

117 2.1. Laplace Prior

118
 119 **THEOREM 2.** *Under assumptions (A1)–(A4), the Laplace prior $f(\beta_{nj} | s_n) =$
 120 $(1/2s_n) \exp(-|\beta_{nj}|/s_n)$ with scale parameter s_n yields a strongly consistent poste-*
 121 *rior if $s_n = C/(\sqrt{p_n n^{\rho/2}} \log n)$ for finite $C > 0$.*
 122

123 2.2. Student's t Prior

124 The density function for the scaled Student's t distribution is

$$125 f(\beta_j | s, d_0) = \frac{1}{s\sqrt{d_0} B(1/2, d_0/2)} \left(1 + \frac{\beta_j^2}{s^2 d_0} \right)^{-(d_0+1)/2},$$

126 with scale s , degrees of freedom d_0 , and $B(\cdot)$ denoting the beta function.
 127
 128

129
 130 **THEOREM 3.** *Under assumptions (A1)–(A3) and (A5), the scaled Student's t prior*
 131 *with parameters s_n and d_{0n} yields a strongly consistent posterior if $d_{0n} = d_0 \in (2, \infty)$*
 132 *and $s_n = C/(\sqrt{p_n n^{\rho/2}} \log n)$ for finite $\rho > 0$ and $C > 0$.*
 133
 134

135 2.3. Generalized Double Pareto Prior

136 As defined by Armagan et al. (2013), the generalized double Pareto density is given
 137 by
 138

$$139 f(\beta_j | \alpha, \eta) = \frac{\alpha}{2\eta} \left(1 + \frac{|\beta_j|}{\eta} \right)^{-(\alpha+1)}, \quad \alpha, \eta > 0.$$

140
 141 **THEOREM 4.** *Under assumptions (A1)–(A3) and (A5), the generalized double Pareto*
 142 *prior with parameters α_n and η_n yields a strongly consistent posterior if $\alpha_n = \alpha \in (2, \infty)$*
 143 *and $\eta_n = C/(\sqrt{p_n n^{\rho/2}} \log n)$ for finite $\rho > 0$ and $C > 0$.*
 144

2.4. Horseshoe-like Priors

As defined in Armagan et al. (2011), generalized beta scale mixtures of normals are obtained by the following three equivalent representations:

$$\beta_j \sim N(0, 1/\varrho_j - 1), f(\varrho_j) = \frac{\Gamma(a_0 + b_0)}{\Gamma(a_0)\Gamma(b_0)} \xi^{b_0} \varrho_j^{b_0-1} (1 - \varrho_j)^{a_0-1} \{1 + (\xi - 1)\varrho_j\}^{-(a_0+b_0)} \quad (1)$$

$$\beta_j \sim N(0, \tau_j), \tau_j \sim \text{Ga}(a_0, \lambda_j), \lambda_j \sim \text{Ga}(b_0, \xi)$$

$$\beta_j \sim N(0, \tau_j), f(\tau_j) = \frac{\Gamma(a_0 + b_0)}{\Gamma(a_0)\Gamma(b_0)} \xi^{-a_0} \tau^{a_0-1} (1 + \tau_j/\xi)^{-(a_0+b_0)}$$

where $a_0, b_0, \xi > 0$. Due to the representation in (1) and the work by Carvalho et al. (2010), we refer to these priors as *horseshoe-like*. The above formulation yields a general family that covers special cases discussed in Johnstone & Silverman (2004), a technical report by Griffin & Brown (2007) and Carvalho et al. (2010). The resulting marginal density on β_j is

$$f(\beta_j | a_0, b_0, \xi) = \frac{\Gamma(b_0 + 1/2)\Gamma(a_0 + b_0)U\{b_0 + 1/2, 3/2 - a_0, \beta_j^2/(2\xi)\}}{(2\pi\xi)^{1/2}\Gamma(a_0)\Gamma(b_0)}, \quad (2)$$

where $U(\cdot)$ denotes the confluent hypergeometric function of the second kind.

THEOREM 5. *Under assumptions (A1)–(A3) and (A5), the prior in (2) with parameters $a_{0n} = a_0 \in (0, \infty)$, $b_{0n} = b_0 \in (1, \infty)$ and ξ_n yields a strongly consistent posterior if $\xi_n = C/(p_n n^\rho \log n)$ for finite $\rho > 0$ and $C > 0$.*

3. FINAL REMARKS

Our analysis is heavily dependent on the construction of good tests. Results can be extended utilizing appropriate tests relying on an estimator with asymptotically vanishing probability of being outside of a *shrinking* neighborhood of the truth. For instance, one could use results similar to Bickel et al. (2009) given additional conditions on X_n . Theorem 7.2 of Bickel et al. (2009) states that

$$\text{pr}_{\beta_n^0} \left(\|\hat{\beta}_{nL} - \beta_n^0\|_2^2 > M \frac{a_n \log p_n}{n} \right) \leq p_n^{1-a_n^2/8} \quad (3)$$

for $a_n > 2\sqrt{2}$ and for some $M > 0$, where $\hat{\beta}_{nL}$ denotes the Lasso estimator. Hence using (3), in a similar fashion to Lemma 1, we can obtain consistent tests with an ϵ -neighborhood contracting at a rate $\mathcal{O}\{(a_n \log p_n)^{1/2}/\sqrt{n}\}$. Assuming $q_n < \infty$ for simplicity and letting $a_n = \mathcal{O}(\log n)$, following Theorems 1, 3, 4 and 5, we anticipate that under the Student's t , generalized double Pareto and horseshoe-like priors, a *near-optimal* contraction rate of $\mathcal{O}\{(\log n \log p_n)^{1/2}/\sqrt{n}\}$ is possible.

As in almost all of the Bayesian asymptotic literature, we have focused on sufficient conditions. Our conditions are practically appealing in allowing priors to be screened for their usefulness in high-dimensional settings. However, it would be of substantial interest to additionally provide theory allowing one to rule out the use of certain classes of priors in particular settings.

4. TECHNICAL DETAILS

Proof of Lemma 1. Noting that $\hat{\beta}_n = (X_n^\top X_n)^{-1} X_n^\top y_n$, $E_{\beta_n^0}(\Phi_n) = \text{pr}_{\beta_n^0}(\|\hat{\beta}_n - \beta_n^0\| > \epsilon/2) \leq \text{pr}_{\beta_n^0}\{\chi_p^2 > \epsilon^2 n \Lambda_{\min}^2 / (4\sigma^2)\}$ where χ_p^2 is a chi-squared distributed random variable with p degrees of freedom. The inequality is attained using assumption (A2). Similarly, $\sup_{\beta_n \in \mathcal{B}_n} E_{\beta_n}(1 - \Phi_n) \leq \sup_{\beta_n \in \mathcal{B}_n} \text{pr}_{\beta_n}(\|\hat{\beta}_n - \beta_n\| - \|\beta_n^0 - \beta_n\| \leq \epsilon/2) \leq \sup_{\beta_n \in \mathcal{B}_n} \text{pr}_{\beta_n}(\|\hat{\beta}_n - \beta_n\| \geq -\epsilon/2 + \|\beta_n^0 - \beta_n\|) = \text{pr}_{\beta_n}(\|\hat{\beta}_n - \beta_n\| \geq \epsilon/2) \leq \text{pr}_{\beta_n^0}\{\chi_p^2 > \epsilon^2 n \Lambda_{\min}^2 / (4\sigma^2)\}$. Simplifying the inequality $\text{pr}\{\chi_p^2 - p \geq 2(px)^{1/2} + 2x\} \leq \exp(-x)$ by Laurent & Massart (2000), we state that $\text{pr}(\chi_p^2 \geq x) \leq \exp(-x/4)$ if $x \geq 8p$. Then, using assumption (A1), as $n \rightarrow \infty$,

$$\begin{aligned} E_{\beta_n^0}(\Phi_n) &\leq \exp\{-\epsilon^2 n \Lambda_{\min}^2 / (16\sigma^2)\}, \\ \sup_{\beta_n \in \mathcal{B}_n} E_{\beta_n}(1 - \Phi_n) &\leq \exp\{-\epsilon^2 n \Lambda_{\min}^2 / (16\sigma^2)\}. \end{aligned}$$

This completes the proof. \square

Proof of Theorem 1. Our proof relies on a technique originally devised by Schwartz (1965). The posterior probability of \mathcal{B}_n is given by

$$\begin{aligned} \Pi_n(\mathcal{B}_n | y_n) &= \frac{\int_{\mathcal{B}_n} \{f(y_n | \beta_n) / f(y_n | \beta_n^0)\} \Pi(d\beta_n)}{\int \{f(y_n | \beta_n) / f(y_n | \beta_n^0)\} \Pi(d\beta_n)} \\ &\leq \Phi_n + \frac{(1 - \Phi_n) J_{\mathcal{B}_n}}{J_n} \\ &= I_1 + I_2 / J_n, \end{aligned}$$

where $J_{\mathcal{B}_n} = \int_{\mathcal{B}_n} \{f(y_n | \beta_n) / f(y_n | \beta_n^0)\} \Pi(d\beta_n)$ and $J_n = J_{\mathbb{R}^{pn}}$. We need to show that $I_1 + I_2 / J_n \rightarrow 0$ $\text{pr}_{\beta_n^0}$ -almost surely as $n \rightarrow \infty$. Let $b = \epsilon^2 \Lambda_{\min}^2 / (16\sigma^2)$. For sufficiently large n , $\text{pr}_{\beta_n^0}\{I_1 \geq \exp(-bn/2)\} \leq \exp(bn/2) E_{\beta_n^0}(I_1) = \exp(-bn/2)$ using Lemma 1. This implies that $\sum_{n=1}^{\infty} \text{pr}_{\beta_n^0}\{I_1 \geq \exp(-bn/2)\} < \infty$ and hence by the Borel–Cantelli lemma $\text{pr}_{\beta_n^0}\{I_1 \geq \exp(-bn/2) \text{ infinitely often}\} = 0$. We next look at the behavior of I_2 :

$$\begin{aligned} E_{\beta_n^0}(I_2) &= E_{\beta_n^0}\{(1 - \Phi_n) J_{\mathcal{B}_n}\} \\ &= E_{\beta_n^0}\left\{(1 - \Phi_n) \int_{\mathcal{B}_n} \frac{f(y_n | \beta_n)}{f(y_n | \beta_n^0)} \Pi_n(d\beta_n)\right\} \\ &= \int_{\mathcal{B}_n} \int (1 - \Phi_n) f(y_n | \beta_n) dy_n \Pi_n(d\beta_n) \\ &\leq \Pi_n(\mathcal{B}_n) \sup_{\beta_n \in \mathcal{B}_n} E_{\beta_n}(1 - \Phi_n) \\ &\leq \exp(-bn) \end{aligned}$$

Then for sufficiently large n , $\text{pr}_{\beta_n^0}\{I_2 \geq \exp(-bn/2)\} \leq \exp(-bn/2)$ using Lemma 1. Again $\sum_{n=1}^{\infty} \text{pr}_{\beta_n^0}\{I_2 \geq \exp(-bn/2)\} < \infty$ and hence by the Borel–Cantelli lemma $\text{pr}_{\beta_n^0}\{I_2 \geq \exp(-bn/2) \text{ infinitely often}\} = 0$.

We have shown that both I_1 and I_2 tend towards zero exponentially fast. Now we analyze the behavior of J_n . To complete the proof, we need to show that $\exp(bn/2) J_n \rightarrow$

241 ∞ $\text{pr}_{\beta_n^0}$ -almost surely as $n \rightarrow \infty$.

$$242 \quad \exp(bn/2)J_n = \exp(bn/2) \int \exp\left\{-n\frac{1}{n} \log \frac{f(y_n|\beta_n^0)}{f(y_n|\beta_n)}\right\} \Pi_n(d\beta_n)$$

$$243 \quad \geq \exp\{(b/2 - \nu)n\} \Pi_n(\mathcal{D}_{n,\nu}) \quad (4)$$

244 where $\mathcal{D}_{n,\nu} = \{\beta_n : n^{-1} \log\{f(y_n|\beta_n^0)/f(y_n|\beta_n)\} < \nu\} = \{\beta_n : n^{-1}(\|y_n - X_n\beta_n\|^2 - \|y_n - X_n\beta_n^0\|^2) < 2\sigma^2\nu\}$ for any $0 < \nu < b/2$. Then $\Pi_n(\mathcal{D}_{n,\nu}) \geq \Pi_n\{\beta_n : n^{-1}|\|y_n - X_n\beta_n\|^2 - \|y_n - X_n\beta_n^0\|^2| < 2\sigma^2\nu\}$. Using the identity $x^2 - x_0^2 = 2x_0(x - x_0) + (x - x_0)^2$ for all $x, x_0 \in \mathfrak{R}$,

$$245 \quad \Pi_n(\mathcal{D}_{n,\nu}) \geq \Pi_n\left\{\beta_n : n^{-1}\left|2\|y_n - X_n\beta_n^0\|(\|y_n - X_n\beta_n\| - \|y_n - X_n\beta_n^0\|)\right.\right.$$

$$246 \quad \left. + (\|y_n - X_n\beta_n\| - \|y_n - X_n\beta_n^0\|)^2\right| < 2\sigma^2\nu\}$$

$$247 \quad \geq \Pi_n\left\{\beta_n : n^{-1}(2\|y_n - X_n\beta_n^0\|\|X_n\beta_n - X_n\beta_n^0\| + \|X_n\beta_n - X_n\beta_n^0\|^2) < 2\sigma^2\nu\right\}$$

$$248 \quad \geq \Pi_n\left(\beta_n : n^{-1}\|X_n\beta_n - X_n\beta_n^0\| < \frac{2\sigma^2\nu}{3\kappa_n}, \|X_n\beta_n - X_n\beta_n^0\| < \kappa_n\right) \quad (5)$$

249 given that $\|y_n - X_n\beta_n^0\| \leq \kappa_n$. For $\kappa_n = n^{(1+\rho)/2}$ with $\rho > 0$ and $\kappa_n^2/\sigma^2 \geq 8n$, $\text{pr}_{\beta_n^0}(y_n : \|y_n - X_n\beta_n^0\|^2 > \kappa_n^2) = \text{pr}_{\beta_n^0}(y_n : \chi_n^2 > \kappa_n^2/\sigma^2) \leq \exp\{-\kappa_n^2/(4\sigma^2)\}$. Since $\sum_{n=1}^{\infty} \text{pr}_{\beta_n^0}(y_n : \|y_n - X_n\beta_n^0\| > \kappa_n) < \infty$, by the Borel–Cantelli lemma $\text{pr}_{\beta_n^0}(y_n : \|y_n - X_n\beta_n^0\| > \kappa_n \text{ infinitely often}) = 0$. Following from (5) and the fact that $\kappa_n \rightarrow \infty$, as $n \rightarrow \infty$, for sufficiently large n , $\Pi_n(\mathcal{D}_{n,\nu}) \geq \Pi_n\{\beta_n : n^{-1}\|X_n\beta_n - X_n\beta_n^0\| < 2\sigma^2\nu/(3\kappa_n)\} \geq \Pi_n(\beta_n : \|\beta_n - \beta_n^0\| < \Delta/n^{\rho/2})$, where $\Delta = 2\sigma^2\nu/(3\Lambda_{\max})$. Hence following (4), $\Pi_n(\mathcal{B}_n|y_n) \rightarrow 0$ $\text{pr}_{\beta_n^0}$ -almost surely as $n \rightarrow \infty$ if $\Pi_n(\beta_n : \|\beta_n - \beta_n^0\| < \Delta/n^{\rho/2}) > \exp(-dn)$ for all $0 < d < b/2 - \nu$. This completes the proof. \square

250 *Proof of Theorem 2.* We need to calculate the probability assigned to the region $\{\beta_n : \|\beta_n - \beta_n^0\| < \Delta/n^{\rho/2}\}$ under the Laplace prior.

$$251 \quad \Pi_n\left(\beta_n : \|\beta_n - \beta_n^0\| < \frac{\Delta}{n^{\rho/2}}\right) = \Pi_n\left\{\beta_n : \sum_{j \in \mathcal{A}_n} (\beta_{nj} - \beta_{nj}^0)^2 + \sum_{j \notin \mathcal{A}_n} \beta_{nj}^2 < \frac{\Delta^2}{n^\rho}\right\}$$

$$252 \quad \geq \prod_{j \in \mathcal{A}_n} \left\{\Pi_n\left(\beta_{nj} : |\beta_{nj} - \beta_{nj}^0| < \frac{\Delta}{\sqrt{p_n n^{\rho/2}}}\right)\right\}$$

$$253 \quad \times \Pi_n\left\{\beta_n^{j \notin \mathcal{A}} : \sum_{j \notin \mathcal{A}_n} \beta_{nj}^2 < \frac{(p_n - q_n)\Delta^2}{p_n n^\rho}\right\}$$

$$254 \quad \geq \prod_{j \in \mathcal{A}_n} \left\{\Pi_n\left(\beta_{nj} : |\beta_{nj} - \beta_{nj}^0| < \frac{\Delta}{\sqrt{p_n n^{\rho/2}}}\right)\right\} \left\{1 - \frac{p_n n^\rho E\left(\sum_{j \notin \mathcal{A}_n} \beta_{nj}^2\right)}{(p_n - q_n)\Delta^2}\right\} \quad (6)$$

255 where $E(\beta_{nj}^2)$ can be verified to be $2s_n^2$. Following from (6)

$$256 \quad \Pi_n\left(\beta_n : \|\beta_n - \beta_n^0\| < \frac{\Delta}{n^{\rho/2}}\right) \geq$$

$$257 \quad \left\{\frac{\Delta}{\sqrt{p_n n^{\rho/2}} s_n} \exp\left(-\frac{\sup_{j \in \mathcal{A}_n} |\beta_{nj}^0|}{s_n} - \frac{\Delta}{s_n \sqrt{p_n n^{\rho/2}}}\right)\right\}^{q_n} \left(1 - \frac{2p_n n^\rho s_n^2}{\Delta^2}\right). \quad (7)$$

289 Taking the negative logarithm of both sides of (7) and letting $s_n = C/(\sqrt{p_n n^{\rho/2}} \log n)$
 290 for some $C > 0$, we obtain

$$291 \quad -\log \Pi_n \left(\beta_n : \|\beta_n - \beta_n^0\| < \frac{\Delta}{n^{\rho/2}} \right) \leq -q_n \log \Delta + q_n \log C - q_n \log \log n$$

$$292 \quad -\log \left\{ 1 - \frac{2C^2}{\Delta^2 (\log n)^2} \right\} + \frac{q_n \Delta \log n}{C} + \frac{q_n \sqrt{p_n n^{\rho/2}} \log n \sup_{j \in \mathcal{A}_n} |\beta_{nj}^0|}{C} \quad (8)$$

293
 294 as $n \rightarrow \infty$. It is easy to see that the dominating term in (8) is the last one and
 295 $-\log \Pi_n(\beta_n : \|\beta_n - \beta_n^0\| < \Delta/n^{\rho/2}) < dn$ for all $d > 0$. This completes the proof. \square

296
 297 *Proof of Theorem 3.* $E(\beta_{nj}^2)$, in this case, is given by $d_0 s_n^2 / (d_0 - 2)$. For the sake of
 300 simplicity, we let $d_0 = 3$. Then following from (6)

$$301 \quad \Pi_n \left(\beta_n : \|\beta_n - \beta_n^0\| < \frac{\Delta}{n^{\rho/2}} \right) \geq \left(1 - \frac{3p_n n^\rho s_n^2}{\Delta^2} \right)$$

$$302 \quad \times \left[\frac{2\Delta}{\sqrt{p_n n^{\rho/2}} s_n \sqrt{3B(1/2, 3/2)}} \left\{ 1 + \frac{2 \sup_{j \in \mathcal{A}_n} (\beta_{nj}^0)^2}{3s_n^2} + \frac{2\Delta^2}{3s_n^2 p_n n^\rho} \right\}^{-2} \right]^{q_n}. \quad (9)$$

303
 304 Taking the negative logarithm of both sides of (9) and letting $s_n = C/(\sqrt{p_n n^{\rho/2}} \log n)$
 305 for some $C > 0$, we obtain

$$306 \quad -\log \Pi_n \left(\beta_n : \|\beta_n - \beta_n^0\| < \frac{\Delta}{n^{\rho/2}} \right) \leq q_n \log \left\{ \frac{\sqrt{3CB(1/2, 3/2)}}{2\Delta} \right\} - q_n \log \log n$$

$$307 \quad -\log \left\{ 1 - \frac{C^2}{\Delta^2 (\log n)^2} \right\} + 2q_n \log \left\{ 1 + \frac{2p_n n^\rho \log n \sup_{j \in \mathcal{A}_n} (\beta_{nj}^0)^2}{3C^2} + \frac{2\Delta^2 (\log n)^2}{3C^2} \right\}$$

$$308 \quad (10)$$

309
 310 as $n \rightarrow \infty$. It is easy to see that the dominating term in (10) is the last one and
 311 $-\log \Pi_n(\beta_n : \|\beta_n - \beta_n^0\| < \Delta/n^{\rho/2}) < dn$ for all $d > 0$. The result can be easily shown
 312 to hold for all $d_0 \in (2, \infty)$. This completes the proof. \square

313
 314 *Proof of Theorem 4.* $E(\beta_{nj}^2)$, in this case, can verified to be $2\eta_n^2 / (\alpha^2 - 3\alpha + 2)$ for $\alpha >$
 315 2. For the sake of simplicity, we let $\alpha = 3$. Then following from (6)

$$316 \quad \Pi_n \left(\beta_n : \|\beta_n - \beta_n^0\| < \frac{\Delta}{n^{\rho/2}} \right) \geq$$

$$317 \quad \left\{ \frac{3\Delta}{\sqrt{p_n n^{\rho/2}} \eta_n} \left(1 + \frac{\sup_{j \in \mathcal{A}_n} |\beta_{nj}^0|}{\eta_n} + \frac{\Delta}{\eta_n \sqrt{p_n n^{\rho/2}}} \right)^{-4} \right\}^{q_n} \left(1 - \frac{p_n n^\rho \eta_n^2}{\Delta^2} \right). \quad (11)$$

318
 319 Taking the negative logarithm of both sides of (11) and letting $\eta_n = C/(\sqrt{p_n n^{\rho/2}} \log n)$
 320 for some $C > 0$, we obtain

$$321 \quad -\log \Pi_n \left(\beta_n : \|\beta_n - \beta_n^0\| < \frac{\Delta}{n^{\rho/2}} \right) \leq -q_n \log 3\Delta - 3q_n \log C - q_n \log \log n$$

$$322 \quad -\log \left\{ 1 - \frac{C^2}{\Delta^2 (\log n)^2} \right\} + 4q_n \log \left(C + \Delta \log n + \sqrt{p_n n^{\rho/2}} \log n \sup_{j \in \mathcal{A}_n} |\beta_{nj}^0| \right) \quad (12)$$

289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336

337 as $n \rightarrow \infty$. It is easy to see that the dominating term in (12) is the last one and
 338 $-\log \Pi_n(\beta_n : \|\beta_n - \beta_n^0\| < \Delta/n^{\rho/2}) < dn$ for all $d > 0$. The result can be easily shown
 339 to hold for all $\alpha \in (2, \infty)$. This completes the proof. \square

340 *Proof of Theorem 5.* Similarly to the previous cases, we can show that $E(\beta_{nj}^2) =$
 341 $\xi_n \Gamma(a_0 + 1) \Gamma(b_0 - 1) / \{\Gamma(a_0) \Gamma(b_0)\}$. Then following from (6)
 342

$$343 \Pi_n \left(\beta_n : \|\beta_n - \beta_n^0\| < \frac{\Delta}{n^{\rho/2}} \right) \geq \left\{ 1 - \frac{p_n n^\rho E(\beta_{nj}^2)}{\Delta^2} \right\} \left(\frac{2\Delta}{\sqrt{p_n n^\rho \xi_n}} \right)^{q_n}$$

$$344 \times \left[\frac{U\{b_0 + 1/2, 3/2 - a_0, \sup_{j \in \mathcal{A}_n} (\beta_{nj}^0)^2 / \xi_n + \Delta / (p_n n^\rho \xi_n)\}}{(2\pi \xi_n)^{1/2} \Gamma(a_0) \Gamma(b_0) \Gamma(b_0 + 1/2)^{-1} \Gamma(a_0 + b_0)^{-1}} \right]^{q_n}. \quad (13)$$

345 We can use the expansion $U(a, b, z) = z^{-a} \{\sum_{m=0}^{R-1} (a)_m (1+a-b)_m (-z)^m / m! +$
 346 $\mathcal{O}(|z|^{-R})\}$ for large z , where $(a)_m = a(a+1)\dots(a+m-1)$ and R th term is the
 347 smallest in the expansion (Abramowitz & Stegun, 1972). Letting $R = 1$, for sufficiently
 348 large n , (13) can be further bounded as

$$349 \Pi_n \left(\beta_n : \|\beta_n - \beta_n^0\| < \frac{\Delta}{n^{\rho/2}} \right) > \left\{ 1 - \frac{p_n n^\rho E(\beta_{nj}^2)}{\Delta^2} \right\}$$

$$350 \times \left[\frac{\sqrt{2\Delta} \Gamma(b_0 + 1/2) \Gamma(a_0 + b_0)}{\sqrt{p_n n^\rho \xi_n} \sqrt{\xi_n} \sqrt{\pi} \Gamma(a_0) \Gamma(b_0) \{\sup_{j \in \mathcal{A}_n} (\beta_{nj}^0)^2 / \xi_n + \Delta / (p_n n^\rho \xi_n)\}^{(b_0 + 1/2)}} \right]^{q_n}. \quad (14)$$

351 Taking the negative logarithm of both sides of (14) and letting $\xi_n = C / (p_n n^\rho \log n)$ for
 352 some $C > 0$, we obtain

$$353 -\log \Pi_n \left(\beta_n : \|\beta_n - \beta_n^0\| < \frac{\Delta}{n^{\rho/2}} \right) <$$

$$354 -q_n \log \left\{ \frac{\sqrt{2\Delta} \Gamma(b_0 + 1/2) \Gamma(a_0 + b_0)}{\sqrt{C} \sqrt{\pi} \Gamma(a_0) \Gamma(b_0)} \right\} - \log \left\{ 1 - \frac{C \Gamma(a_0 + 1) \Gamma(b_0 - 1)}{\log n \Delta \Gamma(a_0) \Gamma(b_0)} \right\}$$

$$355 - \frac{q_n}{2} \log \log n + q_n \left(b_0 + \frac{1}{2} \right) \log \left\{ \frac{p_n n^\rho \log n \sup_{j \in \mathcal{A}_n} (\beta_{nj}^0)^2}{C} + \frac{\Delta \log n}{C} \right\} \quad (15)$$

356 as $n \rightarrow \infty$. It is easy to see that the dominating term in (15) is the last one and
 357 $-\log \Pi_n(\beta_n : \|\beta_n - \beta_n^0\| < \Delta/n^{\rho/2}) < dn$ for all $d > 0$. This completes the proof. \square

376 ACKNOWLEDGEMENTS

377 This work was supported by the National Institute of Environmental Health Sciences.
 378 The content is solely the responsibility of the authors and does not necessarily repre-
 379 sent the official views of the National Institute of Environmental Health Sciences or the
 380 National Institutes of Health. Jaeyong Lee was supported by Advanced Research Center
 381 Program (S/ERC), the National Research Foundation of Korea grant funded by the
 382 Korean government (MSIP). Waheed U. Bajwa was supported in part by the NSF. Nate
 383 Strawn was supported by DARPA Mathematics of Sensing, Exploitation, and Execution
 384 (MSEE) program (managed by Dr. Tony Falcone).

REFERENCES

- 385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
- ABRAMOWITZ, M. & STEGUN, I. A. (1972). *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. New York: Dover.
- ARMAGAN, A., DUNSON, D. B. & CLYDE, M. (2011). Generalized beta mixtures of Gaussians. *Advances in Neural Information Processing Systems (NIPS)* .
- ARMAGAN, A., DUNSON, D. B. & LEE, J. (2013). Generalized double Pareto shrinkage. *Statistica Sinica* **23**, 119–143.
- BICKEL, P. J., RITOV, Y., ALEXANDRE & TSYBAKOV, B. (2009). Simultaneous analysis of Lasso and Dantzig selector. *Annals of Statistics* **37**, 1705–1732.
- BONTEMPS, D. (2011). Bernstein–von Mises theorems for Gaussian regression with increasing number of regressors. *Annals of Statistics* **39**, 2557–2584.
- CARVALHO, C. M., POLSON, N. G. & SCOTT, J. G. (2010). The horseshoe estimator for sparse signals. *Biometrika* **97**, 465–480.
- GHOSAL, S. (1999). Asymptotic normality of posterior distributions in high-dimensional linear models. *Bernoulli* **5**, 315–331.
- JIANG, W. (2007). Bayesian variable selection for high dimensional generalized linear models: Convergence rates of the fitted densities. *The Annals of Statistics* **35**, 1487–1511.
- JOHNSTONE, I. M. & SILVERMAN, B. W. (2004). Needles and straw in haystacks: Empirical Bayes estimates of possibly sparse sequences. *Annals of Statistics* **32**, 1594–1649.
- LAURENT, B. & MASSART, P. (2000). Adaptive estimation of a quadratic functional by model selection. *The Annals of Statistics* **28**, 1302–1338.
- SCHWARTZ, L. (1965). On Bayes procedures. *Zeitschrift für wahrscheinlichkeitstheorie und verwandte gebiete* **4**, 10–26.