

# WAVELET ANALYSIS OF GALACTIC ROTATION CURVES

M. Kuassivi  
BP 170374 Cotonou, Benin

## ABSTRACT

The spatial wavelet spectra of 73 published spiral galaxies's rotation curves are computed and their associated scaleograms are presented. Scaleograms are used to detect and isolate local features observed in spiral galaxies's rotation curves. Although wiggles and bumps are usually interpreted as signs of recent and on—going merging, the analysis of the scaleograms reveals regular patterns consistent with the presence of large—scale modes throughout the disk.

### 1. INTRODUCTION

In recent years, a great deal of focus has been put on the understanding of merger events in the evolution of galaxies. Our own galaxy has a tiny satellite galaxy (the Sagittarius Dwarf Elliptical Galaxy) which is currently being ripped up and eaten by the Milky Way. It is thought these kind of events may be quite common in the evolution of large galaxies. The Sagittarius Dwarf galaxy is orbiting our galaxy at almost a right angle to the disk. It is currently passing through the disk; stars are being stripped off of it with each pass and joining the halo of our galaxy. There are other examples of these minor accretion events, and it is likely a continual process for many galaxies. Such mergers provide gas and dark matter to galaxies. Evidence for this process is often observable as warp and streams coming out of galaxies.

Most of the Early type galaxies show clear sign of interaction and tidal distortion with some of them having bumps and wiggles in the rotation curve (Struve & Conway, 2010). In such galaxies, the rotation curves are fitted best by models with scaled H I disks suggesting that a connection exists between large scale streaming motion and HI density distribution. As a matter of facts,

discontinuities observed within the rotation curves deserve some attention. Analysis of these features is made easy by the use of the wavelet transforms.

The wavelet analyses properties of waves of various scale in real space (Jones 2009). Wavelet transforms are now being adopted for a vast number of applications, often replacing the conventional Fourier transform. Many areas of physics have seen this paradigm shift, including molecular dynamics, astrophysics, density matrix localisation, seismology, optics, turbulence and quantum mechanics. Wavelet transforms have advantages over traditional Fourier transforms for representing functions that have discontinuities and sharp peaks, and for accurately deconstructing and reconstructing, non periodic and non stationnary sinals.


The goal of this paper is to use the wavelet transform for the analysis of perturbations in rotation curves.

In section 2. I describe the data and the method. In section 3. the results are briefly discussed. Scaleograms are presented in the annexe.

### 2. METHODOLOGY

The 97 rotation curves data used in this work come from the paper of Brownstein

& Moffat (2006). Using a short script written in IDL language, I manually extracted the data points from the numerical paper version. Table 1. shows the list of the galaxies finally used in this work. Then, I interpolated the rotation curves on a 0.1 kpc grid and subtracted to each rotation curve the Newtonian model published by Brownstein & Moffat (using the same IDL script). Fig.1 is an example of the resulting output for F563—1 showing remarkable similarities with dissipative oscillations. Note that F563-1 is a low surface brightness (LSB) galaxy dominated by dark matter all the way into its center.



**Fig 1.** Reduced rotation curve of F563-1.

Wavelet transform is applied using the D.O.G mother because it resembles the most bumps often observed. Note that the wavelet software was provided by C. Torrence and G. Compo, and is available at <http://paos.colorado.edu/research/wavelets>.

Names	Vo (km/s)
DD0170	61
F563-1	95
F568-3	109
F571-8	145
F583-1	95
NGC55	85
NGC224	260
NGC247	111
NGC253	200
NGC300	102
NGC598	112
NGC660	145
NGC801	235
NGC891	199
NGC1097	290
NGC1365	240

NGC1417	150
NGC1560	75
NGC2403	135
NGC2708	230
NGC2841	310
NGC2903	200
NGC2998	190
NGC3034	85
NGC3079	200
NGC3109	70
NGC3198	155
NGC3495	150
NGC3521	200
NGC3628	210
NGC3672	220
NGC3726	155
NGC3877	161
NGC3893	180
NGC3972	129
NGC4051	165
NGC4088	170
NGC4096	118
NGC4100	179
NGC4138	160
NGC4157	189
NGC4183	111
NGC4217	190
NGC4303	150
NGC4321	260
NGC4389	119
NGC4527	180
NGC4565	250
NGC4736	150
NGC4945	169
NGC5033	210
NGC5055	199
NGC5457	211
NGC5533	290
NGC5585	85
NGC5907	250
NGC6503	119
NGC6674	270
NGC6946	160
NGC6951	192
NGC7331	250
UGC2259	89
UGC6399	86
UGC6614	190
UGC6667	85
UGC6818	74
UGC6917	102
UGC6923	87
UGC6930	110
UGC6973	175
UGC6983	110
UGC7089	71

**Table.1** list of spiral galaxies used in this work with terminal velocities Vo.

### 3. CONCLUSION

Scaleograms allow to detect and characterize local perturbations in the rotation curves. We can see a variety of spatial frequencies among the scaleograms (from  $1 \text{ kpc}^{-1}$  to  $40 \text{ kpc}^{-1}$ ). Such diversity is well expected if we are observing relief traces of merger events. Surprisingly enough, within each scaleogram the range of the detected frequencies is narrow and the amplitudes of the detection peaks are similar. Some scaleograms show a regular distributions of peaks that could be interpreted as trains of mergers.

However, such regular distributions are more likely related to harmonic modes in the rotation curves. I wish to point out that if a large—scale harmonic velocity structure is embedded in the rotation curve it may indeed be related to or be at the origin of the spiral arm structure. This would be consistent with the presence of modes in amorphous LSB galaxies such as F563-1.

As a conclusion, the interpretation of rotation curve discontinuities as arising from random mergers is put at a test.

### REFERENCES

- Brownstein, J.R., Moffat, J.W., 2006, *ApJ*  
636
- Jones, *Data Analysis in Cosmology*,  
Lecture Notes in Physics, vol. 665,  
, Springer 2009
- Struve, C & Conway, J.E., 2010, *A&A*,  
513

## ANNEXE

The figures below are the scaleograms obtain from Moffat & Brownstein (2006) rotation curves. Prior to the wavelet transformation, the Newtonian Model has been subtracted from the data. The X coordinates are the distance to the center of the galaxy in kpc. The Y coordinates are the spatial frequencies (in  $\text{kpc}^{-1}$ ).

