

# The Herschel Virgo Cluster Survey. IX. Dust-to-gas mass ratio and metallicity gradients in four Virgo spiral galaxies

Laura Magrini<sup>1</sup>, Simone Bianchi<sup>1</sup>, Edvige Corbelli<sup>1</sup>, Luca Cortese<sup>3</sup>, Leslie Hunt<sup>1</sup>, Matthew Smith<sup>2</sup>, Catherine Vlahakis<sup>4</sup>, Jonathan Davies<sup>2</sup>, George J. Bendo<sup>5</sup>, Maarten Baes<sup>6</sup>, Alessandro Boselli<sup>7</sup>, Marcel Clemens<sup>8</sup>, Viviana Casasola<sup>9</sup>, Ilse De Looze<sup>6</sup>, Jacopo Fritz<sup>6</sup>, Carlo Giovanardi<sup>1</sup>, Marco Grossi<sup>10</sup>, Thomas Hughes<sup>11</sup>, Suzanne Madden<sup>12</sup>,  
Ciro Pappalardo<sup>1</sup>, Michael Pohlen<sup>2</sup>, Sperello di Serego Alighieri<sup>1</sup>, Joris Verstappen<sup>6</sup>

<sup>1</sup> INAF–Osservatorio Astrofisico di Arcetri, Largo E. Fermi, 5, I-50125 Firenze, Italy e-mail: laura@arcetri.astro.it

<sup>2</sup> Department of Physics and Astronomy, Cardiff University, The Parade, Cardiff, CF24 3AA, UK

<sup>3</sup> European Southern Observatory, Karl-Schwarzschild-Strasse 2, D-85748 Garching bei Munchen, Germany

<sup>4</sup> Departamento de Astronomia, Universidad de Chile, Casilla 36-D, Santiago, Chile

<sup>5</sup> Jodrell Bank Centre for Astrophysics, Alan Turing Building, School of Physics and Astronomy, University of Manchester, Manchester M13 9PL

<sup>6</sup> Sterrenkundig Observatorium, Universiteit Gent, Krijgslaan 281 S9, B-9000 Gent, Belgium

<sup>7</sup> Laboratoire d'Astrophysique de Marseille, UMR 6110 CNRS, 38 rue F. Joliot-Curie, F-13388 Marseille, France

<sup>8</sup> Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, 35122 Padova, Italy

<sup>9</sup> INAF - Istituto di Radioastronomia, via P. Gobetti 101, 40129 Bologna, Italy

<sup>10</sup> CAAUL, Observatorio Astronomico de Lisboa, Universidade de Lisboa, Tapada de Ajuda, 1349-018, Lisboa, Portugal

<sup>11</sup> The Kavli Institute for Astronomy & Astrophysics, Peking University, Beijing 100871, China

<sup>12</sup> Service d'Astrophysique, CEA/Saclay, l'Orme des Merisiers, 91191 Gif-sur-Yvette, France

Received ; accepted

## ABSTRACT

**Context.** Using Herschel data from the Open Time Key Project *the Herschel Virgo Cluster Survey (HeViCS)*, we investigated the relationship between the metallicity gradients expressed by metal abundances in the gas phase as traced by the chemical composition of HII regions, and in the solid phase, as traced by the dust-to-gas mass ratio.

**Aims.** We derived the radial gradient of the dust-to-gas mass ratio for all galaxies observed by HeViCS whose metallicity gradients are available in the literature. They are all late type Sbc galaxies, namely NGC4254, NGC4303, NGC4321, and NGC4501.

**Methods.** We fitted PACS and SPIRE observations with a single-temperature modified blackbody, inferred the dust mass, and calculated two dimensional maps of the dust-to-gas mass ratio, with the total mass of gas from available HI and CO maps. HI moment-1 maps were used to derive the geometric parameters of the galaxies and extract the radial profiles. We examined different dependencies on metallicity of the CO-to-H<sub>2</sub> conversion factor ( $X_{\text{CO}}$ ), used to transform the <sup>12</sup>CO observations into the amount of molecular hydrogen.

**Results.** We found that in these galaxies the dust-to-gas mass ratio radial profile is extremely sensitive to choice of the  $X_{\text{CO}}$  value, since the molecular gas is the dominant component in the inner parts. We found that for three galaxies of our sample, namely NGC4254, NGC4321, and NGC4501, the slopes of the oxygen and of the dust-to-gas radial gradients agree up to  $\sim 0.6-0.7R_{25}$  using  $X_{\text{CO}}$  values in the range 1/3-1/2 Galactic  $X_{\text{CO}}$ . For NGC4303 a lower value of  $X_{\text{CO}} \sim 0.1 \times 10^{20}$  is necessary.

**Conclusions.** We suggest that such low  $X_{\text{CO}}$  values might be due to a metallicity dependence of  $X_{\text{CO}}$  (from close to linear for NGC4254, NGC4321, and NGC4501 to superlinear for NGC4303), especially in the radial regions  $R_G < 0.6-0.7R_{25}$  where the molecular gas dominates. On the other hand, the outer regions, where the atomic gas component is dominant, are less affected by the choice of  $X_{\text{CO}}$ , and thus we cannot put constraints on its value.

**Key words.** Galaxies: individual: NGC4254, NGC4303, NGC4321, NGC4501 — Galaxies: spiral — ISM:dust, extinction — Galaxies: ISM - Galaxies: abundances — Submillimeter: galaxies

## 1. Introduction

Virgo is one of the best studied galaxy clusters, being the richest cluster nearest to our own Galaxy ( $\sim 17$  Mpc, Gavazzi et al. 1999). It is a relatively populous system, consisting of more than 1000 confirmed members (Binggeli et al. 1985). Galaxies in clusters such as Virgo differ significantly from their field counterparts since interactions with the hostile environment re-

move gas, quenching the star formation (c.f., Boselli & Gavazzi 2006).

A galaxy's metallicity is closely related to the star formation (SF) history by which the interstellar medium (ISM) is enriched with the end-products of stellar evolution, and to the infall process that dilutes the ISM and triggers new SF. As a consequence of their star formation histories, gas stripping and infall events, modified by the cluster environment, galaxies in clusters are expected also to differ in metal content relatively to isolated galax-

ies. A fundamental tool with which tracing the chemical evolution of a galaxy is the study of its radial metallicity gradient. The metallicity gradient tracks indeed the star formation history of galaxies, integrated over time, together with infall and/or outflow events.

The first pioneering work in Virgo was done by Skillman et al. (1996), who analyzed nine spiral galaxies with the aim of seeking correlations among their gas content, locations in the cluster, metallicities and radial gradients, and comparing them with field spirals. Skillman et al. (1996) found weak evidence of shallower gradients in cluster galaxies deficient in HI than gradients in galaxies with a normal HI content. The situation is however very complex because galaxy interactions affect the star formation history and the gas content across the disk, producing metallicity gradients which differ from those measured in isolated galaxies of the same morphological type. Rupke et al. (2010) have shown that gradients in strongly interacting galaxies are flatter than in similar isolated galaxies; on the other hand, the metallicity gradient of M81, the largest member of a small group of galaxies, is steeper than in an isolated counterpart due to gas removal in the outskirts (Stanghellini et al. 2010).

A direct correlation between gas metallicity and the dust-to-gas mass ratio is naturally expected since approximatively half of the metals in the ISM reside in dust grains; thus the dust-to-gas ratio should scale with metal abundance. Such a trend has been obtained theoretically by models computing consistently the evolution of metals and dust, despite the large uncertainties in the yields of both (Dwek 1998; Inoue 2003). The relation of the global dust-to-gas mass ratio with metallicity was investigated by, e.g., James et al. (2002); Draine et al. (2007); Hirashita et al. (2008); Lisenfeld & Ferrara (1998). The radial variation of the dust-to-gas mass ratio was first investigated by Issa et al. (1990) in our Galaxy and in other nearby galaxies (LMC, SMC, M31, M33, and M51) who found evidence for a correlation, with dust-to-gas mass ratio and metallicity decreasing at roughly the same rate with increasing galactocentric radius. More recently, Boissier et al. (2004, 2005) and Thilker et al. (2007) found a clear relationship between metallicity and extinction, thus dust, in several nearby galaxies, suggesting that the variation in extinction is associated with the metallicity gradient. Muñoz-Mateos et al. (2009) found a good correlation between dust-to-gas and metallicity gradients in the Spitzer Infrared Nearby Galaxies Survey (SINGS; Kennicutt et al. 2003). Finally, Bendo et al. (2010a) compared the dust-to-gas ratio and metallicity gradients in NGC2403, finding a similar decreasing behavior with radius.

We have recently obtained observations of the Virgo galaxies with the Herschel Space Observatory (Pilbratt et al. 2010), within the Open Time Key Project HeViCS (Herschel Virgo Cluster Survey) (Davies et al. 2010). HeViCS maps a wide area over the Virgo Cluster at wavelengths from 100 to 500  $\mu\text{m}$ . This spectral range covers the peak of the thermal emission from cold dust ( $T < 30$  K) which enables the detection of the bulk of the dust emission in galaxies. Also, Herschel gives an unprecedented resolution at these wavelengths (ranging from about 10 to 36'', equivalent to 1-3 kpc for Virgo galaxies). Atomic and molecular gas maps are available in the literature at a comparable resolution (Chung et al. 2009a; Kuno et al. 2007), thus providing resolved maps and the possibility of deriving radial gradients of dust-to-gas mass ratios.

In the present paper we assessed the validity of using HeViCS observations to obtain metallicity gradients from radial profiles of the dust-to-gas mass ratio. We investigated the hypothesis that the local dust-to-gas mass ratio is proportional to metallicity, starting from the four spiral galaxy in the Virgo cluster (NGC4254, NGC4303, NGC4321, and NGC4501) whose metallicity gradients are available in the literature. We studied the relation between radial profiles of metallicity and dust-to-gas mass ratios, and how this can be used to constrain the CO-to-H<sub>2</sub> conversion factor ( $X_{\text{CO}}$ ) and its dependence on metallicity.

The paper is structured as follows: in Sect. 2 we describe the new HeViCS observations, and the observations of atomic and molecular gas. We also derive the oxygen abundance and its radial gradient in each galaxy. Sect. 3 discusses the fits to the observed Herschel dust spectral energy distributions (SEDs), how the radial profiles are derived, and how we calculate dust and gas masses. In Sect. 4 we develop the method by which we constrain  $X_{\text{CO}}$ , and Sect. 5 gives our conclusions.

## 2. Data

### 2.1. The sample

In the HeViCS field there are four galaxies for which the oxygen gradient has been well determined in the literature (e.g., Skillman et al. 1996; Moustakas et al. 2010). They are NGC4254 (M99), NGC4303 (M61), NGC4321 (M100), and NGC4501 (M88), all Sbc late-type galaxies.

NGC4254 is a bright spiral galaxy located at the periphery of the Virgo cluster, at a projected distance of  $\sim 1$  Mpc from the cluster center. Optical images show that this galaxy has one-armed structure, also seen in the HI gas distribution (Phookun et al. 1993; Chung et al. 2009a). Such an asymmetric spiral pattern is often observed in tidally galaxies, but there is no apparently massive companion near NGC4254 (Sofue et al. 2003).

NGC4321 is located at a distance of  $\sim 1.1$  Mpc from M87, and has an HI disk that is slightly larger than the optical disk.

NGC4501 is the closest galaxy of our sample to M87, being located at a distance of  $\sim 0.5$  Mpc. It is weakly HI-deficient, following the definition of Chung et al. (2009a). Comparisons with simulations suggest that NGC4501 is in an early stage of ram pressure stripping (Vollmer et al. 2008), entering the high-density region of the cluster for the first time.

NGC4303 is the most isolated galaxy in our sample. It is a barred spiral galaxy with face-on geometry located in the outskirts of the Virgo cluster. We assume for all the galaxies the distance of  $\sim 17$  Mpc (Gavazzi et al. 1999).

### 2.2. Herschel observations

The HeViCS program consists of Herschel observations of an area of about 60 sq. deg. over the denser parts of the Virgo Cluster. The total area is made of 4 overlapping fields, which are observed in parallel scanning mode (*fast* scan rate: 60''/s) with both the PACS and SPIRE instruments, yielding data simultaneously in 5 spectral bands, at 100 and 160  $\mu\text{m}$  (from PACS) and at 250, 350, and 500  $\mu\text{m}$  (from SPIRE). At the completion of the program, each field will be covered with 8 scans done in two perpendicular scan directions. The full width half maximum (FWHM) of the beams is 6''.98  $\times$  12''.7 and 11''.64  $\times$  15''.65 in the two PACS bands (PACS Observers' Manual,

2010), and  $18''2$ ,  $24''9$ ,  $36''3$  in the three SPIRE bands (SPIRE Observers' Manual, 2010). At the time of writing, each field has been observed with at least two scans. In this paper, we use this dataset, whose data reduction and analysis is described in details in PAPER VIII, Davies et al. (2011).

Some papers presenting Herschel observations for these galaxies have been already published. Eales et al. (2010) and Sauvage et al. (2010) analyzed the SPIRE maps of NGC4254 (M99) and NGC4321 (M100) observed within the Herschel Reference Survey (Boselli et al. 2010) to map the ISM using dust emission. Adding these data to archival Spitzer, HI, and CO maps, Pohlen et al. (2010) investigated the spatial distribution of gas and dust in these same galaxies. They also present as a preliminary result, the ratio of the total gas mass ( $\text{HI} + \text{H}_2$ ) to  $500 \mu\text{m}$  flux, an approximation of the dust mass for the two galaxies. They found a decreasing dust-to-gas mass ratio with radius, consistent with results by, e.g., Bendo et al. (2010a) in NGC2403. With the present-time availability of the PACS data the dust SED fitting can be better defined allowing to measure the exact shape of the radial dust-to-gas mass gradient. Finally, Smith et al. (2010) presented a resolved dust analysis of three of the largest (in angular size) spiral galaxies in HeViCS, among them NGC4501.

### 2.3. The calibration of metallicity and the abundance gradients


The metallicity measurements of these galaxies are available in the literature from optical spectroscopy of their HII regions. The most direct method to derive the oxygen abundance is to measure the electron temperature ( $T_e$ ) of the ionized gas using the intensity (relative to a hydrogen recombination line) of one or more temperature-sensitive auroral lines such as  $[\text{O III}] \lambda 4363 \text{\AA}$ ,  $[\text{N II}] \lambda 5755 \text{\AA}$ ,  $[\text{S III}] \lambda 6312 \text{\AA}$ , and  $[\text{O II}] \lambda 7325 \text{\AA}$ , as summarized by Moustakas et al. (2010). However, measurements of the electron temperature were not available from the spectroscopic observations in the original papers of McCall et al. (1985), Shields et al. (1991), Henry et al. (1994), and Skillman et al. (1996). The temperature diagnostic lines are indeed intrinsically faint in metal-rich HII regions. Therefore, oxygen abundance has been derived using the strong-line abundance calibrations which relate the metallicity to one or more line ratios involving the strongest recombination and forbidden lines. In particular, the oxygen excitation index  $R_{23} = ([\text{OII}] + [\text{OIII}]) / \text{H}\beta$  is one of the most often adopted calibrators to estimate the nebular abundances. As discussed by Moustakas et al. (2010), the principal advantage of  $R_{23}$  as an oxygen abundance diagnostic is that it is directly proportional to both principal ionization states of oxygen, whereas one of the major disadvantages is that the relation between  $R_{23}$  and metallicity is degenerate for low and high metallicity.

However, the calibrations of the metallicity by means of the strong-line ratios are not unique. They can be divided in three main categories: those calibrated with photoionization models (e.g., McGaugh 1991; Zaritsky et al. 1994; Kewley & Dopita 2002; Kobulnicky & Kewley 2004); those calibrated directly with the electron temperature, called empirical methods (e.g., Pilyugin 2001; Pettini & Pagel 2004; Pilyugin & Thuan 2005); and those combining both methods (e.g., Denicoló et al. 2002). As discussed widely in Kewley & Ellison (2008), the abundances derived with different methods do not have a common absolute oxygen abundance

scale. The oxygen abundances derived using the theoretical calibration are up to a factor of  $\sim 4$  higher than those based on the empirical calibration. However, despite the significant zero-point offset in the abundance scales, to first order the slope of the abundance gradients agree when calculated with different calibrators (see Moustakas et al. 2010).

We have tested the effect of several metallicity calibrators in NGC4254, the galaxy with the best sampled metallicity gradient. Starting with the abundance estimates from Moustakas et al. (2010), based on literature spectroscopy calibrated with the Kobulnicky & Kewley (2004) formula (hereafter KK04), we used the relationships provided by Kewley & Ellison (2008) to convert to other abundance scales. The relationships of Kewley & Ellison (2008) are obtained for limited oxygen abundance ranges, corresponding to the ranges where they could perform a polynomial fit to transform one abundance scale to another. Because of this, from the abundances of KK04 we were unable to recover the oxygen determination of Pettini & Pagel (2004) and Pilyugin & Thuan (2005). For these cases, we re-computed the oxygen abundance from the original spectra (McCall et al. 1985; Shields et al. 1991; Henry et al. 1994).

In Fig. 1, we show the result of our test: while the slope of the gradient is almost invariant with different calibrations, the zero-point depends on the choice of the calibration. In particular, the empirical calibrations of Pettini & Pagel (2004) and Pilyugin & Thuan (2005) show values of the oxygen abundances roughly 0.5-0.8 dex lower than the other determinations. These two latter empirical calibrations were obtained for HII regions with available electron temperature in a relatively low metallicity regime. They could not be valid for metal rich environments, as the galaxies of our sample. In Sect. 4.1 we will discuss how the dust-to-gas ratio might help in setting a lower limit to the metallicity and to discriminate among different calibrations.



**Fig. 1.** The oxygen abundance gradient of NGC4254 obtained with several metallicity calibrations: Kobulnicky & Kewley (2004) (magenta circles), Pilyugin & Thuan (2005) (yellow squares), Zaritsky et al. (1994) (cyan triangles), McGaugh (1991) (blue stars), Kewley & Dopita (2002) (empty squares), Tremonti et al. (2004) (green asterisks), and Pettini & Pagel (2004) (violet diamond).

## 2.4. HI and CO maps

The radial profile of the gas, including both atomic and molecular components, is necessary to derive the dust-to-gas mass ratio gradient. For NGC4254, NGC4321, and NGC4501, we use the moment-0 HI maps obtained with VLA Imaging survey of Virgo galaxies in Atomic gas (VIVA) survey by Chung et al. (2009a). VIVA observations reach a column density sensitivity of  $3\text{-}5 \times 10^{19} \text{cm}^{-2}$ . The comparison of their total HI fluxes with values in the literature from single dish observations gives a good agreement especially for the large galaxies, indicating no loss of flux in the interferometric observations (see Fig. 5 in Chung et al. 2009a). The beam sizes are:  $30''.86 \times 28''.07$  for NGC4254,  $15''.90 \times 14''.66$  for NGC4321, and  $16''.83 \times 16''.41$  for NGC4501. The HI radial profile of NGC4303 is available from Warmels (1988) and Cayatte et al. (1990). We adopt the combined radial profile of the two, shown in Fig.3 of Skillman et al. (1996).

Maps of molecular gas were available thanks to the  $^{12}\text{CO}$  ( $J = 1-0$ ) mapping survey of 40 nearby spiral galaxies, performed with the Nobeyama 45 m telescope by Kuno et al. (2007).

## 3. Analysis


### 3.1. The mass and temperature of dust

Maps of the dust temperature and mass surface density were obtained as in Smith et al. (2010). The images of the galaxies from the five PACS (100 and 160  $\mu\text{m}$ ) and SPIRE (250, 350, and 500  $\mu\text{m}$ ) bands were all convolved and re-gridded to the lower resolution (FWHM=36''.9, 3 kpc) and pixel size (14''.0, 1.1 kpc) of the 500  $\mu\text{m}$  observations. We used only pixels with  $S/N > 10$  at 500  $\mu\text{m}$  (the *rms* of our maps in this band is about 0.3 MJy/sr, see Davies et al. 2011). The selection of these high surface-brightness pixels was necessary to limit the uncertainties due to background subtraction and avoid the artefacts caused by the high-pass filtering in the PACS data reduction. Despite this limit, we were able to study the dust and gas properties up to at least  $0.7 R_{25}^1$  in all galaxies. For each galaxy we thus considered approximately 150-200 pixels covering about 1/2 of the optically defined area.

We estimated the error on the surface brightness on a pixel by pixel basis by comparing galaxy images from the two scan data used in this paper with those relative to other two scans recently taken by Herschel which cover only part of the HeViCS field. We found errors very similar to those estimated on the total fluxes (see Davies et al. 2011). Including a calibration error of 15% for PACS (PACS ICC, private communication) and 7% for SPIRE (SPIRE Observer' Manual, 2010), the total error is 30%, 20%, 10%, 10%, and 15% of the flux at 100, 160, 250, 350, and 500  $\mu\text{m}$ , respectively.

The SED for each pixel was fitted with a single modified blackbody, using a power law dust emissivity  $\kappa_\lambda = \kappa_0(\lambda_0/\lambda)^\beta$ , with spectral index  $\beta = 2$  and emissivity  $\kappa_0 = 0.192 \text{ m}^2 \text{ kg}^{-1}$  at  $\lambda_0 = 350 \mu\text{m}$ . These values reproduce the average emissivity of models of the Milky Way dust in the FIR-submm (Draine 2003). The fit was obtained with a standard  $\chi^2$  minimization technique. In the pipeline calibration, the flux density observed by the var-

ious instruments, i.e. weighted over each filter passband, is converted into a monochromatic flux density assuming  $F_\nu \propto \nu^{-1}$ . Before fitting a modified blackbody, a color correction should be applied to the data, to account for the real spectral slope of the source. Alternatively, the conversion implemented into the pipeline calibration can be removed from the data (in SPIRE parlance this is equivalent to dividing the pipeline flux densities by the  $K_4$  factor; SPIRE Observer' Manual, 2010); the passband weighted flux thus obtained should then be compared with the mean of the model flux density over the spectral response function for each of the bands. We adopted this second technique, using the appropriate response functions for the PACS and SPIRE bands (for SPIRE, we used the response functions for extended emission). However, color corrections are small for the adopted emissivity and the temperature range derived here (see Davies et al. 2011).



**Fig. 2.** Typical SEDs on three different positions (pixels) on NGC4254. Blue dots are the measured (color-corrected) fluxes and the red (solid) + curves are the modified blackbody fits. The three SED correspond to minimum ( $\approx 21\text{K}$ ), mean ( $\approx 23.5\text{K}$ ), and maximum ( $\approx 26\text{K}$ ) temperature in the galaxy (see Fig. 3). The green (dashed) curves show the fit of the Draine & Li (2007) model to the data, for IRSFs of intensities 2, 4 and 8 times the Local ISRF (see their paper for details).

In Fig. 2 we show typical SEDs and graybody fits for three positions of different dust temperature in NGC4254. As shown by the figure, a single temperature modified blackbody with  $\beta = 2$  is sufficient to obtain reasonable fits of the SED over the wavelength range considered here (see also Davies et al. 2011). When data at shorter wavelength than PACS 100  $\mu\text{m}$  is available, one might want to consider a two-component model, to include emission from warmer dust that might significantly contribute at least to the 100  $\mu\text{m}$  flux. This was done, for example, by Bendo et al. (2010b), who used Herschel data, and by Smith et al. (2010) who used 70  $\mu\text{m}$  data from the Spitzer satellite. However, they found that the inclusion of a warmer temperature component, thought necessary to fit the 70  $\mu\text{m}$  data, improves the fit at  $\lambda > 100 \mu\text{m}$  only slightly, and does not modify

<sup>1</sup>  $R_{25}$  is the radius of the galaxy measured to a B surface brightness of 25 mag arcsec<sup>-2</sup>, and is an indication of the size of the galaxy;  $R_{25}$  were obtained from NED;  $\sim 0.7R_{25}$  is equivalent to the solar radius in our Galaxy.

significantly the estimate of the temperature and mass surface density of cold dust.

### 3.2. Sources of uncertainty in mass and temperature of dust

When only the errors on photometry are considered, the uncertainty in the determination of the temperature is about 2 K, and thus  $\sim 20\%$  on the dust mass surface density.


In principle, fitting the dust SED with a single thermal-equilibrium temperature component could result in larger uncertainties in the dust mass estimates: grains of a given size and material could be exposed to different intensities of the interstellar radiation field (ISRF) and thus attain different equilibrium temperatures which will contribute differently to the SED; conversely, for the same radiation field the SED could depend on the dust distribution, because it results from the emission of a mixture of grains of different size and composition, each with its own equilibrium temperature. We found neither of these to have a strong effects on our mass estimates. In fact, by fitting the SED pixel by pixel, we already take into account the gradients due to the diffuse ISRF, which is more prominent in the radial direction than in the vertical directions (i.e. along the line of sight, for non edge-on disks; Bianchi et al. 2000).

However, the temperature radial gradients found here (see Sect. 3.3) would not produce very large uncertainties, even when the global SED is fitted with a single temperature model. For our targets, the differences between the sum of the dust mass in each pixel, and the dust mass obtained by fitting the sum of the flux densities in each pixel, is smaller than the fit error. The insensitivity of the global dust SED fitting on the shallow diffuse ISRF gradient is clearly shown in the analysis of the the FIR/submm SED of late type galaxies of Draine et al. (2007): though the more complex fitting procedure includes a dust grain model and a range of ISRF intensities (see also Draine & Li 2007), the global SED at  $\lambda \geq 100\mu\text{m}$  is found to be fitted by a dust component that accounts for most of the dust mass (a part from a few percents), heated by an IRSF of constant intensity.

We evaluated the effects of a complex dust mixture by fitting the model of Draine & Li (2007) to our pixel-by-pixel SED. Following Draine et al. (2007), we used a single value of the ISRF for emission at  $\lambda \geq 100\mu\text{m}$ . An example can be seen in Fig. 2. The SED from the dust grain mixtures and our single temperature model fit equally well the data. The dust mass obtained by using the procedure of Draine et al. (2007) is higher by about 10%, that is, within the error we quoted. Thus, the dust mass derived from a simple averaged emissivity, and a single IRSF - or temperature - , is not severely underestimated.

Larger uncertainties can come also from the assumption for the emissivity. For example, the value of the emissivity derived by James et al. (2002) from SCUBA observations of galaxies is equivalent to  $\kappa_0 = 0.41 \text{ m}^2 \text{ kg}^{-1}$ , a factor of two larger than the value we adopted. Adopting this emissivity would result in dust surface densities a factor of two smaller than what we found in this paper.

Also, dust emissivity has been reported to increase by about a factor two in grains associated to denser environments (see, e.g., Bianchi et al. 2003). However, this might be the case for extreme environments and not representative of the bulk of the diffuse dust mass: in the Milky Way, recent results from the Planck satellite show no emissivity variation between dust associated with HI and CO emission, nor emissivity variations with the galactocentric radius (Planck Collaboration et al. 2011). In



**Fig. 3.** Radial variation of dust temperature in NGC4254 (red dot-dashed curve), NGC4303 (green dashed line), NGC4321 (blue dotted curve), and NGC4501 (black solid curve).

Sect. 4.1 we will discuss how the uncertainty on  $\kappa_0$  might affect our discussion.

The limited wavelength coverage does not allow us to investigate in details the effect of variation of the emissivity spectral index. In any case, the modified  $\beta = 2$  blackbody provides good fits for all the SEDs analysed here, as well as for the global SEDs in a larger sample of HeViCS objects (Davies et al. 2011). Local variations of the dust emissivity index with temperature, as those reported by Paradis et al. (2010) cannot be easily verified in our dataset. Variation of  $\beta$  from 1.7 to 2.2, as those measured at a reference temperature of 20K, would result in an underestimate and overestimate of the dust mass of the same order as the quoted errors, respectively. Lower  $\beta$  values, as those found for dust at higher temperatures, will not be able to provide good fits to our SEDs in the central part of the galaxies.

Maps of the temperature and dust mass surface density for NGC4501 have been presented in Smith et al. (2010). For the other galaxies, they will be presented in Vlahakis et al. (2011, in preparation).

### 3.3. Radial profiles

The galaxies in our sample are disturbed by tidal interactions, thus we consider more reliable the determination of the geometric parameters of the galactic disks from kinematics rather than from photometry (see, for example, the discrepancy between parameters derived with the two methods for the case of NGC4321 in Chung et al. 2009a). The task ROTCUR of the package NEMO (Teuben 1995) was used to fit a tilted ring model to the 21-cm moment-1 maps from the VIVA dataset. We obtained values for the rotational velocity, the inclination and the position angle, as a function of the galactocentric radius.

The radial variations of  $i$  and p.a. are shown in Fig. 4. The average values are indicated with solid lines. For NGC4321 and NGC4501,  $i$  and p.a. are quite constant with radius and thus we use the average parameters in the whole radial range. For

NGC4254, which is a one-armed spiral, the variations of p.a. and  $i$  are important due to its asymmetric shape. Since in our comparison between metallicity and dust-to-gas mass ratio gradients we are interested mostly in the inner regions where the metallicity data are available, for NGC4254 we used  $i$  and p.a. averaged in the regions with galactocentric radius  $R_G < R_{25}$  (dashed line in Fig. 4, top panels) where the variations are small. Pohlen et al. (2010) also found different ellipse parameters for the outer parts and for the inner parts of NGC4254. For NGC4303, for which VIVA maps are not available, we adopted the parameters by Koda & Sofue (2006).

The final values are shown in Table 1: in column 1 we show the galaxy name, in columns 2 and 3 the mean  $i$  and p.a., in column 4 the optical diameter  $D_{opt}$  in arcmin, in cols. 6, 7, and 8 the slope of the metallicity gradient, the metallicity at the equivalent solar radius  $0.7R_{25}$ , and their reference. For NGC4254 we report the value we adopt, i.e. the average within the optical radius, and in square brackets the average in the whole radial range. Between square brackets, for all galaxies, we report also inclinations and position angles from the literature, which are generally in good agreement with our values in the radial range considered.

The atomic and molecular gas maps were convolved to the same resolution of the SPIRE  $500 \mu\text{m}$  map and regridded on the same pixel scale. The radial bin size in the profiles correspond to 1 pixel in the maps ( $14''$  corresponding to  $\sim 1.1$  kpc). Adopting the geometric parameters in Table 1, we used a dedicated IDL procedure to obtain radial profiles (azimuthally averaged over elliptical annuli) for HI, CO, and dust mass surface density. The profiles are shown in Figs. 5: the top panels contain PACS and SPIRE profiles at 100, 160, 250, 350, and  $500 \mu\text{m}$  (from top to bottom) for NGC4254, NGC4303, NGC4321, and NGC4501; the bottom panels the profiles of dust (green), HI (red), and  $\text{H}_2$  (blue) surface density, where the CO flux is converted into  $\text{H}_2$  using the standard conversion factor  $X_{\text{CO}} \approx (1.8 \pm 0.3) \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$  (see below). For all galaxies  $\text{H}_2$  is dominant in the inner-most regions, while HI becomes dominant over  $\text{H}_2$  at  $R > 0.5\text{-}0.6 R_{25}$ . HI radial profiles are quite flat within  $R_{25}$ , whereas both  $\text{H}_2$  and dust profiles are decreasing with radius.

The errors on the radial profiles were estimated by a combination in quadrature of the uncertainty in the overall absolute calibration and of the standard deviation of the fluxes in each radial bin. Typical errors are of the order of 10-30%. However, uncertainties for the dust mass are mainly due to the choice of the model parameters and can vary by a factor two.

In the next Section, we report the total mass of  $\text{H}_2$  obtained with the different assumptions on  $X_{\text{CO}}$  (Tab.4).

### 3.4. The $X_{\text{CO}}$ factor

To convert  $^{12}\text{CO}$  observation to measure the amount of molecular hydrogen we assume a CO-to- $\text{H}_2$  conversion factor

$$X_{\text{CO}} = N(\text{H}_2) / \int I(\text{CO}) dv$$

(Lebrun et al. 1983). In our Galaxy  $X_{\text{CO}} \approx (1.8 \pm 0.3) \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$  with excursions of up to a factor of  $\sim 2$  over this value, particularly at high latitudes (Dame et al. 2001). Shetty et al. (2011) and Leroy et al. (2009) showed that  $X_{\text{CO}}$  can vary even more, especially in low metallicity environments where it can reach  $X_{\text{CO}} \sim 100X_{\text{Gal}}$ .

As derived by analysis of observations by Bolatto et al. (2008) and by theoretical models (Glover & Mac Low 2011), this factor can vary due to effects of abundance, excitation, optical depth, and cloud structure averaged over a large area. In particular, the abundance of the heavy elements has an important impact on the value of  $X_{\text{CO}}$  (Israel et al. 1986; Maloney & Black 1988). Metallicity affects cloud structure both directly, as smaller abundances of C and O translate into lower abundance of CO, and indirectly, as a lower dust-to-gas mass ratio diminishes the  $\text{H}_2$  formation rates and the shielding of molecular gas from the photo-dissociation by ultraviolet radiation. As summarized by Bolatto et al. (2008), there are several observational calibrations of  $X_{\text{CO}}$  with the metallicity,  $Z$ , in the literature, showing a range of behaviors; most of the calibrations find an increasing  $X_{\text{CO}}$  with decreasing  $Z$  (e.g., Wilson 1995; Barone et al. 2000; Israel et al. 2003; Strong et al. 2004).

The dependence varies from  $X_{\text{CO}} \sim Z^{-2.5}$  (Israel 2000) to  $X_{\text{CO}} \sim Z^{-1}$  (Wilson 1995; Arimoto et al. 1996; Boselli et al. 2002), while Bolatto et al. (2008) do not find any measurable trend in the range from  $8 < 12 + \log(\text{O}/\text{H}) < 8.8$  on the scales of the individual CO-bright giant molecular clouds, but they did not argue for a Galactic  $X_{\text{CO}}$  on large scales. Strong & Matto (1996) also found a constant  $X_{\text{CO}}$  with metallicity. However, few works were dedicated to the metallicity dependence of  $X_{\text{CO}}$  in super-solar regions, as for example the study of Arimoto et al. (1996) who investigated  $X_{\text{CO}}$  also in two metal-rich galaxies, namely M31 and M51. The spiral galaxies in Virgo are thus of particular importance to verify the assumed dependence on metallicity of  $X_{\text{CO}}$  in metal-rich environments. To evaluate the conversion factor  $X_{\text{CO}}$  at different metallicities and relate it to the dust-to-gas ratio, we compared three different assumptions:

- i) *Uniform*: we used the standard Galactic value from Bolatto et al. (2008)  $X_{\text{CO}} \approx 1.8 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$  (hereafter Galactic  $X_{\text{CO}}$ ) throughout the radial range.
- ii) *Linear metallicity dependence*: we used a linear metallicity dependence as found by Arimoto et al. (1996); Boselli et al. (2002). This is similar to the  $-0.67$  slope derived by Wilson (1995). We adopted the analytic calibration given by Wilson (1995), but rely on the most recent oxygen solar abundance of Asplund et al. (2009) instead of Grevesse & Anders (1989). Assuming that at the new solar oxygen abundance,  $12 + \log(\text{O}/\text{H}) = 8.69$ ,  $X_{\text{CO}} \approx 1.8 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$ , the Boselli et al. (2002)'s relationship becomes

$$\log(X_{\text{CO}}) = -1.0 \times (12 + \log(\text{O}/\text{H})) + 8.94 \quad (1)$$

where  $X_{\text{CO}}$  is expressed in unit of  $10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$ .


- iii) *Super-linear metallicity dependence*: we considered a super-linear dependence on metallicity as found by Israel (2000). Re-scaling to the Asplund et al. (2009) solar oxygen abundance, the Israel (2000) relationship becomes

$$\log(X_{\text{CO}}) = -2.5 \times (12 + \log(\text{O}/\text{H})) + 21.72 \quad (2)$$

where  $X_{\text{CO}}$  is again expressed in unit of  $10^{20}$ .

## 4. The method

We started from the hypothesis that dust-to-gas mass ratio and O/H are tracing the same quantity, i.e. the metal abundance relative to hydrogen. Thus we expected that the slope of the two



**Fig. 4.** Radial variation of position angle and inclination in NGC4254, NGC4321, and NGC4501 from moment-1 H I maps. The continuous lines are the average in the whole radial range, whereas the dashed line is the average in the region  $R_G < R_{opt}$  for NGC4254.

**Table 1.** Parameters of the sample galaxies.

Name	Inclination	Position angle	$D_{opt}^a$ (arcmin)	slope $_{KK04}$ (dex/ $R_{25}$ )	$12+\log(O/H)_{KK04}$ at $R_G=0.7R_{25}$	Ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)
NGC4254	$42^\circ$ ( $53^\circ$ ) <sup>b</sup> [ $42^\circ$ ] <sup>d</sup>	$67^\circ$ ( $61.6^\circ$ ) <sup>b</sup> [ $68^\circ$ ] <sup>d</sup>	5.6	$-0.42\pm 0.06$	$8.84\pm 0.02$	M10 <sup>c</sup>
NGC4303	$30^\circ$ <sup>e</sup>	$135^\circ$ <sup>e</sup>	6.5	$-0.42\pm 0.02^\ddagger$	$9.24\pm 0.02$	S96 <sup>c</sup>
NGC4321	$33^\circ$ [ $26^\circ$ ] <sup>f</sup>	$154.4^\circ$ [ $155^\circ$ ] <sup>f</sup>	7.6	$-0.35\pm 0.13$	$9.04\pm 0.04$	M10 <sup>c</sup>
NGC4501	$63^\circ$ [ $58^\circ$ ] <sup>g</sup>	$141.4^\circ$ [ $140^\circ$ ] <sup>g</sup>	7.1	$-0.07\pm 0.16^\ddagger$	$9.30\pm 0.21$	S96 <sup>c</sup>

**Notes.** *a*  $D_{opt}$  is  $2\times R_{25}$ ; *b* Average values for  $R_G < R_{25}$  and, between brackets, throughout the whole radial range. *c* M10=Moustakas et al. (2010) and S96=Skillman et al. (1996); *d* Phookun et al. (1993); *e* Koda & Sofue (2006); *f* Canzian & Allen (1997); *g* Kenney & Young (1988).  $\ddagger$  We recomputed the gradient slope of NGC4303 and NGC4501 using data from Skillman et al. (1996), converting them with Kewley & Ellison (2008) formula to the KK04 scale, obtaining different values with respect to those quoted in their paper.

radial gradients agree. Within this hypothesis, we attempted to constrain the value of  $X_{CO}$ , and eventually its dependence on metallicity, using the information contained in the slope of the metallicity gradient. In fact, due to the large discrepancy on the zero-point of different calibrations as discussed by, e.g., Kewley & Ellison (2008), it is difficult to place these galaxies on an absolute metallicity scale using only inferences from HII optical spectroscopy.

Here we summarize the steps, while the details are given in the following sections.


I We converted the dust-to-gas mass ratio in oxygen abundances with the relationship of Draine et al. (2007) (see

Eq. 3) (Draine et al. 2007, see Eq. 3), thus obtaining an approximate zero-point for the metallicity scale.

- II We computed the radial profile of the dust-to-gas mass ratio with the Galactic  $X_{CO}$ .
- III Once fixed the approximate abundance scale, we compared the radial profiles of the dust-to-gas mass ratio with the super-linear and linear metallicity dependences of  $X_{CO}$ .

#### 4.1. I. The dust-to-gas ratio and metallicity scales

If we consider that the abundances of all heavy elements are proportional to the oxygen abundance and that all heavy elements condensed to form dust in the same way as in the MW, then the



**Fig. 5.** *Top panels:* radial profiles of dust emission at 100, 160, 250, 350, 500  $\mu\text{m}$  (from top to bottom) for NGC4254, NGC4303, NGC4321, NGC4501. *Bottom panels:* radial profiles of dust mass (green solid line), HI (red long dashed line), H<sub>2</sub> (blue dot-dashed line) surface density using a uniform conversion factor  $X_{\text{CO}} \approx (1.8 \pm 0.3) \times 10^{20} \text{ cm}^{-2}$ , and the O/H radial gradient (dotted line, scaled of a factor 10.5 to match the logarithmic scale of the other quantities).

dust-to-gas mass ratio scales proportionally to the oxygen abundance (Draine et al. 2007) (Draine et al. 2007)

$$\frac{M_{\text{dust}}}{M_{\text{H}}} \approx 0.007 \frac{(\text{O}/\text{H})}{(\text{O}/\text{H})_{\odot}}, \quad (3)$$

where 0.007 is the dust-to-hydrogen ratio of the MW at solar radius,  $\sim 0.7 R_{25}$  with optical radius of our Galaxy of 12 kpc (Reshetnikov 2000). In eq.3, we use the observed dust-to-gas ratio 0.0073 estimated from observed depletions in the solar neighborhood instead of the value from dust models, 0.010, about 40% larger (Draine et al. 2007), and the solar oxygen abundance  $12 + \log(\text{O}/\text{H})_{\odot} = 8.69$  from Asplund et al. (2009).

A similar relationship was obtained also by Boselli et al. (2002) using the data available for the MW (Sodroski et al. 1994), the LMC (Koornneef 1982), and the SMC (Bouchet et al. 1985).

We converted the dust-to-gas ratio, obtained with the Galactic  $X_{\text{CO}}$  at  $R_G = 0.7 R_{25}$ , into  $12 + \log(\text{O}/\text{H})$  with Eq. 3.

Before comparing the metallicity derived from dust-to-gas ratio with that derived from nebular oxygen abundances, we estimated the effect of the assumed values of the mass emissivity coefficient and of the  $X_{\text{CO}}$  on the metallicity derived from Eq. 3. Because the determinations of the total gas mass take place in regions where H<sub>2</sub> is not negligible, the dust-to-gas ratio depends indeed critically on what is assumed about  $X_{\text{CO}}$ . First,

we checked the value of the dust-to-gas ratio we would obtain at  $R_G = 0.7 R_{25}$  adopting different  $X_{\text{CO}}$ : using  $X_{\text{CO}} \approx 4 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$  we would have a dust-to-gas ratio  $\sim 0.2$  dex lower and using  $X_{\text{CO}} \approx 0.5 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$  we would obtain a dust-to-gas ratio  $\sim 0.2$  dex higher. Thus an uncertainty of  $\pm 0.2$  dex is associated to the choice of  $X_{\text{CO}}$ .

Then, we verified how the emission coefficient could affect our discussion. In the most unfavorable conditions we could overestimate (or underestimate) the dust mass by a factor two, corresponding to a variation of O/H derived from Eq.3 of  $\pm 0.3$  dex.

In addition, we have that a typical error on the dust-to-gas mass ratio of  $\sim 30\text{-}50\%$  (including the flux calibration) translates into errors of  $\sim 0.10\text{-}0.15$  dex in O/H. Considering that these sources of uncertainty are independent we combine them in quadrature, obtaining a total uncertainty of  $\sim 0.35$  dex.

We compared with  $12 + \log(\text{O}/\text{H})$  obtained at the same galactocentric distance with three calibrations: KK04, Pettini & Pagel (2004) (PP04), and Pilyugin & Thuan (2005) (P05). The results for NGC4254, NGC4303, NGC4321, and NGC4501 are shown in Table 2.

From the comparison of the metallicity obtained for dust-to-gas ratio with several metallicity calibrations we found that the calibrations of Pilyugin & Thuan (2005) show the largest discrepancy with the metallicity derived from dust also if we



**Table 2.** Oxygen abundance from dust-to-gas mass ratio at  $0.7R_{25}$ .

Name	$O/H_{dust-to-gas}$ $0.7R_{25}$	$O/H_{KK04}$	$O/H_{PP04}$	$O/H_{P05}$
(1)	(2)	(3)	(4)	(5)
NGC4254	8.84	8.95	8.65	8.35
NGC4303	9.24	9.24	8.94	8.64
NGC4321	9.04	9.10	8.80	8.40
NGC4501	9.04	9.30	8.90	8.70

consider all possible sources of errors, i.e., uncertainty on the emission coefficient,  $X_{CO}$ , flux calibrations, and tend to give lower O/H values. The calibrations of KK04 and Pettini & Pagel (2004) are, on the other hand, both consistent within the errors ( $\sim 0.35$  dex  $O/H_{dust-to-gas}$  and  $\sim 0.1$  dex for O/H from nebular abundances) with our findings from the dust-to-gas ratio. We conclude that, at the equivalent solar radius ( $0.7 R_{25}$ ), the four galaxies in analysis are more metal rich than the MW at the same radius (the solar oxygen abundance is 8.69), thus a perfect environment to test if any dependence of  $X_{CO}$  with the metallicity is in place at high metallicity.

Our result is only in apparent disagreement with Draine et al. (2007). Their relationship between dust-to-gas and metallicity is derived theoretically assuming that the interstellar abundances of all heavy elements were proportional to O/H, and their fraction scales as in the MW, and then it is compared with the oxygen abundances obtained with the calibration of Pilyugin & Thuan (2005) and the dust-to-gas mass ratio adopting  $X_{CO} \approx 4 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$ . They found a good agreement with the theoretical relationship within a factor of 2. In our analysis we found a better agreement with KK04 oxygen abundances since we are using a different  $X_{CO}$  factor and a different value of dust-to-gas in the MW at solar radius: the P05 oxygen abundances are indeed on average 2-4 times lower than the KK04's ones, but this is compensated by a  $X_{CO}$  factor two times higher, and the assumed dust-to-gas in the MW for the dust models,  $\sim 1.5$  higher than the one we used, which comes from observations.

With our choice of a Galactic  $X_{CO} \approx 1.8 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$ , a dust-to-gas ratio in the solar neighborhood of 0.0073, and the relation in eq.3, we selected the metallicity scale of KK04, which best agrees with the scale derived from the dust-to-gas ratio, and convert the abundances of our galaxies to a common scale. For NGC4254 and NGC4321 we used the abundance determined by Moustakas et al. (2010) (based on literature spectroscopy calibrated with the formula of Kobulnicky & Kewley 2004). For NGC4303 and NGC4501 we used the original spectroscopy to calculate oxygen abundance on the abundance scale of KK04.

With the calibration of KK04, the abundances at  $0.7R_{25}$  are all super-solar ranging from  $\sim 8.95$  to  $\sim 9.3$ , as shown in Table 2. The central oxygen abundances would range from 9.3 to 9.5, but we have to consider that they are an extrapolation of the metallicity gradient up to the galaxy centre and not a real measurement. In addition, the galaxy disk does not extend up to the central regions where the bulge is present. These central abundances are comparable with recent results obtained from direct electron temperature measurement in the HII regions of M81, where the

extrapolation of the metallicity gradient up to the galactic centre gives an oxygen abundance of 9.37 (Stanghellini et al. 2010), and in our Galaxy where the central O/H from the gradient of Rudolph et al. (2006) is 9.2 (determination of O/H with optical spectroscopy).

#### 4.2. II. The dust-to-gas mass gradient with a standard $X_{CO}$

In Fig. 6 we show the comparison between the dust-to-gas mass ratio gradients, calculated with the Galactic conversion factor (red curves), and the oxygen abundance gradients (magenta circles and solid black lines) calibrated with KK04 and converted in the dust-to-gas mass scale with the relationship discussed in Sect. 4.1.


If we consider a constant Galactic  $X_{CO}$ , the slopes of the dust-to-gas and metallicity gradients are only in marginal agreement. In particular, the dust-to-gas gradients are flatter, with a slight positive slope at large galactocentric radii  $R_G > 0.7R_{25}$ . For NGC4254 and NGC4321 the comparison between the metallicity and dust-to-gas mass ratio gradients is straightforward because their metallicity gradients are well determined in the same radial regions where the dust-to-gas map is obtained (Fig. 6, left panels). For NGC4254 and NGC4321 the dust-to-gas mass ratio gradient are flatter than the metallicity radial distribution, and they are constant within the errors, with a slightly increasing behaviour in the outer regions. The same is true also for NGC4501, while for NGC4303 dust-to-gas mass ratio gradient is positive. For these two galaxies we are planning new spectroscopic observations of HII regions in the inner regions to have a better overlap between dust-to-gas and metallicity.

#### 4.3. III. Can we constrain $X_{CO}$ ?

We used Eqs. 1 and 2 along with the abundance gradients given in Table 1 to calculate the molecular gas surface densities, and then the dust-to-gas mass ratios.

The constant Galactic value of  $X_{CO}$  (dot-dashed red curves in Fig. 6) produces in most cases flat dust-to-gas mass ratio as discussed in Sect. 4.2. The super-linear dependence (dashed blue curves) results in a dust-to-gas radial gradient much steeper than the oxygen gradient, especially in the inner regions. A linear fit of the gradient in the logarithm of the dust-to-gas ratio [ $d \log(\text{dust}/\text{gas})/dR_G$ ] (Table 3) suggests that, within  $\sim 0.7 R_{25}$ , the linear metallicity dependence is able to reproduce, within the errors, the same radial slope of the metallicity gradient for NGC4254 and NGC4321. A flattening of the dust-to-gas mass ratio gradient is appreciable in the outer regions, where the HI component is dominant. For NGC4303 a super-linear metallicity dependence of  $X_{CO}$  is instead necessary to match within the errors the two gradients. For NGC4501 an  $X_{CO}$  value between the Galactic one and the one derived from the linear metallicity dependence is required.

From Fig.5 we can figure out what is happening: for a fixed Galactic  $X_{CO}$  the dust-to-gas ratio would be constant, as in shown by dot-dashed red curves in Fig.6. Once a metallicity dependence of the type described in Eqs. 1 and 2 is introduced in  $X_{CO}$ , that depresses the  $H_2$  in the centers (which have super-solar metallicity) and makes the total gas to look more like the atomic component, which generally has a much more constant distribution with  $R_G$  than the dust. Hence, when adopting the metallicity dependences of  $X_{CO}$ , we favor the outwardly decreasing gradients in dust-to-gas ratio.



**Fig. 6.** Dust-to-gas mass ratio and metallicity radial profiles of NGC4254, NGC4303, NGC4321, and NGC4501 with the oxygen abundance calibrated with KK04. The conversion factors  $X_{\text{CO}}$  are: with the Galactic conversion factor (dot-dashed red curve), with the linear metallicity dependence as in Eq.1 (dotted green curve), with the superlinear dependence on metallicity as in Eq. 2 (dashed blue curve). The magenta circles are the oxygen abundance converted to the dust-to-gas scale with Eq.3. For NGC4254 and NGC4321 we used the recalculation by Moustakas et al. (2010), for NGC4303 and NGC4501 the oxygen abundances by Skillman et al. (1996) converted to the KK04 scale. The continuous black lines show the fit to the abundance data.


**Table 3.** Slopes of the O/H and dust-to-gas gradients within  $0.7R_{25}$ .

Name (1)	slope <sub>O/H</sub> (2)	slope <sub>Gal</sub> (3)	slope <sub>linear</sub> (4)	slope <sub>superlinear</sub> (5)
NGC4254	$-0.42 \pm 0.06$	$0.07 \pm 0.06$	$-0.55 \pm 0.07$	$-1.17 \pm 0.06$
NGC4303	$-0.42 \pm 0.02$	$1.11 \pm 0.15$	$0.27 \pm 0.13$	$-0.44 \pm 0.11$
NGC4321	$-0.35 \pm 0.13$	$0.09 \pm 0.09$	$-0.52 \pm 0.08$	$-1.13 \pm 0.09$
NGC4501	$-0.07 \pm 0.16$	$0.07 \pm 0.07$	$-0.45 \pm 0.07$	$-1.18 \pm 0.12$

Given that these galaxies are dominated by the molecular gas component in their central regions, one could think that the result we found is, in a certain sense, expected: we use a  $X_{\text{CO}}$  that depends on  $1/Z$  and we obtain a dust-to-gas mass ratio decreasing as  $Z$ . However, the radial dependence of  $X_{\text{CO}}$  has only a secondary impact on our results: to obtain a decreasing dust-to-gas mass ratio it is necessary only to have a conversion factor of the order of  $\sim 1/3$ - $1/2$  the Galactic value in the central regions

of the galaxies, while the outer regions, where the atomic gas component is dominant, are less affected by the choice of  $X_{\text{CO}}$ . From Fig. 7 where we show the radial dependence of  $X_{\text{CO}}$  obtained with Eq. 1 (linear dependence) for NGC4254 (green circles), NGC4321 (yellow triangles), NGC4303 (red diamonds), and NGC4501 (magenta squares), it can be seen that no larger radial variations of  $X_{\text{CO}}$  are required. To reproduce a decreasing dust-to-gas mass ratio gradient in the radial regions where the molecular gas dominates the gaseous component, we need for the sample galaxies an  $X_{\text{CO}} \sim 1/3$  of the Galactic value, slightly increasing with radius up to  $\sim 2/3$  the Galactic value. These results agree with  $X_{\text{CO}}$  obtained in the metal rich HII regions of M31 and M51 (e.g., Arimoto et al. 1996).

To check this, we have used a constant  $X_{\text{CO}} : 0.5 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$  for NGC4254,  $0.1 \times 10^{20}$  for NGC4303,  $0.9 \times 10^{20}$  for NGC4321, and  $0.7 \times 10^{20}$  for NGC4501. These values are chosen to produce the best agreement between O/H and dust-to-gas gradients for each galaxy of our sample. They are not related to the  $X_{\text{CO}}$  values derived from the metallicity dependence in Eq.1 and shown in Fig.7. Note that the case of NGC4303 seems to be quite extreme: the  $X_{\text{CO}}$  necessary to reproduce with



**Fig. 7.** Radial variation of  $X_{\text{CO}}$  in the four galaxies with the metallicity computed with the calibration KK04 and the dependence in Eq. 1: NGC4254 (green circles), NGC4303 (red diamonds), NGC4321 (yellow triangles), and NGC4501 (magenta squares).

the dust-to-gas ratio the behaviour of the O/H gradient is much smaller than  $1/3$  Galactic  $X_{\text{CO}}$ , being  $0.1 \times 10^{20}$ .

We have compared these gradients also with the dust-to-gas mass ratio computed with constant "Galactic"  $X_{\text{CO}} = 1.8 \times 10^{20} \text{ cm}^{-2}$  (Bolatto et al. 2008) and  $X_{\text{CO}} = 4.0 \times 10^{20} \text{ cm}^{-2}$  (Draine et al. 2007). The results are shown in Fig. 8: with the low  $X_{\text{CO}}$ , the negative slope of the metallicity gradient is reproduced by the dust-to-gas mass ratio gradient. We note also that the outer regions are marginally affected by the choice of  $X_{\text{CO}}$ , and they are still consistent with a "standard"  $X_{\text{CO}}$  values for normal disk galaxies, while the inner regions need a lower value to reproduce a decreasing dust-to-gas mass ratio. This is in agreement with the evidence found for a lower  $X_{\text{CO}}$  in galaxy centers, e.g., Sodroski et al. (1994), Israel et al. (2006), Israel (2009), Watanabe et al. (2011). On the other hand, the dust-to-gas mass ratio obtained with  $X_{\text{CO}} = 1.8 \times 10^{20} \text{ cm}^{-2}$  and  $X_{\text{CO}} = 4.0 \times 10^{20} \text{ cm}^{-2}$  tend to be flatter or positive, especially with the larger value of  $X_{\text{CO}}$ . Thus, even if it is difficult to prove a dependence of  $X_{\text{CO}}$  on the metallicity due to uncertainties on the O/H and dust mass scales, the comparison between the oxygen and dust-to-gas mass ratio radial gradients in the four galaxies under analysis allows us to put a constraint on their  $X_{\text{CO}}$  conversion factors, favoring lower values of  $X_{\text{CO}}$  in the radial regions  $R_G < 0.6-0.7R_{25}$ .

In Table 4 we report the total masses of HI, H<sub>2</sub>, and dust. We computed the dust masses integrating the radial profile up to  $0.7R_{25}$ . We remind that this limit is due to the requirement of a  $S/N > 10$  at  $500 \mu\text{m}$ , necessary to limit the uncertainties due to background subtraction and avoid the artefacts caused by the high-pass filtering in the PACS data reduction. The comparison with the results of Davies et al. (2011), obtained with an integrated analysis of the galaxies, indicates that only 30% of the dust mass resides in the outer-most regions, with  $R > R_{25}$ .

Bendo et al. (2010a) performed a similar analysis on NGC2403, a late-type galaxy with a relatively low molecular fraction. Because in NGC2403 the atomic gas is the largest component of the ISM, differing assumptions about  $X_{\text{CO}}$  (they used a constant value and a metallicity dependence as in our Eq. 2)

have only a minor impact on the radial profile of the dust-to-gas mass ratio. Unlike NGC2403, in the galaxies of our sample the molecular gas dominates the ISM within  $\sim 0.5 R_{25}$ , and thus a variation of  $X_{\text{CO}}$  has a significant effect on the dust-to-gas ratio. Bendo et al. (2010a) found with both the constant and metallicity dependent  $X_{\text{CO}}$  a good agreement between the O/H gradient ( $-0.084 \pm 0.009 \text{ dex kpc}^{-1}$ ) and the dust-to-gas gradient ( $-0.093 \pm 0.002 \text{ dex kpc}^{-1}$ ), with the dust-to-gas slope slightly steeper. They explain that variations in the ratio of oxygen to the other constituents of dust might originate the difference in the gradients. For example, carbon and oxygen are both produced by supernovae, but carbon is also generated by the nucleosynthesis of low and intermediate mass stars; thus C/O is expected to vary radially, as shown by Garnett et al. (1999).


Thomas et al. (2004) obtained a similar result in their study of the radial distribution of dust and gas in several nearby galaxies for which they had neutral hydrogen and  $850 \mu\text{m}$  images available: their radial profiles of dust-to-gas mass ratio were flatter than the oxygen abundance gradients from HII regions. However in their galaxies CO observations were not available, so they were neglecting an important potentially dominant component in the inner regions.

A similar result was obtained also by Muñoz-Mateos et al. (2009) in the galaxies belonging to the SINGS sample. They have studied the radial variation of the dust-to-gas ratio in a large sample of galaxies of different morphological types. They found that the dust-to-gas mass ratio decreases by an order of magnitude from the center to the edge of the optical disk of each galaxy, similarly to what happens for the gas-phase oxygen abundance. Their relation between dust-to-gas mass ratio and oxygen abundance can be described with a linear scaling law for  $12 + \log(\text{O}/\text{H}) > 8.9$ , as we found for NGC4254, NGC4321, and NGC4501. At lower metallicities, they found that the dust-to-gas ratios are systematically below this simple relation. They explain this fact to be due to the large amounts of gas that has not yet undergone star formation activity and that reside in the outer regions of spiral disks.

#### 4.4. Other evidence for a low $X_{\text{CO}}$

In agreement with our result, a value of  $X_{\text{CO}}$  lower than  $\sim 2 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$  was also expected in NGC4254 and NGC4321 from the analysis of Eales et al. (2010). They calibrated the mass-opacity coefficient ( $\kappa_v$ ) using their dust-to-gas maps. Using a standard  $X_{\text{CO}}$  value they found a mass-opacity coefficient significantly lower than the value ( $0.41 \text{ m}^2 \text{ kg}^{-1}$ ) obtained by James et al. (2002). To solve the discrepancy, they suggested that using only SPIRE bands they are over-estimating the temperature of the dust; if the temperature of the dust in both NGC4254 and NGC4321 were actually  $\approx 10 \text{ K}$  rather than  $20 \text{ K}$ , this would be enough to increase the value of  $\kappa_v$  to that expected from the James et al. study. However, this is not the case since with our fit of PACS and SPIRE bands we found dust temperatures in the range  $17 < T < 25 \text{ K}$  found by Eales et al. (2010) (see Fig. 3).

The other possibility they suggested to reconcile their results with James et al. (2002) was to have values of  $X_{\text{CO}} \approx 6$  times lower than the initial value they assumed,  $\sim 2 \times 10^{20} \text{ cm}^{-2} \text{ K km s}^{-1})^{-1}$ . Considering that we are using a mass-opacity coefficient which is about half of that used by James et al., thus, to have agreement with our results, they would need  $X_{\text{CO}} \approx 3$  times lower than their standard value. This is consistent with our result; val-



**Fig. 8.** Dust-to-gas mass ratio and metallicity radial profiles with the conversion factor  $X_{\text{CO}} = 0.1\text{--}0.9 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$  (see text for the  $X_{\text{CO}}$  adopted for each galaxy—orange solid curves), and with constant  $X_{\text{CO}} = 1.8 \times 10^{20} \text{ cm}^{-2}$  (red dot-dashed curves) and  $X_{\text{CO}} = 4.0 \times 10^{20} \text{ cm}^{-2}$  (green dotted curves). The magenta circles are the oxygen abundance converted to the dust-to-gas scale with Eq.3. The continuous black lines show the fit to the abundance data.

**Table 4.** HI, H<sub>2</sub>, and dust masses

Name	M(HI) 10 <sup>9</sup> M <sub>⊙</sub>	M(H <sub>2</sub> ) <sub>unif.</sub> 10 <sup>9</sup> M <sub>⊙</sub>	M(H <sub>2</sub> ) <sub>linear</sub> 10 <sup>9</sup> M <sub>⊙</sub>	M(H <sub>2</sub> ) <sub>superlinear</sub> 10 <sup>9</sup> M <sub>⊙</sub>	M(dust) 10 <sup>7</sup> M <sub>⊙</sub>
(1)	(2)	(3)	(4)	(5)	(6)
NGC4254	4.4 <sup>a</sup>	5.4	1.8	0.22	7.0
NGC4321	4.8 <sup>a</sup>	4.4	1.3	0.13	8.6
NGC4303	3.3 <sup>b</sup>	4.8	0.8	0.06	5.1
NGC4501	2.7 <sup>a</sup>	3.8	1.0	0.05	7.4

**Notes.** *a* Chung et al. (2009a)

*b* integration of the radial profile shown in Fig.3 of Skillman et al. (1996).

ues of  $X_{\text{CO}}$  in the range  $\sim 0.5\text{--}1 \times 10^{20} \text{ cm}^{-2} \text{ K km s}^{-1})^{-1}$  can reproduce both the correct value of the mass-opacity coefficient and the radial profiles of dust-to-gas mass ratio.

## 5. Summary and conclusions

1. We analyzed FIR observations obtained with PACS and SPIRE, together with CO and HI maps from the literature, of four spiral galaxies in the Virgo cluster (NGC4254,

NGC4303, NGC4321, and NGC4501). We derived the total mass of dust from SED fitting of the FIR images, and used HI moment-1 maps to derive the geometric parameters of the galaxies, which are disturbed by tidal interactions. Finally, we extracted the radial profiles of atomic gas, molecular gas and dust, and compare them with oxygen abundance radial gradients compiled from the literature and placed on a common abundance scale.

2. To avoid the large uncertainties on the zero-points of the oxygen abundance derived with the bright-line methods, we used the dust-to-gas ratio obtained with the Galactic value of  $X_{\text{CO}}$  to fix a lower limit to the oxygen abundance. This allowed us to discard a set of metallicity calibrations and to reduce the uncertainty on the oxygen gradient zero-point. We converted the literature O/H to the scale of KK04.
3. We studied the dependence of  $X_{\text{CO}}$  on metallicity by comparing the radial gradient of metal abundance in gas phase (oxygen abundance in HII regions) and in solid phase (dust in emission from the FIR versus the total gas–atomic and molecular), assuming that these two quantities decrease at the same rate with radius. We considered a constant  $X_{\text{CO}}$  (e.g., Bolatto et al. 2008),  $X_{\text{CO}} \propto Z^{-1}$  (“linear” dependence) (e.g., Boselli et al. 2002), and  $X_{\text{CO}} \propto Z^{-2.5}$  (“super-linear” dependence) (e.g., Israel 2000). A linear fit of the gradient in the logarithm of the dust-to-gas ratio [ $d \log(\text{dust/gas})/dR_G$ ] shows that, within  $\sim 0.7 R_{25}$ , values of  $X_{\text{CO}}$  in the range  $\sim 0.5\text{--}0.9 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$  were able to reproduce negative gradients, similar to the O/H ones for NGC4254, NGC4321, and NGC4501, while NGC4303 needs an extremely low  $X_{\text{CO}}$  value  $\sim 0.1 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$ . For NGC4254, these  $X_{\text{CO}}$  values can be obtained with a linear metallicity dependence and oxygen abundance calibrated by KK04. For NGC4303 a superlinear dependence is instead necessary to match within the errors the two gradients, while for NGC4501 and NGC4321 a  $X_{\text{CO}}$  intermediate between the Galactic one and the linear metallicity dependence would be necessary (always with KK04 O/H calibration). We suggest that a  $X_{\text{CO}}$  lower than the standard Galactic one (e.g., Bolatto et al. 2008) is necessary in these galaxies to obtain a decreasing dust-to-gas mass ratio similar to the O/H gradient of HII regions. These low  $X_{\text{CO}}$  values are favored in the radial regions  $R_G < 0.6\text{--}0.7 R_{25}$  where the molecular gas dominates, while the outer regions, where the atomic gas is the main component, are less affected by the choice of  $X_{\text{CO}}$ , and thus we cannot put constraints on its value. A large sample of galaxies with available metallicity and dust-to-gas mass ratio gradients is necessary to confirm and strengthen these results.

*Acknowledgements.* We thank an anonymous referee for comments and suggestions which have improved the quality of the manuscript and its presentation. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. L.M. and C.P. are supported through an ASI-INAF grant “HeViCS: the Herschel Virgo Cluster Survey” I/009/10/0. C.V. received support from the ALMA-CONICYT Fund for the Development of Chilean Astronomy (Project 31090013) and from the Center of Excellence in Astrophysics and Associated Technologies (PBF 06).

## References

- Arimoto, N., Sofue, Y., & Tsujimoto, T. 1996, PASJ, 48, 275  
 Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, ARA&A, 47, 481  
 Barone, L. T., Heithausen, A., Hüttemeister, S., Fritz, T., & Klein, U. 2000, MNRAS, 317, 649  
 Bendo, G. J., et al. 2010a, MNRAS, 402, 1409  
 Bendo, G. J., et al. 2010b, A&A, 518, L65  
 Bianchi, S., Davies, J. I., & Alton, P. B. 2000, A&A, 359, 65  
 Bianchi, S., Gonçalves, J., Albrecht, M., Caselli, P., Chini, R., Galli, D., & Walmsley, M. 2003, A&A, 399, L43  
 Binggeli, B., Sandage, A., & Tammann, G. A. 1985, AJ, 90, 1681  
 Boissier, S., Boselli, A., Buat, V., Donas, J., & Milliard, B. 2004, A&A, 424, 465  
 Boissier, S., et al. 2005, ApJ, 619, L83  
 Bolatto, A. D., Leroy, A. K., Rosolowsky, E., Walter, F., & Blitz, L. 2008, ApJ, 686, 948  
 Boselli, A., Lequeux, J., & Gavazzi, G. 2002, A&A, 384, 33  
 Boselli, A., & Gavazzi, G. 2006, PASP, 118, 517  
 Boselli, A., et al. 2010, PASP, 122, 261  
 Bouchet, P., Lequeux, J., Maurice, E., Prevot, L., & Prevot-Burnichon, M. L. 1985, A&A, 149, 330  
 Canzian, B., & Allen, R. J. 1997, ApJ, 479, 723  
 Cayatte, V., van Gorkom, J. H., Balkowski, C., & Kotanyi, C. 1990, AJ, 100, 604  
 Chung, A., van Gorkom, J. H., Kenney, J. D. P., Crowl, H., & Vollmer, B. 2009, AJ, 138, 1741  
 Chung, E. J., Rhee, M.-H., Kim, H., Yun, M. S., Heyer, M., & Young, J. S. 2009, ApJS, 184, 199  
 Dame, T. M., Hartmann, D., & Thaddeus, P. 2001, ApJ, 547, 792  
 Davies, J. I., et al. 2010, A&A, 518, L48  
 Davies, J. I., et al. 2011, MNRAS, Paper VIII  
 Denicoló, G., Terlevich, R., & Terlevich, E. 2002, MNRAS, 330, 69  
 Draine, B. T. 2003, ARA&A, 41, 241  
 Draine, B. T., et al. 2007, ApJ, 663, 866  
 Draine, B. T., & Li, A. 2007, ApJ, 657, 810  
 Dwek, E. 1998, ApJ, 501, 643  
 Eales, S. A., et al. 2010, A&A, 518, L62  
 Garnett, D. R., Shields, G. A., Peimbert, M., Torres-Peimbert, S., Skillman, E. D., Dufour, R. J., Terlevich, E., & Terlevich, R. J. 1999, ApJ, 513, 168  
 Gavazzi, G., Boselli, A., Scodreggio, M., Pierini, D., & Belsole, E. 1999, MNRAS, 304, 595  
 Glover, S. C. O., & Mac Low, M.-M. 2011, MNRAS, 110  
 Griffin, M. J., et al. 2010, A&A, 518, L3  
 Grevesse, N., & Anders, E. 1989, Cosmic Abundances of Matter, 183, 1  
 Henry, R. B. C., Pagel, B. E. J., & Chincarini, G. L. 1994, MNRAS, 266, 421  
 Hirashita, H., Kaneda, H., Onaka, T., & Suzuki, T. 2008, PASJ, 60, 477  
 James, A., Dunne, L., Eales, S., & Edmunds, M. G. 2002, MNRAS, 335, 753  
 Kenney, J. D., & Young, J. S. 1988, ApJS, 66, 261  
 Kennicutt, R. C., Jr., et al. 2003, PASP, 115, 928  
 Kewley, L. J., & Dopita, M. A. 2002, ApJS, 142, 35  
 Kewley, L. J., & Ellison, S. L. 2008, ApJ, 681, 1183  
 Kobulnicky, H. A., & Kewley, L. J. 2004, ApJ, 617, 240, KK04  
 Koda, J., & Sofue, Y. 2006, PASJ, 58, 299  
 Koornneef, J. 1982, A&A, 107, 247  
 Kuno, N., et al. 2007, PASJ, 59, 117  
 Inoue, A. K. 2003, PASJ, 55, 901  
 Israel, F. P., de Graauw, T., van de Stadt, H., & de Vries, C. P. 1986, ApJ, 303, 186  
 Israel, F. 2000, Molecular Hydrogen in Space, 293  
 Israel, F. P., Baas, F., Rudy, R. J., Skillman, E. D., & Woodward, C. E. 2003, A&A, 397, 87  
 Israel, F. P., Tilanus, R. P. J., & Baas, F. 2006, A&A, 445, 907  
 Israel, F. P. 2009, A&A, 506, 689  
 Issa, M. R., MacLaren, I., & Wolfendale, A. W. 1990, A&A, 236, 237  
 Lebrun, F., et al. 1983, ApJ, 274, 231  
 Leroy, A. K., et al. 2009, ApJ, 702, 352  
 Lisenfeld, U., & Ferrara, A. 1998, ApJ, 496, 145  
 Maloney, P., & Black, J. H. 1988, ApJ, 325, 389  
 McCall, M. L., Rybski, P. M., & Shields, G. A. 1985, ApJS, 57, 1  
 McGaugh, S. S. 1991, ApJ, 380, 140  
 Moustakas, J., Kennicutt, R. C., Jr., Tremonti, C. A., Dale, D. A., Smith, J.-D. T., & Calzetti, D. 2010, ApJS, 190, 233  
 Muñoz-Mateos, J. C., et al. 2009, ApJ, 701, 1965  
 Paradis, D., et al. 2010, A&A, 520, L8  
 Pettini, M., & Pagel, B. E. J. 2004, MNRAS, 348, L59  
 Phookun, B., Vogel, S. N., & Mundy, L. G. 1993, ApJ, 418, 113  
 Pilbratt, G. L., et al. 2010, A&A, 518, L1  
 Pilyugin, L. S. 2001, A&A, 374, 412  
 Pilyugin, L. S., & Thuan, T. X. 2005, ApJ, 631, 231  
 Planck Collaboration, et al. 2011, arXiv:1101.2032  
 Poglitsch, A., et al. 2010, A&A, 518, L2  
 Pohlen, M., et al. 2010, A&A, 518, L72  
 Reshetnikov, V. P. 2000, Astrophysics, 43, 145  
 Roman-Duval, J., et al. 2010, A&A, 518, L74  
 Rudolph, A. L., Fich, M., Bell, G. R., Norsen, T., Simpson, J. P., Haas, M. R., & Erickson, E. F. 2006, ApJS, 162, 346  
 Rupke, D. S. N., Kewley, L. J., & Chien, L.-H. 2010, ApJ, 723, 1255  
 Sauvage, M., et al. 2010, A&A, 518, L64  
 Shetty, R., Glover, S. C., Dullemond, C. P., & Klessen, R. S. 2011, MNRAS, 11  
 Shields, G. A., Skillman, E. D., & Kennicutt, R. C., Jr. 1991, ApJ, 371, 82

- Skillman, E. D., Kennicutt, R. C., Jr., Shields, G. A., & Zaritsky, D. 1996, *ApJ*, 462, 147
- Smith, M. W. L., et al. 2010, *A&A*, 518, L51
- Sodroski, T. J., et al. 1994, *ApJ*, 428, 638
- Sofue, Y., Koda, J., Nakanishi, H., & Hidaka, M. 2003, *PASJ*, 55, 75
- Stanghellini, L., Magrini, L., Villaver, E., & Galli, D. 2010, *A&A*, 521, A3
- Strong, A. W., & Mattox, J. R. 1996, *A&A*, 308, L21
- Strong, A. W., Moskalenko, I. V., Reimer, O., Digel, S., & Diehl, R. 2004, *A&A*, 422, L47
- Teuben, P. 1995, *Astronomical Data Analysis Software and Systems IV*, 77, 398
- Thilker, D. A., et al. 2007, *ApJS*, 173, 572
- Thomas, H. C., Alexander, P., Clemens, M. S., Green, D. A., Dunne, L., & Eales, S. 2004, *MNRAS*, 351, 362
- Tremonti, C. A., et al. 2004, *ApJ*, 613, 898
- Vollmer, B., Soida, M., Chung, A., van Gorkom, J. H., Otmianowska-Mazur, K., Beck, R., Urbanik, M., & Kenney, J. D. P. 2008, *A&A*, 483, 89
- Warmels, R. H. 1988, *A&AS*, 73, 453
- Watanabe, Y., Sorai, K., Kuno, N., & Habe, A. 2011, *MNRAS*, 411, 1409
- Wilson, C. D. 1995, *ApJ*, 448, L97
- Zaritsky, D., Kennicutt, R. C., Jr., & Huchra, J. P. 1994, *ApJ*, 420, 87