

Characteristics of convection and overshooting in RGB and AGB stars^{*}

X.-J. Lai^{1,2,3} and Y. Li^{1,2}

¹ National Astronomical Observatories/Yunnan Observatory, Chinese Academy of Sciences, Kunming 650011, China; lxj@mail.ynao.ac.cn, ly@ynao.ac.cn

² Key Laboratory for the Structure and Evolution of Celestial Objects, Chinese Academy of Sciences, Kunming 650011, China

³ Graduate School of the Chinese Academy of Sciences, Beijing 100049, China.

Abstract Based on the turbulent convection model (TCM) of Li & Yang (2007), we have studied the characteristics of turbulent convection in the envelopes of 2 and $5M_{\odot}$ stars at the RGB and AGB phases. The TCM has been applied successfully in the whole convective envelopes including the convective unstable zone and the overshooting regions. We find that the convective motions become stronger and stronger when the stellar models are located upper and upper along the Hayashi line. In the convective unstable zone, we find that the turbulent correlations are proportional to functions of a common factor $(\nabla - \nabla_{ad})\bar{T}$, which explains similar distributions of those correlations. For the TCM we find that if the obtained stellar structure of temperature is close to that of the MLT, the convective motion will be at a much larger velocity and thus more violent. However, if the turbulent velocity is adjusted to close to that of the MLT, the superadiabatic convection zone is much more extended inward and a lower effective temperature of the stellar model will be obtained. For the overshooting distance we find that the e-folding lengths of the turbulent kinetic energy k in both the top and bottom overshooting regions decrease as the stellar model is located up along the Hayashi line, but both the extents of the decrease are not obvious. And the overshooting distances of the turbulent correlation $\overline{u'_r T'}$ are almost the same for the different stellar models with a same set of TCM's parameters. For the decay way of the kinetic energy k we find that they are very similar for the different stellar models based on a same set of TCM's parameters, and there is a nearly linear relationship between $\lg k$ and $\ln P$ for the different stellar models. When C_s or α increases while the other parameters are fixed, the obtained linearly decaying distance will become longer.

Key words: RGB and AGB stars: convection of stars: turbulent convection model: convective overshooting

1 INTRODUCTION

Red giant branch (RGB) and asymptotic giant branch (AGB) stars are characterized by the convective motion in their envelopes, which can extend over an enormous range in mass (or radius). Because of huge scale of convection zones and small viscosity of the stellar material, turbulent flow instead of laminar one always occurs in their outer envelopes. Turbulent convection may result in many important effects in stars: mixing different elements to be homogeneous in the convection zone and adding fresh

* Supported by the National Natural Science Foundation of China

fuel to the nuclear burning region, dredging the internal material processed by H or He burning up to the stellar surface, and being as an important means for heat transfer. These effects significantly influence the structure and evolution of stars. Many observational phenomena appearing in the RGB and AGB stars are ascribed to the convection. For the RGB stars as an example, many of them show chemical anomalies (Pilachowski et al. 1993; Charbonnel 1995; Gratton et al. 2000; Kraft et al. 1993; Ramírez & Cohen 2002; Gratton et al. 2004; Busso et al. 2007; Recio-Blanco & de Laverny 2007). And we are still perplexed to the so-called carbon star mystery (Iben 1981; Iben 1975, 1977; Sackmann 1980; Lattanzio 1989; Hollowell & Iben 1988; Straniero et al. 1997; Herwig et al. 1997; Herwig 2005) for the AGB stars. Furthermore, the observed mass loss occurring in the RGB and AGB stars may be due to the turbulent pressure (Jiang & Huang 1997). To solve these problems we need to accurately deal with the convection and to learn details of its characteristics.

For some RGB and AGB stars the convective motion near the surface of the stellar envelope are superadiabtic because of less efficient convective energy transport, and may sometimes become supersonic based on the mixing-length theory (MLT) (Deng & Xiong 2001). The MLT (Böhm-Vitense 1953, 1958) is commonly used to deal with the convection. However, due to its simplicity many shortcomings are found (Renzini 1987; Baker & Kuhfuss 1987; Spruit et al. 1990; Pederson et al. 1990), which thus lead to some confused and debatable problems. For example, convective overshooting (e.g. Saslaw & Schwarzschild 1965), semiconvection (e.g. Castellani et al. 1971; Sreenivasan & Wilson 1978) and the breathing convection (e.g. Castellani et al. 1985). Therefore new convection models have been proposed successively, for example, non-local mixing length theories (Ulrich 1970; Maeder 1975; Bressan et al. 1981) and full-spectrum convection theory (Canuto & Mazzitelli 1991). However, they are still based on the framework of the MLT. A better theory to overcome the MLT's drawbacks is the turbulent convection models (TCMs), which are deduced directly from the Navier-Stokes equations, and can describe many turbulence properties (Xiong 1979, 1981, 1985, 1990; Canuto 1992, 1994, 1997; Xiong, Cheng & Deng 1997; Canuto & Dubovikov 1998; Canuto et al. 1996). However, the dynamic equations for turbulent correlations are expressed by higher order correspondents due to the inherent non-linearity of the Navier-Stokes equations. Therefore these equations are needed to be cut-off with some closure approximations in order to be applicable in calculations of the stellar structure and evolution. Nevertheless, some free parameters have to be introduced by the closure assumptions. Different closure assumptions correspond to different TCMs, and many of them may not be easy to be incorporated into a stellar evolution code. Recently a simple TCM proposed by Li & Yang (2007) was successfully applied to the solar models (Yang & Li 2007), in which the introduced parameters of the TCM are directly related to the corresponding turbulence physics. Some significant changes in the structure of the solar convection zone and better results compared to the solar p-mode observations are obtained. Shortly afterwards, Zhang & Li (2009) applied it successfully to the overshooting region of the solar convection zone.

Up to now, the TCMs have seldom been applied to very large convective envelopes of stars, e.g. in the RGB or AGB stars, which are rather different from the solar environment. Furthermore, in view of the key roles of convection played in these stars, we choose three sites of stellar evolution to test the TCM of Li & Yang (2007). We try to find out convection characteristics for two stars of 5 and $2M_{\odot}$ at the RGB and AGB phases, and to find out their dependence on the TCM's parameter. To analyze the properties of the turbulent correlations among velocity and temperature fluctuations, we separate the whole convection envelope (derived self-consistently from the TCM) into the convective unstable zone and the convective overshooting regions, respectively. We firstly describe the basic equations of the TCM in Section 2. The information of the evolutionary code and input physics are described in Section 3. In Section 4, the convection characteristics in the convective unstable zone and overshooting regions are presented and discussed, respectively. The dependence of the structure of the overshooting regions on the TCM's parameters is given in Section 5. Finally, some concluding remarks are summarized in Section 6.

2 EQUATIONS OF TCM

Using the Reynolds decomposition approximation, and with the aid of some closure approximations, the second-order moment equations of the TCM are (Li & Yang 2007 for details):

$$\frac{1}{\bar{\rho}r^2} \frac{\partial}{\partial \ln \bar{P}} \left(C_s \bar{\rho} r^2 \frac{\overline{u'_r u'_r}}{\sqrt{k}} \frac{\partial \overline{u'_r u'_r}}{\partial \ln \bar{P}} \right) = C_k \frac{\sqrt{k}}{\alpha} \left(\overline{u'_r u'_r} - \frac{2}{3} k \right) + \frac{2}{3} \frac{k^{3/2}}{\alpha} + \frac{2\beta g_r}{\bar{T}} H_P \overline{u'_r T'}, \quad (1)$$

$$\frac{1}{\bar{\rho}r^2} \frac{\partial}{\partial \ln \bar{P}} \left(C_s \bar{\rho} r^2 \frac{\overline{u'_r u'_r}}{\sqrt{k}} \frac{\partial k}{\partial \ln \bar{P}} \right) = \frac{k^{3/2}}{\alpha} + \frac{\beta g_r}{\bar{T}} H_P \overline{u'_r T'}, \quad (2)$$

$$\begin{aligned} \frac{2}{\bar{\rho}r^2} \frac{\partial}{\partial \ln \bar{P}} \left(C_{t1} \bar{\rho} r^2 \frac{\overline{u'_r u'_r}}{\sqrt{k}} \frac{\partial \overline{u'_r T'}}{\partial \ln \bar{P}} \right) &= \left[\frac{\bar{\rho} c_p H_P}{\lambda} \overline{u'_r T'} - \bar{T} (\nabla_R - \nabla_{ad}) \right] \overline{u'_r u'_r} + \\ &\frac{\beta g_r}{\bar{T}} H_P \overline{T'^2} + C_t \left(\frac{\sqrt{k}}{\alpha} + \frac{\lambda}{\bar{\rho} c_p \alpha^2 H_P} \frac{\varepsilon^2}{k^3} \right) \overline{u'_r T'}, \end{aligned} \quad (3)$$

$$\begin{aligned} \frac{1}{\bar{\rho}r^2} \frac{\partial}{\partial \ln \bar{P}} \left(C_{e1} \bar{\rho} r^2 \frac{\overline{u'_r u'_r}}{\sqrt{k}} \frac{\partial \overline{T'^2}}{\partial \ln \bar{P}} \right) &= 2 \left[\frac{\bar{\rho} c_p H_P}{\lambda} \overline{u'_r T'} - \bar{T} (\nabla_R - \nabla_{ad}) \right] \overline{u'_r T'} + \\ &2C_e \left(\frac{\sqrt{k}}{\alpha} + \frac{\lambda}{\bar{\rho} c_p \alpha^2 H_P} \right) \overline{T'^2}. \end{aligned} \quad (4)$$

The terms on the left-hand side of Eqs.(1)-(4) are diffusion ones of the turbulence, representing the non-local characteristics of turbulence and describing mainly the properties of the overshooting regions outside the Schwarzschild boundaries of a convection zone. The terms on the right-hand side of Eqs.(1)-(4) are some production and dissipation ones of the turbulence. All the symbols appeared in above equations have their usual meaning (Li & Yang 2001; Li & Yang 2007). There are seven free parameters in above equations. They are C_t , C_e , C_k , C_{t1} , C_{e1} , C_s , and α . The former three ones are dissipation parameters which describe respectively the dissipation of the turbulent correlation $\overline{u'_r T'}$, the dissipation of the temperature fluctuation $\overline{T'^2}$, and actually the anisotropic degree of the turbulence. The larger the values of them are, the smaller the convective heat fluxes will be (Li & Yang 2007). The next three parameters are diffusion ones related respectively to the three turbulent correlations $\overline{u'_r T'}$, $\overline{T'^2}$ and $\overline{u'_r u'_r}$. Larger values of them will result in lowered and expanded profiles of these correlations (Li & Yang 2007). The last parameter α , the ratio of a typical length l to the local pressure scaleheight, is introduced by the closure assumption similar to the mixing length parameter in the MLT. The effects of these parameters on the properties of turbulence, and then on the structure of the solar convection zone, are investigated by Li & Yang (2007) and Zhang & Li (2009). Here we choose several sets of these parameters' values to study the properties of the turbulent convection in the RGB and AGB stars.

3 INPUT PHYSICS

We calculate the evolution of two stars of 2 and $5M_\odot$ respectively. The initial composition ($Z=0.02$, $X=0.7$) is fixed for both stellar models.

The free parameters adopted in the TCM can be determined by the fluid dynamics experiments in the terrestrial environment (e.g. Hossain & Rodi 1982). Given a set of appropriate values of dissipation and diffusion parameters, the value of α can be determined by calibrating the solar model (Li & Yang 2007; Zhang & Li 2009). The value of α obtained in this way can be used to RGB or AGB calculations, but this approach is at the sacrifice of the uncertainty of its value (Herwig 2005). In this paper, we adopt several sets of the parameters' values, which are derived directly from or mainly based on the works of Li & Yang (2007) and Zhang & Li (2009) for the solar model and are shown in Table 1. In which the values of C_t , C_e and C_k are mainly derived from the terrestrial experiments and their typical values are 3.0, 1.25 and 2.5, respectively. Nevertheless, $C_t = 7.0$ is taken from the work of Canuto (1993)

Table 1 The information of the TCM's parameters

Model Name	C_t	C_e	C_k	C_{t1}	C_{e1}	C_s	α
MLT							1.70
LM	3.0	1.25	2.5				0.90
NLMa	3.0	1.25	2.5	0.03	0.03	0.03	0.90
NLMb1	7.0	0.20	2.5	1.0×10^{-7}	1.0×10^{-7}	0.10	0.76
NLMb2	7.0	0.20	2.5	1.0×10^{-7}	1.0×10^{-7}	0.10	0.74
NLMb3	7.0	0.20	2.5	1.0×10^{-7}	1.0×10^{-7}	0.10	0.15
NLMc1	3.0	1.25	2.5	0.05	0.05	0.05	0.68
NLMc2	3.0	1.25	2.5	0.05	0.05	0.04	0.68
NLMc3	3.0	1.25	2.5	0.05	0.05	0.03	0.68
NLMc4	3.0	1.25	2.5	0.05	0.05	0.02	0.68
NLMc5	3.0	1.25	2.5	0.05	0.05	0.01	0.68
NLMd	0.2	0.10	2.5	0.15	0.25	0.10	0.05

and $C_t = 0.20$, $C_e = 0.10$ from the suggestion that smaller values of them will lead to a better result compared to the solar p-mode observation (Yang & Li 2007). Furthermore, $C_{t1} = C_{e1} = 10^{-7}$ are adopted, meaning a negligible diffusion effect of $\overline{u'_r T'}$ and $\overline{T'^2}$ and a local convection model of them, to assure that an almost same temperature structure in stellar model can be obtained for both the MLT and TCM and then a meaningful comparison between them can be done, which will be discussed in the following subsection 4.3. For simplicity we call the TCM with the diffusion terms in Eqs.(1)-(4) as NLMs and the TCM without the diffusion terms as LM. The MLT with $\alpha = 1.70$ is also adopted to calculate the stellar convection at the RGB and AGB phases for comparisons. In Table 1 both the parameter sets of NLMa and NLMd are corresponding to the standard solar model calibration.

The stellar evolution code used in the present paper was originally described by B. Paczynski and M. Kozłowski and updated by R. Sieniewicz. Nuclear reaction rates are adopted from BP95 (Bahcall, Pinsonneault & Wasserburg 1995). The equation of state is the OPAL EOS from Rogers (1994) and Rogers, Swenson & Iglesias (1996). The OPAL opacities GN93hz series (Rogers & Iglesias 1995; Iglesias & Rogers 1996) are used in the high-temperature region. In the stellar atmosphere, low-temperature opacities from Alexander & Ferguson (1994) are used.

To study the convection properties based on the TCM, we select three evolved models during the RGB and AGB phases derived from the MLT, which are indicated by solid stars in Fig.1.




Fig. 1 The HR diagram for the stellar models of 2 and $5M_{\odot}$ based on the MLT. The three solid stars indicate the certain locations of the stellar models during the RGB ($2M_{\odot}$: $\lg L/L_{\odot} = 1.58$, $T_{eff} = 4750K$; $5M_{\odot}$: $\lg L/L_{\odot} = 3.09$, $T_{eff} = 4190K$) and AGB ($5M_{\odot}$: $\lg L/L_{\odot} = 3.81$, $T_{eff} = 3570K$) phases




Fig.2 Distributions of the turbulent correlations for the $5M_{\odot}$ star at the AGB phase. The solid lines are for NLMa, dashed lines for LM.




Fig.3 The same as Fig.2, but for the $5M_{\odot}$ star at the RGB phase.

4 CHARACTERISTICS OF TURBULENCE

4.1 Profiles of the Turbulent Correlations

The profiles of the turbulent correlations \sqrt{k} , $\overline{u'_r T'}$ and $\overline{T'^2}$ are displayed in Fig.2 for the $5M_{\odot}$ model at the AGB phase. It can be seen that convection widely develops in the stellar envelope ($3.5 \leq \lg T \leq 6.4$) and the turbulent kinetic energy k maintains at a high level in most of the convection zone. Particularly, there are several peaks for all the three correlations appearing at almost the same locations. The appearance of the maximum turbulent kinetic energy is mainly due to ionization of hydrogen and helium which intensively blocks the transfer of heat and results in strong buoyancy to drive the convective motion. The profiles of the correlations $\overline{u'_r T'}$ and $\overline{T'^2}$ vary in a similar way as that of \sqrt{k} , except that the largest peaks are much higher than that of \sqrt{k} .

Diffusion terms in the turbulence equations are taken into account in NLMa but not in LM. As a result the non-local effect is behaved remarkably for NLMa around the boundaries of the convective unstable zone, especially around the upper boundary where the NLMa's solution significantly deviates from the LM's. However, the non-local effect can be safely ignored in the interior of the convective unstable zone because the turbulent correlations distribute quite homogeneously there. These conclusions agree well with those of Yang & Li (2007) and Deng & Xiong (2008).

However, for the other two stellar models of 5 and $2M_{\odot}$ at the RGB phase, the profiles of the three correlations, which are respectively shown in Fig.3 and Fig.4, have some differences for both NLMa and LM. These two models have smaller values of the three correlations. It means that the turbulent motion is less violent and thus transfers less heat. Furthermore, except for the maximum peak of those




Fig. 4 The same as Fig.2, but for the $2M_{\odot}$ star at the RGB phase.

correlations the other peaks become less prominent, especially for the model of $2M_{\odot}$ at the RGB phase. Besides, near the bottom of the convection envelope the turbulent kinetic energy k has a weak peak, and its value seems to decrease monotonically with the stellar luminosity. In another word, the convective motion becomes stronger and stronger when the stellar models are located up along the Hayashi line.

4.2 Turbulence Properties in the Convective Unstable Zone

In order to understand the properties of turbulence in the convective unstable zone, some assumptions are invoked to obtain the explicit relationships between the correlations \sqrt{k} , $\overline{u'_r T'}$ and $\overline{T'^2}$. According to the result stated in the former section, the diffusion terms on the left hand sides of Eqs.(1)–(4) can be approximately ignored in the convective unstable zone. With the aid of Eqs.(1)–(4) we get:

$$\overline{T'^2} = \frac{D(\nabla - \nabla_{ad})\overline{T}^2 k - D(\nabla - \nabla_{ad})\overline{T}^2 k \sqrt{1 + 4 \frac{C_t}{C_e} \frac{k}{D^2 \alpha^2 H_P g_r \beta (\nabla - \nabla)}}}{2H_P g_r \beta}, \quad (5)$$

where $D = \frac{4}{3C_k} + \frac{2}{3}$. In view of the mixing-length theory:

$$k = \frac{g\delta}{8H_P} (\nabla - \nabla_{ad})l^2 \sim c_p \nabla_{ad} (\nabla - \nabla_{ad})\overline{T}, \quad (6)$$

Eq.(5) can be further reduced to be:

$$\overline{T'^2} \sim \frac{D\alpha^2}{8} (\nabla - \nabla_{ad})^2 \overline{T}^2. \quad (7)$$

By utilizing Eq.(6) and Eq.(2), the velocity-temperature correlation can be approximated as:

$$\overline{u'_r T'} \sim \sqrt{c_p \nabla_{ad}} (\nabla - \nabla_{ad})^{3/2} \overline{T}^{3/2}. \quad (8)$$

It can be seen that all the three correlations depend on $(\nabla - \nabla_{ad})\overline{T}$, which explains why the three correlations peak at almost the same places in the convective unstable zone. Fig.5 shows the profiles

of k , $(\overline{u_r T'})^{2/3}$, $(\overline{T'^2})^{1/2}$ and $(\nabla - \nabla_{ad})\overline{T}$ for NLMa of the $5M_\odot$ star at the AGB phase. It can be seen that the profiles of $(\overline{u_r T'})^{2/3}$, $(\overline{T'^2})^{1/2}$ and $(\nabla - \nabla_{ad})\overline{T}$ behave in a very similar way in most of the convective unstable zone. However, the profile of k deviates significantly from the others, but its overall trend is almost consistent with them. It should be noted that the above results are suitable for other stellar models.




Fig. 5 Profiles of k , $(\overline{u_r T'})^{2/3}$, $(\overline{T'^2})^{1/2}$ and $(\nabla - \nabla_{ad})\overline{T}$ at the AGB phase of the $5M_\odot$ star based on NLMa.

4.3 Comparisons with the MLT




Fig. 6 The temperature gradient (including the real, radiative and adiabatic ones) and the square root of the turbulent energy \sqrt{k} as functions of the temperature for the $5M_\odot$ star at the AGB phase (the same as in Fig.2). The dashed line is for MLT, the solid lines in left panels are for NLMb1 and the solid lines in right panels are for NLMb3.

Results of two TCMs (NLMb1 and NLMb3) are compared with that of the MLT. Firstly, for the comparison between NLMb1 and the MLT in the left panel of Fig.6, we can find that the temperature gradient ∇ are almost the same with each other, which means that NLMb1 will result in a similar stellar structure as the MLT does in the convective unstable zone. However, the value of the turbulence velocity ($\sim \sqrt{k}$) of NLMb1 is much larger than that of the MLT. In another word, for a fixed heat flux transferred by the convective motion, a larger value of convective velocity is requested for the TCM. This is partly because that the dissipation effects of the turbulent convection, measured by the parameters C_t , C_e and C_k , are fully taken into account.

When decreasing the value of parameter α , for example $\alpha = 0.15$, however, we obtain an almost equivalent profile of the convective velocity as the result of the MLT, which is shown in the right panel of Fig.6 (namely NLMb3). It can be found that the temperature gradient ∇ of NLMb3 is much larger than that of the MLT near the top of the convection envelope and extending to $\lg T \approx 4.7$ where it approaches the adiabatic temperature gradient ∇_{ad} , a place much deeper than the case of the MLT being approximately at $\lg T \approx 4.1$. It means that the superadiabatic convection region is much more extended inward for the TCM than the case of the MLT and the stellar models based on the TCM will be of lower effective temperature than that of the MLT.

The results obtained above are mainly based on the star of $5M_\odot$ at the AGB phase. Likewise, these conclusions are still valid for the $2M_\odot$ star at the RGB phase and the $5M_\odot$ star at the RGB phase. Fig.7, as an example, shows the case of the $2M_\odot$ star at the RGB phase. It should be noted from the right panel of Fig.7 that the superadiabatic convection region does not extend so much inward as in the case of the $5M_\odot$ star at the AGB phase.




Fig. 7 The same as Fig.6, but for the $2M_\odot$ star at the RGB phase.

4.4 Turbulence Properties in the Overshooting Region

The turbulent correlations in the top and bottom overshooting regions are shown in Figs.8-10 for the considered stellar models. It can be found that the turbulent kinetic energy k monotonously decreases outwards and tends to be zero from the boundaries of the convective unstable zone into the overshooting regions. For the correlation $\overline{T'^2}$, there is a peak in both the top and bottom overshooting regions, its value being positive and much larger than that of Zhang & Li (2009). There is an exception in the top

overshooting region for the $2M_{\odot}$ star at the RGB phase seen in the left panel of Fig.10, the peak of $\overline{T'^2}$ disappearing completely. Nevertheless, the value of the correlation $\overline{u_r' T'}$ is always negative in both the top and bottom overshooting regions. As a result, the temperature gradient ∇ is greater than the radiative temperature gradient ∇_R but smaller than the adiabatic one ∇_{ad} .




Fig.8 Distributions of \sqrt{k} , $\overline{T'^2}$ and the temperature gradients in the top overshooting region (TOV) and the bottom overshooting region (BOV) of the $5M_{\odot}$ star at the AGB phase according to NLMA.

Fig.9 The same as Fig.8, but for the $2M_{\odot}$ star at the RGB phase.

On the other hand, the overshooting extents are different for the different correlations in a certain overshooting region. The profile of \sqrt{k} approximately demonstrates the distance that a convective cell can reach beyond the boundaries of the convective unstable zone. Its overshooting distance and decaying way will be very important to the chemical mixing in the overshooting region, if it is used to construct the diffusion process of chemical mixing (e.g. Freytag et al. 1996, Herwig et al. 1997, Salasnich et al. 1999, Ventura & D'Antona 2005). For the three stellar models, our numerical results show that the e-folding length of \sqrt{k} in the top overshooting region decreases as the stellar model is located upper and upper along the Hayashi line (they are 0.37, 0.34 and 0.34, respectively, in the unit of the local pressure scaleheight H_P), and the case in the bottom overshooting region behaves in the same way (they are 0.24, 0.23, 0.21 H_P , respectively), but the difference among them is not obvious in both the top and bottom overshooting regions. However, for the case of $\overline{u_r' T'}$ that can be inferred by the length of the region satisfying $\nabla_{ad} > \nabla > \nabla_R$, all the overshooting distances in the bottom overshooting regions for the three models are about 0.15 H_P . It means that the bottom convective overshooting has a similar effect on the stellar structure for all of the stellar models. In the top overshooting regions, however, it can be

seen that ∇ and ∇_R nearly completely overlap with each other, except for a tiny difference for the $2M_\odot$ star at the RGB phase. This may be due to very low density in the top overshooting regions, resulting in the convective heat flux comparatively rather low.




Fig. 10 The same as Fig.8, but for the $2M_\odot$ star at the RGB phase.

5 DEPENDENCE OF THE STRUCTURE OF THE OVERSHOOTING REGIONS ON TCM'S PARAMETERS

The dependence of convection characteristics of the sun on the TCM's parameters has been investigated extensively (Li & Yang 2007; Zhang & Li 2009). The turbulent correlations are mainly affected by the corresponding dissipation and diffusion parameters. The ratio of the radial kinetic energy to the horizontal one $\overline{u'_r u'_r} / \overline{u'_h u'_h}$ are shown in Fig.11 for the AGB model of $5M_\odot$, focusing on the effect of value of C_s on the convective structure in the overshooting regions, because C_s is the most sensitive parameter to determine the overshooting distance (Zhang & Li 2009). It can be seen that with the convective cells penetrating more and more into the top overshooting region the convection becomes more and more horizontally dominated. Particularly, the turbulence is isotropic when $\overline{u'_r u'_r} / \overline{u'_h u'_h} = 0.5$. It can be seen that the larger the value of C_s is, the longer the isotropic turbulence will be developed in the top overshooting. With respect to the bottom overshooting region, the ratio $\overline{u'_r u'_r} / \overline{u'_h u'_h}$ shows a much more complicated behavior for smaller values of C_s . When C_s is large enough, this complication disappears and the convection becomes more and more horizontally dominated as the convection penetrates inward into the overshooting region.

The decaying law of the turbulent kinetic energy k is a crucial factor determining the efficiency of convective mixing in the overshooting regions. The decay way of the turbulent kinetic energy k in the




Fig. 11 Ratio of the radial turbulent kinetic energy to the horizontal one $\overline{u'_r u'_r} / \overline{u'_h u'_h}$ with different TCM's parameters for the $5M_\odot$ star at the AGB phase.




Fig. 12 The decay way of the turbulent energy k in the top overshooting zone (TOV) and the bottom overshooting zone (BOV) of the $5M_\odot$ star at the AGB phase according to some non-local models.




Fig. 13 The same as Fig.12, but for the $5M_\odot$ star at the RGB phase.

overshooting regions are given in Figs.12-14 for the three stellar models based on different choices of the TCM's parameters. There is a nearly linear relation between $\lg k$ and $\ln P$ before a turning point on the curve of $\lg k$, beyond which k begins to drop promptly to zero. For example, for NLMa the relation




Fig. 14 The same as Fig.12, but for the $2M_{\odot}$ star at the RGB phase.

can be approximated as $\lg k = 2.72 \ln P + C$ in the top overshooting regions, where $C = -16.8, -12.4$ and -10.2 , respectively, for the three stellar models up along the Hayashi line. On the other hand, in the bottom overshooting regions we can obtain similar relations, for instance, for NLMc3 the result is $\lg k = -4.98 \ln P + C$, where $C = 161, 133.8$ and 140 for the three stellar models. It should be noted that the slope of the decaying law seems only a function of the TCM's parameters, and the stellar parameters only affect its constant.

The extension of these linearly decaying region and the slope of the decaying law are sensitive to the TCM's parameters. It can be found that when C_s increases while the other parameters are fixed (e.g. NLMc1, NLMc3, NLMc5), the turbulent kinetic energy decays slower and slower, and the obtained linearly decaying distance will become longer. For example, for the model of $5M_{\odot}$ star at the AGB phase in Fig.12, the length of linearly decaying distance in the bottom overshooting region for NLMc1, NLMc3, and NLMc5 are respectively about $1.2H_P$, $1.4H_P$ and $1.7H_P$. On the other hand, when the value of parameter α increases, the slope of the decaying law becomes less and the linearly decaying distance is significantly prolonged as seen by comparing the result of NLMd and that of NLMa. From Figs.12-14 it can be found that the decaying ways of the k are very similar for stars of different masses and evolutionary stages with same values of the TCM's parameters.

6 CONCLUDING REMARKS

Based on the TCM of Li & Yang (2007) we have obtained the characteristics of the turbulent convection in the RGB and AGB stars with huge convection zones in their envelopes. Some sets of suitable values of the TCM's parameters that have been used in the solar convection zone are adopted to make an exploratory application in two stars of 2 and $5M_{\odot}$ at the RGB and AGB phases. In practical calculations the TCM has been applied to the whole convective envelope including the overshooting regions. When analyzing the characteristics of the turbulent correlations, we separate the whole convective envelope into the convective unstable zone and the convective overshooting region. In the convective unstable zone, we find an approximation approach to associate the turbulent correlations explicitly, and can be used to explain the turbulent properties. Later, we make a comparison between the results of the TCM and those of the MLT on the velocity profile of turbulence and in influence on stellar structure and evolution. These obtained features of convective motion by the TCM are highly sensitive to the TCM's parameters. The main conclusions are summarized as follows:

- 1) The non-local effect of the turbulent convection is only exhibited significantly around the boundaries of the convective unstable zone, especially around the upper boundary, which is in agreement with the results of Li & Yang (2007). The three correlations \sqrt{k} , $\overline{u_r' T'}$, and $\overline{T'^2}$ are nearly peaked at the same place in the convective unstable zone for a same set of the TCM's parameters. It is because that all of them are directly correlated to a function of $(\nabla - \nabla_{ad})\overline{T}$.

- 2) For a fixed heat flux (closely related by $\overline{u_r T'}$) transferred by the convective motion, compared to the results of the MLT, the TCM will result in a larger turbulent velocity and thus more violent convective motion. Furthermore, for a nearly same profile of the convection velocity obtained by adjusting the value of the mixing length parameter α for both the TCM and MLT, the superadiabatic convection zone will be extended much more inward for the TCM than for the case of the MLT and the stellar models based on the TCM will be of lower effective temperature.
- 3) For the stellar models located up along the Hayashi line, we find that the convective motions become stronger and stronger. However, the e-folding lengths of \sqrt{k} in both the top and bottom overshooting regions decrease as the stellar model is located up along the Hayashi line. The overshooting length of $\overline{u_r T'}$ is found to be shorter than other turbulent quantities for a same set of the TCM's parameters in both overshooting regions, which is consistent with the results of Deng & Xiong (2008) and Zhang & Li (2009). And the overshooting distances of $\overline{u_r T'}$ in the bottom overshooting region are almost the same for different stellar models with a same set of the TCM's parameters. Therefore, the bottom convective overshooting has a similar effect on the stellar structure for all of the considered stellar models.
- 4) The diffusive parameter C_s plays an important role in the development of the isotropic turbulence in the top overshooting region, and the longer the isotropic turbulence will be obtained for larger C_s . In the bottom overshooting region, the convection becomes more and more horizontally dominated for larger value of C_s , and the convection shows a much more complicated behavior when C_s is small enough.
- 5) The decaying ways of the turbulent kinetic energy k are very similar for the different stellar models based on a same set of TCM's parameters. We find that there is a nearly linear relation between $\lg k$ and $\ln P$ in most of the overshooting regions. The slope of the decaying law seems only a function of the TCM's parameters. The values of C_s and α have a similar effect on the decaying distance of the turbulent kinetic energy, and the larger the values of them the slope of the decaying law becomes smaller and therefore the linearly decaying distance becomes longer.

Acknowledgements This work is in part supported by the National Natural Science Foundation of China (Grant No. 10973035 and No. 10673030) and by the Knowledge Innovation Key Program of the Chinese Academy of Sciences under Grant No. KJ CX2-YW-T24. Fruitful discussions with Q.-S. Zhang, C.-Y. Ding and J. Su are highly appreciated.

References

- Alexander, D. R., & Ferguson, J. W. 1994, *ApJ*, 437, 879
Bahcall, J. N., Pinsonneault, M. H., & Wasserburg, G. J. 1995, *Rev. Mod. Phys.*, 67, 781
Baker, N. H., & Kuhfuss, R. 1987, *A&A*, 185, 117
Böhm-Vitense, E. 1953, *Z. Astrophys.*, 32, 135
Böhm-Vitense, E. 1958, *Z. Astrophys.*, 46, 108
Bressan, A. G., Bertelli, G., & Chiosi, C. 1981, *A&A*, 102, 25
Busso, M., Wasserburg, G. J., Nollett, K. M., & Calandra, A. 2007, *ApJ*, 671, 802
Canuto, V. M., Goldman, I., & Mazzitelli, I. 1996, *ApJ*, 473, 550
Canuto, V. M. 1992, *ApJ*, 392, 218
Canuto, V. M. 1993, *ApJ*, 416, 331
Canuto, V. M. 1994, *ApJ*, 428, 729
Canuto, V. M. 1997, *ApJ*, 482, 827
Canuto, V. M., & Dubovikov, M. S. 1998, *ApJ*, 493, 834
Canuto, V. M., & Mazzitelli, I. 1991, *ApJ*, 370, 295
Castellani, V., Chieffi, A., Pulone, L., & Tornamè, A. 1985, *ApJ*, 296, 204
Castellani, V., Giannone, P., & Renzini, A. 1971, *Ap&SS*, 10, 340
Charbonnel, C. 1995, *ApJ*, 453, L41
Deng, L., & Xiong, D.-R. 2008, *MNRAS*, 386, 1979

- Deng, L., Xiong, D.-R., & Chan, K. L. 2006, *ApJ*, 643, 426
- Deng, L. C., & Xiong, D.-R. 2001, *ChJAA* (*Chin. J. Astron. Astrophys.*), 1, 50
- Freytag, B., Ludwig, H. G., & Steffen, M. 1996, *A&A*, 313, 497
- Gratton, R. G., Sneden, C., Carretta, E., & Bragaglia, A. 2000, *A&A*, 354, 169
- Gratton, R., Sneden, C., & Carretta, E. 2004, *ARA&A*, 42, 385
- Herwig, F. 1997, *A&A*, 324, L81
- Herwig, F. 2005, *ARA&A*, 43, 435
- Herwig, F., Blöcker, T., Schönberner, D., & El Eid, M. 1997, *A&A*, 324, L81
- Hollowell, D., & Iben, I. Jr. 1988, *ApJ*, 333, L25
- Hossain, M. S., & Rodi, W., 1982, *Turbulent Buoyant Jets and Plumes*, ed. W. Rodi (Oxford: Pergamon Press), 121
- Iben, I. Jr. 1975, *ApJ*, 196, 525
- Iben, I. Jr. 1977, *ApJ*, 217, 788
- Iben, I. Jr. 1981, *ApJ*, 246, 278
- Iglesias, C. A., & Rogers, F. J. 1996, *ApJ*, 464, 943
- Jiang, S.-Y., & Huang, R.-Q. 1997, *A&A*, 317, 121
- Kraft, R. P., Sneden, C., Langer, G. E., & Shetrone, M. D. 1993, *AJ*, 106, 1490
- Lattanzio, J.C. 1989, *ApJ*, 344, L25
- Li, Y., & Yang, J.-Y. 2001, *ChJAA* (*Chin. J. Astron. Astrophys.*), 1, 66
- Li, Y., & Yang, J.-Y. 2007, *MNRAS*, 375, 388
- Maeder, A. 1975, *A&A*, 40, 303
- Pederson, B. B., Vandenberg, D. A., & Irwin, A. W. 1990, *ApJ*, 352, 279
- Pilachowski, C.A., Sneden C., & Booth, J. 1993, *ApJ*, 407, 699
- Ramírez, S. V., & Cohen, J. G. 2002, *AJ*, 123, 3277
- Recio-Blanco, A., & de Laverny, P. 2007, *A&A*, 461, L13
- Renzini, A. 1987, *A&A*, 188, 49
- Rogers, F. J., 1994, in *IAU Colloquium 147, The Equation of State in Astrophysics*, eds. G. Chabrier, & E. Schatzman (Cambridge: Cambridge Univ. Press), 16
- Rogers, F. J., & Iglesias, C. A. 1995, in *ASP Conf. Ser. 78, Astrophysical Applications of Powerful New Databases*, eds. S. J. Adelman, & W. L. Wiese (San Francisco: ASP), 78
- Rogers, F. J., Swenson, F. J., & Iglesias, C. A. 1996, *ApJ*, 456, 902
- Sackmann, I.J. 1980, *ApJ*, 241, L37
- Salasnich, B., Bressan, A., & Chiosi, C. 1999, *A&A*, 342, 131
- Saslaw, W. C., & Schwarzschild, M. 1965, *ApJ*, 142, 1468
- Spruit, H. C., Nordlund, A., & Title, A. M. 1990, *ARA&A*, 28, 263
- Sreenivasan, S. R., & Wilson, W. J. F. 1978, *Ap&SS*, 53, 193
- Straniero, O., Chieffi, A., Limongi, M., et al. 1997, *ApJ*, 478, 332
- Ulrich, R. K. 1970, *Ap&SS*, 7, 183
- Ventura, P., & D'Antona, F. 2005, *A&A*, 431, 279
- Xiong, D.-R. 1985, *A&A*, 150, 133
- Xiong, D.-R. 1990, *A&A*, 232, 31
- Xiong, D.-R. 1979, *Acta Astron. Sin.*, 20, 238
- Xiong, D.-R., Cheng, Q.-L., & Deng, L. 1997, *ApJS*, 108, 529
- Xiong, D.-R. 1981, *Sci. Sinica.*, 24, 1406
- Yang, J.-Y., & Li, Y. 2007, *MNRAS*, 375, 403
- Zhang, Q.-S., & Li, Y. 2009, *RAA*, 9, 585