

to be submitted to *The Astronomical Journal*,
Version: 9/08/2011

Know the Star, Know the Planet. II. Speckle Interferometry of Exoplanet Host Stars

Brian D. Mason¹, William I. Hartkopf¹

*United States Naval Observatory
3450 Massachusetts Ave., NW, Washington, DC 20392-5420
Electronic mail: (bdm,wih)@usno.navy.mil*

Deepak Raghavan¹

*Georgia State University, Dept. of Physics & Astronomy, Atlanta, GA 30303-3083
Electronic mail: raghavan@chara.gsu.edu*

John P. Subasavage¹

*Cerro Tololo Inter-American Observatory, La Serena, Chile
Electronic mail: jsubasavage@ctio.noao.edu*

Lewis C. Roberts, Jr.

*Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive,
Pasadena CA 91109
Electronic mail: lewis.c.roberts@jpl.nasa.gov*

Nils H. Turner and Theo A. ten Brummelaar

*Center for High Angular Resolution Astronomy, Georgia State University, Mt. Wilson, CA
91023
Electronic mail: (nils,theo)@chara-array.org*

ABSTRACT

¹Visiting Astronomer, Kitt Peak National Observatory and Cerro Tololo Inter-American Observatory, National Optical Astronomy Observatories, operated by Association of Universities for Research in Astronomy, Inc. under contract to the National Science Foundation.

A study of the host stars to exoplanets is important to understanding their environment. To that end, we report new speckle observations of a sample of exoplanet host primaries. The bright exoplanet host HD 8673 (= HIP 6702) is revealed to have a companion, although at this time we cannot definitively establish the companion as physical or optical. The observing lists for planet searches and for these observations have for the most part been pre-screened for known duplicity, so the detected binary fraction is lower than what would otherwise be expected. Therefore, a large number of double stars were observed contemporaneously for verification and quality control purposes, to ensure the lack of detection of companions for exoplanet hosts was valid. In these additional observations, ten pairs are resolved for the first time and sixty pairs are confirmed. These observations were obtained with the USNO speckle camera on the NOAO 4m telescopes at both KPNO and CTIO from 2001 to 2010.

Subject headings: binaries: general — binaries: visual — binaries: orbits — techniques: interferometry — stars:individual (HD 8673)

1. Introduction

As discussed in Paper 1 of this series (Roberts et al. 2011) a study of the host stars to exoplanets is essential if we wish to understand the environment in which those planets formed. Further, the star’s luminosity, distance, mass, and other characteristics are fundamentally related to the determination of the planets mass and size. Determining these parameters directly for the host star as opposed to using a template of the canonical stellar class and type will produce more accurate and precise planetary determinations. As part of this effort we herein report new speckle observations of a large sample of exoplanet host primaries.

Binaries affect the formation and stability of planetary systems, as their long-term relationship must be hierarchical. Generally speaking, based on the precepts of Harrington (1981) if the ratio of semimajor axes is 4:1 or greater, an exoplanet in a stellar binary is dynamically stable. Dynamically permitted systems include the more commonly detected configuration of planet(s) orbiting one stellar component of a sufficiently wide-orbit binary in a hierarchical arrangement, and the harder-to-detect circumbinary configuration of planet(s) in a wide orbit around a close stellar binary (see Raghavan et al. 2006, especially §6.1). That said, Raghavan et al. (2010) in their statistics updating and improving upon Duquennoy & Mayor (1991), find that while the frequency of single stars is the same, the number of companions has increased through instrumental and technique enhancements. Due to the

presence of stellar companions, one might imagine the environment of binary stars to be a rich one for substellar companions. However, dynamical effects should eject companions not found in hierarchical orbits. In any event, as conducive as this environment might be to companions, this is not reflected in the list of known planet hosts, however. This is due entirely to selection effects; because of the complexities of disentangling stellar companions from small planetary signatures, the observing lists for planet searches have for the most part been pre-screened for known duplicity, so the detected binary fraction is lower than what would be expected.

In addition to binary stars that are gravitationally bound, there are optical doubles which are merely chance alignments of unrelated stars. Although they do not contribute dynamically to the system, close optical pairs do contribute light to the system, which should be accounted for in system analysis. While photometric analysis of binary systems can infer “third” light in the system, radial velocities or periodic variation in other astrometric parameters (for example, the Hipparcos acceleration solutions), would not give evidence of these companions. Optical pairs would best be found by direct imaging or interferometric analysis.

2. Speckle Observations

All of these observations were obtained as part of other observing projects, for example, analysis of white, red and subdwarfs (Jao et al. 2009), G dwarfs (§5.3.6 of Raghavan et al. 2010), or Massive stars (Mason et al. 2009), some of which are still in developmental and/or data collection stages. Unpublished observations of exoplanet host stars were extracted from these data and are presented here. The instrument used for these speckle observations was the USNO speckle interferometer, described most recently in Mason et al. (2009).

Speckle Interferometry is a single filled-aperture interferometric technique where the “speckles” of a pair of nearby stars, induced by atmospheric turbulence, constructively interfere. Reduced by simple autocorrelation methods, in the resulting image the binary or double star geometry is the predominant structure when compared with the other uncorrelated pairings. It is capable of resolving pairs to the resolution limit of the telescope in question up to size of the observation field (typically, $\sim 1''.5$), as long as the pairs have magnitude differences of less than about three.

Calibration of the KPNO data was accomplished through the use of a double-slit mask placed over the “stove pipe” of the 4-m telescope during observations of a bright known single star (as described in Hartkopf et al. 2000). This application of the Young’s double-slit

experiment allowed determination of scale and position angle zero points without relying on binaries themselves to determine calibration parameters. Multiple observations through the slit mask (during five separate KPNO runs from 2001 to 2008) yielded mean errors of $0''.11$ in the position angle zero point and 0.165% in the scale error. These “internal errors” are undoubtedly underestimates of the true errors for these observations, because these calibration tests were made on stars that were brighter and nearer the zenith than science stars. Total errors are likely $3\text{--}5\times$ larger than these internal errors.

Because the slit-mask option is not available on the CTIO 4-m telescope, we calibrated the Southern Hemisphere data using observations of numerous well-observed wide equatorial binaries obtained at both the KPNO and CTIO telescopes. Published orbital elements for these pairs were updated as needed, using the recent KPNO and published measures, then predicted ρ and θ values from those orbits deemed of sufficiently high quality were used to determine the CTIO scale and position angle zero points. The calibration errors for these southern observations were (not surprisingly) considerably higher than those achieved using the slit mask. Mean errors for five CTIO runs from 2001 to 2010 (as well as a 2001 KPNO run lacking slit mask data) were $0''.67$ in position angle and 1.44% in scale. These errors are likely overestimated, because we have assumed that the calibration binaries have perfect orbits, and any offsets are incorporated into the errors.

3. Results

Following generation of the Directed Vector Autocorrelation (Bagnuolo et al. 1992), the “DVA” is background subtracted through boxcar subtraction and the sharp central peak which corresponding to the zeroth order speckles correlating with themselves is clipped. Companions in the resulting DVA are then readily apparent as peaks several sigma above the background.

Of the 118 exoplanet hosts we observed only one, HIP 6702 showed signs of a companion and is discussed in Section 3.1 and listed in Table 1. The null results are listed in Table 2, a list of single star detections. In the table, Column (1) gives the Hipparcos number, Column (2) the HD Catalog number, Column (3) lists other common designations, Column (4) is the epoch of observation, and Column (5) identifies the telescope (C = Cerro Tololo 4m, K = Kitt Peak 4m). For all of these observations no companion was detected within the ranges $\Delta m_V < 3$, and $0''.03 < \rho < 1''.5$.

Table 1 lists the observations for this new detection. Column (1) gives the precise position of the system, Column (2) is the Washington Double Star Catalog (hereafter, WDS;

Mason et al. 2001) identifier, and Column (3) lists the discovery designation, here the WSI (= Washington Speckle Interferometry) number. Column (4) gives the Hipparcos number of the primary as a cross-reference. Column (5) gives the epoch of observation, and Columns (6) and (7) provide the relative astrometry. Column (8) lists a crude estimate of the magnitude difference of the pair in the V band (the listed number is paired with the more reliable observation). This estimate is probably only good to $\pm 0.5 \text{ mag}$. Column (9) provides the separation in astronomical units, based on the Hipparcos parallax and this angular separation assuming a face-on orbit.

The resulting multiplicity fraction is extremely low, but artificially so. Observation of known binaries is a prime goal of the USNO speckle program and some of these pairs had been previously published (e.g., HD 28305 in Mason et al. 2009). Others which were known but whose motions were not especially compelling (e.g., HD 50583 in Mason et al. 2011) were observed with our 26in refractor in Washington and those which do have a compelling individual story to tell unrelated to exoplanets are in preparation (Farrington et al. 2012). A simplistic multiplicity determination of this limited result ($= \frac{1}{118}$) is therefore not a meaningful number.

3.1. New Double Star : HIP 6702

Of all the exoplanet hosts which have been serendipitously observed, all were unknown as close visual doubles and only one of the host stars, HIP 6702 (= HD 8673) appeared double in directed vector autocorrelations on both times it was observed. The classification of HIP 6702 as an exoplanet is based upon Hartmann et al. (2010) who, using iodine-cell radial velocity measurements, detected a companion with a $M \sin i$ of $14.2 M_j$ with a period of 1634 ± 17 days and an eccentricity of 0.723 ± 0.016 . The relative astrometric measures of this resolved pair are provided in Table 1. Given the small number of measures presented in Table 2, the pair, while a visual double star, is not necessarily a binary system. Verification of physicality for the new companion to HIP 6702 can be accomplished several ways, among them color-magnitude, proper motion and/or kinematic analysis. The speckle interferometry observable of relative position establishing kinematic-physical (i.e., Keplerian) motion requires at least three measures. So, while close proximity can be a powerful argument for physicality, it is by no means definitive (c.f., ι Ori, §5.1 of Mason et al. 2009). Nevertheless, even a companion which is only nearby in the angular sense should be considered in any detailed analysis of the star, as it will contribute to the photometric signature of the examined target. Such is the case for HIP 6702, which was recently reported as a sub-stellar companion (Hartmann et al. 2010).

Among the possible interpretations of the new speckle companion two stand out: first, the companion is a not-physically associated line-of-sight companion and second, it is the companion detected in Hartmann et al. (2010).

3.1.1. *Physical Companion?*

Hipparcos produced many types of double star solutions. The one which can be most easily compared to other detection techniques and the most common are those where the relative parameters (ρ , θ) are presented. The speckle interferometry measures presented in Table 1 are both near the Hipparcos “C” code double star solution cutoff ($0''.082$ for HDS 446 = HIP 27151). The other Hipparcos double star solutions may not be applicable here. Some depend upon *a priori* orbital information (O code), system dynamics in the plane of the sky (G code), variability (V code) or unknown parameters (X and/or S code). In any event, the lack of Hipparcos detection is a condition which is neither necessary nor sufficient to establish that the Hartmann et al. (2010) companion is not-stellar.

However, if the two Table 1 measures represent relative measures of the Hartmann et al. (2010) pair, the inclination must be extremely low. Assuming a near zero inclination the mean separation of $0''.098$ would approximate the angular semi-major axis ($a'' = 0''.098 \pm 0''.011$). Given this, the Hipparcos parallax of 26.14 ± 0.79 mas and the Hartmann et al. period of 1634 ± 17 days, a mass sum of $2.63 \pm 0.92 M_{\odot}$ is obtained, which is not unreasonable for two similar F dwarf stars, although the error is quite large, primarily due to the uncertainty in a'' . The length of time between the two speckle observations represents $1.47 \times$ the Hartmann et al. period. The two measures of angular position represent $(0.497 \text{ or } 0.503) + n$ revolutions of the system (depending on direction of rotation) which is very similar to the Hartmann et al. period when $n = 1$.

Given the estimated dynamic range ($\Delta m_V = 2.3 \pm 0.5$) and assuming the fainter limit and spectral type of the primary this would make the secondary close to a K2V. Using the canonical mass of a K2V in $M \sin i = 14.2 M_j$ gives an inclination of $1^{\circ}02$. Using this inclination with $a \sin i$ from Hartmann et al. (2010) gives a semi-major axis of $0''.168$ which is consistent with the Table 1 results.

3.1.2. *or Optical Companion?*

Since the interferometric companion to HIP 6702 has been observed so few times, establishing the companion as optical or physical is not possible. The proper motion of the

primary is $0''.25/\text{yr}$ ($\alpha=0''.236/\text{yr}$, $\delta=-0''.085/\text{yr}$). From the relative positions in Table 1, the proper motion of the companion would be an even higher at $0''.276/\text{yr}$. If linear motion is assumed and reasonable errors are applied it is possible to determine where the companion would be at some date in the future. In Figure 1 this determination is performed assuming errors slightly larger than nominal for the two speckle interferometry measures: $\Delta\theta = 1''.0$, $\Delta\rho/\rho = 1.0\%$. The predicted position for 2012 through 2015 are plotted as error boxes. Again, assuming linear motion from the two speckle points, a separation of $0''.37$ and a position angle of 255° is determined for 1991.25, which would be well within the capabilities of the Hipparcos satellite (ESA 1997).




Fig. 1.— A “motion characterization” system plot for HIP 6702 (= WSI 96) with small filled circles indicating the 2001 and 2007 speckle measures from Table 1. Scales are in arcseconds, and in each figure the large shaded circle represents the V band resolution limit of a 4m telescope. The four small error boxes in each figure indicate the predicted location of that pair’s secondary in 2012.0, 2013.0, 2014.0, and 2015.0, assuming the motion is linear and all speckle measures are characterized by errors of $\Delta\theta = 1''.0$, $\Delta\rho/\rho = 1.0\%$. Finding the double within a box appropriate to the observation date would be a strong indication that the relative motion of the pair is linear (that is, just motion from an unrelated field star due to proper motion differences). The **H** indicates where the companion would have been at 1991.25, at the Hipparcos epoch.

All Sky Speckle list




Fig. 2.— Aitoff projection of all 444 targets. Filled circles ($N = 114$) are those listed in Table 2. Open circles ($N = 27$) are those observed by CHARA or USNO with an ICCD and reduced with the DVA method. Asterisks ($N = 11$) are those observed by other interferometry groups, and an “X” ($N = 292$) are those which have yet to be observed. A valid speckle measure is only counted if it was obtained on a 4m class telescope.

4. Future Observing

Due to the relatively even distribution of targets not yet observed by speckle interferometry, one observing run in each semester and each hemisphere will be necessary in order to observe all remaining exoplanet host stars. However, the target list for each of the four runs will be slight, less than one hundred stars each. With an approximately equal number of quality control and equatorial scale calibration pairs, each observing run could easily be completed in 2-3 nights. Priority would obviously be given to targets not observed before. Those observed by other speckle interferometric groups would be next priority so they all have a common reduction algorithm. Figure 2 is an Aitoff plot of targets from the list of known exoplanet host stars taken from the NStED¹ database and gives their observation status.

The USNO speckle interferometry program has been supported by NASA and the SIM preparatory science program through NRA 98-OSS-007, SIM Key Project MASSIF as well as

¹<http://nsted.ipac.caltech.edu/cgi-bin/bgServices/nph-bgExec>, extracted 12 April 2011

No. NNH06AF701 issued through the Terrestrial Planet Finder Foundation Science program. Thanks are also provided to the U.S. Naval Observatory for their continued support of the Double Star Program. A portion of the research in this paper was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (NASA). This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. Grateful acknowledgment is given to USNO interns Dean Kang, Laura Flagg and Ankit Patel for their processing of the speckle pixel data. The telescope operators and observing support personnel of KPNO and CTIO continue to provide exceptional support for visiting astronomers. Thanks to Alberto Alvarez, Skip Andree, Bill Binkert, Gale Brehmer, Bill Gillespie, Angel Guerra, Jim Hutchinson, Hillary Mathis, Oscar Saa, Patricio Ugarte, and the rest of the KPNO and CTIO staff. We also like to thank Richard Green of KPNO, who was able to provide us with two extra nights on the 4-m Mayall telescope during our January 2001 run. While we were hampered by poor weather, this additional allocation of time certainly helped us achieve a greatly enhanced completion fraction. A special thank you is also given to Hal Halbedel, who operated the telescope on all or part of each of these new KPNO runs and was instrumental in the slit-mask work done at KPNO.

Table 1. New Interferometric Double

Coordinates α, δ (2000)	WDS or α, δ (2000)	Discoverer Designation	HIP	BY 2000.0+	θ ($^{\circ}$)	ρ ($''$)	Δm (mag)	sep (AU)
01 26 08.78 +34 34 46.9	01262+3435	WSI	96	6702	1.0193	257.3	0.087	3.3
					7.6049	78.3	0.109	4.2

Table 2. Stars with No Companion Detected

HIP #	HD #	Common Name	BY 2000.0+	Telescope
522	142	...	1.5621	C
1292	1237	GJ 3021	1.5674	C
1499	1461	...	1.5701	C
3391	4113	...	1.5647	C
3479	4208	...	1.5647	C
3497	4308	...	1.5648	C
5054	6434	...	1.5622	C
5529	7199	...	1.5675	C
5806	7449	...	1.5675	C
6379	7924	...	1.0218	K
7513	9826	<i>v</i> And	1.0220	K
7599	10180	...	1.5702	C
7978	10647	...	1.5702	C
8159	10697	...	1.0300	K
9683	12661	...	1.0165	K
10138	13445	GJ 86	1.5676	C
10626	13931	...	1.0219	K
12186	16417	...	1.5676	C
12189	16246	30 Ari	1.0304	K
12653	17051	<i>ι</i> Hor	1.5676	C
14954	19994	...	1.0820	C
15323	20367	...	1.0193	K
15527	20782	...	1.0820	C
16537	22049	<i>ε</i> Eri	10.0652	C
17096	23079	...	1.0738	C
20723	28185	...	1.0764	C
24681	34445	...	1.0306	K
25110	33564	...	1.0222	K
26381	37124	...	1.0307	K

Table 2—Continued

HIP #	HD #	Common Name	BY 2000.0+	Telescope
26394	39091	...	1.0740	C
26394	39091	...	6.1937	C
26664	37605	...	1.0306	K
27887	40307	...	1.0740	C
27887	40307	...	6.1937	C
28767	40979	...	1.0224	K
29804	43848	...	1.0793	C
30034	44627	AB Pic	1.0740	C
30579	45364	...	1.0794	C
30860	45350	...	1.0198	K
30905	45652	...	1.0308	K
31246	46375	...	1.0767	C
31540	47186	...	1.0794	C
32916	49674	...	1.0224	K
32970	50499	...	1.0794	C
33212	50554	...	1.0279	K
33719	52265	...	1.0823	C
36795	60532	...	1.0795	C
37826	62509	...	1.0199	K
38041	63765	...	1.0742	C
38558	65216	...	1.0742	C
40693	69830	...	6.1911	C
40952	70642	...	1.0743	C
43587	75732	55 Cnc	1.0200	K
46076	81040	...	1.0770	C
47007	82943	...	1.0797	C
47202	83443	...	1.0744	C
48235	85390	...	1.0744	C
48739	86226	...	1.0797	C

Table 2—Continued

HIP #	HD #	Common Name	BY 2000.0+	Telescope
49699	87883	...	1.0202	K
50473	89307	...	1.0310	K
50921	90156	...	6.1912	C
52521	93083	...	1.0799	C
53721	95128	47 UMa	1.0202	K
54906	97658	...	1.0203	K
57172	101930	...	1.0746	C
57291	102117	...	1.0746	C
57370	102195	...	1.0311	K
57443	102365	...	6.1915	C
57443	102365	...	10.0659	C
57931	103197	...	1.0746	C
58451	104067	...	6.1915	C
59610	106252	...	1.0312	K
64295	114386	...	1.0827	C
64426	114762	...	1.0232	K
64457	114783	...	5.1915	C
64459	114729	...	1.0775	C
64459	114729	...	1.0802	C
64924	115617	61 Vir	6.1890	C
64924	115617	61 Vir	8.4500	K
64924	115617	61 Vir	10.0688	C
65721	117176	70 Vir	1.0232	K
65721	117176	70 Vir	6.1916	C
67275	120136	...	1.0314	K
67275	120136	τ Boo	6.1916	C
71395	128311	...	1.5664	C
71395	128311	...	6.1916	C
72339	130322	...	1.0829	C

Table 2—Continued

HIP #	HD #	Common Name	BY 2000.0+	Telescope
74500	134987	...	1.0830	C
74500	134987	...	1.5611	C
77740	141937	...	1.5666	C
78459	143761	ρ CrB	8.4503	K
79242	142022A	...	1.5667	C
79248	145675	14 Her	1.4986	K
80250	147018	...	1.5667	C
80337	147513	...	6.1919	C
83389	154345	...	1.4960	K
83949	155358	...	1.4961	K
86796	160691	μ Ara	1.5667	C
87330	162020	...	1.5642	C
88348	164922	...	8.4506	K
90004	168746	...	1.5614	C
90485	169830	...	1.5614	C
91085	171238	...	1.5614	C
94075	178911B	...	1.4990	K
94645	179949	...	1.5614	C
96901	186427	16 Cyg	1.4991	K
97336	187123	...	1.4990	K
97546	187085	...	1.5670	C
98505	189733	...	8.4614	K
98767	190360	...	8.4563	K
99711	192263	...	5.8680	K
99825	192310	GJ 785	1.5616	C
101806	196050	...	1.5615	C
101966	196885	...	8.4481	K
104903	202206	...	1.5618	C
106006	204313	...	1.5618	C

REFERENCES

- Bagnuolo, W.G., Jr., Mason, B.D., Barry, D.J., Hartkopf, W.I., & McAlister, H.A. 1992, AJ 103, 1399
- Duquennoy, A. & Mayor, M. 1991, A&A 248, 485
- ESA 1997, The Hipparcos and Tycho Catalogues, ESA SP-1200
- Farrington, C.D. et al. 2012 (*in preparation*)
- Harrington, R.S. 1981, Planetary orbits in multiple star systems, in *Life in the Universe*, Cambridge, MIT Press, 119
- Hartkopf, W.I. et al. 2000, AJ 119, 3084
- Hartmann, M., Guenther, E.W. & Hatzes, A.P. 2010, ApJ 717, 348
- Jao, W.-C., Mason, B.D., Hartkopf, W.I., Henry, T.J. & Ramos, S.N. 2009, AJ 137, 3800
- Mason, B.D., Hartkopf, W.I., Gies, D.R., Henry, T.J., & Helsel, J.W. 2009, AJ 137, 3358
- Mason, B.D., Hartkopf, W.I. & Wycoff, G.L. 2011, AJ (*in press*)
- Mason, B.D., Wycoff, G.L., Hartkopf, W.I., Douglass, G.G. & Worley, C.E. 2001, AJ 122, 3466²
- Raghavan, D., Henry, T.J., Mason, B.D., Subasavage, J.P., Jao, W.-C., Beaulieu, T.D. & Hambly, N.C. 2006, ApJ 646, 523
- Raghavan, D. et al. 2010, ApJS 190, 1
- Roberts, L.C., Jr., Turner, N.H., ten Brummelaar, T.A., Mason, B.D. & Hartkopf, W.I. 2011, (*in progress*)
- Tokovinin, A., Mason, B.D. & Hartkopf, W.I. 2010, AJ 139, 743

²*The Washington Double Star Catalog*, published in *Second USNO Double Star CD 2006.5* and available online at <http://ad.usno.navy.mil/wds/wds.html>.

Table 2—Continued

HIP #	HD #	Common Name	BY 2000.0+	Telescope
106353	204941	...	1.5618	C
108375	208487	...	1.5672	C
108859	209458	...	8.4615	K
109378	210277	...	1.5645	C
111143	213240	...	1.5618	C
112441	215497	...	1.5646	C
113137	216437	...	1.5647	C
113238	216770	...	1.5646	C
113357	217014	51 Peg	7.5883	K
113357	217014	51 Peg	8.4617	K
116727	222404	γ Cep	1.0218	K
116727	222404	γ Cep	1.4993	K
116906	222582	...	1.5674	C

A. Additional Measures of Known Pairs

Due to the high incidence of single stars among the exoplanet hosts, a substantial number of double stars were observed contemporaneously with the exoplanet host observations to ensure that the observing conditions and detection capabilities given above were met. Additional measures of known or suspected doubles were made as time permitted. Table A1 lists 549 mean positions for 485 known systems. Column (1) is the WDS identification (arcminute coordinate), Column (2) lists the Discovery Designation, and Column (3) provides the Hipparcos number. Column (4) gives the epoch of observation, and Columns (5) and (6) provide the relative astrometry. Column (7) contains the notes for these systems. Also found in in table are ten pairs resolved for the first time and sixty pairs which are here confirmed; estimated magnitude differences for the new pairs (when available) are listed in the Notes column.

Table A1. Other Pairs Measured

WDS or α, δ (2000)	Discovery Designation	HIP	BY 2000.0+	θ ($^{\circ}$)	ρ ($''$)	Note
00024+1047	A 1249 AB	190	7.5994	74.8	0.135	
00026-0829	A 428	210	7.6019	344.5	0.174	
00055-1835	RST 3340	—	1.5701	287.4	0.303	
00095+1907	COU 247	768	7.5992	259.1	0.294	
00117+6145	TDS 1338	—	7.6074	111.2	0.527	1
00167+3629	STT 4	1336	7.6021	99.7	0.232	
00174+0853	A 1803 AB	1392	7.6019	307.2	0.145	
00182+7257	A 803	1461	7.6021	307.9	0.262	
00233+5132	TDS 1431	—	7.6074	102.5	0.227	1
00271-0753	A 431	2143	1.5021	184.5	0.139	
			1.5647	189.0:	0.139:	
			7.6019	13.3	0.201	
00274+3054	HDS 62	2170	7.6074	72.0	0.302	1
00283+6344	HU 1007	2232	7.6021	89.2	0.449	
00290+5848	TDS 1484	—	7.6074	351.9	0.463	1
00310-1005	BU 1158 BC	2444	7.6019	310.6	0.325	
00335+4006	HO 3	2643	7.6021	168.2	0.224	
00337+4310	COU 1354	—	7.6074	64.9:	0.646:	1
00339+5316	TDS 1523	—	7.6075	90.6	0.693	1
00362+7546	TDS 1542	—	7.6075	338.9	0.721	1
00364+3330	HDS 84	2872	7.6075	28.8	0.196	1
00373-2446	BU 395	2941	10.5861	99.4	0.555	
00393+3359	COU 656	—	7.6075	150.6	0.821	1
00429+2047	A 2205	3373	7.6019	8.8	0.344	
00431+7659	HU 1012	—	7.6021	124.0	0.446	
00441+5655	TDS 1589	—	7.6075	315.6	0.788	1
00456+5457	TDS 1602	—	7.6075	70.7	0.409	1
00470+2315	HU 413	3673	7.6019	309.3	0.365	
00492+5720	A 921 AB	—	7.6021	183.2	0.168	
00516+2237	A 1808	4030	7.6019	196.7	0.188	
00520+3154	A 924	—	7.6021	318.5	0.261	
00529+6053	TDS 1655	—	7.6075	27.4	0.560	1
00533+0405	A 2307	4176	7.6019	174.6	0.075	
00533-4530	B 644	4177	10.5862	349.2	0.281	

Table A1—Continued

WDS or α, δ (2000)	Discovery Designation	HIP	BY 2000.0+	θ ($^{\circ}$)	ρ ($''$)	Note
01251+4537	A 939	6633	7.6049	183.5	0.191	
01345+3440	A 1913 AB	7337	7.6049	311.6	0.391	
01361-2954	HJ 3447	7463	10.5862	184.9	0.777	
01409+4952	HU 531 AB	7841	7.6049	358.4	0.467	
01443+5732	BU 870 AB	8115	7.6021	333.6	0.611	
01450+5707	BU 453	8166	7.6021	100.7	0.749	
01450+2703	COU 750	—	7.6021	25.8	0.273	
01462+3343	HU 804	8252	7.6021	67.7	0.285	
01463+4439	A 948 AB	—	7.6021	133.1	0.302	
01495+5645	A 950 AB	—	7.6021	278.3	0.222	
01497-1414	HU 422	8504	7.5994	29.8	0.296	
01499+8053	STT 34	8519	7.6075	284.6	0.510	
01500-0408	A 2602	—	7.5995	178.9	0.632	2
01520+1326	HU 1213	8703	7.5995	84.9	0.105	
01528-0447	RST 4188	—	7.5994	31.3	0.419	
01570+3101	A 819 AB	9087	7.6021	315.3	0.183	
01559+0151	STF 186	8998	5.8592	63.7	0.915	
02020+7054	BU 513 AB	9480	7.6022	287.4	0.705	
02130+0851	MCA 5	10324	7.5995	84.1	0.104	
02157+2503	COU 79	10535	1.0164	301.6	0.100	3
			5.8601	255.0	0.157	
			7.5995	243.6	0.152	
02159+0638	A 2013	10552	7.5995	256.0	0.172	
02174+6121	STF 234 AB	10677	7.6021	223.5	0.719	
02202+2949	A 961	10892	7.6021	278.6	0.345	
02211+4246	STF 248	10952	7.6016	217.3	0.528	
02231+7021	MLR 377 AB	11120	7.6022	140.5	0.674	
02291+6724	CHR 6 Aa,Ab	11569	7.6022	45.8	0.543	
02382+4604	A 1278	—	7.6075	308.9	0.139	
02405-2408	SEE 19	12466	10.5864	241.7	0.117	
02415+4053	COU 1511	12552	7.6075	30.0	0.109	
02442-2530	FIN 379 Aa,Ab	12780	10.5864	343.4	0.118	
02517+4559	A 1281	13336	7.6075	134.8	0.630	
03175+6540	STT 52 AB	15309	7.6022	58.1	0.485	

Table A1—Continued

WDS or α, δ (2000)	Discovery Designation	HIP	BY 2000.0+	θ ($^{\circ}$)	ρ ($''$)	Note
04123+0939	STT 74	—	10.0653	114.4	0.361	
04142-4608	RST 2338	19758	1.0791	197.0	0.135	
			6.1935	269.5	0.187	
04163-6057	GLE 1	19917	10.0677	201.7	0.319	
04215-2544	BU 744 AB	20347	10.0645	189.8	0.297	
04239+0928	HU 304	20522	10.0653	20.7	0.267	
04242-5704	RMK 4	20552	1.0739	246.5	5.426	
04269-2405	BU 311	20765	10.0645	150.3	0.439	
04275-1707	HU 440	20806	1.0822	318.9	2.064	
04286+1558	MCA 15	20885	5.8688	350.5	0.151	
04374-0951	RST 3401	21536	10.0652	260.3	0.399	
04395-4507	I 1489	21698	1.0793	80.6	0.208	
04503+0103	A 2622	—	10.0653	252.3	0.294	
04529+7543	MLR 465	22703	1.0222	208.4	0.866	
04545-0314	RST 5501	22812	10.0652	206.4	0.150	
04584-0344	HDS 644	23116	10.0652	333.0	0.588	
05059-1355	A 3009	23716	1.0767	257.2	1.134	
05079+0830	STT 98	23879	10.0653	299.9	0.883	
05089+0313	A 2636	24227	10.0653	349.8	0.363	
05135+0158	STT 517 AB	24349	10.0653	239.7	0.670	
05167+4600	ANJ 1 Aa,Ab	24608	5.8634	328.8	0.046	3
05190-2159	RST 2375	24800	10.0646	17.7	0.172	
05234-3640	I 275	25190	10.0680	220.7	0.788	
05245-0224	DA 5 AB	25281	10.0652	77.7	1.788	2
05247+6323	STF 677	25300	1.0223	127.6	1.074	
05248-5219	I 345 AB	25303	10.0680	213.1	0.253	2
			10.5863	213.6	0.258	
05255-0033	A 848	25365	10.0653	176.1	0.322	
05387-0236	BU 1032 AB	26549	10.0652	89.0	0.256	
05407-0157	STF 774 AB	26727	10.0653	165.3	2.450	
05418-5000	HU 1568	26830	10.0706	152.6	0.495	
05455+5210	WSI 98	27165	1.0223	156.4	0.086	$\Delta_m = 2.0$
05508-3945	I 1494 AB	27611	10.0680	269.4	0.384	
05525-0217	HDS 787	27758	10.0652	343.6	0.111	

Table A1—Continued

WDS or α, δ (2000)	Discovery Designation	HIP	BY 2000.0+	θ ($^{\circ}$)	ρ ($''$)	Note
06410+0954	CHR 168 Aa,Ab	31978	10.0654	258.3	0.117	
06478+0020	STT 157	32572	10.0654	168.0	0.520	
06481-0948	A 1056	—	10.0656	257.3	0.335	
06545-2734	B 706	33197	10.0656	291.6	0.725	
07013-0906	A 671	—	10.0654	13.4	0.315	
07029-1313	HDS 980	33973	10.0656	220.4	0.488	
07175-4659	I 7	35296	1.0741	208.3	0.822	
			6.1882	205.8	0.775	
			10.0656	204.5	0.745	
07185-5721	HDS 1013 Aa,Ab	35374	10.0656	215.6	0.340	
07305+0743	A 2869	36487	10.0654	124.7	0.188	
07411-0124	A 1968	—	10.0654	314.6	0.301	
07417+0942	STF 1130	37484	10.0654	46.8	0.456	
07430-4511	WSI 99	37606	1.0795	116.2	0.131	$\Delta m = 1.0$
			6.1882	47.2	0.050	
07573+0108	STT 185	38870	10.0654	12.6	0.328	
08017-0836	A 1580	39264	10.0654	291.4	0.294	
08088-7638	RST 1381	39887	1.0742	113.5	0.792	
			10.0657	111.1	0.783	
08122+1739	STF 1196 AB	40167	1.0200	78.8	0.857	3
			1.0768	78.5	0.854	
			5.8636	54.4	0.983	
			6.1910	53.3	0.988	
08125-4616	SEE 96 AB	40183	10.0656	274.6	0.615	
08225-5931	HDS 1190	41037	10.0657	141.1	0.475	
08250-4246	RST 4888 AB	41250	10.0656	104.6	0.519	
08251-4910	RST 321	41261	6.1992	106.8	0.291	
			10.0656	141.9	0.354	
08315-1935	I 489	41817	10.0656	295.1	0.350	
08331-2436	BU 205 AB	41949	10.0656	296.5	0.567	
08380-0844	HDS 1242	42345	10.0654	12.7	0.196	
08391-2240	BU 208 AB	42430	1.0825	33.3	1.312	
08421-5245	B 1624	42695	1.0825	105.9	0.290	
			6.1992	142.0	0.165	

Table A1—Continued

WDS or α, δ (2000)	Discovery Designation	HIP	BY 2000.0+	θ ($^{\circ}$)	ρ ($''$)	Note
09191–4128	CHR 239	45705	1.0825	166.6	0.108	1
09264–4215	B 1122	46290	10.0658	230.8	0.161	
09272–0913	A 1588 AB	46365	10.0660	195.7	0.431	
09278–0604	B 2530	46404	1.0825	153.7	0.362	
09285+0903	STF 1356	46454	1.0200	86.5	0.588	3
			1.0770	86.5	0.589	
			1.0825	87.2	0.583	
			5.8637	95.5	0.663	
			6.1913	96.0	0.667	
09379+4554	A 1765	47250	1.0284	116.4	0.265	
09414+3857	STF 1374 AB	47527	1.0201	307.1	2.851	
09487–2625	I 205	48125	1.0797	332.8	1.097	
09525–0806	AC 5 AB	48437	10.0660	50.2	0.551	
09551–2632	I 843 AB	48645	10.0659	130.8	0.929	
10043–2823	I 292	49336	10.0659	305.7	0.713	
10129–4729	I 361	50036	1.0745	122.6:	5.389:	
10161–5954	HU 1597	50287	10.0658	144.6	0.294	
10183–0326	RST 4454 AB	—	10.0660	222.5	0.294	
10196–5724	FIN 162	—	10.0658	177.1	0.408	
10205+0626	STF 1426 AB	50637	10.0660	311.0	0.942	
10279+3642	HU 879	51233	1.0201	187.6	0.065	
			1.0284	189.9	0.067	
10361–2641	BU 411	51885	10.0659	307.0	1.340	
10397+0851	STT 224 AB	52187	10.0660	140.6	0.512	
10419–7811	HDS 1530	52351	1.0746	113.6	0.278	1
			10.0657	125.1	0.163	
10446+0530	A 2771		10.0660	123.0	0.394	
10452–5944	WSI 56	52547	10.0658	59.2	0.275	
10468–4925	R 155	52727	1.0745	52.5	2.013	
11080–6457	HDS 1588	54405	10.0657	352.3	0.592	
11102–1122	HDS 1590	54580	10.0660	344.5	0.134	
11182+3132	STF 1523 AB	55203	1.0201	268.7	1.827	3
			1.0284	268.8	1.823	
			1.0773	267.1	1.810	

Table A1—Continued

WDS or α, δ (2000)	Discovery Designation	HIP	BY 2000.0+	θ ($^{\circ}$)	ρ ($''$)	Note
12313–4130	I 82	61101	10.0659	11.0	0.737	
12362–4650	RST 5526	61517	10.0658	125.9	0.420	
12446–5717	FIN 65 AB	62179	10.0688	90.3	0.268	
12533+4246	COU 1579	—	8.4611	120.6	0.207	
12595+2814	WSI 1	63407	8.4501	112.5	0.147	
13064+2217	WSI 104	63951	8.4501	142.0	0.061	$\Delta m = 0.3$
13137–6248	HDS 1852	64537	10.0714	120.8	0.161	
13138–6335	WSI 57	—	10.0687	236.4	0.194	
13147–6355	WSI 58 Aa,Ab	64624	10.0688	279.3	0.244	
	MLO 3 AB		10.0688	38.2	1.693	
13198+4747	HU 644 AB	65026	8.4569	92.9	1.424	2
13253+7559	STT 267	65489	8.4554	55.1	0.127	
13372–6142	I 365 AB	66438	10.0688	198.7	0.326	
13461+0507	STF 1781	67186	10.0660	189.4	0.934	
13520–3137	BU 343	67696	10.0715	210.8	0.546	
13539–1910	HU 898	67859	8.4609	285.5:	0.310:	
13563+0517	STT 273 AB	68081	10.0660	112.2	1.019	
14153+0308	STF 1819	69653	10.5907	178.3	0.847	
14165+3334	HU 1266	69742	8.4611	138.2	0.254	
14179+6914	A 1102	69870	8.4611	84.0	0.463	
14462–2111	FIN 309	72217	10.5878	94.3	0.224	
14581–4852	WSI 80	73241	6.1892	327.3	0.236	5, $\Delta m = 0.7$
14588+3551	COU 1136	—	8.4611	64.3	0.085	
14593+4649	COU 1760	73346	8.4480	244.5	0.219	2
15127+6008	HU 1159	74428	8.4611	350.8	0.289	
15210+2104	HU 146	75117	8.4566	124.4	0.696	6
15226–4755	SLR 20	75255	6.1998	27.1	0.783	
15245+3723	CHR 181 Aa,Ab	75411	7.6013	302.7	0.117	
			8.4611	264.2	0.093	
15258+8430	MLR 347	75529	8.4611	213.5:	0.197:	
15278+2906	JEF 1	75695	7.6012	142.2	0.301	
15333+4149	STF 1956 AB	76156	7.6013	63.7:	0.096:	
			8.4611	237.6	0.107	
15347+2655	COU 798	—	7.6012	191.8	0.143	

Table A1—Continued

WDS or α, δ (2000)	Discovery Designation	HIP	BY 2000.0+	θ ($^{\circ}$)	ρ ($''$)	Note
15557–2645	I 977	78002	10.5853	253.4	0.460	2
16044–1122	STF 1998 AB	78727	10.5880	355.8	0.948	
16057–0617	BU 948 AB	78849	10.5907	86.2	0.742	
16057–3252	SEE 264 AB	78842	10.5853	19.4	0.716	
	WSI 84 Ba,Bb		10.5853	58.3	0.105	
16079+1425	A 1798	79034	7.6012	350.4	0.178	
16120–1928	BU 120 AB	79374	10.5907	1.0	1.344	
16161–3037	I 1586	79706	10.5853	204.3	0.337	
16169+0113	A 2181	79774	7.5876	77.1	0.485	
16170–5342	I 987	79787	10.5854	265.6	0.452	2
16199+2238	LAW 4 Aa,Ab	80003	7.6012	27.2	0.176	1
			8.4614	30.6	0.191	
16224–3220	JSP 691	80199	10.5853	23.5	0.529	2
16229+3803	COU 1281	—	7.6013	193.1	0.199	
			8.4613	195.4	0.198	
16238+6142	STF 2054 AB	80309	8.4612	350.5	0.982	
16253–4909	TOK 50 Aa,Ab	80448	10.5854	189.0:	0.229:	1
16263–0943	RST 3945	—	7.6013	335.6	0.237	
16274–3729	B 872	80603	10.5854	217.2	0.125	
16309+0159	STF 2055 AB	80883	10.5879	37.2	1.384	
16318–0701	STF 3105	80954	10.5907	181.9	0.448	
16366+6948	BU 953 AB	81332	7.6067	72.2	0.248	
			8.4612	74.0	0.251	
16385–5728	TOK 51 Aa,Ab	81478	10.5854	61.2	0.286	1,2
	RST 869 AB		10.5854	46.7	0.825	2
16413+3136	STF 2084	81693	1.4961	321.9	0.479	3
			7.6068	197.4	1.038	
16420+7353	MLR 198	81757	7.6067	73.1	0.196	
			8.4611	74.1	0.199	
16446+7145	MLR 182	81961	7.6067	290.9:	0.290:	
			8.4612	284.4	0.297	
16450+3842	COU 1284	—	7.6068	355.2	0.180	
			8.4612	356.1	0.177	
16450+2928	COU 490	—	8.4613	293.9	0.121	

Table A1—Continued

WDS or α, δ (2000)	Discovery Designation	HIP	BY 2000.0+	θ ($^{\circ}$)	ρ ($''$)	Note
17141+5608	STT 327	84300	7.6068	329.1	0.457	
			8.4612	330.2	0.453	
17141-0824	BAR 7	84303	10.5908	51.9	1.443	
17156-3836	FIN 355	84425	10.5856	194.5	0.306	
17157-0949	A 2592 AB	84430	10.5908	144.2	0.191	
17161+2316	COU 315	84468	8.4613	132.6	0.125	
17184+3240	BU 628	84653	7.6041	265.7	0.535	
			8.4613	266.3	0.540	
17191-3849	WSI 60	—	10.5883	333.6	1.331	1
17207-0706	A 2593	84866	7.6014	2.6	0.237	
17215+2845	KUI 80 AB	84934	8.4613	169.6	0.640	
17221+2310	COU 415	84976	7.6014	259.9	0.247	
			8.4613	258.7	0.249	
17240-0921	RST 3972	85141	7.6013	192.7	0.108	
17247+3802	HSL 1 Aa,Ac	85209	7.6041	236.5	0.202	
17248-5913	I 385 AB	85216	10.5854	121.4	0.388	
	WSI 87 AD		10.5854	92.3	0.257	
17272-2957	B 340	85413	1.5614	295.1	0.172	
17305-1446	HU 177	85679	7.6013	199.8	0.205	
			10.5908	184.9	0.228	
17308-3726	B 912	85700	10.5856	168.7	0.257	
17313+1901	COU 499	85740	7.6014	4.7	0.102	
17335+5734	MLR 571	85923	7.6041	195.9	0.088	
17366+0723	A 1156	86174	7.6014	348.1	0.205	
17368-2057	HU 751	—	10.5908	296.7	0.273	
17375+2419	CHR 63	86254	7.6014	61.8	0.098	
17375-3747	B 915	—	10.5856	312.5	0.267	
17390+0240	WSI 88	86374	8.4561	5.2	0.169	5, $\Delta m = 2.5$
17490+3704	COU 1145	87204	7.6041	321.6	0.124	
17530+8354	STT 349	87534	7.6041	47.2	0.477	
17533+2459	A 235	87565	7.6021	280.7	0.749	
17535-0355	TOK 54	—	10.5908	137.4	0.126	
17535-1623	WSI 64	—	10.5883	101.7	2.096	1
17539-3445	B 1871	87616	10.5857	273.9	0.263	

Table A1—Continued

WDS or α, δ (2000)	Discovery Designation	HIP	BY 2000.0+	θ ($^{\circ}$)	ρ ($''$)	Note
18054+6216	HU 1290	88592	7.5876	309.6	0.224:	
18078+2606	CHR 67 Aa,Ab	88818	7.5878	312.9	0.235	
18096+0400	STF 2281 AB	88964	10.5908	288.2	0.644	
18112–1951	BU 132 AB	89114	10.5908	188.0	1.396	
18177–1940	BU 246 AB	89647	10.5909	114.9	0.497	2
18211+7244	LAB 5 Aa,Ab	89937	7.5876	216.3	0.132	
18218–1619	CHR 69	90001	7.5880	209.3	0.076	
18236–2610	HO 566	90128	10.5909	144.5	0.535	2
18253+4846	STT 351 AC	90284	7.5876	22.6	0.742	2
	HU 66 BC		7.5876	27.0	0.962	
18337+5638	TDS 929	—	7.6069	78.0	1.188	1
18338+1744	STF 2239 AB,CD	87407	10.5909	276.5	1.580	
18339+5221	A 1377 AB	91013	7.5876	120.8	0.248	
18374+7741	STT 363	91298	7.5876	336.2	0.446	
18384+0850	HU 198	91389	7.5879	128.0	0.503	
18384–0312	A 88 AB	91394	6.2001	154.3	0.101	3
			7.5879	63.0:	0.063:	
			8.4507	17.6	0.100	
18411+4416	HDS 2647	91615	7.6069	268.0	0.140	1
18443+3940	STF 2382 AB	91919	8.4479	346.8	2.354	
	STF 2383 CD	91926	7.5878	78.7	2.351	
			8.4551	79.8	2.368	
18450+4239	HDS 2657	91994	7.6069	32.2	0.240	1
18475+4926	BU 971 AB	92204	8.4479	34.3	0.450	
18552+3941	TDT 1126	—	7.6071	189.9	0.323	1
18558+0327	A 2192	92909	7.5879	31.0	0.170	
18564+5854	TDS 955	—	7.6071	145.7	0.885	1
18567+6440	TDT 1141	—	7.6069	153.3	0.450	1
18579+4215	TDT 1149	—	7.6071	122.7:	0.456:	1
19006+3952	HDS 2696	93335	7.6071	136.8	0.294	1
19025+5142	TDT 1193	—	7.6071	90.6:	1.168:	1
19029+4930	COU 2626	—	7.6071	84.5	0.893	1
19038+4040	COU 2196	—	7.6071	330.1	0.490	1
19045+6245	TDT 1211	—	7.6069	31.5	0.795	1

Table A1—Continued

WDS or α, δ (2000)	Discovery Designation	HIP	BY 2000.0+	θ ($^\circ$)	ρ ($''$)	Note
19296+1224	A 1653	95845	7.6014	152.9	0.198	
19296-1239	HU 75	95847	7.6014	34.3	0.165	
19302+5639	A 712	95901	7.6016	85.8	0.116	
19308+6337	HU 951	95955	7.6016	197.2	0.187	
19313+4729	A 713	96011	7.6016	296.9	0.236	
			8.4562	300.1	0.230	
19352+0825	BAG 27 Aa,Ab	96339	7.6014	36.6	0.291	1
19356+4002	A 1400	96371	7.6016	115.1	0.320	
19389+3514	HU 953	—	7.6016	19.5	0.237	
19411+1349	KUI 93	96840	10.5909	310.9	0.168	
19426-5901	I 119	96955	1.5642	150.6	2.190	
19471-1953	BU 146	97348	10.5909	237.8	0.539	
19474-0148	A 2993	97367	7.6042	298.8	0.150	
19531-1436	CHR 90	97849	10.5909	20.2	0.266	
19550+4152	HO 581	98001	1.4991	98.0	0.354	3
19553-0644	STF 2597 AB	98038	10.5909	100.7	0.595	
19556+5226	YR 2 Aa,Ab	98055	7.6042	53.1	0.116	
19573+0513	A 604	98203	7.6042	22.5	0.110	
20012-3835	HDO 294	98556	10.5861	29.4	1.176	
20014+1045	TOK 34 Aa,Ab	98578	7.6042	339.0:	0.378:	1
20062+3520	TDT 1981	—	7.6071	0.0	0.420	1
20095+3525	TDT 2034	—	7.6071	120.1	0.339	1
20096+5034	TDT 2035	—	7.6071	90.8	0.783	1
20124+5354	HDS 2880	99573	7.6071	273.7	0.113	1
20154+6412	MLR 60 AB	99832	7.6041	348.8	0.220	
20187+5823	TDT 2157	—	7.6071	35.3	0.765	1
20203+3924	A 1427 AB	100268	7.6042	123.7	0.271	
20213+0250	HLD 158	—	7.6042	43.9	1.090	
20251+5936	A 730	100714	7.6041	288.2	0.143	
20267-4334	I 1420	100836	10.5860	346.6	0.447	2
20274+7245	A 871	—	7.6041	183.2	0.305	
20290+0710	A 610	101046	7.6042	70.2	0.436	
20302+2651	WOR 9 AB	101150	8.4589	261.3	0.518	
20311+1548	A 1675	101223	1.4991	339.5	0.078	

Table A1—Continued

WDS or α, δ (2000)	Discovery Designation	HIP	BY 2000.0+	θ ($^{\circ}$)	ρ ($''$)	Note
21041+0300	WSI 6	103987	7.5880	132.7	0.121	1
			8.4550	167.5	0.095	
21044–1951	FIN 328	104019	7.5880	176.2	0.268	
			10.5861	151.6:	0.321:	
21056+6030	A 758 AB	104113	7.5990	122.7:	0.343:	
21105+4300	VKI 39	—	7.6072	234.3	1.323	1
21115+2144	COU 227	104612	7.6017	105.6	0.523	
21120+4419	TDT 2816	—	7.6072	88.6:	0.289:	1
21125+2821	HO 152	104701	7.6017	131.0	0.204	
21135+1559	HU 767	104771	7.6017	164.5	0.174	
21137+6424	H 1 48	104788	7.5990	238.0	0.276	
21148+4946	COU 2653	—	7.6072	65.1	0.298	1
21148+3803	AGC 13 AB	104887	7.6018	250.1	0.753	
21198–2621	BU 271 AB	105312	10.5861	244.3:	1.704:	
21200+5436	TDT 2908	—	7.6073	295.9	0.301	1
21203+5354	TDT 2910	—	7.6072	284.2	0.666	1
21226+3846	HDS 3044	105543	7.6072	141.0	0.122	1
21239+8317	HDS 3049	105644	7.6073	335.1	0.303	1
21243+3740	WSI 7	105676	7.6018	101.6	0.157	1
			8.4615	94.5	0.166	
21243+0343	A 2288	105682	7.5881	234.7	0.387	
21251+0923	BU 164 AB	105747	7.5881	144.6	0.085	
21268+3732	COU 1821	—	7.6072	93.8	0.651	1
21310–3633	B 1008 AB	106224	10.5861	21.8	0.150	
21373+6604	HDS 3077	106733	7.6073	273.4	0.127	1
21410+2920	STT 448	107060	7.5991	145.6	0.156	
21424+4105	KUI 108	107162	1.5020	190.8	0.101	
			8.4508	59.6	0.146	
21434+2353	HU 374	107254	7.5991	131.0	0.128	
21435+4448	LIN 1 Aa,Ab	107256	8.4509	187.3	0.165	
21446+2539	BU 989 AB	107354	7.5991	319.6	0.155	
21459+1153	A 1223 AB	107461	7.5991	331.0	0.115	
21477–1813	CHR 223	107610	7.5992	299.8	0.169	1
21477–3054	FIN 330 AB	107608	10.5861	43.7	0.096	

Table A1—Continued

WDS or α, δ (2000)	Discovery Designation		HIP	BY 2000.0+	θ ($^\circ$)	ρ ($''$)	Note
22342–1841	HU	389 AB	111406	7.5992	281.6	0.167	
22405+2002	WSI	9	111943	8.4615	133.4:	0.175:	1
22409+1433	HO	296 AB	111974	1.5020	14.0	0.234	3
				1.5645	12.8	0.228	
				5.8626	104.7	0.175	
				7.5883	82.5	0.310	
				8.4617	77.9	0.357	
22519–4647	I	683	112908	1.5646	345.2	1.917	
22550+5132	HU	785	113166	7.5884	102.5	0.147	
22562+7250	STT	484 AB	113273	7.5885	100.8	0.355	
22586–4531	HU	1335	113454	1.5646	109.3	0.350	
22595+4355	COU	2146	113524	7.6042	287.2	0.243	8
23026+4245	BU	1147 AB	113788	1.5020	354.6	0.318	
23114+3813	HO	197 AB	114504	7.5884	286.9	0.197	
23164+6407	BU	992	114898	7.5884	21.3	0.249	
23176+1818	HU	400	115011	7.5883	83.7	0.342	
23191–1328	MCA	74 Aa,Ab	115126	5.8655	25.1	0.100	3
23199+2844	COU	439	115189	7.5884	314.4	0.125	
23209+1643	HEI	88	115279	7.5883	148.2	0.157	
23210+1715	WSI	11	—	7.5883	161.4	0.108	1
23277–1502	HU	295	115404	10.5861	282.1	0.317	
23319+7317	A	897	116133	7.5992	331.4:	0.178:	
23350+0136	MEL	9 AC	116384	7.6018	342.2	0.760	
23374+0737	FOX	102 AB	116571	7.5994	41.6	0.173	
23378+6601	HU	1324	116598	7.5992	175.0	0.372	
23382+5514	A	1493	116634	7.5885	311.7	0.155	
23424+3903	A	1494	116952	7.5992	338.7	0.099	
23498+2741	A	424	117511	7.5992	152.1	0.159	
23502+1940	COU	344	117531	7.5992	332.4	0.188	
23528+3841	A	1496 AB	117752	7.5992	152.4	0.413	2
23561+2520	A	426	118005	1.5021	301.2:	0.370:	
23586–1408	RST	4136 AB	118205	7.6019	215.0	0.115	

¹ Confirming observation.

⁶ Three measurements are averaged to give this mean position.

⁷ First measure of newly resolved pair. Primary is HD 341480.

⁸ Confirming observation or more likely a new component.