The First Three Rungs of the Cosmological Distance Ladder

Kevin Krisciunas,¹ Erika DeBenedictis,² Jeremy Steeger,³ Agnes Bischoff-Kim,⁴ Gil Tabak,⁵ and Kanika Pasricha⁶

ABSTRACT

It is straightforward to determine the size of the Earth and the distance to the Moon without making use of a telescope. The methods have been known since the 3rd century BC. However, few amateur or professional astronomers have worked this out from data they themselves have taken. Here we use a qnomon to determine the latitude and longitude of South Bend, Indiana, and College Station, Texas, and determine a value of the radius of the Earth of 6290 km, only 1.4 percent smaller than the true value. We use the method of Aristarchus and the size of the Earth's shadow during the lunar eclipse of 2011 June 15 to derive an estimate of the distance to the Moon (62.3 R_{\oplus}), some 3.3 percent greater than the true mean value. We use measurements of the angular motion of the Moon against the background stars over the course of two nights, using a simple cross staff device, to estimate the Moon's distance at perigee and apogee. Finally, we use simultaneous CCD observations of asteroid 1996 HW1 obtained with small telescopes in Socorro, New Mexico, and Ojai, California, to derive a value of the Astronomical Unit of $(1.59 \pm 0.19) \times 10^8$ km, about 6 percent too large. The data and methods presented here can easily become part of a beginning astronomy lab class.

Subject headings: Astronomical Techniques

¹George P. and Cynthia Woods Mitchell Institute for Fundamental Physics & Astronomy, Texas A. & M. University, Department of Physics and Astronomy, 4242 TAMU, College Station, TX 77843; krisciunas@physics.tamu.edu

²MSC #227, Caltech, Pasadena, CA 91126; edebened@caltech.edu

³3 Ames Street, Box 62, Cambridge, MA 02142; jsteeger@mit.edu

⁴Department of Chemistry, Physics and Astronomy, Georgia College & State University, Campus Box 82, Milledgeville, GA 31061; agnes.kim@gcsu.edu

⁵6530 El Colegio Road, Box #3101, Santa Barbara, CA 93106; tabak.gil@gmail.com

⁶2269 Frist Campus Center, Princeton University, Princeton, NJ 08544; pasricha@princeton.edu

1. Introduction

As scientists and as educators we ask ourselves, "How do we know what we know?" If we had to derive everything we used from scratch, we would not have the time to make much progress. But it is worthwhile to derive *some* fundamental parameters that we use in our area of expertise. In 1987 MIT physicist Philip Morrison hosted a television series called "Ring of Truth: an inquiry into how we know what we know." One purpose of the show was to demonstrate simple measurements. In the episode called "Mapping" he measured the circumference of the Earth with the "van of Eratosthenes". He and his crew went to the north edge of Kansas and measured the elevation angle of the bright southern star Antares as it transited the meridian. Then they drove 370 miles down Highway 183 to the south edge of Kansas and measured Antares again. They found that it was 5 degrees higher at the second location. The distance corresponded to 1/72 of a circle, so the implied circumference of the Earth was 26,600 miles, which is a bit more than 6 percent greater than the correct value (24,901 miles).

The cosmological distance ladder is a sequence of steps used by astronomers to derive distances within the solar system, throughout the Galaxy, and beyond to the farthest galaxies detectable (Rowan-Robinson 1985). It hinges on simple geometry and the principles of surveying. For example, using the positions of two observers separated by some baseline on the Earth, we can determine the distance to the Moon, a nearby planet or an asteroid.

In this paper we determine the sizes of the first three rungs of the cosmological distance ladder. These are: 1) the radius of the Earth; 2) the distance to the Moon; and 3) the distance to the Sun (i.e. the size of the Astronomical Unit, or AU). The first two can be accomplished without a telescope. The third was attempted by various pre-telescopic astronomers such as Tycho Brahe in 1582 (Gingerich & Voelkel 1998), but was not accomplished until 1672 by Gian Domenico Cassini and John Flamsteed (Van Helden 1995), who made observations of Mars when it was prominently visible in the middle of the night and therefore about as close to the Earth as it gets. (Mars is roughly 0.6 AU distant at such a time. When it is on the other side of the Sun it is 2.6 AU distant.) Measuring the size of the AU requires telescopic measurements or distances to nearby planets or asteroids determined with radar.

Once the scale of the solar system is known, we can determine the distances to nearby stars via trigonometric stellar parallaxes if our positional measurements are good to better

⁷See http://www.viddler.com/explore/jacksmernov/videos/6. Of course, they are referring to the Greek astronomer Eratosthenes (ca. 276-195 BC) who obtained the first estimate of the Earth's circumference from observations of the Sun's elevation on the summer soltice from Alexandria and a town 7 degrees to the south.

than 0.1 arc second. Then, classically, we determine the distance to the Hyades star cluster (Rowan-Robinson 1985, p. 48) and tie other star clusters to the Hyades distance. With the discovery of certain standard candles and standardizable candles in star clusters (e.g. RR Lyrae stars and Cepheids), we can calibrate the mean intrinsic brightness of these pulsating stars. They are useful for distance determinations throughout our Galaxy. With the Hubble Space Telescope we can determine distances to other galaxies using the Cepheid period-luminosity relation as far as 25 million parsecs (Riess et al. 2010). (1 parsec equals 3.086×10^{13} km, or 206265 AU.) If we observe Type Ia supernovae in some galaxies whose distances are known via Cepheids, we can calibrate the intrinsic brightness of these supernovae. Since Type Ia supernovae are typically 4 billion times brighter than the Sun at maximum brightness, they can be used to determine distances halfway across the observable universe with a 4-m class telescope.

The methodology of this paper is to use only our own data, if possible. We make minimal use of the *Astronomical Almanac* and minimal use of telescopes. Many of the results here can be obtained independently by students using very unsophisticated and inexpensive equipment.

2. Determining one's position on the Earth

In Fig. 1 we show the astronomical triangle. The northern sky turns around the North Celestial Pole (NCP), near the direction of the star Polaris. The NCP is ϕ degrees above the horizon. ϕ is the latitude. The azimuth (A) is measured clockwise around the horizon and is equal to 0° deg at the north point on the horizon, 90° deg at the east point on the horizon, etc. The hour angle t divided by 15 deg/hr is the number of hours that an object is west of the celestial merdian. t is negative for objects in the eastern sky.

The declination (δ) of a celestial object is the number of degrees the object is north or south of the celestial equator. If we happen to know the declination of the Sun and we determine how high it is above the horizon when it transits the *celestial meridian*,⁸ we can determine our latitude. At local apparent noontime the elevation angle of the Sun will be:

⁸The celestial meridian is the imaginary line in the sky that separates the east half from the west half. For an observer in the northern hemisphere the meridian extends from the north point on the horizon through the North Celestial Pole, through the zenith and down to the south point on the horizon. An object such as the Sun is at its highest point above the horizon when it crosses the celestial meridian. At that moment the local apparent solar time is, by definition, exactly 12 noon.

$$h_{max} = 90^{\circ} - \phi + \delta. \tag{1}$$

We are concerned here with the Sun and Moon as observed at mid-northern latitudes. These objects transit the celestial meridian between the zenith and the south point on the horizon.

To determine the elevation angle of the Sun we can set up a vertical pointed stick or gnomon. Even better is a vertical stick with a small sphere at the top, like that shown in Fig. 2. It is easier to measure the center of the elliptical shadow of the sphere on the ground than it is to measure the end of the darker part of the shadow of a vertical pointed stick.⁹

The Sun's declination ranges from $-\epsilon$ to $+\epsilon$ over the course of the year. ϵ is the *obliquity* of the ecliptic, the tilt of the Earth's axis of rotation to the plane of its orbit. (The ecliptic is the apparent path of the Sun through the constellations of the zodiac owing to the Earth's orbit around the Sun.) On the first day of summer we have:

$$h_{max}(Jun 21) = 90^{\circ} - \phi + \epsilon . \tag{2}$$

And on December 21st we have:

$$h_{max}(\text{Dec }21) = 90^{\circ} - \phi - \epsilon . \tag{3}$$

Eq. 2 plus Eq. 3 gives:

$$\phi = 90^{\circ} - [h_{max}(Jun\ 21) + h_{max}(Dec\ 21)]/2$$
 (4)

And Eq. 2 minus Eq. 3 gives us:

$$\epsilon = [h_{max}(\text{Jun } 21) - h_{max}(\text{Dec } 21)]/2 .$$
 (5)

In other words, if we determine the maximum elevation angle of the Sun on June 21st and December 21st, the average gives us $90^{\circ} - \phi$, and half the difference gives us ϵ .

In Table 1 we give the raw data from two solstice experiments carried out in College Station, Texas.

⁹If a vertical pointed stick is used, this will give us the elevation angle of the upper limb of the Sun. To find the elevation angle of the center of the Sun then requires subtracting the angular radius of the Sun.

In Fig. 3 we show the X-Y positions of the end of the gnomon's shadow for three key times of the year. Note that when the declination of the Sun is negative the positions of the end of the shadow on the ground trace out an upward pointing curve. On the first day of spring (March 20 or 21) or the first day of autumn (about September 23) the points delineate a straight line. When the declination of the Sun is positive the points trace out a downward curving locus. A graph of this type allows one to determine which points, if any, have been mismeasured. If a student invents data for such an experiment, such a graph allows one to detect such fraud easily.

Plots of the data from Table 1 are shown in Fig. 4. We have converted the time values to the number of minutes since local apparent noontime (when $h = h_{max}$).¹⁰ In the top graph we have added, for graphical purposes only, a hyperbolic fit to the data. In fact, there is no simple function that fits such a data set. The best estimate of the minimum value in the summer solstice plot is obtained from a fourth order polynomial fit to the data within 41 minutes of the meridian transit of the Sun. For the 2010 June 21 observations we used a gnomon of height $g = 632 \pm 1$ mm. The minimum shadow length was $L_{min} = 85.84$ mm with an average scatter of ± 0.64 mm. This average scatter is what is known as a random error, as the points scatter randomly above and below an appropriate function fit to the data. h_{max} is equal to the arctangent of g/L_{min} , or 82° 15′. 9 \pm 3′. 5.

For 2010 December 21 we used a gnomon of height $g = 550 \pm 1$ mm. From a fourth order polynomial fit to all 26 points we find $L_{min} = 753.98 \pm 0.74$ mm. h_{max} is found to be 36° 06′. 6 ± 3 ′. 4.

Eq. 4 gives $\phi = 30^{\circ}$ 48′. 7 ± 3 ′. 5, which is 11′. 5 north of the true value of 30° 37′. 2 from Google Earth. (One arc minute of latitude equals one nautical mile, or 1852 m.) Eq. 5 gives $\epsilon = 23^{\circ}$ 04′. 7 ± 3 ′. 5, which is several standard deviations smaller than the true value of 23° 26′. 2. These values show what kind of uncertainty can be obtained with careful observations.

So far we have used no information from the Astronomical Almanac. However, to determine our longitude requires that we know the amount of time that the apparent Sun is ahead or behind the mean Sun. This difference is known as the equation of time. It ranges from -14.2 minutes to +16.4 minutes over the course of the year.

¹⁰Apparent solar time relates to the hour angle of the actual visible Sun in the sky. Due to the tilt of the Earth's axis of rotation to the plane of its orbit, and due to the ellipticity of the Earth's orbit, apparent solar time ranges from 14 minutes behind to 16 minutes ahead of mean solar time. Our watch time is basically mean solar time adjusted (by longitude difference) to the nearest 15 degree line of longitude west of Greenwich, England. For example, Central Standard Time is for locations that are about 90° longitude west of Greenwich, or 6 hours west.

From four gnomon experiments carried out in College Station, Texas, on 2006 November 12, 2007 September 24, and the two solstices mentioned above we found that the mean Sun transits the meridian at 12:24:26 PM standard time, with an uncertainty of ± 72 sec. Since each degree of longitude corresponds to 4 minutes of time, our longitude is 96° 06′. 5 ± 18 ′. 1. We are within one standard deviation of the true value of 96° 20′. 4.

3. The circumference of the Earth

On 2006 September 3 we carried out a gnomon experiment in South Bend, Indiana, on the campus of the University of Notre Dame. For this we needed a value of the declination of the Sun from the *Astronomical Almanac*. Our location was found to be latitude 41° 51′. 5, longitude 85° 50′. 5.

In late October of 2007 we drove from South Bend through Bella Vista, Arkansas, and on to College Station, Texas, keeping track of our route, mileage, and the length of any side trips. After subtracting the side trips, the elapsed distance on the odometer was 1248 statute miles. For a section of highway through central Illinois we noted that 96.8 miles on the odometer corresponded to 98.0 miles according to the highway markers. The implication is that our odometer exhibited a systematic error. On a subsequent occasion we drove the final 92 percent of the exact same route, but with a different car, and the odometer mileage was 40 miles less (about 4 percent). We conclude that an odometer cannot be explicitly trusted. We adjust the original mileage to $1248 \times \frac{98.0}{96.8} = 1263$ miles.

Next, we used a "map tool" (also known as an *opisometer*) and traced our route on a map from the 1980 edition of *The New International Atlas* by Rand McNally, and determined that if we could have traveled along a great circle route from South Bend to College Station, the direct route had a length that was 0.744 of the length of the route we actually drove. (This means that we did not rely on the stated scale of the map.) Thus, the great circle distance from one place to the other was 940 statute miles.

Consider two locations on the Earth with (latitude, longitude) = (ϕ_1, λ_1) and (ϕ_2, λ_2) , respectively. The length of the great circle arc between them (ρ) can easily be obtained from the law of cosines of spherical trigonometry:

$$\cos(\rho) = \sin(\phi_1)\sin(\phi_2) + \cos(\phi_1)\cos(\phi_2)\cos(\lambda_2 - \lambda_1). \tag{6}$$

Given the latitudes and longitudes determined by gnomon and clock at both locations, the great circle arc was 13.78 deg. Thus, our estimate of the circumference of the Earth is $940 \times \frac{360}{13.78} = 24557$ miles, which is 1.4 percent smaller than the true value. Our value for the radius of the Earth is 6290 km. The true value is 6378 km. Given the simplicity of the tools involved, we feel that we have achieved an accuracy far better than our expectations.

For comparison we refer the reader to Bekeris et al. (2011), which describes projects "Eratosthenes 2009" and "Eratosthenes 2010 America". More than 15,000 students at more than 200 schools determined the radius of the Earth using this method. They obtained 6290 km in 2009 and 6375 km in 2010.

4. The distance to the Moon

4.1. Using the method of Aristarchus

The method used above to determine the circumference of the Earth is in principle the same as that used by Eratosthenes. It turns out that it was a generation *before* Eratosthenes that Aristarchus of Samos (ca. 310-230 BC) determined the distance to the Moon in terms of the radius of the Earth. Inherent in both methods is the acceptance of the idea that the Earth is, for all intents and purposes, spherical.

Aristarchus cleverly deduced that we can determine the distance to the Moon from the geometry of a lunar eclipse. See Evans (1998, pp. 68-73) and Fig. 5. A lunar eclipse can only take place when the Moon is opposite the Sun. The shadow of the (spherical) Earth is circular in any plane perpendicular to the Earth-Sun line. Consider the triangle delineated by points A, C, and H. AC is the radius of the Earth, R_{\oplus} . CH is the distance from the center of the Earth to the Moon, d_{Moon} . The horizontal parallax of the Moon, P_M , is given by:

$$\sin(P_{\rm M}) = \frac{R_{\oplus}}{d_{\rm Moon}} \,. \tag{7}$$

Thus, the distance to the Moon in Earth radii is $1/\sin(P_M)$. In Fig. 5 P_S is the horizontal parallax of the Sun. σ is the angular radius of the Sun. τ is the angular radius of the Earth's shadow at the distance of the Moon. For \triangle XCH it is obvious that \angle XCH + P_S + P_M = 180°. In other words, $180^{\circ} - \angle$ XCH = P_S + P_M . Also, σ + \angle XCH + τ = 180°, which means that $180^{\circ} - \angle$ XCH = σ + τ . It follows that:

$$P_S + P_M = \sigma + \tau . (8)$$

Aristarchus believed that the Sun was 19 times more distant than the Moon. The true

ratio is closer to 400. Our value for the radius of the Earth and our value of the distance to the Sun given in §5 below means that the solar parallax $P_s \approx 0.14$.

Aristarchus knew that the Moon and Sun approximately had the same angular diameter because the duration of a total solar eclipse is never more than a few minutes. He took the angular diameter of the Moon to be 0.5 deg. In the *Almagest*, section V 14, Ptolemy states that the Earth's shadow at the distance of the Moon is 2.6 times the angular diameter of the Moon (Toomer 1984, p. 254). We can use two actual measurements of the Sun made on 2010 May 1 and 2011 June 6 to provide an angular diameter of 30.9, so $\sigma \approx 15.9$.

In a previous paper (Krisciunas 2010) we presented measures of the Moon's angular diameter from sightings of the Moon through a 6.2 mm hole in a piece of cardboard that slides up and down a yardstick. Our mean value for the angular diameter of the Moon was 31'.18. Data in that paper and subsequent data yield a time of lunar perigee of 2011 May 14.70 UT, with a perigee-to-perigee period (i.e. the *anomalistic month*) of 27.4992 \pm 0.0518 d, which is within 1- σ of the correct value of 27.55455 d. Based on our non-telescopic data, the lunar eclipse of 2011 June 15 occurred about 4.6 days after perigee, so the Moon's angular size would have been a bit larger than the mean value.

Using six images obtained from www.slooh.com, we determined that the Earth's shadow was 2.56 ± 0.03 times the angular diameter of the full Moon during the lunar eclipse of 2011 June 15. Thus, $\tau = 0.5 \times 31.18 \times 2.56 \approx 39'.19$. It follows that $P_M = 15'.4 + 39'.19 - 0'.14 \approx 55'.17$. Eq. 7 gives a lunar distance of 62.3 R_{\oplus} with a random error of at least ± 0.7 R_{\oplus} . The true mean distance is 60.27 R_{\oplus} .

4.2. Using the motion of Moon against the stars

We have seen that the geometry of a lunar eclipse allows a determination of the distance to the Moon. Hipparchus (ca. 140 BC) used observations of the *solar* eclipse of 14 March 190 BC to derive a distance to the Moon between 71 and 83 R_{\oplus} (Toomer 1981). But can one determine the Moon's distance without the use of an eclipse? Simultaneous observations of the Moon's position from two widely separated places on the Earth would suffice. Smart (1977, pp. 200-202) shows how this can be done from two observatories widely separated in latitude but on the same meridian of longitude.

Observations of the Moon from the *same* location over the course of a night can also be used to determine the distance to the Moon. In this case one also uses a large fraction of the Earth's diameter as a baseline. The complication is that while we wait for the Earth to rotate, the Moon is also orbiting the Earth.

The Moon's orbital period with respect to the background stars is 27.32 d. Thus, on average, the Moon moves 0.55 deg east per hour against the background of stars as viewed by a hypothetical observer at the Earth's center. However, because of the parallax of the Moon, the observed motion for someone situated on the surface of the Earth is roughly 1/6 deg per hour less (Evans 1998, pp. 252-254).

We need the geocentric vs. topocentric shifts in right ascension and declination, respectively, of the Moon owing to our location on the surface of the Earth. The topocentric coordinate system has the observer, standing on the Earth's surface, at the origin. Let $\Delta \alpha = \alpha' - \alpha = t - t'$. (Right ascension increases to the east, while hour angle increases to the west.) Then Eq. 35 of Smart (1977) gives

$$\tan \Delta \alpha = -\left(\frac{R_{\oplus}}{d_{Moon}}\right) \frac{\sin t \cos \phi'}{\cos \delta - \left(\frac{R_{\oplus}}{d_{Moon}}\right) \cos t \cos \phi'}.$$
 (9)

Letting $\delta' = \delta + \Delta \delta$ and T = tan $\Delta \delta$, Eq. 38 of Smart (1977) is

$$\frac{\tan \delta + T}{1 - T \tan \delta} = \frac{\cos t' \left[\sin \delta - \left(\frac{R_{\oplus}}{d_{Moon}} \right) \sin \phi' \right]}{\cos \delta \cos t - \left(\frac{R_{\oplus}}{d_{Moon}} \right) \cos \phi'} . \tag{10}$$

Here the primed parameters are the topocentric values of declination, hour angle, and latitude, while unprimed values are for a hypothetical observer at the Earth's center. Eqs. 9 and 10 cannot be easily inverted to give d_{Moon}/R_{\oplus} . But since these equations contain this ratio, we now show how single-site observations can be used to obtain the distance to the Moon. What follows is primarily a demonstration of method.

In Fig. 6 we show the rate at which the Moon's apparent position (for an observer situated in College Station, Texas) varies as a function of hour angle for a number of occasions over nearly the full range of distance of the Moon from the Earth. Not surprisingly, the Moon moves against the background of stars more slowly on a night when it is near apogee than when it is near perigee.

In Fig. 7 we show a simple cross staff. The pattern for the device that slides up and down the yardstick was obtained from the University of Washington.¹¹ With such a device we can derive the angular separation of two objects in the sky. For two objects with known right ascensions and declinations ρ is given by an expression similar to Eq. 6:

¹¹http://www.astro.washington.edu/courses/labs/clearinghouse/labs/Skywatch/angles.html

$$\cos(\rho) = \sin(\delta_1)\sin(\delta_2) + \cos(\delta_1)\cos(\delta_2)\cos(\alpha_2 - \alpha_1). \tag{11}$$

If the Moon's angular separation from two stars of known RA and DEC is determined, then it is possible to determine the RA and DEC of the Moon. The easiest way to envision this is to take a star chart and a compass and draw portions of two circles of different radius centered at the locations of the two particular stars. One obtains two numerical solutions of the Moon's position, one on each side of the great circle arc joining the two stars. One of those possibilities can easily be eliminated. Once an approximate location of the Moon's position is found, one can use a computer program to search a box that covers a range of RA and DEC to find the celestial location that is the specified number of degrees from each of the two stars of known position.

In Table 2 we give five sets of observations made on the night of 2011 May 15 (UT), an occasion when the Moon was close to the bright star α Vir and within a day of perigee. On this night the Moon was waxing and 93 percent illuminated. Thus, it was visible almost the entire night. One of our reference "stars" was Saturn. Given the accuracy of our observations we can assume that the position of Saturn was constant on this night. As one can see, the angular separation of Saturn and α Vir was measured to better than 0.2 deg on five occasions. Because the Moon is not a point source, measuring the angular separation of the Moon and Saturn or the Moon and a star is more difficult than measuring the angular separation of two bright stars or one bright star and a planet.

In Table 3 we give the derived right ascensions and declinations of the Moon from our observations of 2011 May 15. At the end of the night the reference stars δ Crv and γ Hya were too low in the sky to be seen. The uncertainties of the right ascension and declination of the Moon at 08:03 and 08:41 UT were derived with the assumption of an uncertainty of \pm 0.3 deg for the angular separation of Saturn vs. the Moon and α Vir vs. the Moon. Because of the large uncertainty in the RA and DEC of these final two observations, we exclude data sets four and five from further analysis. The first, second, and third determinations of the position of the Moon yield an apparent angular rate of 0.508 \pm 0.094 deg/hr. The corresponding range of the Moon's distance is 47.3 to 58.5 R_{\phi} with a most likely value of 52.1 R_{\phi}.

2010 October 21 UT was the day after lunar apogee. On this date we made a series of six measurements of the angular separation of the Moon vs. Jupiter over a 9.1 hour period. (No stars near the Moon were bright enough to be seen with the unaided eye given the quality of the sky.) Thus, we could not use the same method to obtain multiple estimates of the RA and DEC of the Moon at a given time. The separation of the Moon and Jupiter increased from 11.31 to 14.77 deg, with a mean rate of increase of 0.403 ± 0.028 deg/hr.

Under the assumption that the Moon was moving directly away from the position of Jupiter (which was only approximately true), this angular rate is a lower limit to the rate of change of position of the Moon against the background of stars. The implied range of lunar distance was 57.2 to 62.1 R_{\oplus}, with a most likely value of 59.5 R_{\oplus}.

If we take the 1- σ upper limit of the rate of change of position of the Moon on 2011 May 15 and the 1- σ lower limit of the rate of change of the lunar position on 2010 October 21, we get a conservative estimate of the range of the Moon's distance, namely 47 to 62 R_{\oplus} . Obviously, the observations and analysis required for this method of determining an estimate of the Moon's distance are much more complicated than Aristarchus's method using the geometry of a lunar eclipse.

5. The distance to the Sun

In 2008 the authors of this paper were participants in the Summer Science Program (SSP), a residential non-credit enrichment program for incoming high school juniors and seniors. Coauthors Krisciunas and Kim were faculty, while the other coauthors were students. It has been the tradition at SSP to divide the students into teams of three for the observing, but each student then writes computer code to produce a determination of the orbital parameters of a particular asteroid. In Table 4 we give the orbital solution of asteroid 8567 (= 1996 HW1) by one of us (J. S.) along with the parameters obtainable from the Horizons website of the Jet Propulsion Laboratory. The code to determine the orbital parameters was debugged using observations of Ceres and the orbital parameters of Ceres given by JPL Horizons. We note that our orbit solution is based on observations extending over a very small arc of the full orbit. (The orbital period of asteroid 1996 HW1 is 2.93 years.)

It had been an ambition at SSP for some time to observe a near Earth object simultaneously from two sites so that its distance could be derived, the length of the Astronomical Unit could be measured, and the scale of the solar system be determined. We report one such experiment here.

On 2008 July 24 UT we took images of asteroid 1996 HW1 at Etscorn Observatory of the New Mexico Institute of Mining and Technology in Socorro, New Mexico, using a 15 cm Takahashi refractor and CCD camera. The geographic position was latitude +34° 04′ 21″, longitude W 106° 54′ 50″. A simultaneous image was obtained in Ojai, California, at latitude +34° 26′ 04″. longitude W 119° 11′ 22″.6 using a 25 cm Meade reflector and CCD

¹²ssd.jpl.nasa.gov/horizons.cgi#top

camera. Figs. 10 and 11 show the asteroid at 08:17:28 UT on 2008 July 24, as viewed from the two sites.

The right ascensions and declinations of a number of field stars were determined using ds9 and an image of the field obtained from the Space Telescope Science Institute Digital Sky Survey. We then determined a transformation from pixel coordinates to right ascension and declination using the IRAF programs ccmap and cctrans. This allowed us to determine the right ascension and declination of the asteroid for each of our images. From the New Mexico site we determined the asteroid's topocentric position to be $\alpha = 21:27:07.42$, $\delta = +15^{\circ} 53' 02''.77$. From the Etscorn image and the positions of the field stars, the root-mean-square errors in right ascension and declination were ± 0.31 and ± 0.36 arcsec, respectively. Our image obtained in California just barely has the asteroid in the frame, and the guiding was not as good. Still, we find a topocentric position of $\alpha = 21:27:07.77$, $\delta = +15^{\circ} 53' 02''.25$. The RMS errors of the field star positions were ± 0.52 and ± 0.44 arcsec, respectively, for right ascension and declination.

The effect on right ascension and declination due to parallax and the finite size of the Earth can be calculated much more easily than using Eqs. 9 and 10 (above) if the planet or asteroid under consideration has a distance d considerably greater than the distance to the Moon. Smart (1977, pp. 209-210) gives these relevant equations:

$$\Delta \alpha = -\left(\frac{R_{\oplus}}{d}\right) \sin t \cos \phi' \sec \delta. \tag{12}$$

$$\Delta \delta = -\left(\frac{R_{\oplus}}{d}\right) \left(\sin \phi' \cos \delta - \cos \phi' \sin \delta \cos t\right). \tag{13}$$

The values on the left hand sides of these equations are measured in radians.

Given that the latitudes of the New Mexico and California sites were almost the same, we have almost no leverage to use Eq. 13 to determine the distance to the asteroid. We limit ourselves to a consideration of the effect of parallax on the right ascensions. Let the New Mexico site be "position 1" and the California site be "position 2". Consider the seconds part of the observed right ascension of the asteroid. There exist corrections to the right

¹³ http://archive.stsci.edu/cgi-bin/dss_form

¹⁴IRAF is distributed by the National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation (NSF).

ascension such that $7.77 - c_2 = 7.42 - c_1$. The corrections c_i adjust the observed right ascensions to what would be observed by a hypothetical observer at the center of the Earth. So, 0.35 seconds of time equals the difference of the parallactic corrections. Since one second of time in right ascension equals $15 \cos \delta$ arc seconds, we have $5''.05 \pm 0.61 = c_2 - c_1$. The uncertainty comes from the square root of the sum of squares of the uncertainties of the pixel to RA/DEC transformations from **ccmap** in IRAF.

At the New Mexico site at the time Fig. 10 was taken the hour angle of the asteroid was -1.899 deg, while at the California site the hour angle of the asteroid was -14.176 deg when Fig. 11 was taken. Using Eq. 12 we obtain a distance to the asteroid of:

$$d = \frac{R_{\oplus}(0.210010 - 0.028541)}{\left(\frac{5.05 \pm 0.061}{206265}\right)} \,. \tag{14}$$

The reader will know that the number of arc seconds in a radian is 206265. If we adopt our value of the radius of the Earth from §3 (namely, 6290 km), we obtain a distance to the asteroid of $(4.66 \pm 0.56) \times 10^7$ km.

Using the Steeger orbit solution in Table 4, the method of Meeus (1988, p. 125 ff.), and the rectangular coordinates of the Sun on 2008 July 24 from the Astronomical Ephemeris, we determined that asteroid 1996 HW1 was 0.294 AU distant when Figs. 10 and 11 were taken. Our resulting value for the Astronomical Unit is $(1.59 \pm 0.19) \times 10^8$ km, which is roughly 6 percent larger than the true value of 1.496×10^8 km.

Our value of the solar parallax, P_S in Fig. 5, equals $(6290/1.59 \times 10^8) \times 206265 = 8''.2$, or about 0'.14, which we used in §4.1. Our distance to the Sun in Earth radii divided by our distance to the Moon in Earth radii gives a Sun distance that is roughly 406 times the distance to the Moon, which is close to the true mean value of 389.

6. Systematic and Random Errors

In the interests of readability we have not included all statistical details above. However, given the readership of the *American Journal of Physics*, we thought it prudent to have a discussion of various kinds of errors here.

Say you want to answer a simple question such as "How tall am I?" By standing up against the wall, placing a ruler level on the top of your head and making a mark on the wall you can find out the answer. But what if you forgot to take your shoes off? You would have a *systematic error* equal to the height of the heels of your shoes.

What if you measured your height immediately after getting out of bed after a full night's sleep? You would find that first thing in the morning you are about 2 cm taller than you will be later that same day. Why is this so? The disks between the vertebrae in your back expand when you are lying down sleeping. Thus, if you measured your height first thing in the morning, right after lunch, then after a 6 mile run in the afternoon, you could get three different answers. The "mean value of your height during the day" might be 178 \pm 1 cm. This standard deviation of \pm 1 cm is the "mean error of the mean" of the set of measurements. This is referred to as a random error because values that you measure would scatter about some average value.

Very often a set of measurements exhibits a Gaussian (i.e. bell-shaped) distribution. Then 68.3 percent of the measurements are within one standard deviation of the mean value, 95.5 percent are within 2- σ and 99.7 percent are within 3- σ . This allows us to identify *outliers* in the data. If some data value is several (or many) standard deviations from the mean (or expected) value, we have either underestimated the size of our random errors, or there is some unaccounted source of systematic error.

In §2 above we derived a value of the latitude for a particular location in College Station, Texas, which was 11'. 5 north of the true value. This is the systematic error. Usually in science we do not know the "true" value, but in this case we do. From a scatter of the measurements of the shadow length of our gnomon we could obtain a numerical value of the root-mean-square scatter of the data about some best fit line. That gives us an estimate of the uncertainty of the minimum shadow length, which translates into a random error for the maximum elevation angle of the Sun on some given day, which then leads to a random error for our value of the latitude. Our value of the latitude is within 3 of our standard deviations of the true value. Our value of the tilt of the Earth's axis of rotation to the plane of its orbit is 6 standard deviations from the true value. Thus, we have room for improvement in the experiment, should we choose to do it again and again.

Some sources of uncertainty have not yet been mentioned. For the location of the June 21st and December 21st gnomon experiments we have measured the levelness of the spot where the data were taken, finding that a perfectly straight gnomon which was perfectly squarely set in its base would have been tilted 6 ± 1 arc minute north of the zenith. This would have made the elevation angle too low by that amount at local noontime, but less so at other times. To complicate matters, our gnomon is just a wooden dowel rod which is not totally straight, and we do not know just how squarely it sits in the block of wood that is its base.

Regarding the use of the "map tool" in §3, we found that the great circle arc from South Bend, Indiana, to College Station, Texas, was 0.7437 ± 0.0009 of the route driven.

We have rounded this off to 0.744 and have decided to ignore the numerical uncertainty.

In §4.1 we found from six images of the Moon in partial lunar eclipse taken from 21:29:55 to 21:52:59 UT that the angular diameter of the Earth's shadow was 2.46, 2.65, 2.49, 2.62, 2.56, and 2.56 times the angular diameter of the Moon. These were obtained using hard copies of the images, a compass, and a ruler. In all instances we could see more than a 180 degree arc of the illuminated part of the Moon. The mean ratio was 2.56 ± 0.03 . To find angle τ (the angular radius of the Earth's shadow) we used the (more robust) mean value of the angular diameter of the Moon from Krisciunas (2010) rather than an estimated value of the Moon's angular size on 2011 June 15.

In §4.2 and Table 2 we give results of an experiment that is at or beyond the capabilities of a ruler plus cardboard cross piece. We found that the systematic error of the angular separation of two bright point sources (Saturn vs. α Vir) was +0.12 deg. The random error was ± 0.04 deg. These are considerably better than the often quoted uncertainty of 0.20 to 0.25 deg for non-telescopic measurements. But consider our measurements of the angular separation of Saturn vs. the Moon and α Vir vs. the Moon. Our mean systematic errors are as great as 0.39 deg and our random errors are \pm 0.23 to \pm 0.29 deg. Table 3 shows that if we have measured the Moon's position with respect to three or more point sources, we can locate the Moon's location on the sky to within 0.3 deg on average. But if we have measured the Moon's position with respect to only two point sources, the error in position can be too large to be useful for determining the rate of change of the Moon's position over the course of a night.

When it comes to certain kinds of analysis, there are other errors that are neither systematic or random. In the top half of Fig. 4 we have fitted a hyperbola to the full data set. But this is for illustrative purposes only. Why? Careful scrutiny shows that the points do not randomly scatter above and below the curved line. The actual function we need requires the latitude of the site, which is the thing we are trying to derive. In this case, for practical reasons we have fitted a fourth order polynomial to a subset of the data around the time of the minimum shadow length. The bottom line is that we fit some data under the assumption that we have the appropriate function. What the most appropriate function is also depends on the typical size of the error bars of the data points. If the error bars are large, high order polynomial fits would be unjustified.

7. Discussion and Conclusions

Using a vertical stick, a car, watch, map, and map tool, we have measured the size of the Earth from first principles. We used observations of the elevation angle of the Sun over the course of a few hours overlapping local noontime on the summer solstice and winter solstice of 2010 to determine the latitude and longitude of College Station, Texas. We also determined the latitude and longitude of South Bend, Indiana. The two sites were 13.78 deg apart along a great circle arc. We obtained a value of the radius of the Earth of 6290 km, about 1.4 percent too small.

Aristarchus's method of determining the distance to the Moon derives from the geometry of the Earth, Moon, and Sun at the time of a lunar eclipse. One key observable is the angular size of the Earth's shadow at the distance of the Moon compared to the angular size of the Moon. Using images available on the web of the lunar eclipse of 2011 June 15 we found that the Earth's shadow was 2.56 times the angular diameter of the Moon. The corresponding distance of the Moon was 62.3 Earth radii, about 3.3 percent larger than the true mean value.

It is also possible to determine the distance to the Moon from observations of its motion against the background of stars. These observations are made from a single site over the course of one night. This method is far more complicated than Aristarchus's method, but from such observations we showed that the Moon was between 47 and 62 R_{\oplus} distant. The true range of the Moon's distance is 55.9 to 63.8 R_{\oplus} .

Simultaneous observations from Socorro, New Mexico, and Ojai, California allowed us to determine the distance to the asteroid 1996 HW1 on 2008 July 24. A determination of the orbital elements of this asteroid by one of us allowed us to calculate that the asteroid was 0.294 AU distant on that date. The final result was a calibration of the AU equal to $(1.59 \pm 0.19) \times 10^8$ km, which is 6 percent larger than the true value.

Given the seeing and tracking constraints associated with our asteroid observations, our asteroid experiment was at the limits of the technology of our small telescopes. To determine the distance to a *main belt* asteroid (which would have been 10 times more distant) would have required a baseline 10 times bigger than the 1130 km baseline we used if we wanted the parallax to be several arc seconds. Such an experiment would be nearly impossible on the surface of the Earth.

This paper has overflowed with numerical and trigonometrical details. Still, the *qualitative* results are easy to understand and are worth restating. Using very inexpensive equipment we can determine the size of the Earth and the distance to the Moon. To determine the length of the AU we do *not* need to organize an international endeavor like what was done

for the Venus transits of 1761, 1769, 1874, and 1882 (Van Helden 1995). We can measure the length of the AU with carefully timed observations of a near Earth asteroid using telescopes comparable to those owned by many amateur astronomers.

It is also worth reminding ourselves why astronomers have such an obsession with determining cosmic distances. Thanks to Kepler's Third Law the length of the AU gets us the distances to all other objects that orbit the Sun. Once we have the distances to stars like the Sun (converting hydrogen to helium in their cores, but with a range of mass) via trigonometric stellar parallaxes – the next rung of the distance ladder – we can use photometric methods to determine distances to star clusters using the fact that "main sequence stars" of the same mass have comparable intrinsic brightness. A simple equation relates the apparent brightness of a star with its intrinsic brightness and the distance. We can exploit the Cepheid period-luminosity law to calibrate our way across the Galaxy and to nearby galaxies. Knowing the energy budget of a star helps us determine what it is made of, what is its structure, how it will live, and how it will die. Luminosities of Type Ia supernovae have allowed us to address some the largest questions we can ask, such as, "What is the ultimate fate of the Universe?"

The interconnectedness of astronomical topics means that the big questions are related to the Earth size experiment of Eratosthenes carried out more than 22 centuries ago. Starting with an instrument as simple as a vertical stick we can connect basic and profound aspects of the universe we inhabit.

Tom Weimar helped with the construction of the gnomon pictured in Fig. 2. We thank Nick Suntzeff for useful discussions. The NMT observations of asteroid 1996 HW1 were obtained with the help of Jeff Lu, David Oh, Anna Heinz, Peter Combs, and Rebecca Mickol. We thank Dan Klinglesmith and Jason Speights for observing support at Etscorn Observatory. The asteroid observations were made as part of the Summer Science Program, which is funded by private donations and sponsored by New Mexico Institute of Mining and Technology. Elisabeth Button kindly made Fig. 1.

REFERENCES

Bekeris, V., et al. 2011, Astronomy Education Review, 10 (1), 010201

Evans, J. 1998, The History and Practice of Ancient Astronomy (New York & Oxford: Oxford Univ. Press)

- Gingerich, O., & Voelkel, J. R. 1998, J. for the History of Astronomy, 29, 1
- Krisciunas, K. 2010, Amer. J. Phys., 78, 834
- Meeus, J. 1988, Astronomical Formulae for Calculators, 4th ed. (Richmond, Virginia: Willmann-Bell)
- Riess, A. G., et al. 2010, ApJ, 730, 119
- Rowan-Robinson, M. 1985, The Cosmological Distance Ladder: distance and time in the universe (New York: W. H. Freeman)
- Smart, W. M. 1977, Textbook on Spherical Astronomy, 6th ed., revised by R. M. Green (Cambridge: Cambridge Univ. Press)
- Toomer, G. J. 1981, "Hipparchus," in Dictionary of Scientific Biography, C. G. Gillespie, ed. (New York: Scribner), vol. 15, 207
- Toomer, G. J. 1984, Ptolemy's Almagest (New York: Springer-Verlag)
- Van Helden, A. 1995, in Planetary Astronomy from the Renaissance to the Rise of Astrophysics, Part B: The eighteenth and nineteenth centuries, R. Taton and C. Wilson, eds. (Cambridge: Cambridge Univ. Press), 153

This preprint was prepared with the AAS IATEX macros v5.2.

Table 1. Gnomon Data^a

CDT (Jun 21)	L (mm)	CST (Dec 21)	L (mm)
11:25:00	330	11:07:45	844
11:35:00	300.5	11:15:00	829
11:46:30	268.5	11:25:00	807
11:55:00	245.5	11:35:15	787
12:06:17	217	11:49:20	770
12:15:04	196	11:59:20	761.5
12:25:02	172	12:10:00	756.5
12:35:01	151	12:15:00	754.5
12:45:06	131	12:20:00	754
12:55:00	112.5	12:25:15	754
13:00:10	104	12:30:00	755
13:05:00	99	12:36:00	756.5
13:10:47	92.5	12:40:38	759.5
13:15:00	88.5	12:45:18	761.5
13:20:00	86.5	12:50:00	765.5
13:25:00	86	12:55:43	771
13:30:10	86	13:00:00	775
13:37:20	92	13:05:20	782
13:45:00	101	13:10:06	788.5
13:49:30	106	13:20:00	805
13:56:00	117	13:30:00	824.5
14:04:30	132	13:40:00	848
14:15:00	154	13:50:00	875
14:25:00	175.5	14:01:20	910
14:38:12	206	14:10:50	946
14:45:00	224	14:20:06	987
14.55.00	247.5		
15:05:00	275		
15:15:00	303		
15:25:00	331.5		

 $^{\rm a}{\rm Gnomon}$ height was 632 \pm 1 mm for the 2010 June 21 observations, and 550 \pm 1 mm for the 2010 December 21 observations. Column 1 is Central Daylight Time. Columns 2 and 4 gives the shadow lengths in mm. Column 3 is Central Standard Time.

Table 2. Angular separations (2011 May 15)

UT	Object pair	$ ho_{obs}$	$ ho_{true}$	diff ^a
02:21:00	α Vir vs. Moon	3.°41	3°70	+0°29
02:24:00	Saturn vs. Moon	12.85	13.21	+0.36
02:28:00	δ Crv vs. Moon	10.71	10.85	+0.14
02:30:00	Saturn vs. α Vir	13.60	13.71	+0.11
04:04:00	α Vir vs. Moon	3.28	3.40	+0.12
04:06:40	Saturn vs. Moon	13.34	13.86	+0.52
04:08:00	δ Crv vs. Moon	11.33	11.34	+0.01
04:10:00	Saturn vs. α Vir	13.63	13.71	+0.08
05:54:00	α Vir vs. Moon	3.32	3.22	-0.10
05:56:00	Saturn vs. Moon	14.52	14.53	+0.01
05:58:30	δ Crv vs. Moon	11.77	11.91	+0.14
06:01:40	Saturn vs. α Vir	13.55	13.71	+0.16
06:04:00	γ Hya vs. Moon	9.33	9.31	-0.02
08:02:00	α Vir vs. Moon	3.63	3.15	-0.48
08:03:00	Saturn vs. Moon	14.80	15.34	+0.54
08:05:00	Saturn vs. α Vir	13.55	13.71	+0.16
08:31:00	Saturn vs. α Vir	13.60	13.71	+0.11
08:38:00	α Vir vs. Moon	3.35	3.17	-0.18
08:41:00	Saturn vs. Moon	15.06	15.60	+0.54

 $^{^{\}rm a}{\rm In}$ the sense "true" minus "observed".

Table 3. Derived and true topocentric positions of Moon (2011 May 15)

UT	α_{obs}	δ_{obs}	α_{true}	δ_{true}	$ ho^{ m a}$
02:24:00	$198.^{\circ}15 \pm 0.09$	$-12.^{\circ}76 \pm 0.09$	198°27	-13.°20	0.°46
04:04:40	198.76 ± 0.04	-13.14 ± 0.07	198.93	-13.56	0.45
05:56:00	199.53 ± 0.02	-13.94 ± 0.01	199.63	-13.90	0.10
08:03:00	199.50 ± 0.57	-14.31 ± 0.36	200.55	-14.25	1.02
08:41:00	200.03 ± 0.62	-14.27 ± 0.34	200.87	-14.34	0.81

^aAngular distance between derived position of Moon based on observations with cross staff vs. true topocentric position of Moon. For the fourth and fifth determinations the uncertainties in the right ascension and declination derive from the assumption that the angular separations of the Moon vs. α Vir and the Moon vs. Saturn were accurate to \pm 0.3 deg.

Table 4. Steeger orbit determination for asteroid 1996 HW1

Parameter	Description	Steeger value	JPL Horizons value
$egin{array}{c} egin{array}{c} egin{array}{c} egin{array}{c} egin{array}{c} egin{array}{c} B & eta & eta \\ A & eta & eta \\ C & eta & eta \end{array}$	epoch (UT) epoch (Julian Date) mean anomaly at t ₀ semi-major axis eccentricity inclination angle longitude of ascending node	2008 June 29.35 2454646.85 335.°1461 2.0855 AU 0.4575 8.°5033 177.°6887	2007 January 15.00 2454115.5 155.°33678 ^a 2.046041 AU 0.449165 8.°437363 177.°216737
ω	argument of perihelion	176°1618	177°020070

 $^{^{\}mathrm{a}}$ The mean anomaly increases by 0.33676929 deg per day according to JPL Horizons, so it is 334.°2791 at the epoch of the Steeger solution.


- Fig. 1.— The astronomical triangle. The North Celestial Pole is labeled "NCP". The zenith is at Z. The latitude of the site is ϕ . An object in the western sky is marked by an asterisk. The hour angle of the object is t, which is positive in the western sky, negative in the eastern sky. The declination of the object is δ . Its elevation angle above the horizon is h, so the zenith angle is $90^{\circ}-h$. The azimuth A of an object is measured clockwise around the horizon, with north = 0° .
- Fig. 2.— A gnomon. It consists of a wooden base with a hole drilled through it using a drill press, and a vertical stick that fits tightly. It can be a pointed stick. Here we have fashioned a small sphere at the top. It is easier to measure the center of the elliptical shadow of the sphere than the end of the darker part of the shadow of a pointed stick.
- Fig. 3.— The X-Y positions of the end of the shadow of a 632 mm high gnomon used in College Station, Texas. For the 2010 December 21 observations we have scaled the coordinates by 632/550, as a 550 mm high gnomon was used on that date. The curvature of these loci change with the declination of the Sun over the course of the year. In fact, the shadow lengths and the X-positions, along with the hour angle of the Sun obtained from the times of the observations and the time of minimum shadow length, allow us to derive the declination of the Sun. In that case it is best to use observations obtained when the Sun is roughly one hour or more from the meridian.
- Fig. 4.— Upper figure: the shadow length of a 632 mm gnomon on the day of the summer solstice, as measured in College Station, Texas. For illustrative purposes *only* a hyperbola is fit to the data. Lower figure: the shadow length of a 550 mm gnomon on the day of the winter solstice, as measured at the exact same location. A fourth order polynomial is fit to the data.
- Fig. 5.— Geometry of the lunar eclipse (not to scale). At point G the Moon is halfway into the shadow of the Earth. At point H the Moon is halfway out of the shadow. Simple arguments and measurements originating with Aristarchus allow us to estimate the distance to the Moon in terms of the Earth's radius.
- Fig. 6.— The angular motion of the Moon as a function of hour angle for an observer in College Station, Texas.
- Fig. 7.— The cross staff. The cardboard cross piece slides up and down the yardstick. Using simple geometry we can use this device to determine the angular separation of objects in the sky.
- Fig. 8.— The topocentric and geocentric RA and DEC of the Moon on 2011 May 15. The

topocentric values are calculated for College Station, Texas. The zenith angle of the Moon is greater for an observer situated on the surface of the Earth compared to a hypothetical observer at the center of the Earth. In other words, by observing on the surface of the Earth, the Moon appears to be lower in the sky compared to what would be seen by a hypothetical observer at the center of the Earth. This elevation angle offset translates to varying shifts in RA and DEC over the course of the night.


Fig. 9.— Dashed curve: angular motion of an Earth-orbiting object as viewed by a hypothetical observer at the Earth's center, as a function of its distance in R_{\oplus} . Solid curve: mean angular motion of an Earth-orbiting object as viewed from College Station, Texas. The average is taken over an 8 hour period centered on the meridian transit. On 2010 October 21 we measured the Moon to move 0.40 ± 0.03 deg/hr from six observations over 9.1 hours. The implied value of the Moon's distance is between 57 and 62 R_{\oplus} on that occasion. On 2011 May 15 we measured the Moon to move 0.508 ± 0.094 deg/hr from three sets of observations taken over 3.9 hours. The implied distance to the Moon is roughly 47 to 58 R_{\oplus} .


Fig. 10.— Asteroid 1996 HW1 is circled in this unfiltered 90 second image obtained by E. DeBenedictis with a 15 cm Takahashi refractor at Etscorn Observatory, Socorro, New Mexico, on 2008 July 24 at 08:17:27.8 UT. North is up, east is to the left.

Fig. 11.— Asteroid 1996 HW1 is circled in this unfiltered 90 second image obtained by G. Tabak and K. Pasricha with a 25 cm Meade reflector at Besant Hill School in Ojai, California, on 2008 July 24 at 08:17:28 UT, with an uncertainty of no more than \pm 2 seconds. North is up, east to the left. Note that this image was taken at the same time as Fig. 10, but the asteroid is roughly 5 arc seconds to the left (east) as observed at the other end of a baseline of 1130 km.




Krisciunas $et\ al.$ Fig. 1







Krisciunas et al. Fig. 3




Krisciunas et al. Fig. 4




Krisciunas $et\ al.$ Fig. 5




Krisciunas et al. Fig. 6







Krisciunas et al. Fig. 8



Krisciunas $et\ al.$ Fig. 9



Krisciunas $et\ al.$ Fig. 10



Krisciunas et al. Fig. 11