B. R. Dennis¹, A. G. Emslie², and H. S. Hudson³

the date of receipt and acceptance should be inserted later

Abstract In this introductory chapter, we provide a brief summary of the successes and remaining challenges in understanding the solar flare phenomenon and its attendant implications for particle acceleration mechanisms in astrophysical plasmas. We also provide a brief overview of the contents of the other chapters in this volume, with particular reference to the well-observed flare of 2002 July 23.

Keywords Sun: flares; Sun: X-rays; Sun: acceleration; Sun: energetic particles

Contents

1	Historical Perspective			
2	Review of Flare Models			
3	Challenges for Simple Acceleration Models			
4	Importance of Hard X-Rays and γ -Rays as Diagnostics of Accelerated Particles			
5	RHESSI Design and Capabilities			
	5.1 Operations			
	5.2 Collaborations			
6	Outline of this Volume			
Ac	cronyms			

1 Historical Perspective

This volume of *Space Science Reviews* contains a comprehensive review of our current understanding of the high energy aspects of solar flares. It is written with the same philosophy as the book on solar flares (Sturrock 1980b) that grew out of the *Skylab* workshops. The nine chapters are intended to display the accumulated wisdom of the many scientists who have attended the ten *RHESSI* (*Reuven Ramaty High Energy spectroscopic Imager*) science workshops to date. *RHESSI* is a NASA Small Explorer satellite launched in February 2002. The intent was to cover the relevant published literature into 2010, and this succeeded as illustrated in Figure 1. Results are summarized from hard X-ray and γ -ray observations in

¹Code 671, NASA Goddard Space Flight Center, Greenbelt, MD 20771 E-mail: brian.r.dennis@nasa.gov

²Western Kentucky University, Bowling Green, KY 42101E-mail: gordon.emslie@wku.edu

³Space Sciences Laboratory, UC Berkeley, Berkeley, CA 94720 E-mail: hhudson@ssl.berkeley.edu




Fig. 1.1 The distribution by year of the 1,619 (non-unique) citations in Chapters 1-8 back to 1960; the 20 citations prior to that included papers by Alfvén, Carrington, and Giovanelli among other pioneers. The peak in 2002 coincides with the publication of the initial *RHESSI* results and the peak at around 1995 may be recognizable as *Yohkoh* results.

Solar Cycle 23 and the complementary observations at other wavelengths that have provided information on the same flares and the often-associated coronal mass ejections (CMEs). We anticipate that this volume will be a comprehensive reference of our current state of knowledge and relevant published literature up to the onset of renewed solar activity in Hale Cycle 24.

Although great progress has been made in understanding the solar flare phenomenon since the *Skylab* report, the basic concepts were well established at that time from the extensive ground-based observations and early space missions. As summarized by Sturrock (1980a), the basic picture of a flare involved the sudden explosive release of the "free" magnetic energy of a current-carrying magnetic field in the corona. During the flare, the energy would be released by altering (or even destroying) the currents to convert the field to a lower-energy (or even current-free) form. Various energy release mechanisms were considered, including magnetic reconnection, but then, as now, it was recognized that the plasma processes are very complicated and no definitive conclusions could be made on which specific processes were involved.

The acceleration of particles was considered as posing a primary requirement for any flare model. It was recognized though that there was a serious electron "number problem," in that the number of electrons required to explain the measured hard X-ray fluxes was a substantial fraction of the electron content of the corresponding coronal region before the flare. Also, the total power in the 10-100 keV electrons (some $2\times 10^{29}~{\rm erg~s^{-1}}$ in SOL1972-08-04T06:25 or SOL1972-08-04T06:25)¹ and the total energy contained in these electrons assuming thick-target interactions was known to be an unexpectedly large fraction of the total flare energy (Ramaty et al. 1980). Indeed, Lin & Hudson (1976) had shown that the $\sim\!10$ to 100 keV electrons "constitute the bulk of the flare energy," perhaps as high as 10 to

¹ In this volume we identify individual flares using the IAU naming convention; see DOI 10.1007/s11207-010-9553-0

50%. This issue remains one of the key problems in understanding the mechanism of energy release in solar flares as described in chapters (3, 7, and 8) of this volume (Holman et al. 2011; Kontar et al. 2011; Zharkova et al. 2011).

The thermal or nonthermal origin of the hard X-ray emission was an ongoing debate in the *Skylab* workshops (Sturrock 1980b). This debate is still not fully resolved (White et al. 2011) but strong support for the nonthermal electron-beam model came with the hard X-ray imaging of the *Solar Maximum Mission (SMM)*, *Hinotori*, and *Yohkoh* satellites. The Hard X-ray Telescope (HXT) on *Yohkoh* produced images of a great many flares (Sato et al. 1999), with the advantage of simultaneous microwave observations (e.g., Bastian et al. 1998; Hannah et al. 2011). The X-ray images revealed the prevalence of double footpoint sources and the near-simultaneity of their light curves to within a second (Sakao et al. 1996). The stronger X-ray source and the weaker radio source both tend to be located in the weaker magnetic field region, consistent with an electron beam model where magnetic mirroring is significant (e.g., Bastian et al. 1998).

The acceleration of ions was also recognized from the γ -ray observations on OSO-3 (Chupp et al. 1973) and HEAO-1 (Hudson et al. 1980) but their numbers and energy content were not well understood. It required observations made with the Gamma Ray Spectrometer (Forrest et al. 1980) on SMM in the 1980s to reveal that the ions could be accelerated nearly simultaneously with the electrons (Forrest & Chupp 1983) and that the total energy in ions above 1 MeV/nucleon could be comparable to, or even exceed, the total energy in electrons above 20 keV (Ramaty et al. 1995). Similar to electrons (Holman et al. 2011), the total energy content in ions depends to a significant extent on an accurate determination of the low-energy end of the accelerated spectrum, and considerable progress has been made on this front (Vilmer et al. 2011).

The biggest discrepancy between our current understanding of the flare phenomenon and the *Skylab* ideas is in the relationship between flares and coronal mass ejections (CMEs). In the *Skylab* era, CMEs were referred to as coronal transients, and it could still be questioned if they were "incidental to flares or whether they reveal something fundamental about the energy release process" (Rust et al. 1980). It was not until the "solar flare myth" was described by Gosling (1993) that many realized that the CMEs are the major cause of solar effects at the Earth, not the flares as such. This was controversial (Hudson et al. 1995), and with modern data we now understand that flares and energetic CMEs are, in fact, intimately related and may have comparable energy content (Emslie et al. 2004a, 2005) in major events. It is now clear that the largest and fastest CMEs, which have the greatest effect on space weather and pose the greatest danger to satellites, are mostly (possibly always) associated with large flares of comparable total energy. Indeed, the origins and energy sources of flares and CMEs are so intimately entwined that it is impossible to explain one without understanding the other. Thus, if we are to understand these phenomena and develop predictive capabilities, it is imperative that we still consider them as interrelated phenomena.

2 Review of Flare Models

It was established very early on in flare studies that the source of the energy released in a flare is in current-carrying (i.e., "non-potential") magnetic fields. Not only are flares invariably connected with magnetism in the photosphere, but an examination of the various candidate sources of energy reveals magnetic energy to be the only plausible contender (Tandberg-Hanssen & Emslie 1988). Release of the energy stored in the twisted magnetic field configuration can proceed through a process termed *magnetic reconnection*, in which

the connectivity of the magnetic field redefines itself. However, it is far less clear – and this is what to a large extent drives flare research – how the energy can be built up *without* reconnection dissipating it immediately, and what "trigger" initiates the flare process. The energy can build up for many hours, or even days, without significant dissipation of the energy prior to the flare onset.

Sturrock (1980a) provided a concise, yet thorough, review of the various reconnection scenarios that existed through the end of the 1970s. In fairness, it must be conceded that most of these scenarios are still valid, although many have fallen out of favor. Perhaps the most significant "casualty" is the model of Gold & Hoyle (1960), which invoked the self-attraction of magnetic loops carrying parallel, or near-parallel, currents, leading to an energy release at their mutual interface. While this laboratory analogy is rather appealing, it unfortunately fails to recognize that it is not currents that attract *per se*, but rather one current interacts with the magnetic field produced by the other. Thus, in the global force-free field appropriate to the low- β solar corona, the current density $\bf J$ is always nearly parallel to the local magnetic field $\bf B$, and no $\bf J \times \bf B$ force exists. Independent magnetic flux loops therefore have no particular attraction to (or repulsion from) each other, and some external influence (such as a photospheric velocity field; e.g. Heyvaerts et al., 1977) must be postulated in order to drive them together.

Basic Sweet-Parker reconnection (e.g., Sweet 1969), has also fallen out of favor because it is perceived to develop too slowly. In its place Petschek reconnection, which is both much faster and is characterized by energy release at reconnection-driven shocks (Petschek 1964) rather than at the point of reconnection, has appeared in many descriptions. The standard "CSHKP" (Carmichael 1964; Sturrock 1966; Hirayama 1974; Kopp & Pneuman 1976) model, suggested by the growth of soft X-ray "loop prominence systems" accompanying the increasing separation of $H\alpha$ "ribbons," invokes a reconnection site near the apex of the loop system. This injects energy (e.g., as flows) into the underlying loops. Although this model does not naturally account for the impulsive phase of a flare, nor the particle acceleration, a great deal of effort has been expended in quantitative modeling of such structures (density of loop-top source, standing shocks in loop legs, etc.). Aspects of these problems are discussed in Zharkova et al. (2011).

The planning of the *RHESSI* workshop series addressed the need to consider the physics of magnetic reconnection, and the concomitant acceleration of electrons and ions, *in the context of the observations*. For example, it was recognized early on that "test-particle" approaches to particle acceleration, in which the electric and magnetic fields are *prescribed* and constant in time, do not adequately take into account the fact that the mass, momentum, energy, and electrical current carried by the large number of accelerated particles necessary to account for flare observations must have a major feedback on both the electrodynamic and magnetohydrodynamic environments of the acceleration region. A major emphasis of the theory team (Zharkova et al. 2011) at the *RHESSI* workshops was the development of acceleration models that explicitly take these nonlinear aspects of the process into account.

3 Challenges for Simple Acceleration Models

The impulsive phase of a solar flare is characterized, in part, by the emission of a copious flux of hard X-rays (photon energy $\varepsilon \gtrsim 10$ keV). It is generally accepted that these hard X-rays are produced by collisional bremsstrahlung (free-free emission) when accelerated electrons encounter ambient protons and heavier ions in the solar atmosphere, although other emission mechanisms have also been considered (Kontar et al. 2011). The amount of electron

energy required to produce these hard X-rays depends on the model used to characterize the interaction of the accelerated electrons with the target. A lower limit is given by a *collisional thick-target* interpretation (e.g., Brown 1971), in which all of the electron energy is absorbed in the target only through Coulomb collisions with ambient particles (primarily electrons). In this interpretation, the ratio of electron power to emitted hard X-ray power is of order 10^5 , and this gives an order-of-magnitude estimate for the rate of electron acceleration in the flare (Holman et al. 2011). For a large (e.g., *GOES* X-class) flare, the required rate of acceleration of electrons above 20 keV can exceed 10^{37} s⁻¹, a large number with interesting consequences and difficult problems (e.g., Miller et al. 1997).

First, the number of electrons \mathcal{N} confined in the coronal portion of a flare loop is simply the number density n (cm⁻³) multiplied by the volume V (cm³). Inserting typical values of $n\approx 10^{11}$ and $V\approx 10^{27}$ gives $\mathcal{N}\approx 10^{38}$, so that the acceleration process would deplete the store of available electrons in 10 s or so, significantly less than the observed duration of the flare. This simple calculation therefore tends to rule out models in which all the electrons to be accelerated are stored in a coronal volume prior to the onset of the flare. If the acceleration site is indeed in the corona, this requires instead that the electrons "recycle" multiple times or that fresh electrons enter from the chromosphere. Given that electrons in the solar atmosphere have gyroradii of order a centimeter and so are strongly tied to the guiding magnetic field lines, this presents formidable difficulties for current closure of the accelerated electron streams. Although solutions to this problem have been offered (Emslie & Henoux 1995), they require the synergistic interaction of a large number ($\sim 10^{10}$ or so) of separate acceleration regions, and the origin (not to mention stability) of such a system has yet to be adequately explored.

Second, the accelerated electron *number* carries with it an associated *electrical current* of some 10^{18} A (3×10^{27} statamps). In steady-state, such a current, if assumed to propagate unidirectionally in a flux tube of radius 10^9 cm, gives rise, via Ampère's Law, to a magnetic field $B\approx2\times10^8$ Gauss). Not only is such a high magnetic field completely untenable on observational grounds, the associated energy density $B^2/8\pi\approx10^{15}$ erg cm $^{-3}$, for a total energy content $\int (B^2/8\pi)\,dV$ of some 10^{42} ergs, some ten orders of magnitude larger than the energy content in a 1000-second duration beam.

Third, a current of this magnitude cannot appear instantaneously. The self-inductance $\mathscr L$ of a structure scales with ℓ , the characteristic dimension. For a solar flare loop of typical dimensions, we find $\mathscr L\approx 10$ H. Hence, to initiate a current $I\approx 10^{18}$ A in a time $\tau\approx 10$ s requires a voltage $V\approx \mathscr LI/\tau\approx 10^{18}$ V, which is *fourteen* orders of magnitude higher than the typical energy of the accelerated electrons.

Such considerations have led various authors (e.g, Knight & Sturrock 1977; Emslie 1980; Brown & Bingham 1984; Spicer & Sudan 1984; Larosa & Emslie 1989; van den Oord 1990; Zharkova et al. 1995) to consider models in which cospatial return currents locally neutralize the beam current. However, such models can only produce beam-current neutralization in the *propagation* region, and considerable analysis has been performed on the details of the beam/return current interaction in this propagation region (van Oss & van den Oord 1995). In the acceleration region itself, return-current electrons would have to flow in a direction counter to the applied electromotive force and so cannot neutralize the unacceptably large currents therein. Although significant progress on this issue has been made recently (Zharkova et al. 2011), a satisfactory resolution of this issue has yet to be offered. A self-consistent electrodynamic theory would require a description of current closure in the acceleration region as well as in the beam itself. This difficulty has led various authors to reject acceleration models featuring large-scale electric fields in favor either of acceleration by

very large electric fields in localized current sheets, or via acceleration in (stochastic) MHD or plasma waves.

The problems imposed by the return current become more severe as the beam becomes more intense. It has become increasingly clear directly from *RHESSI* imaging (e.g., Dennis & Pernak 2009) that the hypothetical electron beam would occupy only a small area. Closely related emissions such as UV and white light are often unresolved at even higher resolution and suggest areas substantially smaller than 10^{16} cm² (Fletcher et al. 2011). Accordingly, alternatives that replace electron-beam energy transport with Poynting fluxes have been proposed (Emslie & Sturrock 1982; Haerendel 2006; Fletcher & Hudson 2008; Haerendel 2009).

4 Importance of Hard X-Rays and γ-Rays as Diagnostics of Accelerated Particles

It is important to emphasize that the energy released as hard X-rays and γ -rays is – in and of itself – a negligible component of that released in the flare. The importance of this radiation lies not in its energy content *per se*, but rather in the energy in accelerated particles required to produce this diagnostic radiation (Holman et al. 2011; Vilmer et al. 2011).

The process of hard X-ray emission is very inefficient. In order to produce a photon by bremsstrahlung, an electron must suffer a near-direct collision on an ambient ion. Most electrons instead lose their energy in a large number of small-angle scatterings off ambient electrons, and do not contribute to the bremsstrahlung yield. We may compare the energy emitted through bremsstrahlung to that suffered in Coulomb collisions by comparing the cross-sections for the two processes. The nonrelativistic differential cross-section (cm² per unit photon energy) for free-free emission of a photon of energy ε by an electron of energy ε may be, to order-of-magnitude, approximated by the Kramers form

$$\sigma(\varepsilon, E) \approx \alpha \frac{r_o^2 mc^2}{\varepsilon E},$$
 (4.1)

where $\alpha \approx 1/137$ is the fine structure constant, $mc^2 = 511$ keV is the electron rest mass, and r_0 is the classical radius of the electron. From this it follows that the cross-section (cm² keV) for energy loss through bremsstrahlung by an electron of energy E is

$$\sigma_E^B = \int_0^E \varepsilon \sigma(\varepsilon, E) dE \approx \alpha r_o^2 mc^2. \tag{4.2}$$

By contrast, the cross-section for Coulomb energy loss ($cm^2 \text{ keV}$) by an electron of energy E is (Brown 1972; Emslie 1978)

$$\sigma_E^C(\varepsilon, E) = \frac{2\pi e^4 \Lambda}{E} \approx \Lambda \frac{r_o^2 (mc^2)^2}{E}, \tag{4.3}$$

where Λ is the Coulomb logarithm, ε is the electron charge, and we have used $r_o = e^2/mc^2$. Taking the ratio

$$\eta = \frac{\sigma_E^B}{\sigma_F^C} \approx \frac{\alpha}{\Lambda} \frac{E}{mc^2} \approx 4 \times 10^{-4} \frac{E}{mc^2}$$
 (4.4)

gives the energetic efficiency of the bremsstrahlung process relative to Coulomb collisions. For $E\approx 20$ keV, $\eta\approx 1.5\times 10^{-5}$, i.e., for each erg of bremsstrahlung, $\gtrsim 10^5$ ergs of energy in electrons are required.

Gamma-rays in solar flares (Vilmer et al. 2011) are produced principally by the interaction of accelerated protons and heavier ions with nuclei in the ambient atmosphere, although electron-ion bremsstrahlung from accelerated electrons can also contribute. Unlike hard X-ray emission, not all γ -ray emission is prompt – in particular, the capture of neutrons onto ambient protons to create the 2.223 MeV deuterium-formation line can take several minutes because of the need to reduce the momentum of the neutrons to a value where the cross-section for recombination is sufficiently high. The low speed of the deuterium atoms also leads to the extremely small spectral width of the 2.223 MeV line; by contrast, most γ -ray lines (e.g., the prompt nuclear de-excitation lines of 12 C at 4.4 MeV and 16 O at 6.1 MeV) have both narrow and broad spectral profiles depending on whether the heavy ions are the target or the projectile in the interactions with accelerated or ambient protons, respectively.

RHESSI not only provides γ -ray spectra with unprecedented spectral resolution (Smith et al. 2003) but also, on a few occasions, *images* of the γ -ray line emission – in particular in the 2.223 MeV line. Interestingly, the locations of the hard X-ray and γ -ray sources are *not* coincident (Hurford et al. 2003, 2006), indicating a preferential acceleration of electrons vs. protons in different substructures within the flare volume (cf. Emslie et al. 2004b).

Some time after the publication of the *Skylab* volume, studies of observations from the *SMM* Gamma-Ray Spectrometer (GRS) (Share & Murphy 1995) led Ramaty et al. (1995) to realize that the spectra of the accelerated ions could remain steep down to energies as low as 1 MeV, and hence that the *energy content* of accelerated ions in solar flares, as revealed by their γ -ray emission, could rival that of the accelerated electrons, hitherto thought to dominate the energy budget of accelerated particles. Further study of the relative partitioning of energy between accelerated electrons and ions was carried out by Emslie et al. (2004a, 2005), who reached a similar conclusion. Further discussion of the partitioning of flare energy amongst its constituent parts can be found in Fletcher et al. (2011), Holman et al. (2011), and Vilmer et al. (2011).

5 RHESSI Design and Capabilities

5.1 Operations

RHESSI (Lin et al. 2002) uses nine cooled and segmented germanium detectors to achieve high-resolution X-ray and γ -ray spectroscopy across the full energy range from 3 keV to 17 MeV (Smith et al. 2002). The FWHM energy resolution increases from \sim 1 keV at the lowest energies to $\sim 10 \text{ keV}$ at the highest. This has proved adequate to detect the iron-line complex at ~6.7 keV, measure the steep hard X-ray continuum spectra with relative flux accuracies as fine as 1%, measure the width of the positron-annihilation line at 511 keV as it varies with time during a flare, and resolve all of the narrow nuclear γ -ray lines except for the intrinsically narrow neutron-capture line at 2.223 MeV. A bi-grid tungsten collimator over each detector modulates the incident photon flux as the spacecraft rotates at \sim 15 rpm to provide the temporal information needed for the Fourier-transform technique that is used to reconstruct the X-ray and γ -ray images (Hurford et al. 2002). Imaging is possible at all energies up to about 1 MeV, with an angular resolution of $\sim 2''$ (FWHM) up to ~ 100 keV increasing to $\sim 20''$ at 1 MeV. When the count rates are sufficiently high, images can be made with a cadence as short as 4 s. In addition, images can also be made in the neutroncapture γ -ray line at 2.223 MeV in the relatively few flares when the total number of photons detected in this line is sufficiently high (several thousand; Hurford et al. 2006). The field of view is $\sim 1^{\circ}$ such that a flare can be imaged no matter where on the visible disk it occurs.

RHESSI has an effective sensitive area that reaches \sim 60 cm² at 100 keV. Two thin aluminum disks can be automatically moved above each detector to attenuate intense soft X-ray fluxes so that *RHESSI* can operate with minimal detector saturation and pulse pile-up over a wide dynamic range in flux level. This allows coverage of both faint microflares (with the attenuators removed) and the most powerful flares (with both attenuators in place over each detector).

RHESSI was launched on 2002 February 5 and has been in operation almost continuously since 2002 February 11, with brief intervals of pointing away from the Sun for observations of the Crab Nebula and solar global emission (the quiet Sun as a star). The *GOES* class X4.8 flare SOL2002-07-23T00:35 yielded the first γ -ray emission lines detected by *RHESSI* (see Shih et al. 2009) for a summary of all γ -ray events seen with *RHESSI* as of the Cycle 23/Cycle 24 solar minimum). By the time of writing, there have been two successful anneals of the germanium detectors, in November 2007 and again in April 2010. These are month-long procedures needed to restore sensitive volume and energy resolution that become degraded by the accumulation of radiation damage.

5.2 Collaborations

r Because solar flares and other forms of activity are defined by particle acceleration and extreme heating, RHESSI observations are at the heart of many broader studies. Instruments on other spacecraft and at ground-based observatories around the world have been active participants in providing the magnetic, thermal, and dynamic context in which the X-ray and γ -ray sources are produced. In addition, microwave observations of the gyrosynchrotron emission provide additional information on the accelerated electrons themselves. Coronagraph observations of CMEs, and in situ particle-and-field measurements in the near-Earth environment, also provide information that can be used to establish the links between these related phenomena and any associated flares. A partial list of all collaborating spacebased observatories with further information can be found on the Max Millennium Web site at http://solar.physics.montana.edu/max_millennium/obs/SBO.html. They include the ACE, Cluster, CORONAS, GOES, INTEGRAL, SOHO, TRACE, and WIND spacecraft. More recently, complementary observations have been made with the newer solar missions including STEREO. They have provided X-ray, EUV, UV, optical, and in situ particle-andfield measurements relevant to the many events recorded by RHESSI. Thanks to the daily email messages and the coordinating efforts of the Max Millennium program, many collaborative observing campaigns have been conducted to maximize the overlap of the various observatory programs.

6 Outline of this Volume

The contents of this volume center on *RHESSI* capabilities and research flowing from the *RHESSI* data, but this embraces most of flare physics because of the dominating importance of the high-energy processes. The current volume consists of a series of articles, each representing the work of multiple authors. The purpose of each article is to present a review of the pertinent subject matter, linking the results of a variety of published works into a coherent whole, which we hope is useful both for the reader who wishes an overview of contemporary knowledge in the area, and for the experienced researcher to view results in context. Each article presents mid-length reviews of the literature, and provides a comprehensive reference

list for the reader who seeks more detailed developments or information. A composite index appears at the back of this volume.

Taken as a whole, the Editors hope that this volume will be a useful successor for the *Skylab* workshop volume (Sturrock 1980b), and we hope that Zharkova et al. (2011) in particular will play the same role for solar-flare particle acceleration as reviewed earlier by Miller et al. (1997).

Although each of the articles presents a somewhat different aspect of solar flare research, they are all inter-related and should be read in this context. To illustrate the inter-relationship of these articles, we note that much attention has been paid to the characteristics of the first γ -ray line flare detected with *RHESSI* mentioned above (SOL2002-07-23T00:35) This event forms the basis for much of the discussion in each article, namely:

- The temporal, spatial and spectral properties of the intense hard X-ray radiation from this flare are discussed in Kontar et al. (2011); this includes a discussion of the temporal evolution of the hard X-ray spectrum, both for the flare as a whole and for subregions (e.g., coronal sources, chromospheric footpoints) observed within the active region. Using an appropriate cross-section for hard X-ray production, regularized spectral inversion of the observed hard X-ray spectrum then yields the volume-averaged mean source electron spectrum for the event.
- A useful check on the electron spectrum comes from observations of deka-GHz radio emission (White et al. 2011); this radio emission is believed to be produced by the highenergy tail of the same ensemble of electrons that produces the deka-keV hard X-ray emission, but the inferred electron spectra are intriguingly different.
- As discussed in Holman et al. (2011), combining the mean source electron spectrum with an appropriate electron transport model then leads to the *accelerated* electron spectrum. Analysis of this accelerated spectrum (with particular attention to the low-energy end where, due to the typically steep spectra involved, most of the particle energy resides) then yields information on the total energy in the accelerated electrons.
- Vilmer et al. (2011) discuss various aspects of the γ -ray emission from this event. It includes a comparison of the time profiles and spatial locations of hard X-ray and γ -ray sources. The intensity and Doppler shifts of the γ -ray lines are also presented along with the information that these measurements provide on the number and angular distribution of accelerated ions, and even on the magnetic field geometry in the active region.
- Results from all photon energy ranges (hard X-ray, soft X-ray, γ -ray, optical and EUV) are synthesized into a global picture of the energetics of this flare in Fletcher et al. (2011).
- The position of this flare within the statistical ensemble of all flare events detected with *RHESSI*, extending from B-class microflares to large X-class events, is provided in Hannah et al. (2011).
- Zharkova et al. (2011) review the implications of these observational results for theoretical models of particle acceleration and transport in flare plasmas, and in the broader field of acceleration in astrophysical sources.
- Lin (2011) summarizes these results and presents prospects for future research directions.

Acknowledgements H. Hudson was supported by NASA under contract NAS5-98033 for RHESSI.

References

- T. S. Bastian, A. O. Benz, D. E. Gary, Annual Rev Astron. Astrophys. 36, 131 (1998), doi:10.1146/annurev.astro.36.1.131
- J. C. Brown, Solar Phys. 18, 489 (1971)
- J. C. Brown, R. Bingham, Astron. Astrophys. 131, L11 (1984)
- H. Carmichael, NASA Special Publication 50, 451 (1964)
- E. L. Chupp, D. J. Forrest, P. R. Higbie, A. N. Suri, C. Tsai, P. P. Dunphy, Nature 241, 333 (1973)
- B. R. Dennis, R. L. Pernak, Astrophys. J. 698, 2131 (2009), doi:10.1088/0004-637X/698/2/2131
- A. G. Emslie, Astrophys. J. 235, 1055 (1980), doi:10.1086/157709
- A. G. Emslie, B. R. Dennis, G. D. Holman, H. S. Hudson, Journal of Geophysical Research (Space Physics) 110, 11103 (2005), doi:10.1029/2005JA011305
- A. G. Emslie, J.-C. Henoux, Astrophys. J. 446, 371 (1995), doi:10.1086/175796
- A. G. Emslie, H. Kucharek, B. R. Dennis, N. Gopalswamy, G. D. Holman, G. H. Share, A. Vourlidas, T. G. Forbes, P. T. Gallagher, G. M. Mason, T. R. Metcalf, R. A. Mewaldt, R. J. Murphy, R. A. Schwartz, T. H. Zurbuchen, Journal of Geophysical Research (Space Physics) 109, 10104 (2004a), doi:10.1029/2004JA010571
- A. G. Emslie, J. A. Miller, J. C. Brown, Astrophys. J. Lett. 602, L69 (2004b), doi:10.1086/382350
- A. G. Emslie, P. A. Sturrock, Solar Phys. 80, 99 (1982), doi:10.1007/BF00153426
- L. Fletcher, H. S. Hudson, Astrophys. J. 675, 1645 (2008), doi:10.1086/527044
- L. Fletcher et al., Space Sci. Rev. pp. XXX-XXX (2011)
- D. J. Forrest, E. L. Chupp, Nature 305, 291 (1983)
- D. J. Forrest, E. L. Chupp, J. M. Ryan, M. L. Cherry, I. U. Gleske, C. Reppin, K. Pinkau, E. Rieger, G. Kanbach, R. L. Kinzer, Solar Phys. 65, 15 (1980)
- T. Gold, F. Hoyle, Mon. Not. Roy. Astron. Soc. 120, 89 (1960)
- J. T. Gosling, J. Geophys. Res. 98, 18937 (1993), doi:10.1029/93JA01896
- G. Haerendel, Space Sci. Rev. 124, 317 (2006), doi:10.1007/s11214-006-9092-z
- G. Haerendel, Astrophys. J. **707**, 903 (2009), doi:10.1088/0004-637X/707/2/903
- I. Hannah et al., Space Sci. Rev. (2011)
- J. Heyvaerts, E. R. Priest, D. M. Rust, Astrophys. J. 216, 123 (1977)
- T. Hirayama, Solar Phys. 34, 323 (1974), doi:10.1007/BF00153671
- G. Holman et al., Space Sci. Rev. pp. XXX-XXX (2011)
- H. Hudson, B. Haisch, K. T. Strong, J. Geophys. Res. 100, 3473 (1995), doi:10.1029/94JA02710
- H. S. Hudson, T. Bai, D. E. Gruber, J. L. Matteson, P. L. Nolan, L. E. Peterson, Astrophys. J. Lett. 236, L91 (1980), doi:10.1086/183205
- G. J. Hurford, S. Krucker, R. P. Lin, R. A. Schwartz, G. H. Share, D. M. Smith, Astrophys. J. Lett. 644, L93 (2006), doi:10.1086/505329
- G. J. Hurford, E. J. Schmahl, R. A. Schwartz, A. J. Conway, M. J. Aschwanden, A. Csillaghy, B. R. Dennis, C. Johns-Krull, S. Krucker, R. P. Lin, J. McTiernan, T. R. Metcalf, J. Sato, D. M. Smith, Solar Phys. 210, 61 (2002), doi:10.1023/A:1022436213688
- G. J. Hurford, R. A. Schwartz, S. Krucker, R. P. Lin, D. M. Smith, N. Vilmer, Astrophys. J. Lett. 595, L77 (2003), doi:10.1086/378179
- J. W. Knight, P. A. Sturrock, Astrophys. J. **218**, 306 (1977)
- E. Kontar et al., Space Sci. Rev. pp. XXX–XXX (2011)
- R. A. Kopp, G. W. Pneuman, Solar Phys. 50, 85 (1976), doi:10.1007/BF00206193
- T. N. Larosa, A. G. Emslie, Solar Phys. 120, 343 (1989)
- R. Lin, Space Sci. Rev. pp. XXX-XXX (2011)
- R. P. Lin, B. R. Dennis, G. J. Hurford, D. M. Smith, A. Zehnder, P. R. Harvey, D. W. Curtis, D. Pankow, P. Turin, M. Bester, A. Csillaghy, M. Lewis, N. Madden, H. F. van Beek, M. Appleby, T. Raudorf, J. McTiernan, R. Ramaty, E. Schmahl, R. Schwartz, S. Krucker, R. Abiad, T. Quinn, P. Berg, M. Hashii, R. Sterling, R. Jackson, R. Pratt, R. D. Campbell, D. Malone, D. Landis, C. P. Barrington-Leigh, S. Slassi-Sennou, C. Cork, D. Clark, D. Amato, L. Orwig, R. Boyle, I. S. Banks, K. Shirey, A. K. Tolbert, D. Zarro, F. Snow, K. Thomsen, R. Henneck, A. McHedlishvili, P. Ming, M. Fivian, J. Jordan, R. Wanner, J. Crubb, J. Preble, M. Matranga, A. Benz, H. Hudson, R. C. Canfield, G. D. Holman, C. Crannell, T. Kosugi, A. G. Emslie, N. Vilmer, J. C. Brown, C. Johns-Krull, M. Aschwanden, T. Metcalf, A. Conway, Solar Phys. 210, 3 (2002), doi:10.1023/A:1022428818870
- R. P. Lin, H. S. Hudson, Solar Phys. 50, 153 (1976)
- J. A. Miller, P. J. Cargill, A. G. Emslie, G. D. Holman, B. R. Dennis, T. N. LaRosa, R. M. Winglee, S. G. Benka, S. Tsuneta, J. Geophys. Res. 102, 14631 (1997), doi:10.1029/97JA00976
- H. E. Petschek, NASA Special Publication 50, 425 (1964)

R. Ramaty, N. Mandzhavidze, B. Kozlovsky, R. J. Murphy, Astrophys. J. Lett. 455, L193+ (1995), doi:10.1086/309841

- R. Ramaty, C. Paizis, S. A. Colgate, G. A. Dulk, P. Hoyng, J. W. Knight, R. P. Lin, D. B. Melrose, F. Orrall, P. R. Shapiro, in *Skylab Solar Workshop II*, ed. by P. A. Sturrock (1980), pp. 117–185.
- D. M. Rust, E. Hildner, R. T. Hansen, M. Dryer, A. N. McClymont, S. M. P. McKenna-Lawlor, D. J. McLean, E. J. Schmahl, R. S. Steinolfson, E. Tandberg-Hanssen, in *Skylab Solar Workshop II*, ed. by P. A. Sturrock (1980), pp. 273–339.
- T. Sakao, T. Kosugi, S. Masuda, K. Yaji, M. Inda-Koide, K. Makishima, Advances in Space Research 17, 67 (1996)
- J. Sato, T. Kosugi, K. Makishima, Pub. Astron. Soc. Japan 51, 127 (1999)
- G. H. Share, R. J. Murphy, Astrophys. J. 452, 933 (1995), doi:10.1086/176360
- $A.\ Y.\ Shih,\ R.\ P.\ Lin,\ D.\ M.\ Smith,\ Astrophys.\ J.\ Lett.\ \textbf{698},\ L152\ (2009),\ doi: 10.1088/0004-637X/698/2/L152$
- D. M. Smith, R. P. Lin, P. Turin, D. W. Curtis, J. H. Primbsch, R. D. Campbell, R. Abiad, P. Schroeder, C. P. Cork, E. L. Hull, D. A. Landis, N. W. Madden, D. Malone, R. H. Pehl, T. Raudorf, P. Sangsingkeow, R. Boyle, I. S. Banks, K. Shirey, R. Schwartz, Solar Phys. 210, 33 (2002), doi:10.1023/A:1022400716414
- D. M. Smith, G. H. Share, R. J. Murphy, R. A. Schwartz, A. Y. Shih, R. P. Lin, Astrophys. J. Lett. 595, L81 (2003), arXiv:astro-ph/0306292, doi:10.1086/378173
- D. S. Spicer, R. N. Sudan, Astrophys. J. 280, 448 (1984), doi:10.1086/162011
- P. A. Sturrock, Nature 211, 695 (1966), doi:10.1038/211695a0
- P. A. Sturrock, in Skylab Solar Workshop II, ed. by P. A. Sturrock (1980a), pp. 411-449
- P. A. Sturrock (ed.), Solar flares: A monograph from SKYLAB Solar Workshop II (1980b)
- P. A. Sweet, Annual Rev Astron. Astrophys. 7, 149 (1969), doi:10.1146/annurev.aa.07.090169.001053
- E. Tandberg-Hanssen, A. G. Emslie, *The physics of solar flares* (Cambridge and New York, Cambridge University Press, p. 286, 1988)
- G. H. J. van den Oord, Astron. Astrophys. 234, 496 (1990)
- R. F. van Oss, G. H. J. van den Oord, Astron. Astrophys. 299, 297 (1995)
- N. Vilmer et al., Space Sci. Rev. pp. XXX-XXX (2011)
- S. White et al., Space Sci. Rev. pp. XXX–XXX (2011)
- V. V. Zharkova, J. C. Brown, D. V. Syniavskii, Astron. Astrophys. 304, 284 (1995)
- V. V. Zharkova et al., Space Sci. Rev. pp. XXX-XXX (2011)

Appendix: Glossary of Acronyms Used in the Monograph

3DP	3-D Particles, onboard WIND
AAS ACE ACRIM AIA ARTB ATST	American Astronomical Society Advanced Composition Explorer Active Cavity Radiometer Irradiance Monitor, onboard SMM Atmospheric Imaging Assembly, onboard SDO active-region transient brightening Advanced Technology Solar Telescope
BATSE BBSO BCS	Burst And Transient Source Experiment, onboard <i>CGRO</i> Big Bear Solar Observatory Bent or Bragg Crystal Spectrometer, onboard <i>SMM</i> or <i>Yohkoh</i>
CA CCD CDF CDS CGRO CME CoMP	cellular automaton charge-coupled device cumulative distribution function Coronal Diagnostic Spectrometer, onboard SoHO Compton Gamma Ray Observatory coronal mass ejection Coronal Multi-channel Polarimeter Imaging Compton Telescope, onboard CGRO
CORONAS	Complex ORbital ObservatioNs of the Active Sun satellite series

CoSMO Coronal Solar Magnetism Observatory

CSHKP Carmichael, Sturrock, Hirayama, Kopp & Pneuman flare model

DC direct current

DEM differential mission measure DOI Digital Object Identifier DR diffusion region

EDF empirical distribution function

EIS EUV Imaging Spectrometer, onboard Hinode

Extreme ultraviolet Imaging Telescope, onboard SoHO EIT

EM emission measure **ENA** energetic neutral atom EP erupting prominence ESA European Space Agency EST European Solar Telescope **EUV** extreme ultraviolet

EVE EUV Variability Experiment, onboard SDO

FAL Fontenla, Avrett, & Loeser solar atmospheric model

FASR Frequency Agile Solar Radiotelescope

FIP first ionization potential **FMSS** fast-mode standing shock

Focusing Optics hard X-ray Spectrometer Imager **FOXSI**

FP footpoint

FWHM full width at half maximum

GBMGamma-ray Burst Monitor, onboard Fermi

GeDs germanium detectors

GEM Geospace Environment Modeling

GLE ground-level event

GOES Geostationary Operational Environmental Satellite

GONG Global Oscillation Network Group

GRANATGamma Rentgenovskii Astronomicheskii Nauchni ApparaT

GRBgamma-ray burst

Gamma-Ray Imaging Polarimeter for Solar flares **GRIPS** GRS Gamma Ray Spectrometer, onboard SMM **GSFC** [NASA] Goddard Space Flight Center

HEAO High Energy Astrophysical Observatory satellite series HMI Helioseismic and Magnetic Imager, onboard SDO HXIS Hard X-ray Imaging Spectrometer, onboard SMM HXR

hard X-ray

HXRBS Hard X-Ray Burst Spectrometer, onboard SMM **HXRS** Hard X-ray Spectrometer, onboard MTI HXTHard X-ray Telescope, onboard Yohkoh

IAU International Astronomical Union

ICEInternational Cometary Explorer, a.k.a ISEE-3

ICME interplanetary coronal mass ejection IDL Interactive Data Language International Geophysical Year IGY

INTEGRALINTErnational Gamma-Ray Astrophysics Laboratory

IR infrared

ISEE International Sun-Earth Explorer satellite series

KOSMA Kölner Observatorium für SubMillimeter Astronomie

KS Kolmogorov-Smirnov statistic

LASCO Large Angle and Spectrometric Coronagraph, onboard SoHO

LDE long-decay event

LHDI lower hybrid drift instability

LOFAR Low Frequency Array for Radio Astronomy

LOS line-of-sight
LPF large proton flare
LTE local thermal equilibrium

MDI Michelson Doppler Imager, onboard SoHO MEKAL Mewe-Kaastra-Liedahl atomic code

MEM maximum entropy method

MESSENGER MErcury Surface, Space Environment, GEochemistry, and Ranging mission

MHD magnetohydrodynamic(s)

MISS Multichannel Infrared Solar Spectrograph, at PMO

MSFC [NASA] Marshall Space Flight Center

MTI Multi-Thermal Imager

NASA National Aeronautics and Space Administration NOAA National Oceanic and Atmospheric Administration

NoRH Nobeyama Radioheliograph NoRP Nobeyama Radio Polarimeters

NRBH nonrelativistic Bethe-Heitler (bremsstrahlung cross-section)

NRH Nançay Radioheliograph

OSO Orbiting Solar Observatory satellite series

OSPEX Object SPectral EXecutive (IDL-based spectral analysis software)
OSSE Oriented Scintillation Spectrometer Experiment, onboard CGRO

OVSA Owens Valley Solar Array

PA position angle

PASJ Publications of the Astronomical Society of Japan

PDF probability density function

PFL post-flare loop

PHEBUS Payload for High Energy BUrst Spectroscopy, onboard GRANAT

PIC particle-in-cell
PIL polarity inversion line

PMO Purple Mountain solar Observatory POLAR POLAR spacecraft, not an acronym

PVO Pioneer Venus Orbiter

QPP quasi-periodic pulsations

RCS reconnecting current sheet

RESIK REntgenovsky Spektrometr s Izognutymi Kristalami, onboard CORONAS-F

RHESSI Reuven Ramaty High Energy Solar Spectroscopic Imager

RMC rotation modulation collimator

SC (RHESSI) subcollimators
SDO Solar Dynamics Observatory
SEE Solar Eruptive Events
SEP solar energetic particle

SFU Solar Flux Unit (10^{-22} W m⁻² Hz⁻¹)

SHH soft-hard-harder (temporal behavior of spectral index)
SHS soft-hard-soft (temporal behavior of spectral index)

SIGMA Systéme d'Imagerie Gamma à Masque Aléatoire, instrument onboard *GRANAT*

SMART Hida Solar Magnetic Activity Research Telescope

SMM Solar Maximum Mission
SMSS slow-mode standing shock
SOC self-organized criticality

SOHO or SoHO Solar and Heliospheric Observatory

SOLIS Synoptic Optical Long-term Investigations of the Sun magnetograph SONG SOlar Neutrons and Gamma-rays instrument, onboard CORONAS-F

 SORCE
 Solar Radiation and Climate Experiment

 SOT
 Solar Optical Telescope, onboard Hinode

 SOXS
 SOlar X-ray Spectrometer, onboard GSAT-2

 SPI
 SPectrometer on Integral, onboard INTEGRAL

 SPR-N
 Solar Spectropolarimeter, onboard CORONAS-F

SSRT Siberian Solar Radio Telescope SST Solar Submillimeter Telescope ST Hubble Space Telescope

STEREO Solar TErrestrial Relations Observatory

SUMER Solar Ultraviolet Measurements of Emitted Radiation instrument, onboard *SoHO*

SXI Soft X-ray Imager, onboard GOES

SXR soft X-ray

SXT Soft or Solar X-ray Telescope, onboard *Yohkoh* or *Hinotori*, respectively

THEMIS Time History of Events and Macroscale Interactions during Substorms

TIM Total Irradiance Monitor, onboard SORCE

TOF time-of-flight

TRACE Transition Region and Coronal Dynamics Explorer

TS termination shock
TSI total solar irradiance

UV ultraviolet

UVCS Ultraviolet Coronagraph Spectrometer, onboard SoHO

VAL Vernazza, Avrett, & Loeser solar atmospheric model

VLA Very Large Array VUV vacuum ultraviolet

WATCH Wide Angle Telescope for Cosmic Hard X-rays; onboard GRANAT

WAVES Not an acronym; instrument onboard WIND WBS Wide Band Spectrometer, onboard Yohkoh

WIND Spacecraft; not an acronym

WL white light

XBP X-ray bright point

XRT X-Ray Telescope, onboard *Hinode*

XUV X-ray/EUV/UV

Index

accelerated particles	wave energy transport, 6
diagnostics, 6	flares
energy content of, 3	and CMEs, 3
importance of, 2	energy content, 3
low energies, 3	naming convention, 2
number problem, 2, 4	footpoint simultaneity, 3
acceleration	footpoints, 3
test-particle approach, 4	free-free emission, 4
acceleration region	,
return current, 5	gamma-rays
test-particle approach, 4	delayed emission, 7
Ampère's Law, 5	deuterium formation (2.223 MeV), 7
Timpere s Euri, s	nuclear de-excitation, 7
beams	positron annihilation (511 keV), 7
	Giovanelli, R. G., 2
and waves, 6	
area, 6	gyrosynchrotron emission, 8
induced magnetic field, 5	H 1 C 1 24 2
return current, 5, 6	Hale Cycle 24, 2
self-inductance, 5	hard X-rays
bremsstrahlung, 4	emitted power, 5
efficiency, 6	inefficiency of, 6
Kramers approximation, 6	
thick-target, 5	literature citation, 1
	illustration, 2
Carrington, R. C., 2	loop prominence systems, 4
coronal mass ejections (CMEs), 3	
and flares, 3	magnetic field
energy content, 3	force-free, 4
coronal transients, 3	free energy, 3
Crab Nebula, 8	inductive time scale, 5
CSHKP, 4	magnetic structures
current closure, 5	mirror geometry, 3
current systems, 5	microflares
inductive time scale, 5	GOES B class, 9
	sensitive detection of, 8
debates	myths
thermal-nonthermal, 3	solar flare, 3
thermal-nonthermal, 5	solal fiale, 5
electrons	number problem, 2
	number problem, 2
beam energy transport, 6	December - Access 6
number problem, 2	Poynting flux, 6
total energy, 2	1
eras	radio emission
Skylab, 3	gyrosynchrotron, 8
	microwaves, 8
flare (individual)	reconnection, 3
SOL1972-08-04T06:25, 2	Petschek, 4
SOL2002-07-23T00:35 (X4.8), 8	Sweet-Parker, 4
cross-disciplinary analysis, 9	return current, 5
flare models	RHESSI, 1
casualties, 4	annealing operations, 8
collisional thick target, 5	collaborations, 8
CSHKP, 4	design and capabilities, 7
electron beam, 3	launch date, 8
Gold & Hoyle, 4	literature citation, 2
standard, 4	ribbons

```
H\alpha, 4
satellites
      Explorer series, 1 Hinode, 8
       Hinotori, 3
       RHESSI, 1
          launch, 1
       SDO, 8
      Skylab, 1
SMM, 3
       Solar Maximum Mission, 3
       STEREO, 8
       Yohkoh, 2, 3
shocks
      and reconnection, 4
Skylab, 1, 9
solar cycles
      cycle 23
      \gamma-ray events, 8 cycle 24, 2
solar flare myth, 3
Solar Maximum Mission
       Gamma-Ray Spectrometer, 7
space weather
       and extreme events, 3
standard model, 4
test-particle approach, 4
thermal-nonthermal debate, 3
workshops

RHESSI, 1, 4
      theory team, 4 Skylab, 2, 3
Yohkoh
      bibliography, 2
Hard X-ray Telescope (HXT), 3
```