

Spectrum and Morphology of the Two Brightest Milagro Sources in the Cygnus Region: MGRO J2019+37 and MGRO J2031+41.

A. A. Abdo,^{1,2} B. T. Allen,^{3,4} T. Aune,⁵ D. Berley,⁶ E. Bonamente,⁷
 G. E. Christopher,^{8,9} T. DeYoung,¹⁰ B. L. Dingus,¹¹ R. W. Ellsworth,¹²
 J. G. Galbraith-Frew,⁷ M. M. Gonzalez,¹³ J. A. Goodman,⁶ C. M. Hoffman,¹¹
 P. H. Hütemeyer,⁷ B. E. Kolterman,⁸ J. T. Linnemann,¹ J. E. McEnery,¹⁴ A. I. Mincer,⁸
 T. Morgan,¹⁵ P. Nemethy,⁸ J. Pretz,¹¹ J. M. Ryan,¹⁵ P. M. Saz Parkinson,⁵ G. Sinnis,¹¹
 A. J. Smith,⁶ V. Vasileiou,^{6,16} G. P. Walker,^{11,17} D. A. Williams⁵ and G. B. Yodh³
 ebonamen@mtu.edu, petra@mtu.edu

ABSTRACT

The Cygnus region is a very bright and complex portion of the TeV sky, host to unidentified sources and a diffuse excess with respect to conventional cosmic-ray propagation models. Two of the brightest TeV sources, MGRO J2019+37 and MGRO J2031+41, are analyzed using Milagro data with a new technique, and their emission is tested under two different spectral assumptions: a power law and a power law with an exponential cutoff. The new analysis technique is based on an energy estimator that uses the fraction of photomultiplier tubes in the observatory that detect the extensive air shower. The photon spectrum is measured in the range 1 to 200 TeV using the last 3 years of Milagro data (2005-2008), with the detector in its final configuration. MGRO J2019+37 is detected with a significance of 12.3 standard deviations (σ), and is better fit by a power law with an exponential cutoff than by a simple power law, with a probability $> 98\%$ (F-test). The best-fitting parameters for the power law with exponential cutoff model are a normalization at 10 TeV of $7_{-2}^{+5} \times 10^{-10} \text{ s}^{-1} \text{ m}^{-2} \text{ TeV}^{-1}$, a spectral index of $2.0_{-1.0}^{+0.5}$ and a cutoff energy of $29_{-16}^{+50} \text{ TeV}$. MGRO J2031+41 is detected with a significance of 7.3σ , with no evidence of a cutoff. The best-fitting parameters for a power law are a normalization of $2.4_{-0.5}^{+0.6} \times 10^{-10} \text{ s}^{-1} \text{ m}^{-2} \text{ TeV}^{-1}$ and a spectral index of $3.08_{-0.17}^{+0.19}$. The overall flux is subject to an $\sim 30\%$ systematic uncertainty. The systematic uncertainty on the power law indices is ~ 0.1 . A comparison with previous results from TeV J2032+4130, MGRO J2031+41 and MGRO J2019+37 is also presented.

Subject headings: Gamma rays: observations

¹ Department of Physics and Astronomy, Michigan State University, 3245 BioMedical Physical Sciences Building, East Lansing, MI 48824

² Current address: Space Science Division, Naval Research Laboratory, Washington, DC 20375

³ Department of Physics and Astronomy, University of California, Irvine, CA 92697

⁴ Current address: Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138

⁵ Santa Cruz Institute for Particle Physics, University of California, 1156 High Street, Santa Cruz, CA 95064

⁶ Department of Physics, University of Maryland, Col-

lege Park, MD 20742

⁷ Department of Physics, Michigan Technological University, Houghton, MI 49931

⁸ Department of Physics, New York University, 4 Washington Place, New York, NY 10003

⁹ Current address: Department of Physics, Brown University, Providence, RI 02912, USA

¹⁰ Department of Physics, Pennsylvania State University, University Park, PA 16802

¹¹ Group P-23, Los Alamos National Laboratory, P.O. Box 1663, Los Alamos, NM 87545

¹² Department of Physics and Astronomy, George Mason

1. Introduction

The Cygnus region is a part of our Galaxy of active massive star formation and destruction, and has been studied over a broad range of wavelengths, including radio, X-ray, GeV and TeV gamma-ray, as well as in cosmic rays. From GeV up to multi-TeV energies the Cygnus region is the brightest diffuse gamma-ray source in the northern hemisphere (Hunter et al. 1997).

One of the challenges in analyzing the Cygnus region at TeV energies, is the proper separation of the gamma-ray flux that is attributed to the point or extended sources in the region or to the diffuse emission. Previous Milagro analyses computed the diffuse emission from the region using two alternative methods to isolate the contribution from the resolved sources (Abdo et al. 2007, 2008), and found that at TeV energies the flux is still in excess with respect to the predicted flux from the GALPROP model (GALPROP). Milagro also published the discovery of two unidentified TeV sources in the region, MGRO J2019+37 and MGRO J2031+41 (Abdo et al. 2007a, 2009). The location of MGRO J2019+37 was found to be consistent with two EGRET sources, while the best fit position for MGRO J2031+41 was near two EGRET sources and the HEGRA unidentified source TeV J2032+4130. In a correlation study connecting the TeV sources discovered by Milagro with sources detected above 10σ (the so-called Bright Source List, BSL) by the Fermi Large Area Telescope (LAT), the two aforementioned brightest Milagro sources in the region were associated with GeV pulsars (Abdo et al. 2009c,?). MGRO J2019+37 and MGRO J2031+41 are currently associated with two and one pulsars identified by the Fermi LAT, respectively (Abdo et al. 2011c). Milagro also detected two candidate sources in the

Cygnus region ($l \in [65^\circ, 85^\circ]$ and $b \in [-2^\circ, +2^\circ]$) (Abdo et al. 2009).

Recently, several collaborations have presented new surveys of the Cygnus region, confirming the complexity of the region and showing the highly structured diffuse emission. The TeV emission from the position of MGRO J2019+37 has been confirmed by the VERITAS experiment (Aliu et al. 2011). The VERITAS spatial counts map shows a clear structure associated with MGRO J2019+37, a region of extended emission which seems to be produced by previously unresolved sources. At lower energies, the Fermi LAT collaboration has also recently published new results on the diffuse emission from the Cygnus region. An extended source, the so-called Cocoon, overlapping with MGRO J2031+41, has been found and its emission has been explained by freshly accelerated cosmic rays, trapped in a shell of photon-dominated emission formed by stellar winds and supernovae, as shown by mid-infrared observations (Abdo et al. 2011b). The spectrum from this region is hard, extending up to 100 GeV with no evidence of softening, and could be explained as the result of pulsar-accelerated particles or as an active super-bubble. The average diffuse emission from the region has also been analyzed at MeV to GeV energies (Abdo et al. 2011a). Despite the very rich source population, this emission is similar to that of the local interstellar space, and there is no evidence that it is necessary to include an extra contribution in the model, resembling the diffuse excess previously measured by Milagro (Abdo et al. 2007, 2008). Most recently, the ARGO-YBJ collaboration presented the results of a survey of the Cygnus region in the energy range of 600 GeV to 7 TeV. MGRO J2031+41 is observed with a significance of 6.4σ and a flux consistent with previous Milagro results, but there is no significant detection of MGRO J2019+37 (Bartoli et al. 2012).

Here, we present a new analysis of the last three years of data collected with the Milagro experiment (2005-2008). An improved gamma-hadron separation and a newly-developed technique are exploited to reconstruct the energy spectra of gamma rays from the sources in the Cygnus region. The best fits for the spectra of the two brightest Milagro sources are presented: MGRO J2019+37, a source with a post-trials sig-

University, 4400 University Drive, Fairfax, VA 22030

¹³ Instituto de Astronomía, Universidad Nacional Autónoma de México, 74D.F., México, 04510

¹⁴ NASA Goddard Space Flight Center, Greenbelt, MD 20771

¹⁵ Department of Physics, University of New Hampshire, Morse Hall, Durham, NH 03824

¹⁶ CRESST NASA/Goddard Space Flight Center, MD 20771 and University of Maryland, Baltimore County, MD 21250

¹⁷ Current address: National Security Technologies, Las Vegas, NV 89102

nificance in excess of 12σ between 1 and 100 TeV, and MGRO J2031+41 a source with a post-trials significance in excess of 7σ (Abdo et al. 2007a). We compare our spectra with results from the HEGRA, MAGIC, Whipple and ARGO-YBJ experiments (Aharonian et al. 2005; Albert et al. 2008; Lang et al. 2004; Bartoli et al. 2012).

2. Analysis Technique

The Milagro detector was located in the Jemez Mountains in New Mexico, at an altitude of 2,630 m a.l.s. It was operated from 2001 to 2008, and, in its final configuration, consisted of two components: (1) a central pond (60×80 m², 8 m deep) with two photomultiplier tube (PMT) layers, a shallow *air shower* layer consisting of 450 PMTs, and a deep *muon layer* consisting of 273 PMTs and (2) an array of 175 single-PMT tank detectors surrounding the pond covering an area of 40,000 m² (Atkins et al. 2001, 2003).

A detailed description of the analysis method and the parameters used here can be found in the Milagro paper on the spectral measurement of the Crab Nebula (Abdo et al. 2011). Systematic uncertainties of this method are estimated to be $\sim 30\%$ on the overall flux and ~ 0.1 on the spectral index.

The new element in this analysis, with respect to the previous approach (Abdo et al. 2007a, 2009), is the introduction of an estimator, \mathcal{F} , for the energy of the primary particle initiating the extensive air shower, based on the number of PMTs hit for each event. The \mathcal{F} parameter is defined as the sum of two fractions:

$$\mathcal{F} = \frac{N_{A.S.}}{N_{A.S.}^{live}} + \frac{N_{T.A.}}{N_{T.A.}^{live}}, \quad (1)$$

where $N_{A.S.}$ and $N_{T.A.}$ are the number of PMTs detecting the event, while $N_{A.S.}^{live}$ and $N_{T.A.}^{live}$ are the number of functional PMTs in the air shower layer and in the tank array, respectively. Because the typical energy resolution of Milagro is quite broad (50-100%), and the median energy associated with a given \mathcal{F} value is dependent on the spectral assumption for the observed source, the fit is performed in the measured \mathcal{F} space rather than in the energy space.

The reconstructed Milagro data contain information about the direction and \mathcal{F} value of air

shower events. According to their \mathcal{F} value, events are sorted into 9 bins ($0.2 \leq \mathcal{F} \leq 2$), resulting in a total of 9 signal skymaps, binned in $0.1^\circ \times 0.1^\circ$ pixels. For each \mathcal{F} bin, background maps are calculated using the Direct Integration method (Abdo et al. 2011). The two brightest regions of interest in the sky, a $2^\circ \times 2^\circ$ area around the position of the Crab Nebula and a band around the Galactic plane ($-2.5^\circ < b < 2.5^\circ$) are excluded when calculating the background (Abdo et al. 2011). Rather than discriminating between gamma-ray and cosmic-ray initiated air showers, a weight is applied to all measured events, in both signal and background maps, where gamma-ray like events are assigned a higher weight than cosmic-ray like events (Abdo et al. 2011). The gamma-ray excess with respect to the estimated background is calculated as the difference between signal weights and background weights, and is filled into the so-called excess map. Based on these excess maps we compute the \mathcal{F} distributions used in the energy fit.

Excess maps are smoothed according to the detector angular resolution (or point spread function, PSF), which is a function of \mathcal{F} and varies between 0.3° and 0.7° . The measured \mathcal{F} distribution for the target source in a $0.1^\circ \times 0.1^\circ$ bin, as a result, is the average excess coming from a PSF-wide region around the nominal source position. Since the background is measured with the direct integration method removing the Galactic plane ($b > 2.5^\circ$ and $b < -2.5^\circ$), the measured excess from sources in the Galactic plane includes any other diffuse or extended emission possibly present in the vicinity of the source. The Galactic diffuse background was estimated to contribute up to 15% of the flux at 35 TeV for the weakest Galactic BSL source (Abdo et al. 2009).

We test the emission from the two sources for two different spectral hypotheses: a power law, defined as

$$\frac{dN}{dE} = N_{10\text{TeV}} \left(\frac{E}{10 \text{ TeV}} \right)^{-\alpha}, \quad (2)$$

where $N_{10\text{TeV}}$ is the normalization scale set at 10 TeV and α the spectral index, and a power law with an exponential cutoff, defined as

$$\frac{dN}{dE} = N_{10\text{TeV}} \left(\frac{E}{10 \text{ TeV}} \right)^{-\alpha} \exp\left(-\frac{E}{E_c}\right), \quad (3)$$

where E_c is the cutoff energy.

The fit to the data is performed comparing the measured excess to simulations. First, a set of simulated data is generated varying sensitive parameters (i.e. $N_{10\text{TeV}}$, α and E_c). The best spectral parameters and the corresponding fit probability are then found using a χ^2 minimization, comparing the measured and the simulated \mathcal{F} distributions. Uncertainties on fit parameters are computed using $1\text{-}\sigma$ contours of χ^2 histograms (as discussed in sections 3.1 and 3.2). The uncertainty is defined as the distance between the best fit value and the lower and upper edges of the $1\text{-}\sigma$ contour.

3. Results and Discussion

The results presented here are obtained from the analysis of the Milagro data taken with the detector in its final configuration. The start date is 2005-10-22 20:39:16 GMT and the stop date is 2008-04-15 00:02:53 GMT, corresponding to a total of 906 days of observation, resulting in 832 integrated days after data quality cuts.

The positions of MGRO J2019+37 and MGRO J2031+41 are obtained fitting Milagro excess maps with a 2-dimensional Gaussian function with equal widths in right ascension and declination ($\sigma_{RA} = \sigma_{Dec}$), plus a constant base to take into account contribution from diffuse emission. The fit is performed in a square region of $\pm 3^\circ$ around the excess. Table 1 shows the positions used in this paper and a comparison to other surveys.

3.1. MGRO J2019+37 spectrum

In Fig. 1 the \mathcal{F} distribution of data is shown with the two simulated \mathcal{F} distributions for the best fitting power law and power law with exponential cutoff models. These distributions show the weighted excess coming from the source position after background subtraction. The excess from MGRO J2019+37 is compared to the excess measured from the entire Cygnus region ($65^\circ < l < 85^\circ$ and $-2^\circ < b < 2^\circ$). We find that the emission from MGRO J2019+37 accounts for $\sim 12\%$ of the excess from the entire region.

Figs. 2 and 3 show the 1 and $2\text{-}\sigma$ contours from the χ^2 fit for the two tested hypotheses. $1\text{-}\sigma$ contours are used to compute uncertainty on the parameters as described in section 2.

The best fit spectra for the two tested hypotheses (Eqs. 2 and 3) are shown in Fig. 4. The best-fit spectral parameters are summarized in Table 2.

With the power law hypothesis, the best-fit parameters for the normalization and the spectral index are $N_{10\text{TeV}} = (4.1 \pm 0.6) \cdot 10^{-10} \text{ s}^{-1} \text{ m}^{-2} \text{ TeV}^{-1}$ and $\alpha = 2.78 \pm 0.10$ respectively. The χ^2 is 16.12 for 7 degrees of freedom (d.o.f.), which gives a χ^2 fit probability of 0.024.

In the case of the power law with cutoff hypothesis, the best fit parameters are $N_{10\text{TeV}} = 7_{-2}^{+5} \cdot 10^{-10} \text{ s}^{-1} \text{ m}^{-2} \text{ TeV}^{-1}$, $\alpha = 2.0_{-1.0}^{+0.5}$ and $E_c = 29_{-16}^{+50} \text{ TeV}$. The χ^2 is 1.924 (6 d.o.f.), which gives a χ^2 fit probability of 0.93.

The χ^2 fit probabilities suggest that the power law with a cutoff model (χ^2 probability = 93%) fits the data better than a simple power law (χ^2 probability = 2.4%). We use the F-test (Bevington and Robinson 2003) to compute if the improvement in the fit is significant. For MGRO J2019+37, the simple power law is disfavoured at a C.L. $> 98\%$.

Previous Milagro analyses quoted the flux of MGRO J2019+37 at 20 and 35 TeV respectively (Abdo et al. 2007a, 2009). Those values, using a different analysis technique, are in agreement with the results presented here. An independent analysis (Allen 2007), which used a different energy estimator for the particle initiating the air shower and a different parameter to distinguish between gamma and cosmic rays, also is consistent with our results. ARGO-YBJ (Bartoli et al. 2012) has not observed any significant emission from MGRO J2019+37 in the range 600 GeV to 7 TeV. The explanations put forward in (Bartoli et al. 2012) are an insufficient exposure above 5 TeV or a possible time variability of MGRO J2019+37. The 90% C.L. upper limit from ARGO-YBJ is consistent with our best fitting model, a power law with an exponential cutoff (see Fig. 4). On the other hand, the simple power law model ($\alpha = 2.78 \pm 0.10$), already disfavoured by the F-test, does not agree with the ARGO-YBJ results.

3.2. MGRO J2031+41 spectrum

The \mathcal{F} distribution from data and those simulated for the two best fit models are shown in Fig. 5. MGRO J2031+41 accounts for $\sim 6\%$ of the excess from the entire Cygnus region.




Fig. 1.— MGRO J2019+37: distributions of the parameter used to estimate the photon energy, \mathcal{F} . The spectrum fit is performed in the \mathcal{F} space. Black squares represent the data. Red triangles and blue circles represent the simulated distributions for the best fit assuming a power law with a cutoff and a simple power law, respectively. The unit on the y axis is the weighted excess per day, the unit on the x axis is the \mathcal{F} value.




Fig. 2.— MGRO J2019+37: 1 and 2- σ C.L. contours for the power law fit. The point indicates the best fit result (minimum χ^2). The χ^2 increments for the 1 and 2- σ contours are 2.30 and 6.18 respectively.




Fig. 3.— MGRO J2019+37: 1 and 2- σ C.L. contours for the power law with cutoff fit. *left panel* - $N_{10\text{TeV}}$ vs α projection. *right panel* - E_c vs α projection. The points indicate the best fit result (minimum χ^2). The χ^2 increments for the 1 and 2- σ contours are 3.53 and 8.02 respectively.




Fig. 4.— MGRO J2019+37: energy spectra. The best fit is obtained for a power law with cutoff model (in red). The power law model is also shown (in blue). The shaded area represents the 1- σ band, obtained by varying the parameters within the 1- σ contour. ARGO-YBJ 90% C.L. upper limits for MGRO J2019+37 are shown in black (Bartoli et al. 2012).

Fig. 6 and 7 show the 1 and 2- σ contours from the χ^2 fit for the two tested hypotheses.

Best-fit spectral parameters are summarized in Table 2.

For the power law hypothesis, the best fit parameters are $N_{10\text{TeV}} = 2.4_{-0.5}^{+0.6} \cdot 10^{-10} \text{ s}^{-1} \text{ m}^{-2} \text{ TeV}^{-1}$ and $\alpha = 3.08_{-0.17}^{+0.19}$. The χ^2 is 6.98 (7 d.o.f.), with a χ^2 fit probability of 0.43.

For the power law with cutoff hypothesis, the best fit parameters are $N_{10\text{TeV}} = 3.7_{-1.9}^{+90.0} \cdot 10^{-10} \text{ s}^{-1} \text{ m}^{-2} \text{ TeV}^{-1}$, $\alpha = 2.7_{-2.6}^{+0.5}$ and $E_c = 44_{-40}^{+\infty} \text{ TeV}$. In this case we are not able to constrain the upper limit of the 1- σ uncertainty for the cutoff energy (see right panel of Fig. 7). The maximum E_c simulated value is 1000 TeV, almost an order of magnitude above the highest detected energy. The χ^2 is 5.80 (6 d.o.f.), with a χ^2 fit probability of 0.45. The best fit spectra for the two models are shown in Fig. 8.

The improvement in the fit obtained with the more complex model is not significant, according to the F-test, and the simple power law is to be preferred over the power law with the exponential cutoff.

The flux of MGRO J2031+41 at 20 and 35 TeV, measured by previous Milagro analyses (Abdo et al. 2007a, 2009), is in agreement with the results presented here. Results from ARGO-YBJ, MAGIC, HEGRA and Whipple (Bartoli et al. 2012; Albert et al. 2008; Aharonian et al. 2005; Lang et al. 2004) are shown in Fig. 8. The HEGRA and MAGIC models are mutually consistent, but they disagree with our best fit in terms of integral flux in the overlapping energy range. Between 1 and 10 TeV the flux measured by MAGIC accounts for only $\sim 3\%$ of the flux measured by Milagro. The spectrum as measured by the air Cherenkov telescopes (ACTs) is much harder ($\alpha \sim 2$) than the Milagro power law best fit ($\alpha \sim 3$). This discrepancy can be explained by the following two facts. First, the angular resolution of HEGRA and MAGIC ($< 0.1^\circ$) is much better than that of Milagro (0.3° to 0.7°). The Whipple flux was measured at 0.6 TeV, with a PSF of 0.21° , and it agrees better with the extrapolation of the Milagro result at lower energies. The same is true for ARGO-YBJ, which, with an angular resolution between 0.47° and 2.8° , measured an emission consistent with the

results presented here. The measurement presented in this paper accounts for photons coming from a larger region around the nominal source position compared to ACTs. Second, the way the background is computed is also different. Whipple used the ON/OFF method and observed the source with a significance of 3.3σ , while MAGIC primarily was operated in wobble mode (5.6σ). A possible explanation of the presented results is that TeV J2032+4130, whose extension is slightly larger than the HEGRA and MAGIC angular resolution, is surrounded by an extended emission, $\sim 4^\circ \times 3^\circ$ according to the Milagro map (see Fig. 9 discussed in Sec. 3.3). Therefore, Milagro, Whipple and ARGO-YBJ are not able to disentangle the extended emission from the central source and observe a higher flux. MAGIC and HEGRA, on the other hand, take into account the extended emission as a background. As a result, the ACTs measure a fainter emission and a harder spectrum for TeV J2032+4130.

3.3. Morphology

The significance map of the Cygnus region for the last three years of Milagro data (2005-2008) is shown in Fig. 9. An energy-dependent PSF smoothing is applied. The top panel shows the entire region in Galactic coordinates. Milagro positions for MGRO J2019+37 and MGRO J2031+41 are shown, as well as the high-significance Fermi LAT sources ($> 10\sigma$) and the extended Fermi Cocoon.

The Fermi counts map for photon energies above 10 GeV overlaid with the Milagro significance contours in the Cygnus region is shown in Fig. 10. Milagro detects an excess consistent with the position of the two pulsars PSR J2021+3651 and PSR J2032+4127, associated with MGRO J2019+37 and J2031+41 respectively, but no detection of significant emission coincident with PSR J2021+4026.

MGRO J2019+37 is observed with a significance of 12.3σ . Its extension is obtained with the 2-dimensional Gaussian fit discussed in Sec. 3. The fit results in $\sigma = 0.75^\circ$, slightly larger than the angular resolution of the detector in its final configuration. There is evidence from VERITAS that the extended source Milagro observes is the result of the superimposition of close point-like sources (Aliu et al. 2011).




Fig. 5.— MGRO J2031+41: distributions of the parameter used to estimate the photon energy, \mathcal{F} . The spectrum fit is performed in the \mathcal{F} space. Black squares represent the data. Red triangles and blue circles represent the simulated distributions for the best fit assuming a power law with a cutoff and a simple power law, respectively. The unit on the y axis is the weighted excess per day, the unit on the x axis is the \mathcal{F} value.




Fig. 6.— MGRO J2031+41: 1 and 2- σ C.L. contours for the power law fit. The point indicates the best fit result (minimum χ^2).

MGRO J2031+41, observed with a significance of 7.3σ , is an extended source, with $\sigma = 1.1^\circ$. It shows a central core with a higher significance ($> 6\sigma$) surrounded by a broader region (approximately $3^\circ \times 4^\circ$) with a lower significance ($> 4\sigma$). TeV J2032+4130 lies within the central core, and the Fermi Cocoon 1- σ radius ($\sigma = 2^\circ$) completely includes MGRO J2031+41. However, the Fermi analysis (Abdo et al. 2011b) shows that the Cocoon has an elongated shape, poorly correlating with the Milagro contours, and the possible association between the Cocoon and the extended Milagro excess remains unclear.

The spectral fit results do not change if the maximum significance position is chosen instead of the 2-D Gaussian fit position.

4. Conclusions

We present the spectra of the two brightest Milagro sources in the Cygnus region using a new analysis technique applied to the last 3 years of data collected by the Milagro experiment.

MGRO J2019+37 is observed with a significance over 12σ . Its emission is well fitted by a power law with an exponential cutoff ($E_c = 29_{-16}^{+50}$ TeV) and a hard asymptotic spectral index ($\alpha = 2.0_{-1.0}^{+0.5}$). The simple power law hypothesis is disfavoured at a C.L. $> 98\%$. The TeV excess measured by Milagro from MGRO J2019+37, spatially associated with the Fermi LAT pulsars J2018.0+3626 and J2021.0+3651, has been confirmed by VERITAS, and it is likely produced by nearby unresolved sources. ARGO-YBJ does not detect a significant emission from MGRO J2019+37, but the 90% C.L. upper limits do not conflict with the Milagro best fitting model.

The emission from MGRO J2031+41 (7.3σ significance), is well modeled by a power law with $\alpha = 3.08_{-0.17}^{+0.19}$. Our result, in particular the integral flux in the overlapping energy range, is consistent with previous measurements by ARGO-YBJ and Whipple, but it disagrees with HEGRA and MAGIC results for TeV J2032+4130, most likely because of the different PSF of the instruments and different background subtraction methods. MGRO J2031+41 appears to show a structured morphology, produced by the superimposition of a central point-like source, coincident with TeV J2032+4130, and an extended emission, pos-




Fig. 9.— PSF-smoothed significance maps of the Cygnus region. *top panel* - Black triangles mark the positions of the two Milagro sources with the $1\text{-}\sigma$ radius (solid circles), white-filled circles mark Fermi BSL sources (the $1\text{-}\sigma$ radius is consistent with the marker size). The Fermi Cocoon position is marked by the cross and the dashed ellipse is its $1\text{-}\sigma$ radius. *bottom left panel* - zoomed MGRO J2019+37 significance map. The circle represents the $1\text{-}\sigma$ radius. *bottom right panel* - zoomed MGRO J2031+41 significance map. The Milagro position with the $1\text{-}\sigma$ radius is shown in black. The crosses mark the Whipple (green), HEGRA (magenta) and MAGIC (blue) positions with their uncertainty.




Fig. 10.— Fermi LAT counts maps of the Cygnus region above 10 GeV. We used Pass 7 photon data from 2008-08-04 15:43:37 to 2011-12-06 10:10:13 GMT (LAT Data Server), zenith angle $< 107^\circ$, DATA_QUAL=1, LAT_CONFIG=1. A Gaussian smoothing is applied for display purposes. Circles mark the position of the Fermi BSL sources. White lines are the Milagro significance contours (3 , 5 and 11σ).

sibly produced either by unresolved sources or interactions of cosmic rays with the local interstellar medium. The correlation between the TeV extended emission and the overlapping Fermi Cocoon is unclear and needs further studies.

HAWC, the next generation water Cherenkov observatory, will be able to produce a more accurate analysis of the TeV emission from the Cygnus region, with its improved sensitivity (10-15 times better than Milagro) and angular resolution.

We gratefully acknowledge Scott Delay and Michael Schneider for their dedicated efforts in the construction and maintenance of the Milagro experiment. This work has been supported by the National Science Foundation (under grants PHY-0245234, -0302000, -0400424, -0504201, -0601080, -1002445, and ATM-0002744), the US Department of Energy (Office of High-Energy Physics and Office of Nuclear Physics), Los Alamos National Laboratory, the University of California, and the Institute of Geophysics and Planetary Physics.

REFERENCES

- Abdo, A. A. et al., 2007, ApJ, 658, L33
- Abdo, A. A. et al., 2007a, ApJ, 664, L91
- Abdo, A. A. et al., 2008, ApJ, 688, 1078
- Abdo, A. A. et al., 2009, ApJ, 700, L127
- Abdo, A. A. et al., 2009a, Science, 326, 1512
- Abdo, A. A. et al., 2009b, Physical Review Letters, 103, 251101
- Abdo, A. A. et al., 2009c, ApJS, 183, 46
- Abdo, A. A. et al., 2011, arXiv:1110.0409v1
- Abdo, A. A. et al., 2011a, arXiv:1110.6123v1
- Abdo, A. A. et al., 2011b, Science, 334, 1103
- Abdo, A. A. et al., 2011c, arXiv:1108.1435v1
- Aharonian, F. et al., 2005, A&A, 431, 197
- Albert, J. et al., 2008, ApJ, 675, L25
- Aliu, E. for the VERITAS Collaboration, 2011, in Proc. of 32nd ICRC, arXiv:1110.4656v2
- Allen, B. 2007, PhD thesis, University of California, Irvine
- Atkins, R. et al., 2001, in Proc. 27th ICRC, arXiv:astro-ph/0110513
- Atkins, R. et al., 2003, ApJ, 595, 803
- Bartoli, B. et al., 2012, ApJ, 754, L22
- Bevington R. P. and Robinson D. K., 2003, ISBN: 0-07-247227-8
- GALPROP, <http://galprop.stanford.edu>
- Hunter, S. D. et al., 1997, ApJ, 481, 205
- Lang, M. J. et al., 2004, A&A, 423, 415
- LAT Data Server, <http://fermi.gsfc.nasa.gov/ssc/data/access/>

This 2-column preprint was prepared with the AAS L^AT_EX macros v5.2.




Fig. 7.— MGRO J2031+41: 1 and 2- σ C.L. contours for the power law with cutoff fit. *left panel* - $N_{10\text{TeV}}$ vs α projection. *right panel* - E_c vs α projection. The points indicate the best fit result (minimum χ^2). The right panel shows that the present analysis is unable to constrain the upper limit of the energy cutoff within 1 σ .




Fig. 8.— MGRO J2031+41: energy spectra. The power law model is shown in blue and the power law with cutoff model is shown in red. These two hypotheses give the same χ^2 fit probability. The pink cross is the Whipple flux at 0.6 TeV (Lang et al. 2004). The fine-dashed cyan line and the dashed yellow line are the HEGRA and MAGIC best fits (Aharonian et al. 2005; Albert et al. 2008), respectively. The dot-dash black line is the ARGO-YBJ best fit (Bartoli et al. 2012). The shadowed area represents the 1- σ band.

TABLE 1
SOURCE POSITIONS

survey	RA	Dec	width (σ)
MGRO J2019+37			
2-D Gaussian with base	304.646 ± 0.008	36.847 ± 0.006	0.75
Fermi position (0FGL J2020.8+3649) ^a	305.22	36.83	—
Milagro 2007 ^b	304.98	36.66	1.1
MGRO J2031+41			
2-D Gaussian with base	307.41 ± 0.02	41.190 ± 0.015	1.1
Fermi position (0FGL J2032.2+4122) ^a	308.06	41.38	—
Milagro 2007 ^b	307.98	41.51	3.0

^a(Abdo et al. 2009)

^b(Abdo et al. 2007a)

NOTE.—Positions of MGRO J2019+37 (top) and MGRO J2031+41 (bottom) for the 2-D Gaussian with base fit, Fermi position of associated sources and previous Milagro survey.

TABLE 2
FIT RESULTS

Source	$N_{10\text{TeV}} (10^{-10} \text{ s}^{-1} \text{ m}^{-2} \text{ TeV}^{-1})$	α	E_c (TeV)	$\chi^2/\text{d.o.f.}$	prob.
MGRO J2019+37	4.1 ± 0.6	2.78 ± 0.10	∞	16.12/7	0.024
MGRO J2019+37	7^{+5}_{-2}	$2.0^{+0.5}_{-1.0}$	29^{+50}_{-16}	1.924/6	0.93
MGRO J2031+41	$2.4^{+0.6}_{-0.5}$	$3.08^{+0.19}_{-0.17}$	∞	6.980/7	0.43
MGRO J2031+41	$3.7^{+90.0}_{-1.9}$	$2.7^{+0.5}_{-2.6}$	$44^{+\infty}_{-40}$	5.796/6	0.45

NOTE.—Best fit results for MGROJ 2019+37 and MGRO J2031+41 using both the power law and the power law with cutoff hypotheses. The table lists statistical errors only, the systematic flux error is $\sim 30\%$ and the systematic error on the spectral index is ~ 0.1 .