Comment on "First Observation of Ground State Dineutron Decay: 16Be"

F.M. Marqués, ¹ N.A. Orr, ¹ N.L. Achouri, ¹ F. Delaunay, ¹ and J. Gibelin ¹ LPC Caen, ENSICAEN, Université de Caen, CNRS/IN2P3, 14050 Caen cedex, France Published in Physical Review Letters 109, 239201 (2012).

PACS numbers: 21.10.-k, 21.10.Dr, 27.20.+n

In a recent paper by Spyrou et al. [1], an investigation of the unbound system ¹⁶Be via single-proton removal from ¹⁷B was reported. In addition to identifying a structure some 1.5 MeV above the $^{14}\text{Be}+n+n$ threshold, interpreted as the ¹⁶Be ground state, significant enhancements were observed at low n-n relative energy (E_{n-n}) and angle (θ_{n-n}) . Through a comparison with simulations based essentially on direct three-body and "dineutron" decay, Spyrou et al. concluded that only the dineutron mode was consistent with these effects. As such it was claimed that the first case of dineutron decay had been discovered. Here we point out that such an interpretation is, at best, premature as the inclusion of the n-n interaction in the description of direct three-body decay can generate strong enhancements at low n-n relative energy and angle, as observed, without the need to invoke dineutron decay.

An important feature of the interpretation was the treatment of the direct three-body decay mode, whereby the well-known n-n interaction was neglected. By contrast, the dineutron decay was modeled in terms of the two-body decay of 16 Be into 14 Be and a quasi-bound 2n cluster, followed by n+n decay. The neglect, however, of the n-n interaction in the former case is a significant oversight. Indeed, it is well known that the low-energy $^{1}S_0$ n-n interaction invariably leads to a characteristic enhancement near zero relative momentum, a feature which is exploited in determinations of the n-n scattering length [2]. More generally, this enhancement is observed in almost any final state in which two neutrons are emitted over a relatively short time scale (see, for example, Refs. [3–6]).




FIG. 1: (color online) Calculations of $^{14}\text{Be}+n+n$ decay for three-body phase space without (1) and including (2) the n-n FSI, and for dineutron decay (3). The results have been normalized in each panel to the same integrated yield (Y).

It would be surprising, therefore, if such effects were not present in the decay of $^{16}{\rm Be}.$

To put these considerations on a more quantitative footing, we have undertaken three different calculations (Fig. 1). For all three the input is a $^{14}\text{Be}+n+n$ decayenergy distribution following that observed in Ref. [1]. In case (1) the energy is shared by the three particles following phase space considerations alone. In case (3), dineutron decay, it is shared through a sequential process: $^{14}\text{Be}+^2n$ breakup followed by $^2n\rightarrow n+n$, with the 2n decay energy similar to that of Ref. [1] (Fig. 1a).

The new case, denoted (2), takes as its starting point three-body phase space which is then modulated by the n-n final-state interaction (FSI). The calculations follow the formalism of Ref. [7], as described in Ref. [5]. For the example shown here an initial n-n average separation of $r_{nn}^{\rm rms} = 4$ fm was assumed. The results are very similar, to those of the dineutron model: a clear enhancement is observed at low n-n relative angle and energy, together with a $^{14}{\rm Be}$ -n distribution peaking sharply at large angles. While it is beyond the scope of a Comment to attempt to reproduce exactly the data of Spyrou et al., if account were taken of the statistics and experimental response function, it would not be possible to discriminate between these two decay processes.

In summary, the inclusion of the n-n interaction in the description of direct three-body decay of 16 Be generates strong enhancements at low n-n relative energy and angle and large 14 Be-n opening angles, characteristic of those observed by Spyrou $et\ al.\ [1]$, without the need to invoke dineutron decay.

^[1] A. Spyrou et al., Phys. Rev. Lett. 108, 102501 (2012).

^[2] D.E. González Trotter et al., Phys. Rev. C 73, 034001 (2006).

^[3] K. Ieki et al., Phys. Rev. Lett. 70, 730 (1993).

^[4] F.M. Marqués et al., Phys. Lett. B 476, 219 (2000);N.A. Orr, arXiv:0803.0886v1 [nucl-ex].

^[5] F.M. Marqués et~al., Phys. Rev. C ${\bf 64},\,061301$ (2001).

^[6] T. Aumann et al., Phys. Rev. C 59, 1252 (1999).

^[7] R. Lednicky and L. Lyuboshits, Sov. J. Nucl. Phys. 35, 770 (1982).