

Number of Walks and Degree Powers in a Graph

M.A. Fiol, E. Garriga

Universitat Politècnica de Catalunya, BarcelonaTech
 Departament de Matemàtica Aplicada IV
 Barcelona, Catalonia
 (e-mails: {fiol, egarriga}@ma4.upc.edu)

Abstract

This note deals with the relationship between the total number of k -walks in a graph, and the sum of the k -th powers of its vertex degrees. In particular, it is shown that the the number of all k -walks is upper bounded by the sum of the k -th powers of the degrees.

Let $G = (V, E)$ be a connected graph on n vertices, $V = \{1, 2, \dots, n\}$, with adjacency matrix \mathbf{A} . For any integer $k \geq 1$, let $a_{ij}^{(k)}$ denote the (i, j) entry of the power matrix \mathbf{A}^k . Let \mathbf{D} be the diagonal matrix with elements $(\mathbf{D})_{ii} = d_i$ (the degree of vertex i). Here we study the relationship between the sum of all walks of length k in G and the sum of the k -th powers of its degrees. As a main result, and answering in the affirmative a conjecture of Marc Noy [8], we will show that

$$\sum_{i,j} a_{ij}^{(k)} \leq \sum_i d_i^k, \quad (1)$$

with equality if and only if G is regular or $k \leq 2$. In the case $k = 3$ we also provide an exact value of the difference between the above sums in (1). In other line of research, some upper bounds for $\sum_i d_i^k$ have been given by several authors. See, for instance, [3, 6, 7, 10] (for general graphs) and [2, 5] (for graphs not containing a prescribed subgraph).

Let us first begin with the small values of k . The case $k = 0$ is trivial since the number of walks of length 0 equals the number of vertices. Similarly, if $k = 1$, the sum $\sum_{i,j} a_{ij}$ is just the sum of the degrees $d_1 + d_2 + \dots + d_n$. If $k = 2$, we can use that $\mathbf{A}\mathbf{j} = \mathbf{D}\mathbf{j}$ (where \mathbf{j} is the all-1 vector) and the symmetry of the involved matrices to obtain:

$$\sum_{i,j} a_{ij}^{(2)} = \langle \mathbf{j}, \mathbf{A}^2 \mathbf{j} \rangle = \langle \mathbf{A}\mathbf{j}, \mathbf{A}\mathbf{j} \rangle = \|\mathbf{A}\mathbf{j}\|^2 = \|\mathbf{D}\mathbf{j}\|^2 = d_1^2 + d_2^2 + \dots + d_n^2.$$

Assume now that G is regular of degree, say, d . Then, \mathbf{j} is the positive eigenvector corresponding to the eigenvalue d and we get

$$\sum_{i,j} a_{ij}^{(k)} = \langle \mathbf{j}, \mathbf{A}^k \mathbf{j} \rangle = \langle \mathbf{j}, d^k \mathbf{j} \rangle = d^k \|\mathbf{j}\|^2 = nd^k.$$

A similar reasoning shows that, for a general (non-regular) graph, the inequality in (1) always holds for k large enough. Indeed, let $\boldsymbol{\nu} = (\nu_1, \nu_2, \dots, \nu_n)^\top$ be the positive eigenvector of G , normalized in such a way that $\min_{i \in V} \nu_i = 1$. Let λ be its corresponding (positive) eigenvalue, which is known to be smaller than the maximum degree Δ of G (see [1, 4]). Then,

$$\sum_{i,j} a_{ij}^{(k)} \leq \langle \boldsymbol{\nu}, \mathbf{A}^k \boldsymbol{\nu} \rangle = \langle \boldsymbol{\nu}, \lambda^k \boldsymbol{\nu} \rangle = \|\boldsymbol{\nu}\|^2 \lambda^k$$

which, for k large enough, is smaller than the single k -power Δ^k .

To deal with the case $k = 3$, it is more convenient to work with the Laplacian matrix of G ; that is, $\mathbf{L} := \mathbf{D} - \mathbf{A}$. Then, recall that, given any real function defined on V , $f : V \rightarrow \mathbb{R}$, and with \mathbf{f} being the (column) vector with components the values of f on V , we have

$$\langle \mathbf{f}, \mathbf{L} \mathbf{f} \rangle = \sum_{i \sim j} (f(i) - f(j))^2,$$

where the sum is extended over all edges of G (see, for instance, [1]). We are interested in the case when the above function is just the degree of the corresponding vertex: $f(i) = d_i$. In this case, we denote by $\boldsymbol{\delta}$ its corresponding vector. Then, the difference between the two sums in (1) is just

$$\sum_i d_i^3 - \sum_{i,j} a_{ij}^{(3)} = \langle \boldsymbol{\delta}, \mathbf{D} \boldsymbol{\delta} \rangle - \langle \boldsymbol{\delta}, \mathbf{A} \boldsymbol{\delta} \rangle = \langle \boldsymbol{\delta}, \mathbf{L} \boldsymbol{\delta} \rangle = \sum_{i \sim j} (d_i - d_j)^2 \geq 0.$$

To prove the inequality in the general case, note first that, for any two positive numbers a, b with, say, $a \geq b$,

$$a^r b + ab^r = a^{r+1} + b^{r+1} - (a^r - b^r)(a - b) \leq a^{r+1} + b^{r+1}$$

with equality if and only if $a = b$. (The same conclusion is reached when we apply Hölder inequality [8] to the vectors (a, b) and (b^r, a^r) with norms L^{r+1} for the first vector and dual norm $L^{(r+1)/r}$ for the second one.)

Also, notice that all walks of a given length, say $k \geq 1$, can be obtained by considering, for any vertex i , all (i, j) -walks of length $k - 1$ “extended” by each of the d_i edges incident to i . Thus, $\sum_{i,j} a_{ij}^{(k)} = \sum_{i,j} d_i a_{ij}^{(k-1)}$ or, equivalently,

$$\langle \mathbf{j}, \mathbf{A}^k \mathbf{j} \rangle = \langle \mathbf{j}, \mathbf{A}^{k-1} \mathbf{D} \mathbf{j} \rangle = \langle \mathbf{D} \mathbf{j}, \mathbf{A}^{k-1} \mathbf{j} \rangle.$$

Keeping all this in mind, we are now ready to prove (1). Indeed, assuming that $k \geq 3$, we have:

$$\begin{aligned}
\sum_{i,j} a_{ij}^{(k)} &= \sum_{i,j} d_i a_{ij}^{(k-2)} d_j = \sum_i a_{ii}^{(k-2)} d_i^2 + \sum_{i<j} 2a_{ij}^{(k-2)} d_i d_j \\
&\leq \sum_i a_{ii}^{(k-2)} d_i^2 + \sum_{i<j} a_{ij}^{(k-2)} (d_i^2 + d_j^2) \\
&= \sum_{i,j} a_{ij}^{(k-2)} d_j^2 \\
&= \sum_{i,j} d_i a_{ij}^{(k-3)} d_j^2 = \sum_i a_{ii}^{(k-3)} d_i^3 + \sum_{i<j} a_{ij}^{(k-3)} (d_i d_j^2 + d_i^2 d_j) \\
&\leq \sum_i a_{ii}^{(k-3)} d_i^3 + \sum_{i<j} a_{ij}^{(k-3)} (d_i^3 + d_j^3) \\
&= \sum_{i,j} a_{ij}^{(k-3)} d_j^3 \leq \dots \leq \sum_{i,j} a_{ij} d_j^{k-1} = \sum_j d_j^k.
\end{aligned}$$

Moreover, notice that all the above inequalities become equalities if and only if G is regular, as claimed.

As commented by one of the referees, the above prove suggests a wider result: If \mathbf{A} is a real symmetric $n \times n$ matrix, $\mathbf{D} = \text{diag}(d_1, d_2, \dots, d_n)$, where $d_i = \sum_j |a_{ij}|$, and \mathbf{J} is the all-1 matrix, then

$$\text{tr}(\mathbf{A}^k \mathbf{J}) \leq \text{tr}(\mathbf{D}^k \mathbf{J}) = \sum_i d_i^k. \quad (2)$$

In other words, an extremal (not characteristic) property of the positive diagonal matrices is exhibited.

Acknowledgments

Research supported by the Ministerio de Educación y Ciencia, Spain, and the European Regional Development Fund under project MTM2005-08990-C02-01 and by the Catalan Research Council under project 2005SGR00256.

The authors thank one of the referees for the comment about the use of Hölder inequality and the result in (2).

References

- [1] N. Biggs, *Algebraic Graph Theory*, second ed., Cambridge University Press, Cambridge, 1993.

- [2] B. Bollobás and V. Nikiforov, Degree powers in graphs with forbidden subgraphs, *Electron. J. Combin.* 11 (2004), Research paper R42.
- [3] Sebastian M. Cioabă, Sums of powers of the degrees of a graph, *Discrete Math.* 306 (2006) 1959–1964.
- [4] D.M. Cvetković, Graphs and their spectra, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* No. 354–356 (1971), 1–50.
- [5] Y. Caro and R. Yuster, A Turán type problem concerning the powers of the degrees of a graph, *Electron. J. Combin.* 7 (2000), Research paper R47.
- [6] K. Ch. Das, Maximizing the sum of the squares of the degrees in a graph, *Discrete Math.* 285 (2004), 57–66.
- [7] D. de Caen, An upper bound on the sum of the squares of the degrees in a graph, *Discrete Math.* 185 (1998), 245–248.
- [8] G.H. Hardy, J.E. Littlewood and G. Pólya, *Inequalities*, Cambridge Univ. Press, Cambridge, 1934.
- [9] Marc Noy, personal communication (2005).
- [10] U.N. Peled, R. Petreschi and A. Sterbini, (n, e) -graphs with maximum sum of squares of degrees, *J. Graph Theory* 31 (1999), 283–295.