

Empirical parameterization of the high energy behaviour of average charged particle multiplicities in e^+e^- , ep and pp collisions

M.Zavertyaev

Max-Planck Institut für Kernphysik, D-69117 Heidelberg, Germany

P.N.Lebedev Physical Institute, 117924 Moscow B-333, Russia

Abstract

A simple fit of existing data for center-of-mass energies above the b -threshold shows that the average charged particle multiplicities in $p(\bar{p})$ and pp interactions can be parameterized by a simple power law $\langle n_{ch} \rangle = N_0 \cdot (s/s_0)^\gamma$ with γ close to $1/5$. Choosing $s_0 = m_p^2$ one finds $N_0=2.32$. The average charged particle multiplicities in e^+e^- and $e^\pm p$ interactions follows the same power law with an offset of ± 1 unit in N_0 . The observed trend leads to the conclusion that in LHC experiments at energy of 7 TeV one should expect the average charge multiplicity in the region of $\langle n_{ch} \rangle$ close to 80.

Many experiments published their results on average charge multiplicities studies at different energies in e^+e^- , $e^\pm p$ and $p(\bar{p})$ and pp interactions. A compilation of these data may be found in a data files prepared by Particle Data Group ([1] and references therein).

In Fig.1 the data taken from [1] are presented in double logarithmic scale. For $\sqrt{s} > 11$ GeV, just above the mass of the Upsilon resonances, the $p(\bar{p})$ and pp data are fitted by $\langle n_{ch} \rangle = N_0 \cdot (s/m_p^2)^{1/5}$ with m_p the proton mass and one free parameter N_0 . Such a simple power law may find its explanation if the process of multiparticle production has some sort of fractal nature in it. It is intriguing that the power $1/5$ which provides a very good fit to the high energy data appears to be equal to $1/n_f$, the number of active flavours in the process. Furthermore, the best fit value $N_0 = 2.32$ is very close to $\ln(m_\Upsilon/m_p)$.

Another curious observation is related to the e^+e^- and the $e^\pm p$ data points. The green and the red lines are obtained from the fit to the $p(\bar{p})$ and pp data by changing N_0 by ± 1 unit. Surprising is the precision with which the data points from e^+e^- and $e^\pm p$ interactions follow the same power law. One also sees that the $e^\pm p$ measurements, which pertain to the current fragmentation region in the Breit-frame, are about half of the $p(\bar{p})$ and pp multiplicities.

The observed trend leads to the conclusion that in LHC experiments at energy of $\sqrt{s} = 7$ TeV one should expect the average charge multiplicity in the region of $\langle n_{ch} \rangle$ close to 80 - the extrapolation for about one order of magnitude of energy. In Fig.1 the expectation region marked by a big magenta dot.

It would be very interesting to see the results on average charge multiplicity in the future experiments on e^+e^- and $e^\pm p$ new generation accelerators.

Acknowledgments

The author acknowledges very useful discussions with M.Schmelling.

References

- [1] Nakamura, K. *et al.*, *Review of particle physics*, *J.Phys.*, 2010.




Figure 1: The average charge multiplicities in e^+e^- , $e^\pm p$ and $p(\bar{p})$ and pp interactions. The data points are taken from [1]. The blue line is the result of the fit of data in $p(\bar{p})$ and pp interaction with the power fixed to 1/5, the green and red lines are the offset of fit by ± 1 unit. The magenta dot indicates the expected average charge multiplicities at 7 TeV.