

A new approach to the results of Kövari, Sós, and Turán concerning rectangle-free subsets of the grid

Jeremy F. Alm
 Illinois College
 Jacksonville, IL 62650
 alm.academic@gmail.com

Jacob Manske*
 Texas State University
 San Marcos, TX 78666
 jmanske@txstate.edu

November 10, 2018

Abstract

For positive integers m and n , define $f(m, n)$ to be the smallest integer such that any subset A of the $m \times n$ integer grid with $|A| \geq f(m, n)$ contains a rectangle; that is, there are $x \in [m]$ and $y \in [n]$ and $d_1, d_2 \in \mathbb{Z}^+$ such that all four points (x, y) , $(x + d_1, y)$, $(x, y + d_2)$, and $(x + d_1, y + d_2)$ are contained in A . In [12], Kövari, Sós, and Turán showed that $\lim_{k \rightarrow \infty} \frac{f(k, k)}{k^{3/2}} = 1$. They also showed that $f(p^2, p^2 + p) = p^2(p + 1) + 1$ whenever p is a prime number. We recover their asymptotic result and strengthen the second, providing cleaner proofs which exploit a connection to projective planes, first noticed by Mendelsohn in [14]. We also provide an explicit lower bound for $f(k, k)$ which holds for all k .

1 Introduction and motivation

For a positive integer n , let $[n] = \{1, 2, \dots, n\}$. For $m, n \in \mathbb{Z}^+$, define $f(m, n)$ to be the least integer such that if $A \subseteq [m] \times [n]$ with $|A| \geq f(m, n)$, then A contains a rectangle; that is, there is $x \in [m], y \in [n]$, and $d_1, d_2 \in \mathbb{Z}^+$ such that all four points (x, y) , $(x + d_1, y)$, $(x, y + d_2)$, and $(x + d_1, y + d_2)$ are contained in A . For ease in notation, let $f(k) = f(k, k)$. For $c \in \mathbb{Z}^+$, a c -coloring of a set S is a surjective map $\chi : S \rightarrow [c]$. If χ is constant on a set $A \subset S$, we say that A is *monochromatic*.

We will write $g(k) \sim h(k)$ to mean that functions g and h are *asymptotically equal*; that is, $\lim_{k \rightarrow \infty} \frac{g(k)}{h(k)} = 1$. Also, notice that $f(m, n) = f(n, m)$ for any choice of n and m .

*Corresponding author.

The problem of finding bounds or exact values of $f(m, n)$ finds its roots in the famous theorem of van der Waerden from [21], which states that given any positive integers c and d , there exists an integer N such that any c -coloring of $[N]$ contains a monochromatic arithmetic progression of length d . Szemerédi proved a density version of this theorem in [20], using the now well-known Regularity Lemma. Progress in this area is still being made. For instance, in [3], Axenovich and the second author try to find the smallest k so that in any 2-coloring of $[k] \times [k]$ there is a monochromatic *square*; i.e., a rectangle with $d_1 = d_2$. While the upper bounds are enormous, they proved $k \geq 13$; in [4], Bacher and Eliahou show that $k = 15$. In [10], the authors are interested in finding OBS_c , which is the collection of $[m] \times [n]$ grids which cannot be colored in c colors without a monochromatic rectangle, but every proper subgrid can be; see also [7]. For a more complete survey on van der Waerden type problems, see [11].

Zarankiewicz introduced the problem of finding $f(m, n)$ in [22] using the language of minors of (0,1)-matrices. In [12], Kővari, Sós, and Turán show that $f(k) \sim k^{3/2}$ and that whenever p is a prime number, we have $f(p^2 + p, p^2) = p^2(p + 1) + 1$. In this manuscript, we will recover this asymptotic result and strengthen the second result.

In [17], Reiman achieved the bound of

$$f(m, n) \leq \frac{1}{2} \left(m + \sqrt{m^2 + 4mn(n-1)} \right) + 1. \quad (1)$$

Notice that by setting $m = p^2 + p$ and $n = p^2$, the right hand side of (1) becomes $p^2(p + 1) + 1$, so the result of Kővari, Sós, and Turán implies that the inequality is sharp. Reiman showed equality in (1) in the case that $m = n = q^2 + q + 1$, provided q is a prime power. In [14], Mendelsohn recovers and strengthens the equality result of Reiman by noticing the connection of the Zarankiewicz problem to projective planes.

A $k \times k$ (0,1)-matrix A corresponds to a subset $S_A \subset [k] \times [k]$ by

$$(i, j) \in S \text{ if and only if the } (i, j) \text{ entry of } A \text{ is } 1.$$

Notice that the set S_A contains a rectangle if and only if the matrix $A^T A$ has an entry off the main diagonal which is not equal to 0 or 1. Also notice that $\text{tr}(A^T A) = |S_A|$.

Such (0,1)-matrices arise in the study of projective planes. A projective plane of order n is an incidence structure consisting of $n^2 + n + 1$ points and $n^2 + n + 1$ lines such that

- (i) any two distinct points lie on exactly one line;
- (ii) any two distinct lines intersect in exactly one point;
- (iii) each line contains exactly $n + 1$ points; and
- (iv) there is a set of 4 points such that no 3 of these points lie on the same line.

It is not known for which positive integers n there exists a projective plane of order n ; projective planes have been constructed for all prime-power orders, but for no others. In the well-known paper [5], Bruck and Ryser show that if the square-free part of n is divisible by a prime of the form $4k + 3$, and if n is congruent to 1 or 2 modulo 4, then there is no projective plane of order n ; see also [6]. More recently, the authors in [8] draw a connection between the existence of projective planes of order greater than or equal to 157 and the number of cycles in $n \times n$ bipartite graphs of girth at least 6. In 1989, a computer search conducted by the authors in [13] showed that there is no projective plane of order 10. The smallest order for which it is still not known whether there is a projective plane is 12, although the results in [15, 19, 16, 1, 2] suggest that there is no such structure.

Next we state a lemma which appears in [14] connecting projective planes to the Zarankiewicz problem.

Lemma 1. *If n is a positive integer such that there exists a projective plane of order n , then $f(n^2 + n + 1) = (n + 1)(n^2 + n + 1) + 1$.*

We will include a proof of Lemma 1 both for completeness and since we will reference the lower bound construction in the proof of Theorem 2.

Proof of Lemma 1. Let n be a positive integer such that there is a projective plane of that order. For ease in notation, set $N = n^2 + n + 1$. First we will show that $f(N) \geq (n + 1)N + 1$.

We begin by constructing a $N \times N$ $(0, 1)$ -matrix A . There exists a projective plane P of order n ; so let A be the $N \times N$ matrix whose rows correspond to the points of P and whose columns correspond to the lines of P where the (i, j) entry of A is equal to 1 if and only if the point indexed by i lies on the line indexed by j . Since any two distinct lines have exactly one point in common, the scalar product of any two distinct columns must be 1; hence, S_A does not contain a rectangle. Since each line contains exactly $(n + 1)$ points, $|S_A| = \text{tr}(A^T A) = (n + 1)N$, so $f(N) \geq (n + 1)N + 1$.

Now, suppose A is any $N \times N$ $(0, 1)$ -matrix with $(n + 1)N + 1$ nonzero entries, and let a_i denote the number of 1s in row i . The number of pairs of 1s in row i is $\binom{a_i}{2}$, so the total number of pairs of 1s from each row is $\sum_{i=1}^N \binom{a_i}{2}$.

The number of pairs of distinct column indices is $\binom{N}{2}$. If $\sum_{i=1}^N \binom{a_i}{2} > \binom{N}{2}$, the pigeonhole principle implies that there is a pair of column indices such that there are two distinct rows which have 1s in both of those columns; i.e., S_A contains a rectangle.

To see that $\sum_{i=1}^N \binom{a_i}{2} > \binom{N}{2}$, recall that the Cauchy-Schwarz inequality

gives

$$\left(\sum_{i=1}^N a_i\right)^2 \leq \sum_{i=1}^N a_i^2 \sum_{i=1}^N 1^2. \quad (2)$$

Since $\sum_{i=1}^N a_i = (n+1)N + 1$ by assumption, the bound in (2) gives

$$(n+1)^2 N + 2(n+1) + \frac{1}{N} \leq \sum_{i=1}^N a_i^2. \quad (3)$$

Since $\sum_{i=1}^N a_i^2 = \sum_{i=1}^N a_i(a_i - 1) + \sum_{i=1}^N a_i = 2 \sum_{i=1}^N \binom{a_i}{2} + (n+1)N + 1$, inequality (3) gives

$$N((n+1)^2 - (n+1)) + 2(n+1) + \frac{1}{N} - 1 \leq 2 \sum_{i=1}^N \binom{a_i}{2}. \quad (4)$$

Since $(n+1)^2 - (n+1) = n^2 + n + 1 - 1 = N - 1$, inequality (4) can be rewritten as

$$\frac{N(N-1)}{2} + n + \frac{1}{N} + \frac{1}{2} \leq \sum_{i=1}^N \binom{a_i}{2}, \quad (5)$$

and since $n > 0$, the left hand side of (5) is bound from below by $\binom{N}{2}$, as desired. \square

It is interesting to note that we have equality in (2) just in case all of the a_i are equal; that is, each row and column contain the same number of 1s.

2 Main results

Our main lemma is below, a useful proposition for dealing with asymptotic behavior of functions when some explicit values of the functions are known. A version of this lemma is used in [12], but it is neither proved nor explicitly stated.

Lemma 2. *Suppose g and h are monotonically increasing functions. If a_n is a strictly increasing sequence of positive integers such that*

$$(i) \quad \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = 1;$$

$$(ii) \quad \lim_{n \rightarrow \infty} \frac{h(a_{n+1})}{h(a_n)} = 1; \text{ and}$$

$$(iii) \quad g(a_n) = h(a_n) \text{ for all } n,$$

then $g \sim h$.

Theorem 1 recovers the asymptotic result of Kövari, Sós, and Turán. Theorem 2 strengthens another of their results. The proofs exploit the connection to projective planes, cleaning up the arguments found in [12]. Theorem 3 is an explicit lower bound for $f(k)$, which holds for all k .

Theorem 1. $f(k) \sim k^{3/2}$.

Theorem 2. Let n be a positive integer. If there is a projective plane of order n , then $f(n^2, n^2 + n) = n^2(n + 1) + 1$.

Theorem 3. If $k \in \mathbb{Z}$ with $k \geq 3$, then $f(k) \geq \frac{1}{16}((k + 4)\sqrt{4k - 3} + 5k + 22)$.

3 Proof of Lemma 2

Now we prove Lemma 2.

Proof. Let g and h be monotonically increasing functions. Suppose a_n is a strictly increasing sequence of positive integers such that $\lim_{n \rightarrow \infty} \frac{h(a_{n+1})}{h(a_n)} = 1$ and that $g(a_n) = h(a_n)$ for all n . Let $\varepsilon > 0$. Choose N so that

$$\left| \frac{h(a_{n+1})}{h(a_n)} - 1 \right| < \varepsilon \text{ and } \left| \frac{h(a_n)}{h(a_{n+1})} - 1 \right| < \varepsilon \quad (6)$$

whenever $n > N$. Next, choose m large enough so that for some $n > N$, we have $a_n \leq m \leq a_{n+1}$. Since g is increasing and g and h agree on the sequence a_n , we have

$$h(a_n) = g(a_n) \leq g(m) \leq g(a_{n+1}) = h(a_{n+1}). \quad (7)$$

Since h is monotone increasing, $h(a_n) \leq h(m) \leq h(a_{n+1})$, so we may transform (7) into

$$\frac{h(a_n)}{h(a_{n+1})} \leq \frac{g(m)}{h(m)} \leq \frac{h(a_{n+1})}{h(a_n)}. \quad (8)$$

Subtracting 1 from every term in (8) and taking absolute values gives that either

$$\left| \frac{g(m)}{h(m)} - 1 \right| \leq \left| \frac{h(a_{n+1})}{h(a_n)} - 1 \right| \text{ or } \left| \frac{g(m)}{h(m)} - 1 \right| \leq \left| \frac{h(a_n)}{h(a_{n+1})} - 1 \right|.$$

Without loss of generality, say $\left| \frac{g(m)}{h(m)} - 1 \right| \leq \left| \frac{h(a_{n+1})}{h(a_n)} - 1 \right|$. By (6), we have

$$\left| \frac{g(m)}{h(m)} - 1 \right| < \varepsilon,$$

so $\frac{g}{h} \rightarrow 1$ and $g \sim h$, as desired. □

4 Proof of Theorem 1

Now we prove Theorem 1.

Proof. For a positive integer k , set

$$h(k) = \left(\sqrt{k - \frac{3}{4}} + \frac{1}{2} \right) k + 1.$$

Notice that $h(k) \sim k^{3/2}$ and that $h(n^2 + n + 1) = (n + 1)(n^2 + n + 1) + 1$, so by Lemma 1, we have $f(n^2 + n + 1) = h(n^2 + n + 1)$ whenever there is a projective plane of order n . Since there is a projective plane of order p for every prime p , we have that f and h agree on an infinite sequence of integers a_n for which $\frac{a_{n+1}}{a_n} \rightarrow 1$ (see [18, 9]). Notice that $\frac{h(a_{n+1})}{h(a_n)} \rightarrow 1$, so we may apply Lemma 2 to achieve $f \sim h$, and thus $f \sim k^{3/2}$, as desired. \square

5 Proof of Theorem 2

Proof. Let n be a positive integer such that there is a projective plane of order n . Set $N = n^2 + n + 1$. As in the proof of Lemma 1, we can construct an $N \times N$ matrix A such that $\text{tr}(A^T A) = (n + 1)N$ and that $A^T A$ has only 1s off the main diagonal; hence, the corresponding subset S_A of the $N \times N$ grid has no rectangle.

To construct an $n^2 \times (n^2 + n)$ matrix B from A , we delete the first column of A along with all rows having a 1 in the first column. Since each row and column of A contains exactly $n + 1$ nonzero entries, we have deleted $n + 1$ rows and 1 column. The resulting matrix B is thus an $n^2 \times (n^2 + n)$ matrix. Since $A^T A$ has no entries off the main diagonal greater than 1, $B^T B$ has no entries off the main diagonal greater than 1. Since we have deleted $(n + 1)^2$ nonzero entries from A , we have that

$$|S_B| = (n + 1)N - (n + 1)^2 = (n + 1)(n^2 + n + 1) - (n + 1)^2 = n^2(n + 1),$$

so $f(n^2, n^2 + n) \geq n^2(n + 1) + 1$.

Using the inequality from Reiman (1),

$$f(n^2, n^2 + n) \leq n^2(n + 1) + 1,$$

and hence $f(n^2, n^2 + n) = n^2(n + 1) + 1$, as desired. \square

The structure obtained by taking a projective plane and deleting a line together with all of the points on that line is called an *affine plane*. Our result is stronger than that of the authors in [12], since we need only that there is a projective plane of order n , not that n is a prime number.

6 Proof of Theorem 3

Proof. Suppose k is an integer with $k \geq 3$. There exists a nonnegative integer α such that

$$2^{2\alpha} + 2^\alpha + 1 \leq k \leq 2^{2\alpha+2} + 2^{\alpha+1} + 1. \quad (9)$$

By focusing on the upper bound from (9), this gives $k \leq (2^{\alpha+1} + 1/2)^2 + 3/4$, or

$$\frac{\sqrt{k - 3/4} - 1/2}{2} \leq 2^\alpha. \quad (10)$$

Let $g(n) = (n+1)(n^2 + n + 1) + 1$, and let $h(k) = \frac{\sqrt{k - 3/4} - 1/2}{2}$. Since g is an increasing function, inequality (10) gives

$$g(h(k)) \leq g(2^\alpha). \quad (11)$$

By Lemma 1, we have $g(n) = f(n^2 + n + 1)$ whenever there exists a projective plane of order n . Since there is a projective plane of any prime power order, (11) gives

$$g(h(k)) \leq f(2^{2\alpha} + 2^\alpha + 1). \quad (12)$$

But since f is increasing, the lower bound in (9) gives $g(h(k)) \leq f(k)$, and since $g(h(k)) = \frac{1}{16}((k+4)\sqrt{4k-3} + 5k + 22)$, we have the desired result.

We also note that while $g(h(k)) \sim \frac{1}{8}k^{3/2}$, which is worse than the result in Theorem 1, this lower bound holds for every choice of k , and not just those k for which there exists a projective plane of order k . \square

7 Further Research

Trying to find the exact value of $f(m, n)$ without conditions on m and n (that is, removing the extra hypotheses from the results in [12]) would be attractive, although this problem has been open for years, and likely requires a new idea.

The next attractive direction is to take the approach of the authors in [10], and consider colorings of rectangular grids.

Recall that OBS_c is the collection of $[m] \times [n]$ grids which cannot be colored in c colors without a monochromatic rectangle, but every proper subgrid can be. An open problem from [10] is the *rectangle-free conjecture*: if there exists a rectangle-free subset of $[m] \times [n]$ of size $\lceil mn/c \rceil$, then it is possible to color $[m] \times [n]$ in c colors so there is no monochromatic rectangle. Since the authors in [10] have theorems which depend on the rectangle-free conjecture, resolving this conjecture either in the affirmative or the negative would result in progress for obtaining $|\text{OBS}_c|$ or even OBS_c .

Acknowledgments. The authors wish to thank Jim Marshall for carefully reading a draft of this manuscript. The authors also thank the anonymous referees for their helpful comments made toward improving the paper.

References

- [1] K. Akiyama and C. Suetake. The nonexistence of projective planes of order 12 with a collineation group of order 8. *J. Combin. Des.*, 16(5):411–430, 2008.
- [2] K. Akiyama and C. Suetake. On projective planes of order 12 with a collineation group of order 9. *Australas. J. Combin.*, 43:133–162, 2009.
- [3] M. Axenovich and J. Manske. On monochromatic subsets of a rectangular grid. *Integers*, 8:A21, 14, 2008.
- [4] R. Bacher and S. Eliahou. Extremal binary matrices without constant 2-squares. *J. Comb.*, 1(1, [ISSN 1097-959X on cover]):77–100, 2010.
- [5] R. H. Bruck and H. J. Ryser. The nonexistence of certain finite projective planes. *Canadian J. Math.*, 1:88–93, 1949.
- [6] S. Chowla and H. J. Ryser. Combinatorial problems. *Canadian J. Math.*, 2:93–99, 1950.
- [7] J. Cooper, S. Fenner, and S. Purewal. Monochromatic boxes in colored grids. *SIAM J. Discrete Math.*, 25(3):1054–1068, 2011.
- [8] S. De Winter, F. Lazebnik, and J. Verstraëte. An extremal characterization of projective planes. *Electron. J. Combin.*, 15(1):Research Paper 143, 13, 2008.
- [9] P. Erdős. On a new method in elementary number theory which leads to an elementary proof of the prime number theorem. *Proc. Nat. Acad. Sci. U. S. A.*, 35:374–384, 1949.
- [10] S. Fenner, W. Gasarch, C. Glover, and S. Purewal. Rectangle free coloring of grids. [arXiv:1005.3750](https://arxiv.org/abs/1005.3750) [math.CO], 2010.
- [11] R. L. Graham, B. L. Rothschild, and J. H. Spencer. *Ramsey theory*. Wiley-Interscience Series in Discrete Mathematics and Optimization. John Wiley & Sons Inc., New York, second edition, 1990. A Wiley-Interscience Publication.
- [12] T. Kövari, V. T. Sós, and P. Turán. On a problem of K. Zarankiewicz. *Colloquium Math.*, 3:50–57, 1954.
- [13] C. W. H. Lam, L. Thiel, and S. Swiercz. The nonexistence of finite projective planes of order 10. *Canad. J. Math.*, 41(6):1117–1123, 1989.
- [14] N. S. Mendelsohn. Packing a square lattice with a rectangle-free set of points. *Math. Mag.*, 60(4):229–233, 1987.
- [15] A. R. Prince. Projective planes of order 12 and $PG(3, 3)$. *Discrete Math.*, 208/209:477–483, 1999. *Combinatorics (Assisi, 1996)*.

- [16] A. R. Prince. Ovals in finite projective planes via the representation theory of the symmetric group. In *Finite groups 2003*, pages 283–290. Walter de Gruyter GmbH & Co. KG, Berlin, 2004.
- [17] I. Reiman. Über ein Problem von K. Zarankiewicz. *Acta. Math. Acad. Sci. Hungar.*, 9:269–273, 1958.
- [18] A. Selberg. An elementary proof of the prime-number theorem. *Ann. of Math. (2)*, 50:305–313, 1949.
- [19] C. Suetake. The nonexistence of projective planes of order 12 with a collineation group of order 16. *J. Combin. Theory Ser. A*, 107(1):21–48, 2004.
- [20] E. Szemerédi. On sets of integers containing no k elements in arithmetic progression. *Acta Arith.*, 27:199–245, 1975. Collection of articles in memory of Juriĭ Vladimirovič Linnik.
- [21] B.L. van der Waerden. Beweis einer Baudetchen Vermutung. *Nieuw Arch. Wiskunde*, 15:212–216, 1927.
- [22] K. Zarankiewicz. Problem P101. *Colloq. Math.*, 3:19–30, 1954.