1 LINEAR AND PROJECTIVE BOUNDARY OF NILPOTENT GROUPS

BERNHARD KRÖN, JÖRG LEHNERT, NORBERT SEIFTER, AND ELMAR TEUFL

ABSTRACT. We define a pseudometric on the set of all unbounded subsets of a metric space. The Kolmogorov quotient of this pseudometric space is a complete metric space. The definition of the pseudometric is guided by the principle that two unbounded subsets have distance 0 whenever they stay sublinearly close. Based on this pseudometric we introduce and study a general concept of boundaries of metric spaces. Such a boundary is the closure of a subset in the Kolmogorov quotient determined by an arbitrarily chosen family of unbounded subsets.

Our interest lies in those boundaries which we get by choosing unbounded cyclic sub-(semi)groups of a finitely generated group (or more general of a compactly generated, locally compact Hausdorff group). We show that these boundaries are quasi-isometric invariants and determine them in the case of nilpotent groups as a disjoint union of certain spheres (or projective spaces).

In addition we apply this concept to vertex-transitive graphs with polynomial growth and to random walks on nilpotent groups.

CONTENTS

4	1.	Introduction	1
5	2.	General construction	3
6	3.	Quasi-isometries	8
7	4.	Boundary at infinity and angular metric in a $CAT(0)$ space	9
8	5.	Boundaries of groups	10
9	6.	Boundaries of nilpotent groups	13
10	7.	Boundaries of vertex-transitive graphs with polynomial growth	15
11	8.	Attaching the boundary	20
12	9.	Random walks on nilpotent groups	23
13	Ap	pendix A. Compactly generated groups	24
14	Ap	pendix B. Nilpotent Lie groups	27
15	Re	ferences	33

16

1. Introduction

There are numerous boundary notions of graphs, groups, manifolds, metric spaces and other geometric objects. The literature on the subject is extensive and boundary notions proved to be a useful tool in studying the underlying space. An early instance is the theory of ends which was developed in the first half of the twentieth century by Freudenthal (see e.g. [Fre42]) and others. Various geometric ideas were used to refine the notion of ends:

In 1973 Eberlein and O'Neill [EO73] constructed the boundary at infinity of a CAT(0) space by considering equivalence classes of non-compact geodesic rays. The equivalence notion of geodesic rays uses the natural parametrization, i.e. two geodesic rays are equivalent if they stay at bounded distance as the parameter tends to ∞ . A different description is given by

2

Date: 15 September 2013.

²⁰¹⁰ Mathematics Subject Classification. 20F65 (54E35,20F18,22E25,05C63).

Key words and phrases. metric spaces, boundaries, nilpotent groups.

 $\mathbf{2}$

Gromov in [Gro81] which uses an embedding into the set of continuous functions relying on the metric only.

In graph theory in the 1990s Jung [Jun93] and Jung, Niemayer [JN95] introduced a re-28 finement of ends of graphs called b-fibers and d-fibers, respectively. The basic idea behind 29 fibers is to consider points at infinity as equivalence classes of rays (infinite paths) which 30 stay at bounded distance "up to linear reparametrization". In 2005 Bonnington, Richter and 31 Watkins [BRW07] modified this concept by considering rays as equivalent whenever they stay 32 at sublinear distance "up to linear reparametrization". They were able to use this concept 33 to prove some nice results on infinite planar graphs, but the boundary, whose elements have 34 been called "bundles", was not topologized and not considered for groups or vertex-transitive 35 graphs. 36

Another instance, where the concept of staying at sublinear distance is used, is given 37 by Kaimanovich in [Kai91, Theorem 5.5]. The so-called "ray approximation" is used to 38 determine, whether a given probability space is the Poisson boundary of a random walk on a 39 countable group G defined by a probability measure μ on G. A proposal space (B, λ) is the 40 Poisson boundary of (G, μ) , if compatibility conditions between μ and λ hold and if there 41 exist measureable "projections" $\pi_n: B \to G$, such that almost every trajectory (g_1, g_2, \dots) 42 stays sublinear close to $(\pi_1(g_\infty), \pi_2(g_\infty), \dots)$, where g_∞ is the limit point of the trajectory 43 (g_1, g_2, \dots) in *B*. 44

In these examples the "parametrization" of rays or sequences is used in the definition of staying (sublinearly) close. In the following we relax this and work with general subsets and not only with rays or sequences. Let (X, d) be a metric space, let $o \in X$ be a fixed reference point and denote by B(x, r) the closed ball in (X, d) with center x and radius r. If R, S are two unbounded subsets of X, their distance t(R, S) is defined to be the square root of the infimum over all $\alpha \geq 0$, such that

$$S \subseteq \bigcup_{x \in R} B(x, \alpha d(o, x) + a) \qquad \text{and} \qquad R \subseteq \bigcup_{y \in S} B(y, \alpha d(o, y) + a)$$

for some $a \ge 0$. The sets R, S are sublinearly close, if t(R, S) = 0. We show that the set of all unbounded subsets of (X, d) equipped with the distance t is a pseudometric space, whose Kolmogorov quotient is a complete metric space (Proposition 2.6 and Theorem 2.9). Given some family \mathcal{E} of unbounded subsets the associated "boundary" of X is the closure of all equivalence classes which contain an element from \mathcal{E} in the Kolmogorov quotient. Interesting families of unbounded subsets include: geodesics, horoballs, cyclic sub(semi)groups (in the case of groups), one-parameter sub(semi)groups (in the case of topological groups).

We mainly focus on the group case. Let G be a finitely generated (or more general com-58 pactly generated, locally compact Hausdorff) group and let d be a word metric on G. If the 59 family \mathcal{E} is given by all unbounded cyclic subsemigroups or by all unbounded cyclic sub-60 groups, we call the associated boundary *linear boundary* in the former case and *projective* 61 boundary in latter case. We prove that these two boundaries are quasi-isometric invariants 62 (Lemma 5.3). In our main result (Theorem 6.1) we identify the linear and projective bound-63 ary for nilpotent groups. Let G be either a connected, nilpotent Lie group or a finitely 64 generated, nilpotent group with descending central series 65

$$G = \gamma_1(G) \supseteq \gamma_2(G) \supseteq \cdots \supseteq \gamma_c(G) \supseteq \gamma_{c+1}(G) = 1.$$

66 Let $\nu(i)$ denote the compact-free dimension or the torsion-free rank of the commutative 67 group $\gamma_i(G)/\gamma_{i+1}(G)$. Then the linear boundary is homeomorphic to the disjoint union of *c* 68 spheres

$$\mathbb{S}^{\nu(1)-1}
in \mathbb{S}^{\nu(2)-1}
in \dots
in \mathbb{S}^{\nu(c)-1}$$

- and the projective boundary is homeomorphic to the disjoint union of projective spaces $\mathbb{P}^{\nu(1)-1} \uplus \mathbb{P}^{\nu(2)-1} \uplus \cdots \uplus \mathbb{P}^{\nu(c)-1}.$
- ⁷⁰ Here \mathbb{S}^d is the *d*-dimensional sphere and \mathbb{P}^d is the *d*-dimensional projective space.

The following facts about the boundary notion introduced above must be stressed: Com-71 pact elements of a group G do not contribute to the boundaries $\mathcal{P}G$ and $\mathcal{L}G$. Hence, whenever 72 G only contains compact elements, these boundaries are empty. In particular, this means in 73 the discrete case, that $\mathcal{P}G$ and $\mathcal{L}G$ are empty for torsion groups. We also emphasize that we 74 compare unbounded sets using the distance function t and not sequences or rays using their 75 parametrizations, as it is e.g. done in [BRW07]. For instance, two sequences or rays might 76 be distant in the sense of [BRW07] using parametrizations $a_n = n$ and $b_n = \sqrt{n}$, respectively, 77 although the underlying unbounded sets are the same hence sublinearly close. 78

79 The paper is organized as follows:

- The general framework for metric spaces is studied in Section 2. The distance t and some auxiliary quantities are introduced and several basic results are proved. For instance we show that the distance t has all properties stated above.
 - In Section 3 we investigate the relationship to quasi-isometries. It is shown that the distance t is preserved up to bi-Lipschitz-equivalence under quasi-isometries of the underlying space (Theorem 3.3).
 - In Section 4 we show that the boundary at infinity of a complete CAT(0) space equipped with the angular metric can be obtained by the boundary construction outlined above using the set of unbounded geodesics up to bi-Hölder equivalence.
- In Section 5 we apply this concept to groups using unbounded cyclic sub(semi)groups as families of unbounded subsets. Some general results are obtained and the case of abelian groups is discussed in detail. In the latter case the projective boundary is homeomorphic to a projective space and the linear boundary is homeomorphic to a sphere.
 - Section 6 is devoted to the formulation and proof of the main result (Theorem 6.1). Most technical parts of the proof are deferred to Appendix B.
- Section 7 discusses the situation for graphs. The projective boundary of a graph is defined by the above procedure, using the family of unbounded orbits generated by cyclic subgroups of the automorphism group and the linear boundary is defined analogously. For connected, vertex-transitive graphs with polynomial volume growth we obtain the same description of the projective (respectively linear) boundary as in the case of nilpotent groups (Corollary 7.12).
- In Section 8 we construct a topology on the disjoint union of the base space Xand some boundary which is obtained by the construction above. The definition is reminiscent of the cone topology of the boundary at infinity of CAT(0) spaces. The subspace topology on X of this topology is always induced by the metric d, but the subspace topology on the boundary is neither induced by t nor Hausdorff in general. We discuss criteria (Lemma 8.2 and Proposition 8.3) which guarantee both: the subspace topology on the boundary is Hausdorff and induced by t.
 - In Section 9 we show that every boundary point in the linear boundary of a nilpotent Lie group is obtained as a limit of a random walk with drift and vice versa.
- Appendix A collects some known results on compactly generated groups and word metrics which are used in the previous sections.
 - Appendix B mostly contains the technical lemmas used in the proof of Theorem 6.1 and the necessary notions from Lie theory.

115

109

110

113

114

83

84

85

86

87

88

94

95

2. General construction

Let (X, d) be a metric space. We write \mathcal{U} to denote the family of unbounded subsets of (X, d). The closed and open ball with center $x \in X$ and radius $r \geq 0$ in (X, d) are denoted by

$$B(x,r) = \{ y \in X : d(y,x) \le r \} \quad \text{and} \quad U(x,r) = \{ y \in X : d(y,x) < r \},\$$

respectively. Let o be a fixed reference point, let $R \subseteq X$, and let α and a be nonnegative real numbers. We set

$$\alpha R + a = \bigcup_{x \in R} B(x, \alpha d(o, x) + a)$$

121 and write αR instead of $\alpha R + 0$.

122 Remark. The notation $\alpha R + a$ is unusual, but turns out to be convenient for computations 123 involving sets of this form. Mostly this notation will be used if X is a metric space, so no 124 confusion should occur. However, if X is a linear space too, $\alpha R + a$ will always be used in 125 the above meaning and never means a linearly scaled and translated set. Furthermore, it 126 should be the stressed that 0R = R and

$$0R + a = \bigcup_{x \in R} B(x, a),$$

127 which is often called *a*-neighborhood of R or generalized ball of radius a around R.

128 Lemma 2.1. Let $R \in \mathcal{U}$ and $\alpha > 1$. Then $\alpha R = X$.

129 Proof. Let x be any point in X. Since R is unbounded, there is an element $y \in R$ such that 130 $(\alpha - 1)d(o, y) \ge d(o, x)$. Hence

$$d(x,y) \le d(x,o) + d(o,y) \le (\alpha - 1)d(o,y) + d(o,y) = \alpha d(o,y),$$

131 and $x \in B(y, \alpha d(o, y)) \subseteq \alpha R$.

Lemma 2.2. Let R, S, and T be subsets of X. If $T \subseteq \beta S + b$ and $S \subseteq \alpha R + a$ then 133 $T \subseteq (\alpha + \alpha \beta + \beta)R + \beta a + a + b$.

134 Proof. Let z be in T. Since the sets $\alpha R + a$ and $\beta S + b$ are defined as unions of balls, z 135 is in $B(y,\beta d(o,y) + b)$ for some $y \in S$ and y is in $B(x,\alpha d(o,x) + a)$ for some $x \in R$. Set 136 c = d(o,x). Then $d(x,y) \leq \alpha c + a$. By the triangle inequality,

$$d(o, y) \le d(o, x) + d(x, y) \le (\alpha + 1)c + a.$$

137 Hence

$$d(y,z) \le \beta d(o,y) + b \le (\alpha\beta + \beta)c + \beta a + b.$$

138 Finally,

$$d(x,z) \le d(x,y) + d(y,z) \le (\alpha + \alpha\beta + \beta)c + \beta a + a + b.$$

139 This means that $z \in (\alpha + \alpha\beta + \beta)R + \beta a + a + b$.

Definition 2.3. For two subsets $R, S \subseteq X$ let $s^+(R, S)$ be the infimum of all $\alpha \ge 0$ such that $S \subseteq \alpha R + a$ for some $a \ge 0$. Set

$$s(R, S) = \max\{s^+(R, S), s^+(S, R)\}$$

and $t(R,S) = \sqrt{s(R,S)}$. If $R, S \in \mathcal{U}$ and s(R,S) = 0 then R and S are called *linearly equivalent* and we write $R \sim S$.

Remark. The functions s^+ , s, t depend on the metric space (X, d). In order to emphasize the underlying metric space (X, d) we write s_X^+ or $s_{(X,d)}^+$ and analogously for s and t. Similarly, we write \mathcal{U}_X or $\mathcal{U}_{(X,d)}$ instead of \mathcal{U} , if it is necessary to specify the metric space.

Lemma 2.4. Let R, S be two subsets of X. Then $s^+(R, S)$ and therefore s(R, S) and t(R, S)do not depend on the reference point o in X.

Proof. Let $o, p \in X$ and set c = d(o, p). We write $s_o^+(R, S)$ in order to emphasize the reference point o. Furthermore, write $C_o(R, \alpha, a)$ to denote the set $\alpha R + a$ with respect to the reference point o. For $\alpha > s_o^+(R, S)$ there is a number a > 0 such that $S \subseteq C_o(R, \alpha, a)$. Hence for $y \in S$ we can find a point $x \in R$ such that $d(y, x) \leq \alpha d(o, x) + a$. The triangle inequality implies that

$$d(y,x) \le \alpha(d(p,x) + d(p,o)) + a = \alpha d(p,x) + \alpha c + a.$$

4

Therefore $S \subseteq C_p(R, \alpha, \alpha c + a)$ and thus $s_p^+(R, S) \leq s_o^+(R, S)$. The reversed inequality is obtained by changing the rôle of o and p.

Lemma 2.5. Let $R, S \in \mathcal{U}$. Then $s^+(R, S)$ is the infimum of all $\alpha \ge 0$ such that $S \setminus U(o, r) \subseteq \alpha R$ for some $r \ge 0$.

Proof. We write $\sigma^+(R,S)$ to denote the infimum of all $\alpha \geq 0$ such that $S \setminus U(o,r) \subseteq \alpha R$ for 158 some $r \ge 0$. First we show that $s^+(R, S) \le \sigma^+(R, S)$. Assume that $\alpha > \sigma^+(R, S)$ and $r \ge 0$ 159 such that $S \setminus U(o, r) \subseteq \alpha R$. Set a = r + d(o, R), where $d(o, R) = \inf\{d(o, x) : x \in R\}$. Then, 160 by triangle inequality, $S \subseteq \alpha R + a$. Therefore $s^+(R, S) \leq \alpha$ and hence $s^+(R, S) \leq \sigma^+(R, S)$. 161 Now we prove the reversed inequality: Let $\alpha > s^+(R, S)$ and set $\varepsilon = \frac{1}{2}(\alpha - s^+(R, S)) > 0$. 162 Then, by definition of $s^+(R, S)$, there exists a constant $a \ge 0$ such that $S \subseteq (\alpha - \varepsilon)R + a$. 163 We claim that $S \setminus U(o, r) \subseteq \alpha R$ holds for $r = \frac{a}{\varepsilon}(1 + \alpha)$. Let $y \in S$. Then there is a $x \in R$ 164 with $d(y,x) \leq (\alpha - \varepsilon)d(o,x) + a$. Using the triangle inequality yields 165

$$d(o,y) \leq d(o,x) + d(y,x) \leq d(o,x) + (\alpha - \varepsilon)d(o,x) + a \leq (1 + \alpha - \varepsilon)d(o,x) + a,$$

166 which implies

$$d(o, x) \ge \frac{d(o, y) - a}{1 + \alpha - \varepsilon}$$

167 If $d(o, y) \ge r$ then we obtain

$$a \leq \varepsilon \cdot \frac{d(o, y) - a}{1 + \alpha - \varepsilon} \leq \varepsilon d(o, x)$$

168 and

177

$$d(y,x) \le (\alpha - \varepsilon)d(o,x) + a \le (\alpha - \varepsilon)d(o,x) + \varepsilon d(o,x) = \alpha d(o,y).$$

169 Therefore $S \setminus U(o, r) \subseteq \alpha R$ and $\sigma^+(R, S) \leq s^+(R, S)$.

Proposition 2.6. The function s^+ is a premetric on \mathcal{U} satisfying a weak form of the triangle inequality, i.e. if R, S, T are unbounded subsets of X, then

172 • $s^+(R,S) \in [0,1] \text{ and } s^+(R,R) = 0,$

173 • $s^+(R,T) \le s^+(R,S) + s^+(R,S)s^+(S,T) + s^+(S,T).$

Similarly, s is a symmetric premetric on \mathcal{U} satisfying the same weak triangle inequality, i.e. $s(R,S) \in [0,1]$ and s(R,R) = 0,

176 • s(R, S) = s(S, R),

•
$$s(R,T) \le s(R,S) + s(R,S)s(S,T) + s(S,T)$$

178 Finally, t is a pseudometric on \mathcal{U} , i.e.

179 • $t(R, S) \in [0, 1]$ and t(R, R) = 0,

180 • t(R, S) = t(S, R),

181 • $t(R,T) \le t(R,S) + t(S,T).$

182 *Proof.* The statements for s^+ and s follow from the definition and from the Lemmas 2.1 and 183 2.2.

It remains to show that t satisfies the triangle inequality. Let R, S be unbounded subsets of X. Then

$$\begin{aligned} s(R,T) &\leq s(R,S) + s(R,S)s(S,T) + s(S,T) \\ &\leq s(R,S) + 2\sqrt{s(R,S)s(S,T)} + s(S,T) \end{aligned}$$

186 which implies

$$t(R,T) = \sqrt{s(R,T)} \le \sqrt{s(R,S)} + \sqrt{s(S,T)} = t(R,S) + t(S,T).$$

187 Corollary 2.7. Assume that R, S, T are unbounded subsets of X. If $s^+(S,T) = 0$ then 188 $s^+(R,T) \leq s^+(R,S)$ and $s^+(S,R) \leq s^+(T,R)$. Therefore, if $S \sim T$, then $s^+(R,S) =$ 189 $s^+(R,T), s^+(S,R) = s^+(T,R)$, and s(R,S) = s(R,T), t(R,S) = t(R,T).

190 Proof. Using Proposition 2.6 and $s^+(S,T) = 0$ we get

$$s^+(R,T) \le s^+(R,S) + s^+(R,S)s^+(S,T) + s^+(S,T) = s^+(R,S)$$

191 and

192

$$s^+(S,R) \le s^+(S,T) + s^+(S,T)s^+(T,R) + s^+(T,R) = s^+(T,R).$$

The remaining claims follow, since $S \sim T$ implies $s^+(S,T) = s^+(T,S) = 0.$

193 Corollary 2.8. Linear equivalence is an equivalence relation on unbounded subsets and the

194 functions s^+ , s, t are well-defined on the quotient space \mathcal{U}/\sim .

Proof. Reflexivity and symmetry follow immediately from the definition. Transitivity follows from Corollary 2.7. Let R_1, R_2, S_1, S_2 be unbounded subsets and suppose $s(R_1, R_2) =$ $s(S_1, S_2) = 0$. Corollary 2.7 implies that $s^+(R_1, S_1) = s^+(R_2, S_1) = s^+(R_2, S_2)$, whence s^+ and therefore s, t are well-defined on equivalence classes.

Theorem 2.9. $(\mathcal{U}/\sim, t)$ is a complete metric space.

200 Proof. By Proposition 2.6 and Corollary 2.8 $(\mathcal{U}/\sim, t)$ is a metric space. It remains to prove 201 that it is also complete.

Let $(\xi_n)_{n\geq 0}$ be a Cauchy sequence in \mathcal{U}/\sim . Without loss of generality we may assume that $s(\xi_n, \xi_m) \leq 1/2$ for all n, m. Choose representatives $R_n \in \xi_n$. Then for any $\varepsilon > 0$ there is an index N such that $s(R_n, R_m) < \varepsilon$ for $n, m \geq N$. Therefore there exists a function $\varepsilon^* \colon \mathbb{N} \to (0, 1/2]$ such that ε^* is decreasing, $\varepsilon^*(n) \to 0$ as $n \to \infty$, and $s(R_m, R_n) < \varepsilon^*(m)$ for $m \leq n$.

According to Lemma 2.5 there are $r(m, n) \ge 0$, for $m \le n$, such that

$$R_n \setminus U(o, r(m, n)) \subseteq \varepsilon^*(m) R_m$$
 and $R_m \setminus U(o, r(m, n)) \subseteq \varepsilon^*(m) R_n$.

Hence there is an increasing function $r^* \colon \mathbb{N} \to [0, \infty)$ such that $r^*(n) \geq r(m, n)$ for $m \leq n$. Applying Lemma 2.5 to $R_m \setminus U(o, r^*(n))$ and $R_n \setminus U(o, r^*(n))$ for $m \leq n$ implies that there are $q(m, n) \geq 0$ such that

$$R_n \setminus U(o, q(m, n)) \subseteq \varepsilon^*(m) (R_m \setminus U(o, r^*(n))),$$

$$R_m \setminus U(o, q(m, n)) \subseteq \varepsilon^*(m) (R_n \setminus U(o, r^*(n))).$$

Thus there is an increasing function $q^* \colon \mathbb{N} \to [0,\infty)$ such that $q^*(n) \to \infty$ as $n \to \infty$ and $q^*(n) \ge q(m,n)$ for $m \le n$.

Let $x \in R_m$ with $q^*(n) \le d(o, x) < q^*(n+1)$ for some $n \ge m$. Then there is a $y \in R_n$ such that

$$d(o, y) \ge r^*(n)$$
 and $d(x, y) \le \varepsilon^*(m)d(o, y)$

Using the triangle inequality and $\varepsilon^*(m) \leq 1/2$ we get

$$d(o, y) \le d(o, x) + d(x, y) \le d(o, x) + \varepsilon^*(m)d(o, y) \le d(o, x) + d(o, y)/2$$

216 and

(1)
$$d(o,y) \le 2d(o,x) < 2q^*(n+1).$$

217 We write x^* to denote this element y and define the set S by

$$S = \bigcup_{m \ge 1} \{ x^* : x \in R_m, \, d(o, x) \ge q^*(m) \}.$$

Then S is an unbounded subset of X. Note that if $x \in S$ and $d(o, x) \ge 2q^*(m)$ for some m then $x \in R_n$ for some $n \ge m$ due to the estimate in (1). We claim that $s(S, R_m) \le \varepsilon^*(m)$ for $m \ge 1$.

• Let x be an element of R_m with $d(o, x) \ge q^*(m)$. Then, by construction of S, there is a $y \in S$ with $d(x, y) \le \varepsilon^*(m)d(o, y)$. This implies

$$R_m \setminus U(o, q^*(m)) \subseteq \varepsilon^*(m)S$$

 $\mathbf{6}$

• Let x be an element of S with $d(o, x) \ge 2q^*(m)$. Then $x \in R_n$ for some $n \ge m$. This implies the lower bound $d(o, x) \ge r^*(n)$. By definition of r^* there is a $y \in R_m$ such that $d(x, y) \le \varepsilon^*(m)d(o, y)$. Hence

 $S \setminus U(o, 2q^*(m)) \subseteq \varepsilon^*(m)R_m.$

²²⁶ This implies the claim. Let ζ be the equivalence class of S. Then

$$s(\xi_m, \zeta) = s(R_m, S) \le \varepsilon^*(m)$$

for $m \ge 1$. Therefore ξ_m converges to ζ proving Cauchy completeness.

Definition 2.10. We call t angle metric of unbounded sets (see Example 2.11).

If Ξ is a subset of \mathcal{U}/\sim , we write $\mathsf{cl}(\Xi)$ to denote the closure of Ξ in the metric space $(\mathcal{U}/\sim, t)$. Let $\mathcal{E} \subseteq \mathcal{U}$ be a family of unbounded subsets of (X, d). Define \mathcal{E}/\sim to be the set of equivalence classes in \mathcal{U}/\sim which contain at least one element from \mathcal{E} , this is

$$\mathcal{E}/{\sim} = \{[R] : R \in \mathcal{E}\} \subseteq \mathcal{U}/{\sim}$$

where [R] is the equivalence class of R with respect to linear equivalence \sim . Note that cl(\mathcal{E}/\sim) is a well-defined subset of \mathcal{U}/\sim which is closed and hence Cauchy complete. Thus up to isometry (cl(\mathcal{E}/\sim), t) is the Cauchy completion of (\mathcal{E}/\sim , t).

Remark. The definition of the set \mathcal{E}/\sim depends on the underlying metric space (X, d). However, no confusion should occur, since the underlying metric space will be clear from the context. Moreover, the above definition of \mathcal{E}/\sim is somewhat unusual, since $\mathcal{E}/\sim \subseteq \mathcal{U}/\sim$. The reason for this definition is that we will use topological notions of $(\mathcal{U}/\sim, t)$ for the subset \mathcal{E}/\sim . Furthermore, note that, if $\sim_{\mathcal{E}}$ denotes the restriction of \sim to the set \mathcal{E} then

$$\mathcal{E}/\sim \to \mathcal{E}/\sim_{\mathcal{E}}, \quad \zeta \mapsto \zeta \cap \mathcal{E}$$

240 is a canonical bijection.

Example 2.11. Consider \mathbb{R}^n equipped with the usual ℓ^2 -metric. For a nonzero vector 242 $x \in \mathbb{R}^n$ let L_x denote the line $\{\lambda x : \lambda \in \mathbb{R}\}$ and H_x the half-line $\{\lambda x : \lambda \geq 0\}$. Set 243 $\mathcal{L} = \{L_x : x \in \mathbb{R}^n, x \neq 0\}$ and $\mathcal{H} = \{H_x : x \in \mathbb{R}^n, x \neq 0\}$. Then $\mathsf{cl}(\mathcal{L}/\sim)$ is the projective 244 space \mathbb{P}^{n-1} and $\mathsf{cl}(\mathcal{H}/\sim)$ is the sphere \mathbb{S}^{n-1} . If $x, y \in \mathbb{R}^n \setminus \{0\}$ then

$$s(L_x, L_y) = \sin(\angle(L_x, L_y)) \quad \text{and} \quad s(H_x, H_y) = \sin(\min\{\frac{1}{2}\pi, \angle(H_x, H_y)\})$$

where $\angle(L_x, L_y)$ is the smaller angle between the lines L_x and L_z and $\angle(H_x, H_y)$ is the angle between the half-lines H_x, H_y .

The following examples show that the function s is not always a metric and that geodesics do not always yield a nice structure.

Example 2.12. Consider the 2-dimensional space \mathbb{R}^2 with ℓ^1 -metric d_1 . Let $x_1 = (1,0)$, 250 $x_2 = (2,1)$, and $x_3 = (1,1)$. Set $L_i = \{\lambda x_i : \lambda \in \mathbb{R}\}$ for $i \in \{1,2,3\}$. Then

$$s(L_1, L_2) = \frac{1}{2}, \qquad s(L_2, L_3) = \frac{1}{3}, \qquad s(L_1, L_3) = 1,$$

²⁵¹ and the triangle inequality is not satisfied.

Example 2.13. Consider the metric space (\mathbb{Z}^2, d_1) , where d_1 is the ℓ^1 -metric. In this discrete 252 setting a geodesic ray is an infinite sequence $(x_0, x_1, ...)$ in \mathbb{Z}^2 such that $d(x_i, x_j) = |i - j|$. 253 Let \mathcal{G} be the family of all geodesic rays emanating from the origin. Furthermore, let \mathcal{E} 254 be the family of all sets $\{nx : n \in \mathbb{N}_0\}$ for $x \in \mathbb{Z}^2, x \neq 0$. Then the space $\mathsf{cl}(\mathcal{G}/\sim)$ 255 contains much more elements than $cl(\mathcal{E}/\sim)$. To see this set $x_{2n} = (2^n - 1, 2^n - 1)$ and 256 $x_{2n+1} = (2^{n+1} - 1, 2^n - 1)$ for $n \in \mathbb{N}_0$. Join x_m and $x_{m+1}, m \in \mathbb{N}_0$, by a geodesic path and 257 let R denote the ray consisting of the union of these finite geodesic paths. Obviously R is a 258 geodesic ray and there is some $\varepsilon > 0$ such that $s(R, S) \ge \varepsilon$ for all $S \in \mathcal{E}$. 259

3. Quasi-isometries

Definition 3.1. Let (X, d_X) and (Y, d_Y) be metric spaces and let q > 0. A function $f: X \to Y$ is called a q-quasi-isometry if

$$q^{-1}d_X(x,x') - q \le d_Y(f(x), f(x')) \le qd_X(x,x) + q$$

for all $x, x' \in X$ and such that every closed ball in Y with radius q contains an element of f(X). We say that two metrics d_1 and d_2 on X are quasi-isometrically equivalent, if the identity is a quasi-isometry from (X, d_1) to (X, d_2) .

Lemma 3.2. Let $f: X \to Y$ be a q-quasi-isometry of the metric spaces (X, d_X) and (Y, d_Y) . Let R, S be unbounded subsets of X. Then f(R), f(S) are unbounded subsets of Y and

$$q^{-2}s_X^+(R,S) \le s_Y^+(f(R),f(S)) \le q^2s_X^+(R,S)$$

268 Proof. We fix reference points o and f(o) in X and Y, respectively. First of all note that

$$q^{-1}d_X(x,x') - q \le d_Y(f(x), f(x')) \le qd_X(x,x') + q$$

269 implies

$$q^{-1}d_Y(f(x), f(x')) - 1 \le d_X(x, x') \le qd_Y(f(x), f(x')) + q^2$$

270 for all $x, x' \in X$.

Let $\alpha > s_X^+(R, S)$. Then there is a number a such that $S \subseteq \alpha R + a$. Hence for $x' \in S$ there is a point $x \in R$ with $d_X(x', x) \leq \alpha d_X(o, x) + a$. Since f is a q-quasi-isometry, we get $d_X(o, x) \leq q d_Y(f(o), f(x)) + q^2$. This implies

$$d_Y(f(x'), f(x)) \le q d_X(x', x) + q \le q \alpha d_X(o, x) + q a + q$$
$$\le q^2 \alpha d_Y(f(o), f(x)) + q^3 \alpha + q a + q,$$

274 proving that

$$f(S) \subseteq q^2 \alpha f(R) + q^3 \alpha + qa + q$$

275 holds. Thus $s_Y^+(f(R), f(S)) \le q^2 s_X^+(R, S)$.

If $s_X^+(R,S) = 0$ then $s_Y^+(f(R), f(S)) \ge q^{-2}s_X^+(R,S)$ trivially holds. Hence we assume that $s_X^+(R,S) > 0$. Then, for $\alpha < s_X(R,S)$, $S \subseteq \alpha R + a$ fails to be true for all $a \ge 0$. Hence for every $a \ge 0$ there exists a point $x' \in S$ which is not contained in $\alpha R + a$. Thus $d_X(x',x) > \alpha d(o,x) + a$ for all $x \in R$. This implies

$$d_Y(f(x'), f(x)) \ge q^{-1} d_X(x', x) - q > q^{-1} \alpha d_X(o, x) + q^{-1} a - q$$

$$\ge q^{-2} \alpha d_Y(f(o), f(x)) + q^{-1} (a - \alpha) - q.$$

Thus f(x') is not contained in $q^{-2}\alpha f(R) + q^{-1}(a-\alpha) - q$. Since $a \ge 0$ was arbitrary, this means that $s_Y^+(f(R), f(S)) \ge q^{-2} s_X^+(R, S)$.

Theorem 3.3. Let $f: X \to Y$ be a q-quasi-isometry of the metric spaces (X, d_X) and (Y, d_Y). Then f induces a bijection $f: \mathcal{U}_X/\sim \to \mathcal{U}_Y/\sim$ which is bi-Lipschitz continuous:

$$q^{-1}t_X(\zeta,\xi) \le t_Y(f(\zeta),f(\xi)) \le qt_X(\zeta,\xi)$$

for all $\zeta, \xi \in \mathcal{U}_X/\sim$. In particular, if \mathcal{E} is a family of unbounded subsets in X, then $f(\mathcal{E}/\sim) = f(\mathcal{E})/\sim$ and $f(cl(\mathcal{E}/\sim)) = cl(f(\mathcal{E})/\sim)$.

Proof. Of course $f(\mathcal{U}_X)$ is a subset of \mathcal{U}_Y . By Lemma 3.2 the function $f: \mathcal{U}_X/\sim \to \mathcal{U}_Y/\sim$ which maps the equivalence class of an unbounded $R \subseteq X$ to the equivalence class of f(R)is well-defined, one-to-one, and satisfies

$$q^{-1}t_X(\zeta,\xi) \le t_Y(f(\zeta),f(\xi)) \le qt_X(\zeta,\xi)$$

for all $\zeta, \xi \in \mathcal{U}_X/\sim$. Thus it remains to show that f is also onto. Let S be an unbounded subset of Y. Since f is a q-quasi-isometry, the set $R = f^{-1}((0S+q) \cap f(X))$ is an unbounded subset of X and $f(R) = (0S+q) \cap f(X) \sim S$.

8

4. Boundary at infinity and angular metric in a CAT(0) space

Let us recall the definitions of the boundary at infinity and the angular metric in a 293 CAT(0) space. For more details we refer to the book of Bridson and Haefliger [BH99], 294 see especially Chapter II.8 and Chapter II.9 therein. A *geodesic ray* in a metric space (X, d)295 is a curve $c: [0,\infty) \to X$ such that d(c(x),c(y)) = |x-y| for all $x,y \ge 0$. The boundary 296 at infinity ∂X of X is defined to be the set of equivalence classes of geodesic rays, where 297 geodesic rays c, c' are equivalent whenever they stay at bounded distance, that is, if there is 298 a constant K, such that $d(c(x), c'(x)) \leq K$ for all $x \in [0, \infty)$. In the sequel we assume that 299 X is a complete CAT(0) space. 300

For each point p in X and ξ in ∂X there is precisely one geodesic ray belonging to ξ which emanates from p. Then $\angle_p(\xi, \zeta)$ for $\xi, \zeta \in \partial X$ is defined to be the angle at p between the uniquely determined rays in ξ and ζ which emanate from p. The angle between ξ and ζ is defined by

$$\angle(\xi,\zeta) = \sup\{\angle_p(\xi,\zeta) : p \in X\}.$$

This yields a metric on ∂X called *angular metric* and $(\partial X, \angle)$ is a complete metric space. For our purposes the following description of the angular metric is useful. Fix a reference point o in X. If $\xi \in \partial X$, we write c_{ξ} for the uniquely determined geodesic ray in ξ which emanates from o and R_{ξ} for the image of c_{ξ} in X, i.e. $R_{\xi} = c_{\xi}([0, \infty))$. Then, see [BH99, Proposition 9.8 (4)],

$$2\sin\left(\frac{1}{2}\angle(\xi,\zeta)\right) = \lim_{x \to \infty} \frac{1}{x} d(c_{\xi}(x), c_{\zeta}(x)).$$

310 Lemma 4.1. Let ξ, ζ be elements in ∂X . Then

$$s(R_{\xi}, R_{\zeta}) \le 2\sin\left(\frac{1}{2}\angle(\xi, \zeta)\right) \le 4s(R_{\xi}, R_{\zeta}).$$

³¹¹ Proof. Note that $d(o, c_{\xi}(x)) = d(o, c_{\zeta}(x)) = x$ for all $x \in [0, \infty)$, since $c_{\xi}(0) = c_{\zeta}(0) = o$. Sup-³¹² pose that $\alpha > 2\sin(\frac{1}{2}\angle(\xi,\zeta))$. Then there exists a constant $a \ge 0$, such that $d(c_{\xi}(x), c_{\zeta}(x)) \le \alpha x$ for all $x \ge a$. This implies that

$$R_{\xi} \subseteq \alpha R_{\zeta} + a$$
 and $R_{\zeta} \subseteq \alpha R_{\xi} + a$

Therefore $s(R_{\xi}, R_{\zeta}) \leq \alpha$ which yields the lower bound.

If $s(R_{\xi}, R_{\zeta}) \geq \frac{1}{2}$ then the upper bound is trivially true. Hence assume that $s(R_{\xi}, R_{\zeta}) < \frac{1}{2}$ and fix some α , such that $s(R_{\xi}, R_{\zeta}) < \alpha \leq \frac{1}{2}$. By Lemma 2.5 there is a constant $r \geq 0$, such that $R_{\zeta} \setminus U(o, r) \subseteq \alpha R_{\xi}$. Hence, for any $x \geq r$, there is a $y = y(x) \geq 0$, such that

$$d(c_{\zeta}(x), c_{\xi}(y)) \le \alpha d(o, c_{\xi}(y)) = \alpha y.$$

Using the triangle inequality the estimate above yields $y \le x + \alpha y$ and $x \le y + \alpha y$. It follows that $|y - x| \le \alpha y$ and $y \le 2x$, since $\alpha \le \frac{1}{2}$. Collecting the pieces we get

$$d(c_{\zeta}(x), c_{\xi}(x)) \leq d(c_{\zeta}(x), c_{\xi}(y)) + d(c_{\xi}(y), c_{\xi}(x))$$
$$\leq \alpha d(o, c_{\xi}(y)) + |y - x|$$
$$\leq 2\alpha y \leq 4\alpha x.$$

320 and thus

292

$$2\sin\left(\frac{1}{2}\angle(\xi,\zeta)\right) = \lim_{x \to \infty} \frac{1}{x} d(c_{\zeta}(x), c_{\xi}(x)) \le 4\alpha.$$

As a consequence of the previous lemma we get that two geodesic rays c and c' stay at bounded distance if and only if the subsets $c([0,\infty))$ and $c'([0,\infty))$ are linearly equivalent. Write \mathcal{G} to denote the family of all subsets of the form $c([0,\infty))$, where c is some geodesic ray in X.

Proposition 4.2. Let X be a complete CAT(0) space and equip ∂X with the angular metric \angle . Then

$$\partial X \to \mathcal{G}/\sim, \quad \xi \mapsto [R_{\xi}],$$

where $[R_{\xi}]$ is the equivalence class of R_{ξ} with respect to linear equivalence, is one-to-one, onto, and bi-Hölder continuous:

$$\frac{1}{\pi} \angle (\xi, \zeta) \le \left(t([R_{\xi}], [R_{\zeta}]) \right)^2 \le \angle (\xi, \zeta)$$

for all $\xi, \zeta \in \partial X$. Furthermore, \mathcal{G}/\sim is a closed subset of $(\mathcal{U}/\sim, t)$, since $(\partial X, \angle)$ is a complete metric space.

331 Proof. Since $\angle(\xi,\zeta) \in [0,\pi]$ and $\frac{2}{\pi}x \leq \sin(x) \leq x$ for all $x \in [0,\frac{\pi}{2}]$, Lemma 4.1 yields 332 $\frac{1}{\pi}\angle(\xi,\zeta) \leq s(R_{\xi},R_{\zeta}) \leq \angle(\xi,\zeta)$

333

5. Boundaries of groups

Let G be a group and d be a metric on G. Fix the identity element $1 \in G$ as reference point. From an algebraic point of view it is natural to consider the families of unbounded cyclic subgroups and unbounded cyclic subsemigroups of the group G. Hence define

$$\mathcal{C}G = \{ \langle g \rangle \, : \, g \in G, \, \langle g \rangle \in \mathcal{U} \}$$

337 and

$$\mathcal{C}^+G = \{ \langle g \rangle^+ : g \in G, \, \langle g \rangle^+ \in \mathcal{U} \},\$$

where $\langle g \rangle^+ = \{g^n : n \in \mathbb{N}_0\}$ is the semigroup generated by $g \in G$. Note that, if $\langle g \rangle^+ \in \mathcal{C}^+G$, then $\langle g \rangle \in \mathcal{C}G$.

Definition 5.1. We define the projective boundary of G by $\mathcal{P}G = \mathsf{cl}(\mathcal{C}G/\sim)$ and the linear boundary by $\mathcal{L}G = \mathsf{cl}(\mathcal{C}^+G/\sim)$

Remark. Both, $\mathcal{P}G$ and $\mathcal{L}G$, depend on the metric d. If it is necessary to emphasize this dependence, we write $\mathcal{P}(G, d)$ and $\mathcal{L}(G, d)$, respectively.

Lemma 5.2. If $g \in CG$ and $h \in C^+G$ then $\langle g^n \rangle \sim \langle g \rangle$ and $\langle h^n \rangle^+ \sim \langle h \rangle^+$ for all $n \in \mathbb{N}$. Furthermore, if d is left-invariant or right-invariant, then $\langle g \rangle^+ \in C^+G$ if and only if $\langle g \rangle \in CG$.

There are two interesting sources for metrics on a group G. If G is finitely generated (or more generally compactly generated), it is natural to consider word metrics on G. If G is a connected Lie group, it is natural to consider left-invariant Riemannian metrics on G. In this case G is also compactly generated and Corollary A.7 implies that any left-invariant Riemannian metric is quasi-isometrically equivalent to any word metric on G. Hence for our purposes it is sufficient to study the setting of compactly generated groups in more detail.

A topological group is called *compactly generated*, if there is a compact generating set $K \subseteq G$. In this case $S = K \cup K^{-1}$ is a compact, symmetric (i.e. $S = S^{-1}$), generating set. Set $S^0 = \{1\}$ and $S^n = \{s_1 \cdots s_n : s_1, \ldots, s_n \in S\}$ for $n \ge 1$. Note that S^n is compact and symmetric for all $n \ge 0$ and

$$G = \bigcup_{n \ge 0} S^n.$$

The word metric d of G with respect to S is defined by $d(g,h) = \inf\{n : g^{-1}h \in S^n\}$. The metric d is left-invariant and induces the discrete topology on G which is in general different from the group topology. In the sequel we consider the class of compactly generated, locally compact Hausdorff groups. Some facts about such groups and their word metrics are provided by Appendix A. Finitely generated groups fit in this setting (in this case a finitely generated group is equipped with the discrete topology). If not stated otherwise, all topological notions refer to the group topology (except for boundedness which refers to the word metric d).

We fix some compactly generated, locally compact Hausdorff group G and a word metric d on G. Notice that a subset of G is bounded with respect to d if and only if it is relatively compact (see Lemma A.1). Suppose that d' is another word metric on G or (more general) a metric which is quasi-isometrically equivalent to d. Then, by Theorem 3.3, $t_{(G,d)}$ and $t_{(G,d')}$ are bi-Lipschitz-equivalent. Hence linear equivalence and all notions which only depend

on the topological or uniform structure of \mathcal{U}/\sim (like closure or Cauchy completeness for instance), do not depend on the generating set. In particular, we obtain the following statement.

Lemma 5.3. If a compactly generated, locally compact Hausdorff group G is equipped with a word metric d then the (topological) spaces $\mathcal{L}G$ and $\mathcal{P}G$ do not depend on the choice of the word metric (or of the generating set).

A group element g is called *compact*, if $\langle g \rangle$ is relatively compact, and *non-compact* otherwise. Thus g is non-compact if and only if $\langle g \rangle \in CG$. Notice that, if G is finitely generated, a group element g is non-compact if and only if g is non-torsion. Furthermore, by Weil's lemma (see [HR79, Theorem 9.1]) g is non-compact, if and only if $\langle g \rangle$ is the image of a monomorphism $\mathbb{Z} \to G$ which is a topological isomorphism onto $\langle g \rangle$ (a topological isomorphism is a group isomorphism which is also a homeomorphism). Hence, Weil's lemma implies the following.

Lemma 5.4. If $g \in G$ is non-compact then $d(1, g^n) \to \infty$ for $n \to \infty$.

Remark. Notice, that compact group elements of a group G do not contribute to the boundaries $\mathcal{P}G$ and $\mathcal{L}G$. Especially, if G only contains compact group elements, then these boundaries are empty. In the discrete case this means that torsion groups have empty boundaries.

Remark. Let g and h be non-compact group elements. We have seen that $s(\langle g \rangle^+, \langle h \rangle^+) \leq 1$ 386 and $s(\langle g \rangle^+, \langle g^n \rangle^+) = 0$ for all $n \in \mathbb{N}$. Now it is natural to ask, what can be said about 387 $s(\langle g \rangle^+, \langle g^{-1} \rangle^+)$. Often $s(\langle g \rangle^+, \langle g^{-1} \rangle^+) = 1$, but in [KLS12] Krön, Lehnert and Stein give 388 an example of a finitely generated group constructed by iterated HNN-extensions with a non-torsion element g such that $s(\langle g \rangle^+, \langle g^{-1} \rangle^+) \leq \frac{12}{17}$. They also show that in general this 389 390 value cannot be arbitrarily close to zero. Indeed, $s(\langle g \rangle^+, \langle g^{-1} \rangle^+)$ is always greater or equal 391 $\frac{1}{2}$. The infimum of these values (for all groups) is unknown. In [KLS12] there is also an 392 example of a finitely generated group with non-torsion elements g, h for which $\langle g \rangle^+ \sim \langle h \rangle^+$ 393 but $\langle g^{-1} \rangle^+ \not\sim \langle h^{-1} \rangle^+$. 394

The following lemma yields a useful alternative to compute $s^+(\langle g \rangle^+, \langle h \rangle^+)$ and $s^+(\langle g \rangle, \langle h \rangle)$.

Lemma 5.5. Let g and h be non-compact group elements. Then

$$s^+(\langle g \rangle^+, \langle h \rangle^+) = \limsup_{n \to \infty} \inf \left\{ \frac{d(h^n, g^m)}{d(1, g^m)} \, : \, m \in \mathbb{N}_0
ight\}$$

397 and

$$s^+(\langle g \rangle, \langle h \rangle) = \limsup_{|n| \to \infty} \inf \left\{ \frac{d(h^n, g^m)}{d(1, g^m)} : m \in \mathbb{Z} \right\}.$$

Proof. We only prove the first claim, since the proof of the second is analogous. Suppose that $\alpha > s^+(\langle g \rangle^+, \langle h \rangle^+)$. Hence $\langle h \rangle^+ \subseteq \alpha \langle h \rangle^+ + a$ for some $a \ge 0$. Thus, for each $n \in \mathbb{N}_0$, there is an integer $k = k(n) \ge 0$, such that $d(h^n, g^k) \le \alpha d(1, g^k) + a$. Then

$$\inf\left\{\frac{d(h^{n}, g^{m})}{d(1, g^{m})} : m \in \mathbb{N}_{0}\right\} \leq \frac{d(h^{n}, g^{k})}{d(1, g^{k})} \leq \alpha + \frac{a}{d(1, g^{k})}.$$

401 Using the triangle inequality we get

$$(1 - \alpha)d(1, g^k) - a \le d(1, h^n) \le (1 + \alpha)d(1, g^k) + a$$

402 If $n \to \infty$ then $d(1, h^n) \to \infty$ by Lemma 5.4 and therefore $d(1, g^k) \to \infty$. This implies

$$\limsup_{n \to \infty} \inf \left\{ \frac{d(h^n, g^m)}{d(1, g^m)} : m \in \mathbb{N}_0 \right\} \le \limsup_{n \to \infty} \alpha + \frac{a}{d(1, g^k)} = \alpha.$$

403 In order to prove the reversed inequality assume that

$$\alpha > \limsup_{n \to \infty} \inf \left\{ \frac{d(h^n, g^m)}{d(1, g^m)} : m \in \mathbb{N}_0 \right\}.$$

Then there is an integer $N \ge 0$, such that 404

$$\inf\left\{\frac{d(h^n, g^m)}{d(1, g^m)} : m \in \mathbb{N}_0\right\} \le \alpha$$

for all $n \ge N$. Let $\varepsilon > 0$. Then, for each $n \ge N$, we can find an integer $k = k(n) \ge 0$, such 405 that 406

$$\frac{d(h^n, g^k)}{d(1, g^k)} \le \alpha + \varepsilon$$

Set $a = \max\{d(1, h^n) : 0 \le n < N\}$. Then we obtain $\langle h \rangle^+ \subseteq (\alpha + \varepsilon) \langle g \rangle^+ + a$. 407

Using Corollary A.6 and its notation, we obtain the following: 408

Lemma 5.6. Let G be a compactly generated, locally compact group. The following state-409 ments are true up to bi-Lipschitz-equivalence of the metric t: 410

• Suppose that N is a compact group and H is a topological Hausdorff group. If $\{1\} \longrightarrow$ 411 $N \longrightarrow H \xrightarrow{\pi} G \longrightarrow \{1\}$ is a topological short exact sequence, such that $\pi \colon H \to G$ is 412 also open, then H and G have the same linear and projective boundaries, respectively. 413 414

• If H is a closed subgroup of G and $(H \setminus G, d_{H \setminus G})$ is bounded then

$$\mathcal{L}H \subseteq \mathcal{L}G$$
 and $\mathcal{P}H \subseteq \mathcal{L}G$.

If
$$H$$
 is of finite index in G then equality holds.

Proof. In order to prove the first statement note that, by Corollary A.6 the homomorphism 416 $\pi: H \to G$ is a quasi-isometry. Assume that $h \in H$ and $\langle \pi(h) \rangle$ is bounded in G then 417 $\pi^{-1}(\langle \pi(h) \rangle)$ is bounded by Lemma A.3. Hence $\langle h \rangle \subseteq \pi^{-1}(\langle \pi(h) \rangle)$ is bounded. Thus un-418 bounded cyclic sub(semi)groups of H are mapped onto unbounded cyclic sub(semi)groups 419 of G. This implies the first statement using Theorem 3.3. 420

Now suppose that H is a closed subgroup of G and $H \setminus G$ is bounded. By Corollary A.6 421 the inclusion is a quasi-isometry. In order to emphasize the dependence on H and G, we use 422 subscripts H and G. By Theorem 3.3 we have 423

$$\mathcal{L}H = \mathsf{cl}_H(\mathcal{C}^+H/\sim_H) = \mathsf{cl}_G(\mathcal{C}^+H/\sim_G) \subseteq \mathsf{cl}_G(\mathcal{C}^+G/\sim_G) = \mathcal{L}G$$

and analogously for $\mathcal{P}H \subseteq \mathcal{P}G$. Assume that H has finite index in G. If $g \in G$ then $H \setminus H \langle g \rangle$ 424 is finite. Thus there are $k \in \mathbb{Z}$ and n > 0, such that $Hg^{k+n} = Hg^k$. This implies $g^n \in H$. 425 Hence in this case Lemma 5.2 implies 426

$$\mathcal{C}^+H/\sim_H = \mathcal{C}^+G/\sim_G$$
 and $\mathcal{C}H/\sim_H = \mathcal{C}G/\sim_G$

which yields the assertion. 427

In the setting of finitely generated groups the previous lemma implies that two weakly 428 commensurable finitely generated groups G and H (i.e. there is a group Q and homomor-429 phisms $Q \to G$ and $Q \to H$ both having finite kernels and images of finite index) have 430 the same linear and projective boundaries. In the continuous setting the situation is more 431 complicated: In general it is possible that 432

$$\mathcal{C}^+H/\sim_H \subsetneq \mathcal{C}^+G/\sim_G$$

(consider for instance $\mathbb{Z}^2 \leq \mathbb{R}^2$). However, equality may hold after taking closures on both 433 sides, i.e. $\mathcal{L}H = \mathcal{L}G$. The problem here is to find for each non-compact $g \in G$ a sequence 434 $(h_n)_{n\geq 0}$ in H, such that $t(\langle g \rangle^+, \langle h_n \rangle^+) \to 0$ for $n \to \infty$. Notice that there is always an unbounded subset $R \subseteq H$ with $\langle g \rangle^+ \sim R$, if $H \setminus G$ is bounded. 435 436

With this preparations we can settle the commutative case completely. Recall that, if G437 is a commutative, compactly generated, locally compact Hausdorff group, then by [HR79, 438 Theorem 9.8] there are integers $a, b \ge 0$ and a commutative, compact Hausdorff group C, 439 such that G is topologically isomorphic to $\mathbb{R}^a \times \mathbb{Z}^b \times C$. 440

12

441 Corollary 5.7. Assume that G is a commutative, compactly generated, locally compact
 442 Hausdorff group.

- If G is topologically isomorphic to $\mathbb{R}^a \times \mathbb{Z}^b \times C$ for some integers $a, b \ge 0$ and some compact, commutative group C then $\mathcal{L}G = \mathbb{S}^{a+b-1}$ and $\mathcal{P}G = \mathbb{P}^{a+b-1}$.
- If H is a closed subgroup, such that G/H is compact then $\mathcal{L}H = \mathcal{L}G$ and $\mathcal{P}H = \mathcal{P}G$.

446 Proof. As $\mathbb{R}^a \times \mathbb{Z}^b$ is a quotient of G with compact kernel, the linear and projective boundaries 447 of G and $\mathbb{R}^a \times \mathbb{Z}^b$ are the same, respectively. Since any word metric on $\mathbb{R}^a \times \mathbb{Z}^b$ is quasi-448 isometrically equivalent to the ℓ^2 -metric on $\mathbb{R}^a \times \mathbb{Z}^b$, we may use the ℓ^2 -metric. It is then 449 easy to see that $\mathbb{R}^a \times \mathbb{Z}^b$ and \mathbb{R}^{a+b} have the same boundaries. Hence the assertion follows 450 from Example 2.11.

Suppose that H is a closed subgroup, such that G/H is compact. As before, let G be topologically isomorphic to $\mathbb{R}^a \times \mathbb{Z}^b \times C$. It follows that H is topologically isomorphic to $\mathbb{R}^{a-c} \times \mathbb{Z}^{b+c} \times D$ for some integer c and some commutative, compact Hausdorff group D. Thus the first assertion implies the second.

In the setting of topological groups it is natural to consider also unbounded one-parameter 455 subgroups and unbounded one-parameter subsemigroups, as well. A one-parameter subgroup 456 in G is the image of a continuous homomorphism $\mathbb{R} \to G$ and a one-parameter subsemi-457 group is the image of a continuous semigroup homomorphism $[0,\infty) \to G$. Obviously, if 458 $\varphi \colon [0,\infty) \to G$ is a continuous semigroup homomorphism, then there is a canonical exten-459 sion to a continuous group homomorphism $\bar{\varphi} \colon \mathbb{R} \to G$ and φ has unbounded image, if and 460 only if $\bar{\varphi}$ has. Define $\mathcal{C}_{\mathbb{R}}G$ and $\mathcal{C}^+_{\mathbb{R}}G$ to be the family of unbounded one-parameter subgroups 461 and unbounded one-parameter subsemigroups, respectively. Again, by Weil's lemma a con-462 tinuous homomorphism $\varphi \colon \mathbb{R} \to G$ has unbounded image if and only if φ is a topological 463 isomorphism onto its image. 464

Lemma 5.8. Suppose that $\varphi \colon \mathbb{R} \to G$ is a continuous homomorphism with unbounded image. Then

$$\varphi([0,\infty)) \sim \langle \varphi(t) \rangle^+$$
 and $\varphi(\mathbb{R}) \sim \langle \varphi(t) \rangle$

467 for all t > 0. Hence

$$\mathcal{C}_{\mathbb{R}}G/\sim \subseteq \mathcal{C}G/\sim \quad and \quad \mathcal{C}^+_{\mathbb{R}}G/\sim \subseteq \mathcal{C}^+G/\sim.$$

468 Proposition 5.9. Let G be a connected, nilpotent Lie group. Then

$$\mathcal{C}_{\mathbb{R}}G/\sim = \mathcal{C}G/\sim$$
 and $\mathcal{C}_{\mathbb{R}}^+G/\sim = \mathcal{C}^+G/\sim$.

469 Proof. Let \mathfrak{g} be the Lie algebra of G and $\exp: \mathfrak{g} \to G$ be the exponential map. Then \exp is 470 surjective. Thus, if g is a non-compact group element, then there is an element $x \in \mathfrak{g}$ with 471 $\exp(x) = g$. Then $\mathbb{R} \to G$, $t \mapsto \exp(tx)$ is a continuous homomorphism with unbounded 472 image which proves the statement.

473

6. Boundaries of nilpotent groups

In the following we determine the linear and projective boundary of connected, nilpotent Lie groups and their discrete counterparts, finitely generated nilpotent groups. A commutative, connected Lie group G is isomorphic to $\mathbb{R}^a \times (\mathbb{R}/\mathbb{Z})^b$ for some integers a and b. In analogy to the discrete case we call the integer a the compact-free dimension of G. For convenience we define \mathbb{S}^{-1} and \mathbb{P}^{-1} to be the empty set.

Theorem 6.1. Let G be a nilpotent group which is either a connected Lie group or a finitely generated group. Suppose that G has descending central series

$$G = G_1 \supseteq G_2 \supseteq \cdots \supseteq G_c \supseteq G_{c+1} = \{1\},\$$

where $c \geq 1$ is the nilpotency class of G. Let $\nu(i)$ denote the compact-free dimension or torsion-free rank of G_i/G_{i+1} , respectively. Then the linear boundary $\mathcal{L}G$ is homeomorphic BERNHARD KRÖN, JÖRG LEHNERT, NORBERT SEIFTER, AND ELMAR TEUFL

483 to the disjoint union of c spheres:

$$\mathcal{L}G = \mathbb{S}^{\nu(1)-1} \uplus \mathbb{S}^{\nu(2)-1} \uplus \cdots \uplus \mathbb{S}^{\nu(c)-1}.$$

Analogously, the projective boundary PG is homeomorphic to the disjoint union of projective spaces:

$$\mathcal{P}G = \mathbb{P}^{\nu(1)-1} \uplus \mathbb{P}^{\nu(2)-1} \uplus \cdots \uplus \mathbb{P}^{\nu(c)-1}.$$

If two finitely generated, nilpotent groups G and H are weakly commensurable then previous result yields a new proof of the fact that the multisets

 $\{\nu_1(G), \nu_2(G), \dots\}$ and $\{\nu_1(H), \nu_2(H), \dots\}$

of torsion-free ranks are equal, since the boundaries of G and H are bi-Lipschitz-equivalent. Notice that there is no information on the ordering and it is unclear, whether it is possible to deduce the ordering from the angle metrics of G and H, respectively. It is a corollary of Pansu's theorem (see [Pan89, Théorème 3]), that even the tuples

$$(\nu_1(G), \nu_2(G), \dots)$$
 and $(\nu_1(H), \nu_2(H), \dots)$

492 are equal.

First we prove the theorem for connected Lie groups and then use the Mal'tsev completion to deduce the statement for finitely generated groups. For both cases we use the notation and results of Appendix B.

Proof of Theorem 6.1 in the Lie case. We prove that statement for $\mathcal{L}G$, as the other case is completely analogous. Let G be a connected, nilpotent Lie group with word metric d_G . Set $t_G = t_{(G,d_G)}$ and write \sim_G to denote linear equivalence in (G,d_G) . By Lemma B.1 and Lemma 5.6 we may assume that G is also simply connected. Set $t_a = t_{(\mathfrak{g},d_a)}$ and write \sim_a to denote linear equivalence in (\mathfrak{g}, d_a) . By the Lemmas B.7, B.8, B.11 the map

$$\varphi \colon \mathcal{C}^+(\mathfrak{g},+)/\sim_a \to \mathcal{C}^+(G,\cdot)/\sim_G$$

which maps the equivalence class of $\langle x \rangle^+ \in \mathcal{C}^+(\mathfrak{g}, +)$ to the equivalence class of $\langle \exp(x) \rangle^+ \in \mathcal{C}^+(G, \cdot)$, is well-defined and bi-Hölder continuous with respect to the metrics t_a and t_G , respectively. Hence φ extends to a bi-Hölder continuous map from $\mathcal{L}(\mathfrak{g}, d_a)$ to $\mathcal{L}(G, d_G)$. Then the assertion follows from the first part of Lemma B.8.

Proof of Theorem 6.1 in the discrete case. Let Γ be a finitely generated, nilpotent group. 505 We only show the assertion for $\mathcal{L}\Gamma$ for the same reason as above. By Lemma B.1 and 506 Lemma 5.6 we may assume that Γ is also torsion-free. Then the (real) Mal'tsev completion 507 of Γ yields a connected, simply connected, nilpotent Lie group G, such that Γ is a uniform 508 subgroup of G, see [Mal51]. Using Lemma 5.6 it follows that $\mathcal{L}\Gamma \subseteq \mathcal{L}G$. Let d_G be a word 509 metric on G and set $t_G = t_{(G,d_G)}$. In order to prove equality, it is sufficient to construct for 510 each $g \in G$ a sequence $h_1, h_2, \ldots \in \Gamma$, such that $t_G(\langle g \rangle^+, \langle h_m \rangle^+) \to 0$ if $m \to \infty$. Suppose 511 that g is an element of G_n . Set $\Lambda = \log(\Gamma)$ and $x = \log(g)$. Then $\Lambda \cap \mathfrak{g}_k$ is a uniform subgroup 512 in (\mathfrak{g}_k, \cdot) for all k. Hence $\pi_n(\Lambda \cap \mathfrak{g}_n)$ is a uniform subgroup of $(V_n, +)$, since π_n is a continuous 513 epimorphism from (\mathfrak{g}_n, \cdot) to $(V_n, +)$. As V_n is isomorphic to $\mathbb{R}^{\nu(n)}$, $\pi_n(\Lambda \cap \mathfrak{g}_n)$ is isomorphic 514 to $\mathbb{Z}^{\nu(n)}$. Thus there is a sequence $y_1, y_2, \ldots \in \Lambda \cap \mathfrak{g}_n$, such that $t_a(\langle \pi(x) \rangle^+, \langle \pi(y_m) \rangle^+) \to 0$ if 515 $m \to \infty$. Since $\langle x \rangle^+ \sim_a \langle \pi(x) \rangle^+$ and $\langle y_m \rangle^+ \sim_a \langle \pi(y_m) \rangle^+$, we infer that $t_a(\langle x \rangle^+, \langle y_m \rangle^+) \to 0$ if $m \to \infty$. Set $h_m = \exp(z_m) \in \Gamma$. Then $t_G(\langle g \rangle^+, \langle h_m \rangle^+) \to 0$ for $m \to \infty$ using Lemma B.11 516 517 as required. 518

Remark. We have carried out an alternative proof for the discrete case which avoids the use of Mal'tsev completion and tools from Lie theory and employs techniques from combinatorial group theory—mainly commutator calculus and careful analysis of word lengths'. This proof follows similar lines compared to the proof for the Lie case given here.

523 7. Boundaries of vertex-transitive graphs with polynomial growth

Let G be a group and let S be a finite generating set of G. Then the Cayley graph X of G with respect to S is given by VX = G and $EX = \{\{g,gs\} : g \in G, s \in S\}$. If we define a Cayley graph in this way, namely by right multiplication, then G acts as a vertex-transitive group of automorphisms on X by left multiplication. Hence Cayley graphs of finitely generated groups give rise to a particular class of locally finite, vertex-transitive graphs.

Since we have defined our notion of boundary for metric spaces in general, it is natural 530 to consider $\mathcal{L}G$ and $\mathcal{P}G$ not only for groups G (and thus for their Cayley graphs), but also 531 for vertex-transitive graphs in general. But as Example 2.13 shows, even for simple struc-532 tures, as Cayley graphs of \mathbb{Z}^d , for Cayley graphs of groups G the space \mathcal{U}/\sim is much richer 533 than $\mathcal{L}G$ or $\mathcal{P}G$. Hence it seems rather difficult to characterize our boundaries for graphs 534 without involving group actions. Therefore, we define—roughly speaking—the projective 535 (linear) boundary of a graph as the projective (linear) boundary induced by the action of 536 its automorphism group. Then, at least for graphs with polynomial growth, it is possible to 537 obtain results similar to the above. 538

Furthermore, we emphasize that the concepts defined in the sequel are not restricted to locally finite graphs. In addition the results up to Corollary 7.8 also hold without the assumption of local finiteness. From Theorem 7.9 to the end of this section we consider graphs with polynomial growth which of course implies that they are locally finite. Hence, although the main results of this section only hold for locally finite graphs, this assumption is never explicitly stated.

In the following we always endow a graph X with the graph metric d, i.e., for any two vertices $u, v \in VX$, the distance d(u, v) is the infimum of all numbers k such that there is a path of length k connecting u and v.

Definition 7.1. Let X = (VX, EX) be an infinite, connected graph and let $\operatorname{Aut} X$ be the automorphism group of X. For $v \in VX$, we write $\operatorname{Unb}_v X \subseteq \operatorname{Aut} X$ to denote the set of group elements $g \in \operatorname{Aut} X$ for which the set $\langle g \rangle v = \{g^n v : n \in \mathbb{Z}\}$ is unbounded.

Lemma 7.2. Let X be an infinite, connected graph, $v \in VX$, and let $q \in Unb_v X$.

- The set $Unb_v X$ is symmetric and both, $g^{\infty}v = \{g^nv : n \in \mathbb{N}_0\}$ and $(g^{-1})^{\infty}v = \{g^{-n}v : n \in \mathbb{N}_0\}$, are unbounded.
- If n is a nonzero integer then $g^n \in Unb_v X$. Furthermore, $\langle g \rangle v$, $\langle g^n \rangle v$ are linearly equivalent and $g^{\infty}v$, $(g^n)^{\infty}$ are linearly equivalent, too.

⁵⁵⁶ *Proof.* The first part is immediate. The second part can be proved in the same way as Lemma 5.2.

Definition 7.3. Let X be an infinite, connected graph and let $G \leq \text{Aut } X$. For $v \in VX$ we define

$$\mathcal{C}_{G,v}X = \{ \langle g \rangle v : g \in \mathsf{Unb}_v X \cap G \}, \qquad \mathcal{C}_{G,v}^+ X = \{ g^\infty v : g \in \mathsf{Unb}_v X \cap G \}.$$

Lemma 7.4. Let X be an infinite, connected graph. Then, for $u, v \in VX$, we have

$$\mathsf{Unb}_u X = \mathsf{Unb}_v X$$

561 If $G \leq \operatorname{Aut} X$ then

$$\mathcal{C}_{G,u}X/\sim = \mathcal{C}_{G,v}X/\sim$$
 and $\mathcal{C}_{G,u}^+X/\sim = \mathcal{C}_{G,v}^+X/\sim$

562 up to isometric isomorphy.

From f Assume that $g \in Unb_u X$. Then $d(u, g^n u) \to \infty$ as $n \to \infty$. As X is connected, $d(u, v) < \infty$ for all $v \in VX$. The triangle inequality implies

$$d(u, g^{n}u) \leq d(u, v) + d(v, g^{n}v) + d(g^{n}v, g^{n}u) = 2d(u, v) + d(v, g^{n}v),$$

hence $d(v, g^n v) \ge d(u, g^n u) - 2d(u, v) \to \infty$ as $n \to \infty$. Thus $\mathsf{Unb}_u X \subseteq \mathsf{Unb}_v X$ and the reversed inclusion follows by means of symmetry. In order to prove the second part of our assertion, note that, for $g \in \mathsf{Unb}_u X \cap G = \mathsf{Unb}_v X \cap G$,

$$\langle g \rangle u \subseteq 0 \langle g \rangle v + d(u, v)$$
 and $\langle g \rangle v \subseteq 0 \langle g \rangle u + d(u, v)$

which means that $\langle g \rangle u$ and $\langle g \rangle v$ are linearly equivalent, thus implying $\mathcal{C}_{G,u}X/\sim = \mathcal{C}_{G,v}X/\sim$ up to isometric isomorphy. An analogous reasoning yields $\mathcal{C}^+_{G,u}X/\sim = \mathcal{C}^+_{G,v}X/\sim$.

In the light of Lemma 7.4 we may drop dependence on the vertex v. This motivates the following definition:

Definition 7.5. Let X be an infinite, connected graph and fix a reference vertex v. Then we define Unb $X = \text{Unb}_v X$. If G is a subgroup of Aut X then we set

$$\mathcal{P}_G X = \mathsf{cl}(\mathcal{C}_{G,v} X/\sim)$$
 and $\mathcal{L}_G X = \mathsf{cl}(\mathcal{C}_{G,v}^+ X/\sim).$

The spaces $\mathcal{P}X = \mathcal{P}_{\operatorname{Aut} X}X$ and $\mathcal{L}X = \mathcal{L}_{\operatorname{Aut} X}X$ are called *projective boundary* and *linear boundary* of X, respectively.

Let X be a graph and let σ be a partition of the vertex set VX. The quotient graph X_{σ} 576 is defined as follows: the vertex set VX_{σ} is σ , and two vertices $x, y \in VX_{\sigma}$ are adjacent, 577 if there are adjacent vertices $v, w \in X$ with $v \in x$ and $w \in y$. Let $G \leq \operatorname{Aut} X$ be a group 578 of automorphisms such that σ is G-invariant, i.e. $g(b) \in \sigma$ for all $b \in \sigma$ and all $g \in G$. 579 Then G naturally induces a group action on X_{σ} . The subgroup of the automorphism group 580 Aut X_{σ} corresponding to this action is denoted by G_{σ} . Also, there is a homomorphism 581 $\varphi \colon G \to G_{\sigma}$ such that the kernel of φ consists of all those $g \in G$ with g(b) = b for all $b \in \sigma$. 582 If $G \leq \operatorname{Aut} X$ acts vertex-transitively on X and σ is a G-invariant partition of VX then σ is 583 called *imprimitivity system of* G on X. The elements of an imprimitivity system are called 584 blocks. 585

Let σ be an Aut X-invariant partition of VX. In order to avoid ambiguity we write (Aut X) $_{\sigma}$ to denote the subgroup of Aut X_{σ} corresponding to the natural action of Aut X on X $_{\sigma}$. Notice that $(\operatorname{Aut} X)_{\sigma} \subseteq \operatorname{Aut} X_{\sigma}$, but these two groups are not necessarily equal as the next example shows.

Example 7.6. Consider the graph X depicted in Figure 1. It consists of two disjoint infinite double-rays $\{v_i : i \in \mathbb{Z}\}$ and $\{w_i : i \in \mathbb{Z}\}$ and additional "crossed rungs": For even i, v_i is connected to w_{i+1} and for odd i, v_i is connected to w_{i-1} . This graph is vertex-transitive,



FIGURE 1. An example graph X for $(\operatorname{Aut} X)_{\sigma} \subsetneq \operatorname{Aut} X_{\sigma}$.

592

16

and the sets $\{v_i, w_i\}$, $i \in \mathbb{Z}$, give rise to an imprimitivity system σ of Aut X on X. The quotient graph X_{σ} is an infinite double-ray $\{x_i : i \in \mathbb{Z}\}$, where the vertices x_i correspond to the sets $\{v_i, w_i\}$ for $i \in \mathbb{Z}$. The mapping g_{σ} which fixes x_0 and maps x_i onto x_{-i} for $i \in \mathbb{Z}$ is obviously an automorphism of X_{σ} . But there exists no automorphism $g \in \operatorname{Aut} X$ with

$$g(\{v_i, w_i\}) = \{v_{-i}, w_{-i}\}$$

597 for $i \in \mathbb{Z}$. Hence, for this graph X, $(\operatorname{Aut} X)_{\sigma} \subsetneq \operatorname{Aut} X_{\sigma}$ holds.

Let X be an infinite, connected graph and $H \leq G \leq \operatorname{Aut} X$. As the underlying metric space (VX, d) is fixed, the inclusion $H \leq G$ implies, that $\mathcal{L}_H X$ and $\mathcal{P}_H X$ are up to isometric isomorphy subspaces of $\mathcal{L}_G X$ and $\mathcal{P}_G X$, respectively: Fix some reference vertex $v \in VX$ and notice that $\mathcal{C}^+_{H,v} \subseteq \mathcal{C}^+_{G,v}$. Hence the map

$$\mathcal{C}^+_{H,v} \to \mathcal{C}^+_{G,v}, \quad h^\infty v \mapsto h^\infty v$$

induces an isometric embedding $\mathcal{C}_{H,v}^+/\sim \to \mathcal{C}_{G,v}^+/\sim$ which extends naturally to the topological closures $\mathcal{L}_H X$ and $\mathcal{L}_G X$. Similarly, there is an isometric embedding $\mathcal{P}_H X \to \mathcal{P}_G X$.

Lemma 7.7. Let X be an infinite, connected graph.

• If $H \leq G \leq \operatorname{Aut} X$ and H has finite index in G then $\mathcal{L}_H X$ and $\mathcal{L}_G X$ ($\mathcal{P}_H X$ and $\mathcal{P}_G X$) are isometrically isomorphic.

607

• Let $G \leq \operatorname{Aut} X$ and let σ be a G-invariant partition of VX such that

$$\sup\{d(x,y)\,:\,x,y\in b,\,b\in\sigma\}<\infty$$

608

Then $\mathcal{L}_G X$ and $\mathcal{L}_{G_{\sigma}} X_{\sigma}$ ($\mathcal{P}_G X$ and $\mathcal{P}_{G_{\sigma}} X_{\sigma}$) are bi-Lipschitz-equivalent.

Proof. In order to prove the first statement, we may assume that H is a normal subgroup of G with finite index, as the intersection of all conjugates of H forms a normal subgroup with finite index. Let n be the finite index of H in G. Then, for any $g \in \text{Unb}_v X \cap G$, $g^n \in$ Unb_v $X \cap H$ and the unbounded subsets $g^{\infty}v \in \mathcal{C}^+_{G,v}$, $(g^n)^{\infty}v \in \mathcal{C}^+_{H,v}$ are linearly equivalent. Therefore, the isometric embedding $\mathcal{C}^+_{H,v}/\sim \to \mathcal{C}^+_{G,v}/\sim$ is an isometric isomorphism which extends naturally to $\mathcal{L}_H X$ and $\mathcal{L}_G X$. Analogous reasoning yields the statement for $\mathcal{P}_H X$ and $\mathcal{P}_G X$.

We now prove the second assertion. For $x \in VX$ we write \bar{x} to denote the element of $\sigma = VX_{\sigma}$ containing x. Similarly, we write $\bar{g} \in G_{\sigma}$ for the automorphism of X_{σ} induced by the group element $g \in G$. Fix some reference vertex v and set

$$a = \sup\{d(x, y) : x, y \in b, b \in \sigma\} < \infty.$$

619 The map $\pi: VX \to VX_{\sigma}, x \mapsto \bar{x}$, is a quasi-isometry, since

$$d_{X_{\sigma}}(\bar{x},\bar{y}) \le d_X(x,y) \le (a+1)d_{X_{\sigma}}(\bar{x},\bar{y}) + a$$

620 for $x, y \in VX$. Furthermore, π induces a map from $\mathcal{C}^+_{G,v}X$ onto $\mathcal{C}^+_{G_{\sigma},\bar{v}}X_{\sigma}$: if $g^{\infty}v =$ 621 $\{v_0, v_1, \ldots\} \in \mathcal{C}^+_{G,v}X$ then $\pi(g^{\infty}v) = \{\bar{v}_0, \bar{v}_1, \ldots\} = \bar{g}^{\infty}\bar{v} \in \mathcal{C}^+_{G_{\sigma},\bar{v}}X_{\sigma}$. Theorem 3.3 implies 622 that

$$\mathcal{L}_G X = \mathsf{cl}(\mathcal{C}^+_{G,v} X/\sim) \quad \text{and} \quad \mathcal{L}_{G_\sigma} X_\sigma = \mathsf{cl}(\mathcal{C}^+_{G_\sigma,\bar{v}} X_\sigma/\sim)$$

are bi-Lipschitz-equivalent. Again the statement for $\mathcal{P}_G X$ and $\mathcal{P}_{G_{\sigma}} X_{\sigma}$ follows along the same lines.

625 Corollary 7.8. Let X be an infinite, connected graph.

- If $G \leq \operatorname{Aut} X$ acts vertex-transitively on X and σ is an imprimitivity system of Gon X with finite blocks then $\mathcal{L}_G X$ and $\mathcal{L}_{G_{\sigma}} X_{\sigma}$ ($\mathcal{P}_G X$ and $\mathcal{P}_{G_{\sigma}} X_{\sigma}$) are bi-Lipschitzequivalent.
- If $G \leq \operatorname{Aut} X$ acts freely and with finitely many orbits on VX then $\mathcal{L}G$ and $\mathcal{L}_G X$ ($\mathcal{P}G$ and $\mathcal{P}_G X$) are bi-Lipschitz-equivalent.

631 *Proof.* The first statement is immediate:

$$\sup\{d(x,y) : x, y \in b, b \in \sigma\} < \infty$$

follows from the fact that G acts vertex-transitively on X and the blocks of σ are finite.

To prove the second statement we apply the ideas of the proof of the so-called Contraction Lemma (see [Bab77]): Since G acts freely and with finitely many orbits on X, there is a finite tree T in X which contains exactly one vertex of each orbit of G on X. Furthermore, the sets gVT for $g \in G$ form a partition of VX. Set $\sigma = \{gVT : g \in G\}$. Then X_{σ} is isomorphic to a Cayley graph of G, and the groups G and G_{σ} are isomorphic, as VT contains exactly one vertex of each orbit. Hence $\mathcal{L}G$ is (by definition) equal to $\mathcal{L}_{G_{\sigma}}X_{\sigma}$ and the spaces $\mathcal{L}_{G_{\sigma}}X_{\sigma}, \mathcal{L}_{G}X$ are bi-Lipschitz-equivalent by the previous lemma. **Theorem 7.9.** Let X be an infinite, connected, vertex-transitive graph with polynomial growth. Then there is a finitely generated, torsion-free, nilpotent group N which has the same growth rate as X, and $\mathcal{L}N$ and $\mathcal{P}N$ are bi-Lipschitz-equivalent to $\mathcal{L}X$ and $\mathcal{P}X$, respectively.

To prove this result about graphs with polynomial growth, the following two results of Trofimov [Tro84] are essential.

Theorem 7.10 (Theorem 1 in [Tro84]). Let X be an infinite, connected, vertex-transitive graph with polynomial growth. Then there exists an imprimitivity system σ of Aut X on VX with finite blocks such that Aut X_{σ} is a finitely generated virtually nilpotent group and the stabilizer in Aut X_{σ} of a vertex of X_{σ} is finite.

Theorem 7.11 (Theorem 2 in [Tro84]). Let X be an infinite, connected graph with polynomial growth and let a group $G \leq \operatorname{Aut} X$ act vertex-transitively on VX. Then there exists an imprimitivity system σ of G on VX with finite blocks such that G_{σ} is a finitely generated virtually nilpotent group and the stabilizer in G_{σ} of a vertex of X_{σ} is finite.

Proof of Theorem 7.9. Let $G = \operatorname{Aut} X$ and let σ and G_{σ} as in Theorem 7.11. Then G_{σ} contains a finitely generated, nilpotent, normal subgroup N of finite index. By [Sei91b, Corollary 2.7] we can furthermore assume that N is torsion-free. Since the finite index of N in G_{σ} implies that N acts with finitely many orbits on X, we can assume by [Sei91a, Theorem 2.3] that all $n \in N, n \neq 1$, act with infinite orbits on X_{σ} .

Since the vertex stabilizers of Aut X_{σ} and G_{σ} are both finite (by Theorems 7.10 and 7.11), both groups have the same growth rate as the graph X_{σ} which is of course equal to the growth rate of X. Hence G_{σ} has finite index in Aut X_{σ} . As N has finite index in G_{σ} , it has also finite index in Aut X_{σ} . Therefore Lemma 7.7 implies that the projective (linear) boundary induced by N on X_{σ} is bi-Lipschitz-equivalent to the projective (linear) boundary induced by Aut X_{σ} which we defined to be the projective (linear) boundary of X_{σ} .

Since X_{σ} is a quotient graph of X with respect to the finite blocks of σ , Corollary 7.8 implies that the projective (linear) boundaries of X and X_{σ} which are induced by Aut X and $G_{\sigma} = (\operatorname{Aut} X)_{\sigma}$, respectively, are bi-Lipschitz-equivalent.

To conclude the proof we show that $\mathcal{L}N$ and $\mathcal{P}N$ are bi-Lipschitz-equivalent to $\mathcal{L}_N X_{\sigma}$ and $\mathcal{P}_N X_{\sigma}$, respectively. As N is torsion-free and the stabilizer of a vertex is finite, N acts freely on X_{σ} . Since N also acts with finitely many orbits on X_{σ} , the claim follows directly from Corollary 7.8.

As a consequence of Theorem 6.1 we obtain the following result.

Corollary 7.12. Let X be an infinite, connected, vertex-transitive graph with polynomial growth and let N be a finitely generated, torsion-free, nilpotent group supplied by Theorem 7.9. Then the linear boundary $\mathcal{L}X$ is homeomorphic to a disjoint union of spheres:

$$\mathcal{L}X = \mathbb{S}^{\nu(1)-1} \uplus \mathbb{S}^{\nu(2)-1} \uplus \cdots \uplus \mathbb{S}^{\nu(c)-1},$$

where c is the nilpotency class of N and $\nu(i)$ is the torsion-free rank of the *i*-th quotient in the descending central series of N. Analogously, the projective boundary $\mathcal{P}X$ is homeomorphic to a disjoint union of projective spaces:

$$\mathcal{P}X = \mathbb{P}^{\nu(1)-1} \uplus \mathbb{P}^{\nu(2)-1} \uplus \cdots \uplus \mathbb{P}^{\nu(c)-1}.$$

Having these characterizations of the linear and projective boundaries of vertex-transitive graphs with polynomial growth, immediately the following question arises: When are the linear (projective) boundary of an infinite, connected, vertex-transitive graph X with polynomial growth and the linear (projective) boundary of its automorphism group Aut X bi-Lipschitz-equivalent? Using the concept of bounded automorphisms we are able to present a partial answer to this question.

An automorphism $b \in \operatorname{Aut} X$ is called *bounded* if there is an integer k, depending on b, such that $d(x, b(x)) \leq k$ holds for all $x \in VX$. Of course the bounded automorphisms of ⁶⁸⁶ X give rise to a normal subgroup B(X) of Aut X. As was shown in [GIS⁺89], the same ⁶⁸⁷ holds for the bounded automorphisms of finite order of X. We denote the normal subgroup ⁶⁸⁸ of Aut X generated by all bounded automorphisms of finite order by $B_0(X)$. As was also ⁶⁸⁹ shown in [GIS⁺89], $B_0(X)$ is locally finite, periodic and has finite orbits on X. Furthermore, ⁶⁹⁰ in [Sei91b] the following result concerning $B_0(X)$ was proved:

Proposition 7.13 (Corollary 2.7 in [Sei91b]). Let X be an infinite, connected graph with polynomial growth and let $G \leq \text{Aut } X$ act vertex-transitively on X. Then the orbits of $B_0(X) \cap G$ on X give rise to an imprimitivity system σ of G on VX such that G_{σ} satisfies the assertions of Theorem 7.11.

Together with the following result of Sabidussi [Sab64], Proposition 7.13 now immediately implies a partial answer to the above formulated question. To formulate Sabidussi's result we need another definition.

If X is a graph and m is a cardinal then the graph mX is defined on the Cartesian product of VX by a set M of cardinality m, and

$$E(mX) = \Big\{ \{ (x,i), (y,j) \} : \{ x, y \} \in EX, \, i, j \in M \Big\}.$$

Theorem 7.14 (Theorem 4 in [Sab64]). Let X be a connected graph and let $G \leq \operatorname{Aut} X$ act vertex-transitively on X. Furthermore, let m denote the cardinality of the stabilizer in G of a vertex of X. Then mX is a Cayley graph of G.

Corollary 7.15. Let X be an infinite, connected, vertex-transitive graph with polynomial growth. Then \mathcal{L} Aut X and \mathcal{P} Aut X are bi-Lipschitz-equivalent to \mathcal{L} X and \mathcal{P} X, respectively, if $B_0(X)$ is finite.

Proof. $B_0(X)$ is a normal subgroup of Aut X. If it is in addition finite then it follows from 707 7.13 and 7.10 that the stabilizer of a vertex of X in Aut X has some finite cardinality m. 708 Then, by Theorem 7.14, mX is a Cayley graph of Aut X and arguments quite similar to 709 those in the proof of Theorem 7.9 immediately complete the proof.

In [Tro83] Trofimov defined a *lattice* as a connected locally finite graph X, such that for one of the groups G, acting vertex-transitively on X, there exists an imprimitivity system σ with finite blocks, such that G_{σ} is a finitely generated, commutative group. As was shown in [Tro83], in this case $G \leq B(X)$ holds. Furthermore, it is obvious that lattices have polynomial growth with the same growth rate as G_{σ} . In addition lattices can be characterized as follows:

Theorem 7.16 (Theorem 1 in [Tro83]). Let X be a connected locally finite graph. Then X is a lattice if and only if a group $G \leq B(X)$ acts vertex-transitively on X.

717 This immediately leads to the following:

Theorem 7.17. Let X be a connected locally finite graph of polynomial growth with growth rate r and let a group $G \leq B(X)$ act vertex-transitively on X. Then

$$\mathcal{L}X = \mathbb{S}^{r-1}$$
 and $\mathcal{P}X = \mathbb{P}^{r-1}$.

⁷²⁰ *Proof.* Applying Theorem 7.16 this result can be shown analogously to the proof of Theo-⁷²¹ rem 7.9.

Let X now be a Cayley graph of a group G. Then any group element $g \in G$ gives rise to a bounded automorphism of X if and only if the conjugacy class of g in G is finite (see e.g. [GIS⁺89, page 335]). So the boundedness of an element $g \in G$ is independent of whatever Cayley graph represents G.

A group G is called FC-group if for every $g \in G$ the conjugacy class of g in G is finite. Hence for FC-groups G each $g \in G$ acts as a bounded automorphism on any Cayley graph of G. Therefore Cayley graphs of finitely generated FC-groups are lattices and Theorem 7.17 immediately implies: **Corollary 7.18.** Let G be a finitely generated FC-group with polynomial growth of growth rate r. Then

$$\mathcal{L}G = \mathbb{S}^{r-1}$$
 and $\mathcal{P}G = \mathbb{P}^{r-1}$.

732

747

8. Attaching the boundary

Let Ξ be any subset of \mathcal{U}/\sim . In the following we describe a topology τ on the disjoint union \bar{X} of X and Ξ , such that two requirements hold:

• The subspace topology of τ on X is induced by the metric d.

• If x_1, x_2, \ldots is a sequence in X, which eventually leaves any ball in X, and ξ is an equivalence class in Ξ , such that $x_1, x_2, \ldots \in R$ for some $R \in \xi$ then x_1, x_2, \ldots converges to ξ in τ .

Due to the second requirement the subspace topology of τ on Ξ is in general neither induced by the metric t nor Hausdorff, see Lemma 8.2.

Fix some reference point o in X and let $\xi \in \Xi$ be an equivalence class. If $R \in \xi$ and $\alpha > 0$ and $r \ge 0$ then we set

$$N(R,\alpha,r) = \mathsf{int}(\alpha R \setminus U(o,r)) \uplus \{\zeta \in \Xi : s^+(\xi,\zeta) < \alpha\}$$

where $\operatorname{int}(A)$ is the interior of the set $A \subseteq X$. Note that $N(R, \alpha, p) \subseteq N(S, \beta, q)$ if $R \subseteq S$, $\alpha \leq \beta, p \geq q$. We define the topology τ on $\overline{X} = X \uplus \Xi$ by assigning to each $x \in \overline{X}$ a family \mathcal{V}_x of sets which serves as an open neighborhood base for x:

• If $x \in X$ then \mathcal{V}_x is the family of open balls centered at x.

• If $\xi \in \Xi$ then \mathcal{V}_{ξ} is the family of sets $N(R, \alpha, r)$ with $R \in \xi$, $\alpha > 0$, and $r \ge 0$.

Lemma 8.1. The families \mathcal{V}_x , $x \in \overline{X}$, are open neighborhood bases of a topology τ on \overline{X} . Its subspace topology on X is induced by the metric d, X is dense and open in \overline{X} , and the subspace topology on Ξ is T_0 .

Proof. By Theorem 4.5 in [Wil04] we have to check the following three conditions for all $x \in \overline{X}$:

• If $V \in \mathcal{V}_x$ then $x \in V$.

• If $V_1, V_2 \in \mathcal{V}_x$ then $V_3 \subseteq V_1 \cap V_2$ for some $V_3 \in \mathcal{V}_x$.

• If $V \in \mathcal{V}_x$ and $z \in V$ then $W \subseteq V$ for some $W \in \mathcal{V}_z$.

The first condition is immediate for all $x \in \overline{X}$ and the second and third condition hold for all $x \in X$. Hence let $\xi \in \Xi$. In order to prove the second condition for ξ consider $N(R, \alpha, p), N(S, \beta, q) \in \mathcal{V}_{\xi}$ with $R, S \in \xi, \alpha, \beta > 0$, and $p, q \ge 0$. Choose ε in $(0, \beta)$ and set

$$\gamma = \min\{\alpha, \frac{\beta - \varepsilon}{1 + \varepsilon}\}.$$

Since $R, S \in \xi$, it follows that s(R, S) = 0 and by Lemma 2.5 there is a number $r \ge \max\{p, q\}$ such that $R \setminus U(o, r) \subseteq \varepsilon S$. Using Lemma 2.2 this yields

$$\gamma R \subseteq \gamma(R \setminus U(o, r)) \cup \gamma U(o, r) \subseteq (\gamma + \varepsilon \gamma + \varepsilon)S \cup U(o, (1 + \gamma)r) \subseteq \beta S \cup U(o, (1 + \gamma)r)$$

⁷⁶¹ by the choice of γ . Therefore

$$N(R, \gamma, (1+\gamma)r) \subseteq N(R, \alpha, p) \cap N(S, \beta, q),$$

whence the second condition holds for ξ . The third condition holds for ξ , if $z \in V \cap X$ or $z = \xi$. Hence consider $V = N(R, \alpha, p)$ with $R \in \xi$, $\alpha > 0$, $p \ge 0$, and let $\zeta \neq \xi$ be an element in $V \cap \Xi$. Choose an element S in ζ and choose β in $(s^+(R, S), \alpha)$, which is possible, since $s^+(R, S) = s^+(\xi, \zeta) < \alpha$. There is a number $r \ge p$, such that $S \setminus U(o, r) \subseteq \beta R$. Set $\gamma = \frac{\alpha - \beta}{1 + \beta} > 0$. Then

$$\gamma S \subseteq \gamma(S \setminus U(o, r)) \cup \gamma U(o, r) \subseteq (\gamma + \beta \gamma + \beta)R \cup U(o, (1 + \gamma)r) = \alpha R \cup U(o, (1 + \gamma)r)$$

⁷⁶⁷ by the choice of β and γ . Hence we obtain

$$N(S, \gamma, (1+\gamma)r) \subseteq N(R, \alpha, p).$$

The last three assertions follow from the construction of τ .

20

matric space. Then (\bar{Y}, σ) is cor

Remark. Let X be an unbounded, locally compact, metric space. Then (\bar{X}, τ) is compact if the equivalence class of the unbounded set X is an element of Ξ . If, apart from the equivalence class of X, Ξ contains further elements then (\bar{X}, τ) is not Hausdorff.

Lemma 8.2. Let Ξ be any subset of \mathcal{U}/\sim and let (\bar{X}, τ) be defined as above.

• The space
$$(X, \tau)$$
 is Hausdorff if and only if

$$s^+(\xi,\zeta) = 0 \Longleftrightarrow s^+(\zeta,\xi) = 0$$

for all $\xi, \zeta \in \Xi$. In this case, the subspace topology of τ on Ξ is induced by the metric t.

776

77

777

• Suppose that $\Xi = \mathsf{cl}(\mathcal{E}/\sim)$ for some family $\mathcal{E} \subseteq \mathcal{U}$. If there exists a function

 $f: [0,1] \rightarrow [0,\infty)$, such that f(0) = 0, f is continuous at 0, and $s^+(S,R) \leq 0$

778 $f(s^+(R,S))$ for all $R, S \in \mathcal{E}$ then (\bar{X}, τ) is Hausdorff and the subspace topology 779 of τ on $cl(\mathcal{E}/\sim)$ is induced by the metric t.

Proof. The first assertion is a direct consequence of the definition of the open neighborhood bases \mathcal{V}_{ξ} for $\xi \in \Xi$. The second statement is a consequence of the first, since the hypotheses imply that

$$s^+(\xi,\zeta) = 0 \Longleftrightarrow s^+(\zeta,\xi) = 0$$

for all $\xi, \zeta \in \mathsf{cl}(\mathcal{E}/\sim)$: If $s^+(\xi, \zeta) = 0$ and $\varepsilon > 0$ is given then there are $\xi', \zeta' \in \mathcal{E}/\sim$, such that $s(\xi, \xi') \leq \varepsilon$ and $s(\zeta, \zeta') \leq \varepsilon$. Thus

$$s^{+}(\zeta,\xi) \leq 2\varepsilon + \varepsilon^{2} + s^{+}(\zeta',\xi')(1+\varepsilon)^{2}$$

$$\leq 2\varepsilon + \varepsilon^{2} + f(s^{+}(\xi',\zeta'))(1+\varepsilon)^{2}$$

$$\leq 2\varepsilon + \varepsilon^{2} + f(2\varepsilon + \varepsilon^{2})(1+\varepsilon)^{2}.$$

785 This shows that $s^+(\zeta,\xi) = 0$.

With these preparations we are able to provide a criterion which ensures that the topology defined above on the disjoint union of a compactly generated, locally compact Hausdorff group G and its linear boundary $\mathcal{L}G$ (projective boundary $\mathcal{P}G$) is Hausdorff and the subspace topology on $\mathcal{L}G$ ($\mathcal{P}G$) is induced by the angle metric t.

Proposition 8.3. Let G be a compactly generated, locally compact Hausdorff group. Assume that there exists a constant $C \ge 1$, such that for every group element $g \in G$ with $\langle g \rangle^+ \in C^+G$ there is an element $\tilde{g} \in G$ with the following two properties:

793 • $\langle \tilde{g} \rangle^+ \sim \langle g \rangle^+$ and

794

• $d(1, \tilde{g}^m) \leq Cd(1, \tilde{g}^n) + C$ for all m, n with $0 \leq m \leq n$.

Then the topology τ on $G \uplus \mathcal{L}G$ defined by Lemma 8.1 is Hausdorff and the subspace topology of τ on $\mathcal{L}G$ is induced by the metric t. An analogous statement holds for the projective boundary.

Proof. We check that the function $f: [0,1] \to [0,\infty), x \mapsto 2(1+4C)x$ satisfies the conditions of the second part of Lemma 8.2 which implies the statement.

Of course, f is continuous and f(0) = 0. Furthermore, if $x \ge \frac{1}{2}$, then $f(x) \ge 1 + 4C \ge$ 1. Hence $s^+(\langle h \rangle^+, \langle g \rangle^+) \le f(s^+(\langle g \rangle^+, \langle h \rangle^+))$ is trivially true if $\langle g \rangle^+, \langle h \rangle^+ \in \mathcal{C}^+G$ and $s^+(\langle g \rangle^+, \langle h \rangle^+) \ge \frac{1}{2}$, since $s^+(\langle h \rangle^+, \langle g \rangle^+) \le 1$. Hence we may assume that $s^+(\langle g \rangle^+, \langle h \rangle^+) <$ $\frac{1}{2}$. Additionally, after replacing g by \tilde{g} if necessary, we may assume that $d(1, g^m) \le Cd(1, g^n) +$ C for all m, n with $m \le n$. Choose a number α which satisfies $s^+(\langle g \rangle^+, \langle h \rangle^+) < \alpha < \frac{1}{2}$. Then there is a constant $a \ge 0$, such that $\langle h \rangle^+ \subseteq \alpha \langle g \rangle^+ + a$. Hence, for each $n \in \mathbb{N}_0$ there is an integer $\nu(n) \ge 0$ such that

$$d(h^n, g^{\nu(n)}) \le \alpha d(1, g^{\nu(n)}) + a$$

Now we define the function $\kappa \colon \mathbb{N}_0 \to \mathbb{N}_0$ by

$$\kappa(n) = \min\{m \in \mathbb{N}_0 : \nu(m) \le n \le \nu(m+1)\}.$$

808 We claim that

$$d(g^n, h^{\kappa(n)}) \le 2(1+4C)\alpha d(1, h^{\kappa(n)}) + 2Cd(1, h) + 2a + 8Ca + C$$

for all $n \in \mathbb{N}_0$. Once this claim is established then, by the second assertion of Lemma 8.2, the proof is finished. Let $n \ge 0$ be an integer and set $k = \kappa(n)$. Since

$$d(g^n, h^k) \le d(g^n, g^{\nu(k)}) + d(g^{\nu(k)}, h^k)$$

we need to find upper bounds for $d(g^n, g^{\nu(k)})$ and $d(g^{\nu(k)}, h^k)$, see Figure 2. Then



FIGURE 2. The positive powers of two elements g, h and constellation used in the proof of Proposition 8.3.

811

$$d(1, g^{\nu(k)}) \le d(1, h^k) + d(h^k, g^{\nu(k)}) \le d(1, h^k) + \alpha d(1, g^{\nu(k)}) + \alpha d(1, g$$

812 yields

$$d(1,g^{\nu(k)}) \leq \frac{1}{1-\alpha}(d(1,h^k) + a) \leq 2d(1,h^k) + 2a$$

using the bound $\alpha \leq \frac{1}{2}$. Thus

$$d(g^{\nu(k)}, h^k) \le \alpha d(1, g^{\nu(k)}) + a \le 2\alpha d(1, h^k) + 2a.$$

814 We obtain

$$d(g^{\nu(k)}, g^{\nu(k+1)}) \le d(g^{\nu(k)}, h^k) + d(h^k, h^{k+1}) + d(h^{k+1}, g^{\nu(k+1)})$$

$$\le \alpha d(1, g^{\nu(k)}) + a + d(1, h) + \alpha d(1, g^{\nu(k+1)}) + a$$

815 Then $d(1, g^{\nu(k+1)}) \le d(1, g^{\nu(k)}) + d(g^{\nu(k)}, g^{\nu(k+1)})$ implies $d(g^{\nu(k)}, g^{\nu(k+1)}) \le \alpha d(g^{\nu(k)}, g^{\nu(k+1)}) + 2\alpha d(1)$

$$d(g^{\nu(k)}, g^{\nu(k+1)}) \le \alpha d(g^{\nu(k)}, g^{\nu(k+1)}) + 2\alpha d(1, g^{\nu(k)}) + d(1, h) + 2\alpha d(1, g^{\nu(k)}) + d(1, g^{\nu(k)}) + d(1, g^{\nu(k)}) + d(1, g^{\nu(k)}) +$$

and by rearranging the last inequality we get

$$\begin{aligned} d(g^{\nu(k)}, g^{\nu(k+1)}) &\leq \frac{1}{1-\alpha} (2\alpha d(1, g^{\nu(k)}) + d(1, h) + 2a) \\ &\leq 4\alpha d(1, g^{\nu(k)}) + 2d(1, h) + 4a \\ &\leq 8\alpha d(1, h^k) + 2d(1, h) + 8a \end{aligned}$$

using the bound $\alpha \leq \frac{1}{2}$ twice. The assumption on g implies

$$d(g^n, g^{\nu(k)}) \le Cd(g^{\nu(k)}, g^{\nu(k+1)}) + C \le 8C\alpha d(1, h^k) + 2Cd(1, h) + 8Ca + C.$$

818 Collecting the pieces yields

$$d(g^n, h^k) \le 2(1+4C)\alpha d(1, h^k) + 2Cd(1, h) + 2a + 8Ca + C.$$

Lemma 8.4. Let G be a connected, nilpotent Lie group or a finitely generated, nilpotent group. Then the assumption of the previous proposition on G holds.

Proof. Without loss of generality we may assume that G is simply connected in the Lie case or torsion-free in the discrete case, see Lemma B.1 and Corollary A.6. Furthermore, it is sufficient to prove the statement in the Lie case, as the discrete case follows by embedding G in its real Mal'tsev completion.

Hence suppose that G is a connected, simply connected, nilpotent Lie group and let d_G be a word metric on G. We use the notation of Appendix B. By Lemma B.6 there exists a constant q, such that

$$q^{-1}|x| \le d_G(1, \exp(x)) \le q|x| + q$$

for all $x \in \mathfrak{g}$. We claim that the assumption of the previous proposition holds for $C = q^2$. Let g be a group element of G. Then $g \in G_i$ but $g \notin G_{i+1}$ for some $i \ge 1$. Set $y = \pi_i(\log(g)) \in V_i$ and $h = \exp(y)$. Then, for $0 \le m \le n$, we have $|y^m| = m^{1/i}|y| \le n^{1/i}|y| = |y^n|$ and therefore

$$d_G(1, h^m) \le q \|y^m\| + q \le q \|y^n\| + q \le q^2 d_G(1, h^n) + q.$$

831

9. Random walks on nilpotent groups

Many aspects of random walks on nilpotent groups were studied, see for instance [Ale02, Gui73, Gui80, Kai91, Tan11]. In the sequel we give a simple corollary of some results of Kaimanovich in [Kai91]. Let G be a connected, simply connected, nilpotent Lie group with descending central series

$$G = G_1 \supseteq G_2 \supseteq \dots G_c \supseteq G_{c+1} = \{1\}$$

and let d_G be a word metric on G. A random walk $(S_k)_{k\geq 0}$ on G has finite first moment, whenever

$$E(d_G(1,S_1)) = \int_G d_G(1,g)d\mu(g) < \infty,$$

where μ is the law of S_1 . Note that this notion does not depend on the choice of the word metric. We say that $(S_k)_{k\geq 0}$ has *drift* if there is an integer $n \geq 1$, such that $S_1 \in G_n$ almost surely and $(S_k G_{n+1})_{k\geq 0}$ is a random walk in the commutative group G_n/G_{n+1} with drift, i.e. if we identify G_n/G_{n+1} with $\mathbb{R}^{\nu(n)}$, where $\nu(n)$ is the dimension of G_n/G_{n+1} , then the expected direction $E(S_1G_{n+1}) \in G_n/G_{n+1} = \mathbb{R}^{\nu(n)}$ is non-zero.

Theorem 9.1. Let $(S_k)_{k\geq 0}$ be a random walk with finite first moment and drift on a connected, simply connected, nilpotent Lie group G. Then there is a deterministic group element g, such that $\{S_k : k \geq 0\} \sim \langle g \rangle^+$ holds almost surely. In terms of the topology on $G \uplus \mathcal{L}G$, of Lemma 8.1, this means that almost surely $(S_k)_{k\geq 0}$ converges to the equivalence class of $\langle g \rangle^+$ in $\mathcal{L}G$. On the other hand, every point in $\mathcal{L}G$ is limit point of a random walk with drift (in the sense above).

Proof. Let $n \ge 1$ be the integer, such that $S_1 \in G_n$ almost surely and $(S_k G_{n+1})_{k\ge 0}$ is a random walk with drift. By triviality of Poisson boundary and a result of Kaimanovich (see Theorem 4.2 and the following Remark in [Kai91]) there is a deterministic group element $g \in G_n \ (g \notin G_{n+1}$ by the assumptions), such that $d_{G_n}(S_k, g^k) = o(k)$ almost surely. This implies $d_G(S_k, g^k) = o(k^{1/n})$. Since $d_G(1, g^k) \ge C(g)k^{1/n}$ for some constant C(g) > 0, we obtain $\{S_k : k \ge 0\} \sim \langle g \rangle^+$ almost surely. And the other statements follow.

⁸⁵⁵ On the other hand, every point in $\mathcal{L}G$ is limit point of a corresponding deterministic ⁸⁵⁶ random walk.

Remark. As pointed out by Tanaka [Tan12] the deterministic group element g in the previous theorem is given by $gG_{n+1} = E(S_1G_{n+1})$, where n is the integer, such that $S_1 \in G_n$ almost surely and $(S_kG_{n+1})_{k>0}$ is a random walk with drift.

Remark. A similar statement holds for finitely generated, torsion-free, nilpotent groups. Suppose that G is such a group and consider a random walk $(S_k)_{k\geq 0}$ on G with drift. Then $(S_k)_{k\geq 0}$ converges to an element in $\mathcal{L}G$ with respect to the topological space $(G \uplus \mathcal{L}G, \tau)$, where τ is the topology of Lemma 8.1. On the other hand, every point in $\mathcal{L}G$ is limit point of a random walk with drift. 24

BERNHARD KRÖN, JÖRG LEHNERT, NORBERT SEIFTER, AND ELMAR TEUFL

865

875

876

877

878

908

Appendix A. Compactly generated groups

We provide some results on word metrics of compactly generated, locally compact groups and related issues which are completely analogous to the case of finitely generated groups. The books of Hewitt and Ross [HR79], Stroppel [Str06], and de la Harpe [dlH00] provide a good background on topological and finitely generated groups. We recall some basics from [Gui80].

Lemma A.1 (Proposition 1 in [Gui80]). Let G be a compactly generated, locally compact Hausdorff group.

- If S is a compact, symmetric, generating set then, for some $n \ge 0$, the set S^n contains a neighborhood of 1.
 - A subset of G is compact, if and only if it is closed and bounded with respect to some word metric. Consequently, a subset is bounded if and only if it is relatively compact.
 - If S and S' are two compact symmetric generating sets then the associated word metrics d and d' are bi-Lipschitz-equivalent, i.e. there is a constant q > 0, such that

$$q^{-1}d(x,y) \le d'(x,y) \le qd(x,y)$$

for all $x, y \in G$.

Proof. For sake of completeness we provide a short proof: Since G is Hausdorff, the sets S^n , $n = 0, 1, \ldots$, are closed and their union is equal to G. Hence, as locally compact Hausdorff spaces are Baire spaces (see [Wil04, Corollary 25.4]), there is an integer $n \ge 0$, such that S^n contains a non-empty open subset. Since S^n is symmetric, S^{2n} contains a neighborhood of 1.

Let S be any compact, symmetric, generating subset of G and d be the associated word metric. Choose $n \ge 0$ such that S^n contains some open neighborhood U of 1. Suppose that A is a compact subset of G. Then there are finitely many elements a_1, \ldots, a_r of A such that $A \subseteq a_1 U \cup \cdots \cup a_r U$. Thus

$$d(1,a) \le n \max\{d(1,a_1), \dots, d(1,a_r)\}\$$

for all $a \in A$. Hence A is bounded with respect to d and, since G is assumed to be Hausdorff, the set A is also closed. Now suppose that A is a closed subset of G and bounded with respect to d. Then $A \subseteq S^m$ for some $m \ge 0$ which implies that A is compact.

By the second statement, there is a constant q > 0, such that $d'(1, x) \le q$ for all $x \in S$ and $d(1, y) \le q$ for all $y \in S'$. This implies the third assertion.

A metric space (X, d) is called *q*-quasi-geodesic, if for all $x, y \in X$ there is an integer $n \ge 0$ and points $x = x_0, x_1, \ldots, x_{n-1}, x_n = y$ in X, such that

 $n \leq qd(x, y) + q$ and $d(x_{i-1}, x_i) \leq q$

for all $1 \le i \le n$. We remark that similar notions are used in the literature (see for instance [BH99, Definition 8.22] and [Gro93, Section 0.2.D]). Of course, any geodesic metric space is 1-quasi-geodesic and any word metric on a compactly generated, locally compact group is 1-quasi-geodesic.

In the following we give a straightforward generalization of the classical Milnor-Švarc lemma (see for instance [BH99, Proposition 8.19] or [dlH00, Theorem IV.B.23]) to the continuous case. Before stating the lemma we give a precise description of the setting: Let Gbe a locally compact group and X be a Hausdorff space. Furthermore, let d_X be a quasigeodesic metric on X (we do not assume that d_X induces the topology on X). If not stated otherwise, all topological notions concerning X refer to the topology on X with the exception of boundedness, which refers to the metric d_X . An action $G \times X \to X$, $(g, x) \mapsto gx$ is called

- continuous, if it is a continuous mapping from $G \times X$ to X,
 - q-cobounded, if for all $x, y \in X$ there is a $g \in G$ with $d_X(gx, y) \leq q$,
- proper, if $\{g \in G : d_X(gx, x) \le r\}$ is compact for all $x \in X$ and all $r \ge 0$.

We say that G acts by isometries, if $x \mapsto gx$ is an isometry with respect to d_X for all $g \in G$. Note that if the action is continuous and $K \subseteq G$ is compact then, for any $x \in X$, the set $Kx = \{gx : g \in K\}$ is compact and hence bounded. With these preparations we are ready to state the lemma:

Lemma A.2. Let G be a locally compact Hausdorff group and X be a Hausdorff space which is additionally endowed with a quasi-geodesic metric d_X , such that all compact subsets are bounded. Suppose that there is a continuous, cobounded, proper action of G by isometries on X. Then G is compactly generated and for any $x \in X$ the map $G \to X$, $g \mapsto gx$ is a quasi-isometry from (G, d_G) to (X, d_X) , where d_G is some word metric of G.

919 Proof. Except for minor modifications the proof is the same as in [BH99, dlH00].

For simplicity we assume that the constant q involved in the quasi-geodesic metric is the same as the constant q of the cobounded action. Fix $x \in X$. Since the action is proper, the set $\{g \in G : d(gx, x) \leq 3q\}$ is compact. Let S be the union of this set and its inverse. Then S is compact and symmetric and $1 \in S$.

We show that S generates G. Let $g \in G$. Since (X, d_X) is q-quasi-geodesic, there are $x = x_0, x_1, \ldots, x_n = gx$, such that $n \leq qd(x, gx) + q$ and $d(x_{i-1}, x_i) \leq q$ for $1 \leq i \leq n$. Since the action is q-cobounded, there are group elements $g_0 = 1, g_1, \ldots, g_n = g$, such that $d_X(g_ix, x_i) \leq q$ for all $0 \leq i \leq n$. Then

$$d_X(g_{i-1}^{-1}g_ix, x) = d_X(g_ix, g_{i-1}x) \le d_X(g_ix, x_i) + d_X(x_i, x_{i-1}) + d_X(x_{i-1}, g_{i-1}x) \le 3q.$$

It follows that $s_i = g_{i-1}^{-1}g_i \in S$ and thus $g = g_n = s_1 \cdots s_n \in S^n$. Hence S is a generating set. Let d_G be the word metric on G with respect to S. Then the estimate above for $g \in G$ yields

$$d_G(1,g) \le n \le q d_X(x,gx) + q.$$

Now we prove that $G \to X$, $g \mapsto gx$ is a quasi-isometry from (G, d_G) to (X, d_X) . Let $g, h \in G$. Then we obtain

$$d_G(g,h) = d_G(1,g^{-1}h) \le q d_X(x,g^{-1}hx) + q = q d_X(gx,hx) + q.$$

For the reversed bound, note that Sx is bounded, since S is compact. Hence

$$M = \sup\{d_X(x,y) : y \in Sx\}$$

is finite. Suppose that $d_G(g,h) = n \ge 1$ and $g^{-1}h = s_1 \cdots s_n$ for some $s_1, \ldots, s_n \in S$. Then

$$d_X(gx, hx) = d_X(x, g^{-1}hx) = d_X(x, s_1 \cdots s_n x)$$

$$\leq d_X(x, s_1 x) + d_X(s_1 x, s_1 s_2 x) + \dots + d_X(s_1 \cdots s_{n-1} x, s_1 \cdots s_n x)$$

$$= d_X(x, s_1 x) + d_X(x, s_2 x) + \dots + d_X(x, s_n x)$$

$$\leq Mn = Md_G(g, h).$$

In order to have a handy reference we formulate the following well-known results, see [HR79, Section 5] and [Bou66, Section I.10.2].

937 Lemma A.3. Let G be a Hausdorff group.

- Suppose that H is a subgroup. We write $H \setminus G$ to denote the set of right cosets Hg, g \in G, and equip $H \setminus G$ with the quotient topology. Then the projection $\pi: G \to H \setminus G$ is open (i.e. images of open sets are open). If H is compact then π is also proper (i.e. preimages of compact sets are compact).
- Suppose that H is a Hausdorff group and $\pi: H \to G$ is a continuous and open homomorphism which is onto. If the kernel of π is compact then π is proper.

Example A.4. Let G be a compactly generated, locally compact Hausdorff group with word metric d_G , N a compact Hausdorff group, and H a Hausdorff group. Suppose that

$$\{1\} \longrightarrow N \longrightarrow H \xrightarrow{\pi} G \longrightarrow \{1\}$$

is a topological exact sequence (i.e. all involved homomorphisms are continuous). The action $H \times G \to G$, $(h,g) \mapsto \pi(h)g$ is continuous and it acts by isometries. As π is onto, this action is obviously cobounded. Furthermore, the action is proper, if and only if

$$\{h \in H : d_G(hg,g) \le r\} = \pi^{-1}(gB(1,r)g^{-1})$$

is compact for all $g \in G$ and all $r \geq 0$. Here B(1, r) is the closed ball in G with respect to d_G. If π is an open map, it follows that the action is proper (Lemma A.3) and H is locally compact, since this is an extension property.

Example A.5. Consider a compactly generated, locally compact Hausdorff group G with word metric d_G and let H be a subgroup of G. Then $H \times G \to G$, $(h, g) \to hg$ is a continuous action which acts by isometries. The set $H \setminus G$ inherits a metric $d_{H \setminus G}$ from G:

$$d_{H\setminus G}(Hg_1, Hg_2) = \min\{d_G(h_1g_1, h_2g_2) : h_1, h_2 \in H\}$$

for $g_1, g_2 \in G$, which is well-defined, since d_G is discrete. By left-invariance the action is cobounded, if and only if $(H \setminus G, d_{H \setminus G})$ is bounded. Notice that $(H \setminus G, d_{H \setminus G})$ is bounded, if $H \setminus G$ is compact with respect to the quotient topology of G. To see this, choose $n \geq 1$, such that S^n contains an open neighborhood U of 1. Since the projection $\pi : G \to H \setminus G$ is open (Lemma A.3), $\{\pi(gU) : g \in G\}$ is an open cover of $H \setminus G$. Hence there is a finite subcover $\{\pi(g_1U), \ldots, \pi(g_mU)\}$. Thus any coset of $H \setminus G$ is of the form Hg_iu for some $1 \leq i \leq m$ and some $u \in U$. This yields the bound

$$d_{H\setminus G}(H, Hg_i u) \le d_G(1, g_i u) \le d_G(1, g_i) + d_G(1, u) \\ \le \max\{d_G(1, g_i) : 1 \le i \le m\} + n$$

If H is a closed subgroup then H is locally compact and this action is proper. To see this let $g \in G$ and $r \ge 0$ be given. Then

$${h \in H : d_G(hg,g) \le r} = gB(1,r)g^{-1} \cap H$$

is compact, since $gB(1,r)g^{-1}$ is compact and H is closed.

By an application of the generalized Milnor-Švarc lemma to the situations described in the two previous examples we obtain the following:

Corollary A.6. Consider a compactly generated, locally compact Hausdorff group G with word metric d_G .

969

26

 $\{1\} \longrightarrow N \longrightarrow H \xrightarrow{\pi} G \longrightarrow \{1\}$

• Suppose that N is a compact Hausdorff group and H is a Hausdorff group and that

- is a topological exact sequence, such that $\pi: H \to G$ is open. Then H is compactly generated and locally compact and π is a quasi-isometry from (H, d_H) to (G, d_G) for any word metric d_H on H.
- 973 If H is a closed subgroup of G and $(H \setminus G, d_{H \setminus G})$ is bounded then H is compactly 974 generated and locally compact and the inclusion is a quasi-isometry from (H, d_H) 975 to (G, d_G) for any word metric d_H on H. Furthermore, if $H \setminus G$ is compact, then 976 $(H \setminus G, d_{H \setminus G})$ is bounded.

Finally, we note the following consequence of the Milnor-Švarc lemma, which says, that any reasonable metric on a compactly generated, locally compact Hausdorff group is quasiisometrically equivalent to any word metric on the group.

Corollary A.7. Let G be a locally compact Hausdorff group. Suppose that d_Q is a leftinvariant, q-quasi-geodesic metric on G with the property, that compact subsets are bounded with respect to d_Q and closed balls with respect to d_Q are compact. Then G is compactly generated and d_Q is quasi-isometrically equivalent to any word metric on G. Note that it is not assumed that the metric d_Q induces the group topology. However, the assumptions guarantee some compatibility between the metric d_Q and the group topology. For example, the assumptions on d_Q are satisfied, if d_Q is left-invariant, geodesic, proper and induces the group topology.

Appendix B. Nilpotent Lie groups

The purpose of the appendix is to provide some background on nilpotent Lie groups, see for instance [CG90, Goo76, Hoc65], and, mainly, to prove several technical results, which are used in the proof of Theorem 6.1.

Let G be a group. We denote by $[g,h] = g^{-1}h^{-1}gh$ the commutator in G and define the k-fold commutator inductively by $[g_1] = g_1$ and $[g_1, \ldots, g_k] = [g_1, [g_2, \ldots, g_k]]$. The descending central series of G is inductively defined by

$$\gamma_1(G) = G$$
 and $\gamma_{n+1}(G) = \langle [G, \gamma_n(G)] \rangle$

for $n \ge 1$. A group G is called *nilpotent* if $\gamma_{n+1}(G) = \{1\}$ for some integer n and the least integer n with this property is called *nilpotency class* of G. If A is a subset of G then the set

 $I(A) = \{ g \in G : g^n \in A \text{ for some } n \in \mathbb{N} \}$

997 is called *isolator* of A.

If G is commutative and finitely generated, we denote its torsion-free rank by $\mathsf{rk}(G)$. If G is a commutative, connected Lie group then G is isomorphic to $\mathbb{R}^a \times (\mathbb{R}/\mathbb{Z})^b$ for some integers a, b. In analogy to the discrete case we call a the compact-free dimension of G and denote it by $\mathsf{dim}(G)$.

Lemma B.1. Let G be a nilpotent group and set $G_n = \gamma_n(G)$ for $n \in \mathbb{N}$.

• If G is additionally a connected Lie group then the set C of all compact elements in G is a characteristic, connected, compact subgroup, G/C is simply connected and

$$\dim(\gamma_n(G/C)/\gamma_{n+1}(G/C)) = \dim(G_n/G_{n+1})$$

1005 for all $n \in \mathbb{N}$. 1006 • If G is finitely generated the

• If G is finitely generated then the set T of torsion elements in G is a characteristic, finite subgroup, G/T is torsion-free and

$$\mathsf{rk}(\gamma_n(G/T)/\gamma_{n+1}(G/T)) = \mathsf{rk}(G_n/G_{n+1})$$

1008 for all $n \in \mathbb{N}$.

• If G is finitely generated and torsion-free then $G = I(G_1) \supseteq I(G_2) \supseteq \cdots$ is a central series of G with torsion-free quotients, G_n has finite index in $I(G_n)$ and

$$\mathsf{rk}(I_n(G)/I_{n+1}(G)) = \mathsf{rk}(G_n/G_{n+1})$$

1011 for all $n \in \mathbb{N}$

1012 Proof. Let G be a connected, nilpotent Lie group. Theorem 5.1 in [Glu55] implies the 1013 statements concerning C and G/C. It remains to show the equality concerning dimensions. 1014 By induction we have $\gamma_n(G/C) = G_n C/C$ and it is easy to check that

$$G_n/G_{n+1} \to (G_nC/C)/(G_{n+1}C/C), \quad gG_{n+1} \mapsto gC \cdot (G_{n+1}C/C)$$

1015 is a continuous epimorphism with compact kernel which implies the equality.

Now let G be a finitely generated, nilpotent group. Corollary 1.10 in [Seg83] yields the first part and the assertion concerning ranks follows mutatis mutandis.

Finally, assume that G is a finitely generated, torsion-free, nilpotent group. By Lemma 3.4 in [Seg83] $I(G_1) \supseteq I(G_2) \supseteq ...$ is a central series with torsion-free quotients. Furthermore, it is easy to see that $I(G_n)/G_n = T(G/G_n)$, where $T(G/G_n)$ is the characteristic, finite subgroup of all torsion elements in G/G_n . Consider the map

$$G_n/G_{n+1} \to I(G_n)/I(G_{n+1}), \quad gG_{n+1} \mapsto gI(G_{n+1}).$$

988

1003

1022 This is a homomorphism which has finite kernel and an image of finite index. This yields 1023 the claim concerning ranks. \Box

In the following we fix a connected, simply connected, nilpotent Lie group G with nilpotency class c and set $G_n = \gamma_n(G)$ for $n \in \mathbb{N}$. We denote by \mathfrak{g} the associated Lie algebra and by (x, y) the Lie bracket of \mathfrak{g} . Furthermore, we define the k-fold Lie bracket inductively by $(x_1) = x_1$ and $(x_1, \ldots, x_k) = (x_1, (x_2, \ldots, x_k))$. The descending central series of \mathfrak{g} is

$$\mathfrak{g}_1 = \mathfrak{g}$$
 and $\mathfrak{g}_{n+1} = \operatorname{span}_{\mathbb{R}}(\mathfrak{g}, \mathfrak{g}_n)$

for $n \ge 1$. The Lie algebra of G_n is \mathfrak{g}_n . Let $\nu(n)$ be the compact-free dimension of G_n/G_{n+1} . Then

$$G_n/G_{n+1} \simeq \mathfrak{g}_n/\mathfrak{g}_{n+1} \simeq \mathbb{R}^{\nu(n)}$$

as commutative groups. The *exponential map* exp: $\mathfrak{g} \to G$ is a diffeomorphism from \mathfrak{g} to Gand its inverse is log: $G \to \mathfrak{g}$. The Baker-Campbell-Hausdorff formula yields a multiplicative group structure on \mathfrak{g} :

$$xy = x + y + \frac{1}{2}(x, y) + \frac{1}{12}(x, x, y) - \frac{1}{12}(y, x, y) - \frac{1}{24}(y, x, x, y) \pm \cdots$$

for $x, y \in \mathfrak{g}$. Then the exponential map exp is a group isomorphism from (\mathfrak{g}, \cdot) to (G, \cdot) and it is common to identify the Lie group G with its Lie algebra \mathfrak{g} .

1035 A subgroup Γ is called *uniform* in G, if Γ is discrete and the quotient $\Gamma \setminus G$ is compact. In 1036 the following lemma we study uniform subgroups. Its proof depends on well-known results 1037 on such subgroups which can be found in [CG90, Chapter 5].

1038 Lemma B.2. Let Γ be a uniform subgroup in G and set $\Gamma_n = \gamma_n(\Gamma)$ for $n \in \mathbb{N}$. Then 1039 $\Gamma \cap G_n = I(\Gamma_n)$ and

$$\mathsf{rk}(\Gamma_n/\Gamma_{n+1}) = \mathsf{dim}(G_n/G_{n+1})$$

1040 for all $n \in \mathbb{N}$

1041 *Proof.* First we show that $\Gamma \cap \gamma_n(G) = I(\gamma_n(\Gamma))$ for all $n \in \mathbb{N}$ by backward induction on n:

• Suppose that n = c: Obviously, $I(\Gamma_c) \subseteq \Gamma$ and $I(\Gamma_c) \subseteq G_c$, hence $I(\Gamma_c) \subseteq G_c \cap$ 1042 Γ . To prove the reversed inclusion, note that exp is a group homomorphism from 1043 $(\mathfrak{g}_c, +)$ to (G_c, \cdot) . Let $X \subseteq \mathfrak{g}$ be a strong Mal'tsev basis strongly based on Γ and 1044 set $Z = \exp(X)$. Then $\Gamma_c = \langle [Z, \ldots, Z] \rangle$ (see [MKS04, Theorem 5.4]) and thus 1045 $\log(\Gamma_c) = \operatorname{span}_{\mathbb{Z}}(X, \ldots, X)$, since $\exp((x_1, \ldots, x_c)) = [\exp(x_1), \ldots, \exp(x_c)]$ for all 1046 $x_1, \ldots, x_c \in \mathfrak{g}$. Furthermore, we have $\mathfrak{g}_c = \operatorname{span}_{\mathbb{R}}(X, \ldots, X)$. This implies that Γ_c 1047 and $G_c \cap \Gamma$ are uniform subgroups in G_c . Therefore $(G_c \cap \Gamma)/\Gamma_c$ is finite, whence 1048 $G_c \cap \Gamma \subseteq I(\Gamma_c).$ 1049

• Assume that the claim holds for $n \ge 2$: Consider the groups G/G_n and $\Gamma G_n/G_n$. Then $\Gamma G_n/G_n$ is (topologically) isomorphic to $\Gamma/(\Gamma \cap G_n)$. By φ we denote the canonical isomorphism $\Gamma G_n/G_n \to \Gamma/(\Gamma \cap G_n)$. Since $\Gamma G_n/G_n$ is a uniform subgroup in G/G_n and G/G_n is nilpotent with nilpotency class n-1, using the initial step for the nilpotent group G/G_n yields

$$(\Gamma \cap G_{n-1})G_n/G_n = \Gamma G_n/G_n \cap \gamma_{n-1}(G/G_n) = I(\gamma_{n-1}(\Gamma G_n/G_n)).$$

1055

Applying the isomorphism φ on both sides we obtain

$$(\Gamma \cap G_{n-1})/(\Gamma \cap G_n) = I(\gamma_{n-1}(\Gamma/(\Gamma \cap G_n)))$$
$$= I(\Gamma_{n-1}(\Gamma \cap G_n))/(\Gamma \cap G_n)$$
$$= I(\Gamma_{n-1})/(\Gamma \cap G_n)$$

using the induction hypothesis $\Gamma \cap G_n = I(\Gamma_n)$ once more. It follows that $\Gamma \cap G_{n-1} = I(\Gamma_{n-1})$.

Now we prove the assertion concerning ranks. Since $\Gamma \cap G_k$ is uniform in G_k for all $k \ge 1$, it follows that $(\Gamma \cap G_n)G_{n+1}/G_{n+1}$ is uniform in G_n/G_{n+1} . This implies that

$$\mathsf{rk}((\Gamma \cap G_n)/(\Gamma \cap G_{n+1})) = \mathsf{rk}((\Gamma \cap G_n)G_{n+1}/G_{n+1}) = \dim(G_n/G_{n+1})$$

1060 which yields the statement using the last part of Lemma B.1.

Since \mathfrak{g} is a real vector space of finite dimension $\nu(1) + \cdots + \nu(c)$, there are linear subspaces $V_n \subseteq \mathfrak{g}$ of dimension $\nu(n)$, such that $\mathfrak{g}_n = V_n \oplus \mathfrak{g}_{n+1}$. Hence

$$\mathfrak{g}_n=V_n\oplus\cdots\oplus V_c.$$

Write $\pi_n: \mathfrak{g} \to V_n$ to denote the canonical projection. Then π_n is a continuous epimorphism from (\mathfrak{g}_n, \cdot) to $(V_n, +)$ with kernel \mathfrak{g}_{n+1} . Let $\|\cdot\|_n$ be some ℓ^2 -norm on V_n . Then

$$||x|| = \max\{||\pi_n(x)||_n : 1 \le n \le c\}$$

is a norm on \mathfrak{g} . Notice that $\|\pi_n(x)\| = \|\pi_n(x)\|_n$. Since the Lie bracket (\cdot, \cdot) is bilinear, we have the following simple statement.

1067 Lemma B.3. There is a constant $M \ge 1$, such that $||(x,y)|| \le M||x|| ||y||$ for all $x, y \in \mathfrak{g}$. 1068 Consequently,

$$||(x_1,\ldots,x_k)|| \le M^{\kappa-1}||x_1||\cdots||x_k||$$

1069 for all $x_1, \ldots, x_k \in \mathfrak{g}$.

1070 For $x \in \mathfrak{g}$ set

$$|x| = \max\{\|\pi_n(x)\|^{1/n} : 1 \le n \le c\}.$$

1071 Then $|\cdot|$ is called (homogeneous) gauge or quasi-norm (see for instance [Bre12, Goo76, 1072 Gui73]). Note that $|\cdot|$ is homogeneous with respect to the dilation $\delta_t(x) = t\pi_1(x) + \cdots +$ 1073 $t^c\pi_c(x)$, i.e. $|\delta_t(x)| = t|x|$, and it satisfies a weak form of the triangle inequality with respect 1074 to the Lie group structure on \mathfrak{g} (see Lemma B.5).

1075 **Lemma B.4.** For all $x, y \in \mathfrak{g}$ the following holds:

1076 • |-x| = |x|,

1077 • $|x+y| \leq |x|+|y|$,

1078 • if $x \in \mathfrak{g}_n$ and $\alpha \ge 1$ then $|\alpha x| \le \alpha^{1/n} |x|$,

1079 • if
$$0 \le \alpha \le 1$$
 then $|\alpha x| \le \alpha^{1/c} |x|$.

1080 In any case, $|\alpha x| \leq \max\{1, \alpha\} |x|$ for all $\alpha \geq 0$.

¹⁰⁸¹ The following lemma is a crucial observation due to Guivarc'h [Gui73, Lemme II.1], see ¹⁰⁸² also [Bre12, Lemma 2.5].

Lemma B.5. Let $\alpha > 0$. Then, by appropriately rescaling the norms $\|\cdot\|_n$, we have

$$|xy| \le |x| + |y| + c$$

1084 for all $x, y \in \mathfrak{g}$.

In the sequel we assume that the norms $\|\cdot\|_n$ are chosen appropriately, so that the previous lemma holds with $\alpha = 1$. As a simple consequence we obtain $|(x, y)| \leq 2|x| + 2|y| + 2$ and it follows by induction, that

(2)
$$|(x_1, \dots, x_k)| \le 2^{k-1} (|x_1| + \dots + |x_k|) + 2^k$$

1088 for all $x_1, \ldots, x_k \in \mathfrak{g}$.

Since $(G, \cdot) \simeq (\mathfrak{g}, \cdot)$ is a connected, locally compact group, it is compactly generated. Let d_w be some word metric on the group (\mathfrak{g}, \cdot) . The following result shows a fundamental connection between the gauge $|\cdot|$ and the word metric d_w .

1092 Lemma B.6 (Theorem 2.7 in [Bre12]). There is a constant $q \ge 1$, such that

$$q^{-1}|x| \le d_w(0,x) \le q|x| + q$$

1093 for all $x \in \mathfrak{g}$.

After providing the basic setup and important tools from Lie theory, we now apply the notions of Section 2 to this setting. We write s_w^+ instead of $s_{(\mathfrak{g},d_w)}^+$. The quantity d_a defined by $d_a(x,y) = |-x+y|$ yields by Lemma B.4 a metric on \mathfrak{g} , and as before we write s_a^+ instead of $s_{(\mathfrak{g},d_a)}^+$. Although $(x,y) \mapsto |x^{-1}y|$ is not a metric, we define

$$s_m^+(\langle x \rangle^+, \langle y \rangle^+) = \limsup_{n \to \infty} \inf \left\{ \frac{|y^{-n} x^m|}{|x^m|} : m \in \mathbb{N}_0 \right\}$$

1098 and

$$s_m^+(\langle x \rangle, \langle y \rangle) = \limsup_{|n| \to \infty} \inf \left\{ \frac{|y^{-n} x^m|}{|x^m|} \, : \, m \in \mathbb{N}_0 \right\}$$

for $x, y \in \mathfrak{g} \setminus \{0\}$. Using Lemma 5.5 and Lemma B.6 we get the following comparison of s_w^+ and s_m^+ .

1101 Lemma B.7. Let $x, y \in \mathfrak{g}$ with $x \neq 0$ and $y \neq 0$. Then

$$q^{-2}s_m^+(\langle x \rangle^+, \langle y \rangle^+) \le s_w^+(\langle x \rangle^+, \langle y \rangle^+) \le q^2s_m^+(\langle x \rangle^+, \langle y \rangle^+)$$

1102 and

$$q^{-2}s_m^+(\langle x \rangle, \langle y \rangle) \le s_w^+(\langle x \rangle, \langle y \rangle) \le q^2 s_m^+(\langle x \rangle, \langle y \rangle),$$

1103 where q is the constant of Lemma B.6.

1104 Our goal is the comparison of s_a^+ and s_m^+ . We restrict this comparison to elements of $\mathcal{C}^+\mathfrak{g}$ 1105 and $\mathcal{C}\mathfrak{g}$. Note that $\mathcal{C}^+(\mathfrak{g},\cdot) = \mathcal{C}^+(\mathfrak{g},+)$ and $\mathcal{C}(\mathfrak{g},\cdot) = \mathcal{C}(\mathfrak{g},+)$, since $x^n = nx$ for all $x \in \mathfrak{g}$ and 1106 $n \in \mathbb{Z}$. Before we provide the necessary tools for this comparison, let us identify $\mathcal{L}(\mathfrak{g},d_a)$ 1107 and $\mathcal{P}(\mathfrak{g},d_a)$.

1108 Lemma B.8. Up to homeomorphism we have

$$\mathcal{L}(\mathfrak{g}, d_a) = \mathbb{S}^{\nu(1)-1} \uplus \cdots \uplus \mathbb{S}^{\nu(c)-1}, \qquad \mathcal{P}(\mathfrak{g}, d_a) = \mathbb{P}^{\nu(1)-1} \uplus \cdots \uplus \mathbb{P}^{\nu(c)-1},$$

1109 Moreover, the following three statements yield a precise description of $\mathcal{L}(\mathfrak{g}, d_a)$ and $\mathcal{P}(\mathfrak{g}, d_a)$. 1110 (a) If $x, y \in \mathfrak{g}_i$ and $x + \mathfrak{g}_{i+1} = y + \mathfrak{g}_{i+1} \neq \mathfrak{g}_{i+1}$ then

$$s_a^+(\langle x \rangle^+, \langle y \rangle^+) = 0$$
 and $s_a^+(\langle x \rangle, \langle y \rangle) = 0.$

1111 (b) If $x \in \mathfrak{g}_i$, $x \notin \mathfrak{g}_{i+1}$, and $y \in \mathfrak{g}_{i+1}$ then

$$s_a^+(\langle x \rangle^+, \langle y \rangle^+) = 1$$
 and $s_a^+(\langle x \rangle, \langle y \rangle) = 1.$

1112 (c) If $x, y \in V_i$ and $x, y \neq 0$ then, using the notation of Example 2.11,

$$s_a^+(\langle x \rangle^+, \langle y \rangle^+) = \left(\sin(\min\{\frac{1}{2}\pi, \angle(H_x, H_y)\})\right)^{1/i}$$

1113

$$s_a^+(\langle x \rangle, \langle y \rangle) = \left(\sin(\angle(L_x, L_y))\right)^{1/i}$$

1114 Proof. Once we have proved (a), (b), (c) the statement of the lemma follows. We only prove 1115 these three statements for $s_a^+(\langle x \rangle^+, \langle y \rangle^+)$ the other case being analogous.

1116 Statement (a). By assumption $-y + x \in \mathfrak{g}_{i+1}$, whence

$$|-ny + nx| = |n(-y + x)| \le n^{1/(i+1)} |-y + x|.$$

1117 Since $x \in \mathfrak{g}_i \setminus \mathfrak{g}_{i+1}$, it follows that $\pi_i(x) \neq 0$ and

$$|nx| \ge |\pi_i(nx)| = n^{1/i} |\pi_i(x)|.$$

1118 From this we infer that

and

$$s_a^+(\langle x \rangle^+, \langle y \rangle^+) \le \limsup_{n \to \infty} \frac{|-ny + nx|}{|nx|} \le \limsup_{n \to \infty} \frac{n^{1/(i+1)}|-y + x|}{n^{1/i}|\pi_i(x)|} = 0.$$

1119 Statement (b). Using (a), we may assume that $x \in V_i$. Then $\pi_i(-ny+mx) = mx$ and so

$$|-ny+mx| \ge |\pi_i(-ny+mx)| = |mx|.$$

1120 This implies

$$\inf\left\{\frac{|-ny+mx|}{|mx|}: m \in \mathbb{N}_0\right\} \ge 1$$

1121 and therefore $s_a^+(\langle x \rangle^+, \langle y \rangle^+) \ge 1$.

1121 and therefore $s_a(\langle x \rangle^+, \langle y \rangle^+) = 1$. 1122 Statement (c). Note that $\|v\| = \|v\|^{1/i}$ for all $v \in V_i$. Since $s_a^+(\langle x \rangle^+, \langle y \rangle^+) = s_a^+(H_x, H_y)$, 1123 the statement follows from Example 2.11.

We now compare s_a^+ and s_w^+ . Let y, z be elements in \mathfrak{g} and consider the product $y^{-1}(y + 1)$ z = (-y)(y + z). Then, using the Baker-Campbell-Hausdorff formula,

(3)

$$y^{-1}(y+z) = (-y) + (y+z) + \frac{1}{2}(-y,y+z) + \frac{1}{12}(-y,-y,y+z) + \frac{1}{12}(-y,-y,y+z) + \frac{1}{12}(y+z,-y,y+z) \pm \cdots = z - \frac{1}{2}(y,z) + \frac{2}{12}(y,y,z) + \frac{1}{12}(z,y,z) \pm \cdots$$

1126 Of course in the last expression above at most c-fold Lie brackets occur and, for each $1 \leq 1127$ $k \leq c$, there are finitely many k-fold Lie brackets, say $v_{k,1}, \ldots, v_{k,m(k)}$, whose entries are 1128 either y or z, and each of which contains at least one y and at least one z. If $1 \leq k \leq c$ and 1129 $1 \leq j \leq m(k)$ then write $q_{k,j}$ for the rational coefficient in front of the k-fold Lie bracket 1130 $v_{k,j}$. Then

$$y^{-1}(y+z) = \sum_{1 \le k \le c} \sum_{1 \le j \le m(k)} q_{k,j} v_{k,j}$$

1131 Note that the constants $q_{k,j}$ depend on the Baker-Campbell-Hausdorff formula only. For 1132 convenience we set $Q_{k,j} = \max\{1, q_{k,j}\}$ and

$$Q = \sum_{1 \le k \le c} \sum_{1 \le j \le m(k)} Q_{k,j}$$

1133 Lemma B.9. Suppose that $x, y \in \mathfrak{g}_i$ and $x\mathfrak{g}_{i+1} = y\mathfrak{g}_{i+1} \neq \mathfrak{g}_{i+1}$. Then

$$|y^{-n}x^{n}| \le 2^{c-1}Q(c|x| + c|y| + 2)n^{(1-1/c)/i}$$

1134 for all $n \geq 0$.

1135 Proof. Set z = x - y and m = |x| + |y|. By assumption $z \in \mathfrak{g}_{i+1}$ and obviously $|x|, |y|, |z| \le m$. 1136 m. Using the representation (3) of the product $y^{-1}(y+z)$ we obtain

$$y^{-n}x^n = y^{-n}(y+z)^n = \sum_{1 \le k \le c} \sum_{1 \le j \le m(k)} q_{k,j}n^k v_{k,j}.$$

1137 Since each k-fold Lie bracket $v_{k,j}$ contains at least one z, we get $v_{k,j} \in \mathfrak{g}_{ki+1}$. Using (2) 1138 yields $|v_{k,j}| \leq 2^{k-1}km + 2^k = 2^{k-1}(km+2)$ for all k, j and therefore

$$|q_{k,j}n^k v_{k,j}| \le Q_{k,j}n^{k/(ki+1)}2^{k-1}(km+2).$$

1139 Collecting the pieces, we obtain

$$\begin{aligned} |y^{-n}x^{n}| &\leq \sum_{1 \leq k \leq c} \sum_{1 \leq j \leq m(k)} |q_{k,j}n^{k}v_{k,j}| \\ &\leq \sum_{1 \leq k \leq c} \sum_{1 \leq j \leq m(k)} Q_{k,j}n^{k/(ki+1)}2^{k-1}(km+2) \\ &\leq 2^{c-1}Q(cm+2)n^{(1-1/c)/i} \end{aligned}$$

1140 Lemma B.10. Suppose that $x, y \in V_i$ and $|x| \ge |y| = 1$ and $|x - y| = \alpha |x|$ for some 1141 $\alpha \in [0, 1]$. Then

$$|y^{-n}x^n| \le MQ\alpha^{i/c}|x^n|$$

1142 for all $n \geq 0$.

Proof. Set $z = x - y \in V_i$. Of course $||x|| \ge ||y|| = 1$, and $||z|| = \alpha^i ||x||$. Using the 1143 representation (3) we get as in the proof above 1144

$$y^{-n}x^n = y^{-n}(y+z)^n = \sum_{1 \le k \le c} \sum_{1 \le j \le m(k)} q_{k,j}n^k v_{k,j}.$$

Each k-fold Lie bracket $v_{k,j}$ contains at least one z, but this time $v_{k,j} \in \mathfrak{g}_{ki}$. An application 1145 of Lemma B.3 implies 1146

$$\begin{aligned} \|v_{k,j}\| &= \max\{\|\pi_j(v_{k,j})\|^{1/l} : ik \le l \le c\} \\ &\le \max\{\|v_{k,j}\|^{1/l} : ik \le l \le c\} \\ &\le \max\{(M^{k-1}\alpha^i \|x\|^k)^{1/l} : ik \le l \le c\} \\ &\le M\alpha^{i/c} \|x\|^{1/i} = M\alpha^{i/c} \|x\|. \end{aligned}$$

1147 Hence we obtain

$$\begin{aligned} |y^{-n}x^{n}| &\leq \sum_{1 \leq k \leq c} \sum_{1 \leq j \leq m(k)} |q_{k,j}n^{k}v_{k,j}| \\ &\leq \sum_{1 \leq k \leq c} \sum_{1 \leq j \leq m(k)} Q_{k,j}n^{1/i}M\alpha^{i/c}|x| \\ &= MQ\alpha^{i/c}|x^{n}| \end{aligned}$$

Lemma B.11. The following three statements hold. 1148

(a) If $x, y \in \mathfrak{g}_i$ and $x\mathfrak{g}_{i+1} = y\mathfrak{g}_{i+1} \neq \mathfrak{g}_{i+1}$ then 1149

$$s_m^+(\langle x \rangle^+, \langle y \rangle^+) = 0$$
 and $s_m^+(\langle x \rangle, \langle y \rangle) = 0.$

1150 (b) If
$$x \in \mathfrak{g}_i$$
, $x \notin \mathfrak{g}_{i+1}$, and $y \in \mathfrak{g}_{i+1}$ then

$$s_m^+(\langle x \rangle^+, \langle y \rangle^+) = 1$$
 and $s_m^+(\langle x \rangle, \langle y \rangle) = 1.$

1151 (c) If
$$x, y \in V_i$$
 and $x, y \neq 0$ then

$$s_a^+(\langle x \rangle^+, \langle y \rangle^+) \le s_m^+(\langle x \rangle^+, \langle y \rangle^+) \le MQ \left(s_a^+(\langle x \rangle^+, \langle y \rangle^+) \right)^{i/c}$$

1152

$$s_a^+(\langle x \rangle, \langle y \rangle) \le s_m^+(\langle x \rangle, \langle y \rangle) \le MQ(s_a^+(\langle x \rangle, \langle y \rangle))^{i/c}.$$

Proof. Statement (a). By assumption $\pi_i(x) \neq 0$ and we get 1153

$$|x^n| \ge |\pi_i(x^n)| = n^{1/i} |\pi_i(x)|.$$

On the other hand Lemma B.9 implies 1154

$$|y^{-n}x^{n}| \le 2^{c-1}Q(c|x| + c|y| + 2)n^{(1-1/c)/i}$$

1155 for all $n \ge 0$. Hence

and

$$s_m^+(\langle x \rangle^+, \langle y \rangle^+) \le \limsup_{n \to \infty} \frac{|y^{-n} x^n|}{|x^n|}$$
$$\le \limsup_{n \to \infty} \frac{2^{c-1}Q(c|x| + c|y| + 2)n^{(1-1/c)/i}}{n^{1/i}|\pi_i(x)|} = 0.$$

Statement (b): By the first claim we may assume that $x \in V_i$. Using the Baker-Campbell-1156 Hausdorff formula we obtain $\pi_i(y^{-n}x^m) = x^m$ and thus 1157

$$|y^{-n}x^{m}| \ge |\pi_{i}(y^{-n}x^{m})| = |x^{m}|.$$

This implies 1158

$$\inf\left\{\frac{|y^{-n}x^m|}{|x^m|}: m \in \mathbb{N}_0\right\} \ge 1$$

1159 and $s_m^+(\langle x \rangle^+, \langle y \rangle^+) \ge 1$.

32

1160 Statement (c): To prove the lower bound, note that

$$|y^{-n}x^{m}| \ge |\pi_{i}(y^{-n}x^{m})| = |-ny + mx|$$

1161 for all $n, m \in \mathbb{N}_0$. This implies $s_m^+(\langle x \rangle^+, \langle y \rangle^+) \ge s_a^+(\langle x \rangle^+, \langle y \rangle^+)$. 1162 Now we prove the upper bound. Set $\alpha = s_a^+(\langle x \rangle^+, \langle y \rangle^+)$. Without loss of generality

1162 Now we prove the upper bound. Set $\alpha = s_a^+(\langle x \rangle^+, \langle y \rangle^+)$. Without loss of generality 1163 we may assume that $\alpha < 1$. Furthermore, we may scale x and y by positive constants 1164 without changing the value of $s_a^+(\langle x \rangle^+, \langle y \rangle^+)$ or of $s_m^+(\langle x \rangle^+, \langle y \rangle^+)$. Hence we may assume 1165 that ||y|| = 1 and y is orthogonal to x - y with respect to the inner product on V_i associated 1166 with $||\cdot||$, see Figure 3. As a consequence we get $1 = ||y|| \le ||x||$ and $||x - y|| = \alpha^i ||x||$ (due to



FIGURE 3. The constraints for the choice of x and y.

1167 Lemma B.8). Then $1 = |y| \le |x|$ and $|x - y| = \alpha |x|$. By Lemma B.10 we get

 $|y^{-n}x^n| < MQ\alpha^{i/c}|x^n|$

1168 for all $n \ge 0$. Thus

$$s_m^+(\langle x \rangle^+, \langle y \rangle^+) \le \limsup_{n \to \infty} \frac{|y^{-n} x^n|}{|x^n|} \le MQ\alpha^{i/c}.$$

1169

1166

References

Georgios K. Alexopoulos, Random walks on discrete groups of polynomial volume growth, Ann. [Ale02]1170 Probab. 30 (2002), no. 2, 723-801, doi:10.1214/aop/1023481007, MR1905856 (2003d:60010), 1171 Zbl1023.60007.23 1172 László Babai, Some applications of graph contractions, J. Graph Theory 1 (1977), no. 2, 125-[Bab77] 1173 130, Special issue dedicated to Paul Turán, doi:10.1002/jgt.3190010207, MR0460171 (57 #167), 1174 Zbl0381.05029.17 1175 Martin R. Bridson and André Haefliger, Metric spaces of non-positive curvature, Grundlehren [BH99] 1176 der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 319, 1177 Springer-Verlag, Berlin, 1999, MR1744486 (2000k:53038), Zbl0988.53001. 9, 24, 25 1178 [Bou66] Nicolas Bourbaki, Elements of mathematics. General topology. Part 1, Hermann, Paris, 1966, 1179 MR0205210 (34 #5044a), Zbl0301.54001. 25 1180 Emmanuel Breuillard, Geometry of locally compact groups of polynomial growth and shape of large [Bre12] 1181 balls, preprint, 2012. 29 1182 [BRW07] C. Paul Bonnington, R. Bruce Richter, and Mark E. Watkins, Between ends and fibers, J. 1183 Graph Theory 54 (2007), no. 2, 125–153, doi:10.1002/jgt.20202, MR2285455 (2007k:05100), 1184 Zbl1118.05017.2,3 1185 Lawrence J. Corwin and Frederick P. Greenleaf, Representations of nilpotent Lie groups and their [CG90] 1186 applications. Part I, Cambridge Studies in Advanced Mathematics, vol. 18, Cambridge University 1187 Press, Cambridge, 1990, Basic theory and examples, MR1070979 (92b:22007), Zb10704.22007. 27, 1188 1189 Pierre de la Harpe, Topics in geometric group theory, Chicago Lectures in Mathematics, University [dlH00] 1190 of Chicago Press, Chicago, IL, 2000, MR1786869 (2001i:20081), Zbl0965.20025. 24, 25 1191 P. Eberlein and B. O'Neill, Visibility manifolds, Pacific J. Math. 46 (1973), [EO73] 45 - 109.1192 doi:10.2140/pjm.1973.46.45, MR0336648 (49 #1421), Zbl0264.53026.1 1193 Hans Freudenthal, Neuaufbau der Endentheorie, Ann. of Math. (2) 43 (1942), 261–279, [Fre42] 1194 doi:10.2307/1968869, MR0006504 (3,315a), Zbl0060.40006.1 1195 $[GIS^+89]$ Chris D. Godsil, Wilfried Imrich, Norbert Seifter, Mark E. Watkins, and Wolfgang Woess, A 1196 note on bounded automorphisms of infinite graphs, Graphs Combin. 5 (1989), no. 4, 333-338, 1197 doi:10.1007/BF01788688, MR1032384 (91c:05092), Zbl0714.05029.19 1198 1199 [Glu55] Viktor M. Gluškov, Locally nilpotent locally bicompact groups, Trudy Moskov. Mat. Obšč. 4 (1955), 1200 291-332, MR0072422 (17,281b), Zbl0068.01901. 27

- 1201[Goo76]Roe W. Goodman, Nilpotent Lie groups: structure and applications to analysis, Lecture Notes in1202Mathematics, Vol. 562, Springer-Verlag, Berlin, 1976, MR0442149 (56 #537), Zbl0347.22001. 27,120329
- I204 [Gro81] Mikhael Gromov, Hyperbolic manifolds, groups and actions, Riemann surfaces and related topics:
 Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y.,
 1206 1978), Ann. of Math. Stud., vol. 97, Princeton Univ. Press, Princeton, N.J., 1981, pp. 183–213,
 1207 MR624814 (82m:53035), Zbl0467.53035. 2
- 1208[Gro93]_____, Asymptotic invariants of infinite groups, Geometric group theory, Vol. 2 (Sussex, 1991),1209London Math. Soc. Lecture Note Ser., vol. 182, Cambridge Univ. Press, Cambridge, 1993, pp. 1–295,1210MR1253544 (95m:20041), Zbl0841.20039. 24
- 1211 [Gui73]
 Yves Guivarc'h, Croissance polynomiale et périodes des fonctions harmoniques, Bull.

 1212
 Soc. Math. France 101 (1973), 333-379, MR0369608 (51 #5841), Zbl0294.43003,

 1213
 http://www.numdam.org/item?id=BSMF_1973_101_333_0.23, 29
- 1214 [Gui80]..., Sur la loi des grands nombres et le rayon spectral d'une marche aléatoire, Conference on1215Random Walks (Kleebach, 1979) (French), Astérisque, vol. 74, Soc. Math. France, Paris, 1980,1216pp. 47–98, 3, MR588157 (82g:60016), Zbl0448.60007. 23, 24
- 1217 [Hoc65] Gerhard P. Hochschild, *The structure of Lie groups*, Holden-Day Inc., San Francisco, 1965,
 1218 MR0207883 (34 #7696), Zbl0131.02702. 27
- 1219[HR79]Edwin Hewitt and Kenneth A. Ross, Abstract harmonic analysis. Vol. I, second ed., Grundlehren1220der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 115,1221Springer-Verlag, Berlin, 1979, Structure of topological groups, integration theory, group represen-1222tations, MR551496 (81k:43001), Zbl0416.43001. 11, 12, 24, 25
- 1223 [JN95]
 Heinz A. Jung and Peter Niemeyer, Decomposing ends of locally finite graphs, Math. Nachr. 174

 1224
 (1995), 185–202, doi:10.1002/mana.19951740113, MR1349044 (96h:05120), Zbl0833.05052. 2
- 1225[Jun93]Heinz A. Jung, Notes on rays and automorphisms of locally finite graphs, Graph structure theory1226(Seattle, WA, 1991), Contemp. Math., vol. 147, Amer. Math. Soc., Providence, RI, 1993, pp. 477-1227484, MR1224725 (94d:05075), Zbl0787.05048. 2
- 1228 [Kai91] Vadim A. Kaimanovich, Poisson boundaries of random walks on discrete solvable groups, Prob1229 ability measures on groups, X (Oberwolfach, 1990), Plenum, New York, 1991, pp. 205–238,
 1230 MR1178986 (94m:60014), Zbl0823.60006. 2, 23
- [KLS12] Berhnard Krön, Jörg Lehnert, and Maya J. Stein, *Linear boundary and HNN-extensions*, preprint, 2012. 11
- 1233 [Mal51] Anatolii I. Mal'tsev, On a class of homogeneous spaces, Amer. Math. Soc. Translation 1951 (1951),
 1234 no. 39, 33, MR0039734 (12,589e), ZblZbl 0034.01701. 14
- [MKS04] Wilhelm Magnus, Abraham Karrass, and Donald Solitar, *Combinatorial group theory*, second ed.,
 Dover Publications Inc., Mineola, NY, 2004, Presentations of groups in terms of generators and
 relations, MR2109550 (2005h:20052), Zbl1130.20307. 28
- 1238 [Pan89]
 Pierre Pansu, Métriques de Carnot-Carathéodory et quasiisométries des espaces symétriques de rang

 1239
 un, Ann. of Math. (2)
 129 (1989), no. 1, 1–60, doi:10.2307/1971484, MR979599 (90e:53058),

 1240
 Zbl0678.53042. 14
- 1241 [Sab64]
 Gert Sabidussi, Vertex-transitive graphs, Monatsh. Math. 68 (1964), 426–438,

 1242
 doi:10.1007/BF01304186, MR0175815 (31 #91), Zbl0136.44608. 19
- 1243 [Seg83] Daniel Segal, *Polycyclic groups*, Cambridge Tracts in Mathematics, vol. 82, Cambridge University
 1244 Press, Cambridge, 1983, doi:10.1017/CB09780511565953, MR713786 (85h:20003), Zbl0516.20001.
 1245 27
- 1246 [Sei91a]
 Norbert Seifter, Groups acting on graphs with polynomial growth, Discrete Math. 89 (1991), no. 3,

 1247
 269–280, doi:10.1016/0012-365X(91)90120-Q, MR1112445 (92g:05099), Zbl0739.05041. 18
- 1248 [Sei91b]
 ______, Properties of graphs with polynomial growth, J. Combin. Theory Ser. B 52 (1991), no. 2,

 1249
 222-235, doi:10.1016/0095-8956(91)90064-Q, MR1110471 (92i:05106), Zbl0668.05034. 18, 19
- 1250 [Str06]Markus Stroppel, Locally compact groups, EMS Textbooks in Mathematics, European Mathematical1251Society (EMS), Zürich, 2006, doi:10.4171/016, MR2226087 (2007d:22001), Zbl1102.22005. 24
- 1252 [Tan11]
 Ryokichi Tanaka, Large deviation on a covering graph with group of polynomial growth, Math. Z.

 1253
 267 (2011), no. 3-4, 803–833, doi:10.1007/s00209-009-0647-z, MR2776059 (2012i:60051). 23

 1254 [Tan12]
 ______, 2012, private communication. 23
- 1255 [Tro83]
 Vladimir I. Trofimov, Bounded automorphisms of graphs and the characterization of grids, Izv.

 1256
 Akad. Nauk SSSR Ser. Mat. 47 (1983), no. 2, 407–420, MR697303 (84h:05067), Zbl0519.05037. 19

 1257
 [Tro84]
 ______, Graphs with polynomial growth, Mat. Sb. (N.S.) 123(165) (1984), no. 3, 407–421, doi:10.1070/SM1985v051n02ABEH002866, MR735714 (85m:05041), Zbl0548.05033. 18
- 1259 [Wil04]Stephen Willard, General topology, Dover Publications Inc., Mineola, NY, 2004, Reprint of the 19701260original [Addison-Wesley, Reading, MA; MR0264581], MR2048350, Zbl1052.54001. 20, 24

- 1261 BERNHARD KRÖN, FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN, NORDBERGSTRASSE 15, 1090 VI-
- 1262 ENNA, AUSTRIA
- 1263 E-mail address: bernhard.kroen@univie.ac.at
- 1264 JÖRG LEHNERT, MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES, INSELSTRASSE 22, 04103 1265 LEIPZIG, GERMANY
- 1265 LEIPZIG, GERMANY
- 1266 *E-mail address*: lehnert@mis.mpg.de
- 1267 NORBERT SEIFTER, DEPARTMENT MATHEMATIK UND INFORMATIONSTECHNOLOGIE, MONTANUNIVERSITÄT
- 1268 Leoben, Franz-Josef-Strasse 18, 8700 Leoben, Austria
- 1269 E-mail address: seifter@unileoben.ac.at
- 1270 Elmar Teufl, Mathematisches Institut, Universität Tübingen, Auf der Morgenstelle 10, 72076
- 1271 TÜBINGEN, GERMANY
- 1272 E-mail address: elmar.teufl@uni-tuebingen.de