Note on the bondage number of graphs on topological surfaces

Vladimir Samodivkin

Department of Mathematics, UACEG, Hristo Smirnenski Blv. 1, 1046 Sofia, Bulgaria E-mail: vl.samodivkin@uacg.bg

August 30, 2012

Abstract

The bondage number b(G) of a graph G is the smallest number of edges whose removal from G results in a graph with larger domination number. In this paper we present new upper bounds for b(G) in terms of girth, order and Euler characteristic.

Keywords: Bondage number, girth, genus, Euler characteristic.

MSC Classification: 05C35 (Primary).

1 Introduction and main results

We shall consider graphs without loops and multiple edges. An orientable compact 2-manifold \mathbb{S}_h or orientable surface \mathbb{S}_h (see [8]) of genus h is obtained from the sphere by adding h handles. Correspondingly, a non-orientable compact 2-manifold \mathbb{N}_k or non-orientable surface \mathbb{N}_k of genus k is obtained from the sphere by adding k crosscaps. The Euler characteristic is defined by $\chi(\mathbb{S}_h) = 2 - 2h$, $h \geq 0$, and $\chi(\mathbb{N}_k) = 2 - k$, $k \geq 1$. Compact 2-manifolds are called simply surfaces throughout the paper. If a graph G is embedded in a surface \mathbb{M} then the connected components of $\mathbb{M} - G$ are called the faces of G. If each face is an open disc then the embedding is called a 2-cell embedding. For such a graph G, we denote its vertex set, edge set, face set, maximum degree, and minimum degree by V(G), E(G), F(G), $\Delta(G)$, and $\delta(G)$, respectively. Set |G| = |V(G)|, ||G|| = |E(G)|, and f(G) = |F(G)|. We call

|G| and |G| the order and the size of G. For a 2-cell embedding in a surface \mathbb{M} the (generalized) Euler's formula states $|G| - ||G|| + f(G) = \chi(\mathbb{M})$ for any multigraph G that is 2-cell embedded in \mathbb{M} [8, p. 85]. The Euclidean plane \mathbb{S}_0 , the projective plane \mathbb{N}_1 , the torus \mathbb{S}_1 , and the Klein bottle \mathbb{N}_2 are all the surfaces of nonnegative Euler characteristic. The degree of a face is the length of its boundary walk. A face of degree i is called a i-face. For $i \geq 3$, let $f_i(G)$ denote the number of i-faces in the embedded graph G. We say that two faces are intersecting or adjacent if they share a common vertex or a common edge, respectively. The girth of a graph G, denoted as g(G), is the length of a shortest cycle in G. If G has no cycle then $g(G) = \infty$.

A dominating set for a graph G is a subset $D \subseteq V(G)$ of vertices such that every vertex not in D is adjacent to at least one vertex in D. The minimum cardinality of a dominating set is called the domination number of G. The concept of domination in graphs has many applications in a wide range of areas within the natural and social sciences. One measure of the stability of the domination number of G under edge removal is the bondage number b(G), defined in [2] (previously called the domination line-stability in [1]) as the smallest number of edges whose removal from G results in a graph with larger domination number. In general it is hard to determine the bondage number b(G) (see Hu and Xu [6]), and thus useful to find bounds for it.

The main result of the paper is the following theorem.

Theorem 1. Let G be a graph embeddable on a surface whose Euler characteristic χ is as large as possible and let $g(G) = g < \infty$.

- (i) Then $b(G) \le 3 + \frac{8}{g-2} \frac{4\chi g}{|G|(g-2)}$.
- (ii) If G contains no intersecting g-faces, then $b(G) \leq 3 + \frac{8g+4}{g^2-g} 4(1 + \frac{2}{g-1})\frac{\chi}{|G|}$.
- (iii) If G contains no adjacent g-faces, then $b(G) \leq \frac{4g(g+1)}{q^2-q-1}(1-\frac{\chi}{|G|})-1$.

Remark 1. If G is a planar graph with girth $g \ge 4 + i$, $i \in \{0, 1, 2\}$ then Theorem 1(i) leads to $b(G) \le 6 - i$. These results were first proved by Fischermann et al.[3].

Recently, the following results on bondage number of graphs on surfaces were obtained.

Theorem 2 (Gagarin and Zverovich [4]). Let G be a graph embeddable on an orientable surface of genus h and a non-orientable surface of genus k. Then $b(G) \leq \min\{\Delta(G) + h + 2, \Delta(G) + k + 1\}$.

Theorem 3 (Jia Huang [7]). Let G be a graph embeddable on a surface whose Euler characteristic χ is as large as possible. If $\chi \leq 0$ then $b(G) < \Delta(G) + \sqrt{12-6\chi}+1/2$. If $\chi \leq 0$ then $b(G) \leq \Delta(G) + \frac{\sqrt{8g(2-g)\chi+(3g-2)^2}-g+6}{2(g-2)}$.

Remark 2. In many cases the bound stated in Theorem 1(i) is better than those given by Theorems 2 and 3. Indeed, it is easy to see that if $\chi \leq 0$ then:

- (a) $s(\chi,g,|G|) < z(\Delta,h,k)$ at least when both $\Delta(G) \geq \frac{8}{g-2}$ and $|G| > 8 + \frac{16}{g-2}$
- (b) $s(\chi, 3, |G|) < j_1(\Delta, \chi)$ at least when both $\Delta(G) \ge 11$ and $-\frac{|G|^2}{24} \le \chi$ hold;
- (c) $s(\chi, g, |G|) < j_2(\Delta, \chi, g)$ at least when both $\Delta(G) \ge \frac{7}{2} + \frac{6}{g-2}$ and $-\frac{|G|^2}{8}(1-g)$ $\frac{2}{a}$) $\leq \chi \ hold$,

where (under the notation of Theorems 1, 2 and 3): $s(\chi, g, |G|) = 3 + \frac{8}{g-2} - \frac{4\chi g}{|G|(g-2)}$, $z(\Delta, h, k) = \min\{\Delta(G) + h + 2, \Delta(G) + k + 1\}$, $j_1(\Delta, \chi) = \Delta(G) + \sqrt{12 - 6\chi} + 1/2$ and $j_2(\Delta, \chi, g) = \Delta(G) + \frac{\sqrt{8g(2-g)\chi + (3g-2)^2 - g + 6}}{2(g-2)}$.

2 Proof of the main result

The average degree of a graph G is defined as ad(G) = 2||G||/|G|. For the proof of Theorem 1 we needs the following lemmas.

Lemma 4 (Hartnell and Rall [5]). For any graph G, $b(G) \leq 2ad(G) - 1$.

Lemma 5. Let G be a connected graph embeddable on a surface whose Euler characteristic χ is as large as possible and let $g(G) = g < \infty$.

- (i) Then $ad(G) \leq \frac{2g}{g-2}(1-\frac{\chi}{|G|})$.
- (ii) If G contains no intersecting g-faces, then $ad(G) \leq 2 + \frac{4g+2}{g^2-g} 2(1+\frac{2}{g-1})\frac{\chi}{|G|}$.
- (iii) If G contains no adjacent g-faces, then $ad(G) \leq \frac{2g(g+1)}{g^2-g-1}(1-\frac{\chi}{|G|})$.

- *Proof.* By Euler's formula, $f(G) = \chi |G| + \frac{1}{2}ad(G)|G|$. (i) We have $ad(G)|G| = 2||G|| = \sum_{i \geq g} i f_i(G) \geq gf(G) = g(\chi |G| + \frac{1}{2}ad(G)|G|)$ and the result easy follows.
- (ii) Since G contains no intersecting g-faces, each vertex is incident to at most one g-face. This implies $gf_g(G) \leq |G|$. Hence $ad(G)|G| = 2||G|| = \sum_{i \geq g} if_i(G) \geq i$

 $(g+1)f(G)-f_g(G) \ge (g+1)(\chi-|G|+\frac{1}{2}ad(G)|G|)-\frac{|G|}{g}$. After a short computation, the result follows.

(iii) Since G contains no adjacent g-faces, it follows that $gf_g(G) \leq ||G|| = \frac{1}{2}ad(G)|G|$. Hence $ad(G)|G| \geq (g+1)f(G) - f_g(G) \geq (g+1)(\chi - |G| + \frac{1}{2}ad(G)|G|) - \frac{1}{2g}ad(G)|G|$. After some obvious manipulations we obtain the result.

It remains to note that Theorem 1 follows by combining Lemmas 4 and 5.

References

- [1] D. Bauer, F. Harary, J. Nieminen, C.L. Suffel, Domination alteration sets in graphs, Discrete Math. 47 (1983) 153–161.
- [2] J.F. Fink, M.J. Jacobson, L.F. Kinch and J. Roberts, The bondage number of a graph, Discrete Math. 86 (1990) 47–57.
- [3] M. Fischermann, D. Rautenbach and L. Volkmann, Remarks on the bondage number of planar graphs, Discrete Math. 260 (2003) 57–67.
- [4] A. Gagarin and V. Zverovich, Upper bounds for the bondage number of graphs on topological surfaces, Discrete Math. http://dx.doi.org/10.1016/j.disc.2011.10.018
- [5] B.L. Hartnell and D.F. Rall, A bound on the size of a graph with given order and bondage number, Discrete Math. 197/198 (1999) 409–413.
- [6] F.-T. Hu and J.-M. Xu, On the complexity of the bondage and reinforcement problems, Journal of Complexity. 28(2012) 192-201.
- [7] J. Huang, An improved upper bound for the bondage number of graphs on surfaces, Discrete Math. 312(2012) 2776–2781.
- [8] G. Ringel, Map Color Theorem, Springer-Verlag, Berlin, 1974.