# THE LIFETIME OF SHAPE OSCILLATIONS OF A BUBBLE IN AN UNBOUNDED, INVISCID AND COMPRESSIBLE FLUID WITH SURFACE TENSION

O. COSTIN, S. TANVEER, AND M.I. WEINSTEIN

Abstract. General perturbations of a spherical gas bubble in a compressible and inviscid fluid with surface tension were proved in [\[5\]](#page-13-0), in the linearized approximation, to decay exponentially,  $\sim e^{-\Gamma t}$ ,  $\Gamma > 0$ , as time advances. Formal asymptotic and numerical evidence led to the conjecture that  $\Gamma \approx$  $\frac{A}{\epsilon} \frac{We}{\epsilon^2}$  exp  $\left(-B\frac{We}{\epsilon^2}\right)$ , where  $0 < \epsilon \ll 1$  is the Mach number and A and B are positive constants. In this paper, we prove this conjecture and calculate A and  $B$  to leading order in  $\epsilon$ .

## 1. Introduction and outline

The detailed non-spherical deformations of gas bubbles in a liquid is a problem of great physical interest in fundamental and applied physics; see, for example, [\[3,](#page-13-1) [4,](#page-13-2) [5\]](#page-13-0) and references cited therein. We consider the dynamics of a gas bubble in a compressible, inviscid and irrotational fluid with surface tension. This physical system has an equilibrium state, consisting of: a spherically symmetric gas bubble at constant pressure inside the bubble, and a fluid at constant (lower) pressure and vanishing velocity, outside the bubble.

In [\[5\]](#page-13-0) the linearized dynamics about such equilibria was studied and it was shown that general finite energy perturbations of the spherical equilibrium damp out as time advances. In particular, the  $L^{\infty}$  and local energy norms of the perturbation tend to zero exponentially,  $\sim e^{-\Gamma t}$ , as  $t \to \infty$ . Formal asymptotic and numerical evidence led to the conjecture that

<span id="page-0-1"></span>(1) 
$$
\Gamma(\epsilon) \approx \frac{A}{\epsilon} \frac{We}{\epsilon^2} \exp\left(-B\frac{We}{\epsilon^2}\right)
$$

Here,  $0 \leq \epsilon \ll 1$  denotes the Mach number, a dimensionless ratio of speeds, and We denotes the Weber number, a dimensionless measure of surface tension. A and B are positive constants determined in [\[5\]](#page-13-0) by computer simulation.  $(1)$ 

.

The asymptotics [\(1\)](#page-0-1) are in marked contrast to the decay rate of perturbations which are spherically symmetric:

$$
\Gamma_{\text{radial}}(\epsilon) = \mathcal{O}(\epsilon)
$$

Thus, non-spherical deformations excite shape modes which lose their energy to the fluid and radiate sound waves very slowly.

<span id="page-0-0"></span><sup>(1)</sup> The Mach number,  $\epsilon$ , is the ratio of the bubble-wall radial velocity to the sounds speed in the fluid, exterior to the bubble. We shall later set  $We$  equal to one and consider the asymptotics for small  $\epsilon.$ 

Our goal in this article is to present a proof of the above conjecture and a calculation of A and B to leading order in  $\epsilon$ .

A systematic and detailed discussion of the physical problem, the full nonlinear compressible equations and the appropriate linearization is presented in [\[5\]](#page-13-0). The dynamics of the linearized velocity potential,  $\Psi(x, t)$ ,  $x \in \mathbb{R}^3$ , and the bubble shape perturbation,  $\beta(\Omega, t)$ ,  $\Omega \in S^2$  is governed by the wave-system:

<span id="page-1-7"></span><span id="page-1-1"></span>
$$
(3a)
$$

<span id="page-1-2"></span>
$$
\epsilon^2 \partial_t^2 \Psi - \Delta \Psi = 0, \qquad \qquad r = |x| > 1
$$
  
(3b) 
$$
\partial_r \Psi = \partial_t \beta, \qquad \qquad r = 1
$$

<span id="page-1-3"></span>(3c) 
$$
\partial_t \Psi = 3\gamma \left(\frac{1}{2} + \frac{2}{We}\right) \langle \beta, Y_0^0 \rangle - \frac{1}{We} (2 + \Delta_S) \beta,
$$
  $r = 1$ 

<span id="page-1-4"></span>(3d) 
$$
\langle \beta, Y_1^m \rangle = 0
$$
.  $|m| \le 1$ 

[\(2\)](#page-1-0) Equation [\(3a\)](#page-1-1) follows from linearization of the continuity and Euler momentum equations, exterior to the equilibrium bubble. Equation [\(3b\)](#page-1-2) is the linearization of the kinematic boundary condition. Equation [\(3c\)](#page-1-3) is the linearization of the Laplace-Young boundary condition, stating that the pressure jump across the gas - fluid interface is proportional to the mean curvature. Finally, [\(3d\)](#page-1-4) is the linearization of the statement that the origin of coordinates (the center of the bubble) is in a frame of reference moving with the bubble center of mass. The equilibrium bubble has been normalized to have unit radius. Here,  $Y_l^m = Y_l^m(\Omega)$ ,  $\Omega \in S^2$  are spherical harmonics, which satisfy  $-\Delta_S Y_l^m = l(l+1)Y_l^m$ ,  $l \geq 0$ ,  $|m| \leq l$ , where  $\langle Y_l^m, Y_{l'}^{m'} \rangle_{L^2(S^2)} = \delta_{ll'} \delta_{mm'}$ . In particular,  $Y_0^0(\Omega) \equiv (4\pi)^{-1}$ .

As explained in [\[5\]](#page-13-0) and now outlined, the  $L^{\infty}$  and local-energy decay, for  $\epsilon$  small, is controlled by a non-selfadjoint eigenvalue problem, which arises by seeking timeharmonic solutions of  $(3a)-(3d)$  $(3a)-(3d)$ :

$$
\Psi(\Omega, t) = e^{-i\lambda t} \Psi_{\lambda}(r, \Omega), \quad \beta(\Omega, t) = e^{-i\lambda t} \beta_{\lambda}(\Omega) ,
$$

which satisfy an outgoing radiation condition as  $r \to \infty$ :

<span id="page-1-5"></span>The Scattering Resonance Spectral Problem

(4a)

$$
\left(\Delta + (\epsilon \lambda)^2\right) \Psi_{\lambda} = 0, \qquad \qquad r > 1
$$

(4b) 
$$
\partial_r \Psi_\lambda = -i\lambda \beta_\lambda, \qquad r = 1
$$

(4c) 
$$
-i\lambda\Psi_{\lambda} = 3\gamma \left(\frac{1}{2} + \frac{2}{We}\right) \langle \beta_{\lambda}, Y_0^0 \rangle Y_0^0 - \frac{1}{We} (2 + \Delta_S)\beta_{\lambda}, \quad r = 1
$$

<span id="page-1-6"></span>(4d) 
$$
\Psi_{\lambda}
$$
 outgoing  $r \to \infty$ .

If  $\lambda$  is such that [\(4\)](#page-1-5) has a non-trivial solution, then we call  $\lambda$  a (deformation) scattering resonance energy, and  $(\Psi_{\lambda}, \beta_{\lambda})$  a corresponding scattering resonance mode. Due to the radiation condition [\(4d\)](#page-1-6), the eigenvalue problem [\(4\)](#page-1-5) is non-selfadjoint. The eigenvalues of [\(4\)](#page-1-5) all lie in the open lower half complex plane and for each  $\epsilon > 0$  are uniformly bounded away from the real axis. In particular, for any  $\epsilon > 0$ 

<span id="page-1-0"></span><sup>(2)</sup> For simplicity, we have set  $Ca$ , the cavitation number, appear in [\[5\]](#page-13-0) equal to one.

there is a scattering resonance eigenvalue  $\lambda_{\star}(\epsilon)$  with Im  $\lambda_{\star}(\epsilon) < 0$  and such that any scattering resonance eigenvalue of  $(4)$ ,  $\lambda$ , satisfies

$$
\text{Im }\lambda \leq \text{Im }\lambda_{\star}(\epsilon) < 0 ;
$$

see Theorem 3.2 of [\[5\]](#page-13-0).

The following theorem, a consequence of Theorem 5.1 in [\[5\]](#page-13-0), relates  $\lambda_{\star}(\epsilon)$  to the rate of decay of perturbations about the spherical bubble equilibrium:

**Theorem 1.** There exists  $\epsilon_0 > 0$ , such that for all  $0 < \epsilon \leq \epsilon_0$  the following holds. Consider the initial boundary value problem [\(3\)](#page-1-7). Assume the initial conditions:

(a) 
$$
\beta(t, \Omega) = \sum_{l \geq 0} \sum_{|m| \leq l} \beta_l^m(0) Y_l^m(\Omega)
$$
, where  

$$
\|\beta(t=0)\| = \sum_{l \geq 0} \sum_{|m| \leq l} (1+l)^{2+\frac{1}{6}} |\beta_l^m(0)| < \infty
$$

(b) 
$$
\Psi(t, x), \ \partial_t \Psi(t, x) \equiv 0, \ t = 0, \ |x| > 1,
$$

Then, there exists a unique solution  $\Psi(r, \Omega, t)$ ,  $\beta(\Omega, t)$ , defined for  $r > 1$ ,  $\Omega \in S^2$ , which solves the the initial-boundary value problem.

Furthermore, define  $\Gamma(\epsilon)$  to be the minimum distance of a scattering resonance (in the lower half plane) to the real axis. That is,

(5) 
$$
\Gamma(\epsilon) = |\text{Im } \lambda_{\star}(\epsilon)|
$$

Then, the solution  $\Psi(x,t)$ ,  $\beta(\Omega,t)$  satisfies the

#### <span id="page-2-1"></span>Decay Estimate:

(6) 
$$
|\beta(\Omega, t)| \le C \|\beta(t=0)\| e^{-\Gamma(\epsilon)t}, \quad \Omega \in S^2
$$
  
\n(7)  $|\Psi(x, t)| \le \begin{cases} C \frac{1}{|x|} e^{-\Gamma(\epsilon)} (t - \epsilon(|x|-1)} \|\beta(t=0)\|, & 1 < |x| < 1 + \epsilon^{-1}t \\ 0, & |x| > 1 + \epsilon^{-1}t \end{cases}$ 

Our main result is the following precise asymptotic statement about  $\Gamma(\epsilon)$ :

<span id="page-2-0"></span>**Theorem 2.** For  $0 < \epsilon < \epsilon_0$ , there exist positive constants  $A_{\epsilon}$  and  $B$ , such that

(8) 
$$
\Gamma(\epsilon) = \frac{A_{\epsilon}We}{\epsilon^3} \exp\left(-\frac{BWe}{\epsilon^2}\right)
$$

where  $B \approx 0.26924$  and where  $A_{\epsilon}$  has asymptotic expansion

(9) 
$$
A_{\epsilon} = A_0 + \mathcal{O}\left(\frac{\epsilon^2}{We}\right), \text{ where } A_0 \approx e^{-2.1465}
$$

Acknowledgement: O.C. and S.T. were supported in part by NSF grant DMS-1108794. M.I.W. was supported in part by NSF grant DMS-10-08855.

#### 2. Proof of Theorem [2](#page-2-0)

We begin by deriving a characterization of the scattering resonance energies of the eigenvalue problem  $(4)$  as the zeros of an analytic function in  $\mathbb{C}$ .

We first note that outgoing solutions of the three-dimensional Helmholtz equation are linear combinations of solutions of the form  $h_l^{(1)}$  $\binom{1}{l}(r)Y_l^m(\Omega), |m| \leq l$ , where  $h_l^{(1)}$  $\ell_l^{(1)}$  denotes the *outgoing* spherical Hankel function of order *l*. Thus, we seek solutions of [\(4\)](#page-1-5) of the form:

$$
\Psi_{\lambda}(r,\Omega)=a\ Y_{l}^{m}(\Omega)\ h_{l}^{(1)}\left(\epsilon\lambda r\right),\ \ \beta_{\lambda}(\Omega)=b\ Y_{l}^{m}(\Omega),\ \ r\geq 1,\ \ \Omega\in S^{2}.
$$

where a and b are constants to be determined. This choice of  $\Psi_{\lambda}$  solves the Helmholtz equation and satisfies the outgoing radiation condition. To impose the boundary conditions at  $r = 1$  we substitute the expressions for  $\Psi_{\lambda}$  and  $\beta_{\lambda}$  into [\(4\)](#page-1-5) and obtain the following two linear homogeneous equations for the unknown constants a and b:

(10) 
$$
\begin{pmatrix} \epsilon \lambda h_l^{(1)'}(\epsilon \lambda) & i\lambda \\ -i\lambda h_l^{(1)}(\epsilon \lambda) & -(l+2)(l-1)/We \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, l \ge 2.
$$

Setting the determinant equal to zero yields

<span id="page-3-1"></span>
$$
\lambda^2 h_l^{(1)}(\epsilon \lambda) + \frac{(l+2)(l-1)}{We} \epsilon \lambda h_l^{(1)'}(\epsilon \lambda) = 0.
$$

Finally, multiplying through by  $\epsilon^2$  and defining

(11) 
$$
Q(l,\epsilon) = (l+2)(l-1)\frac{\epsilon^2}{We}
$$

yields the following transcendental equation

<span id="page-3-0"></span>(12) 
$$
zh_l^{(1)}(z) + Q(l, \epsilon)h_l^{(1)'}(z) = 0, \ z = \epsilon \lambda \neq 0.
$$

Here  $h_l^{(1)}$  $\ell_l^{(1)}$  denotes spherical Hankel function of the first kind of order l.

<span id="page-3-2"></span>**Remark 1.** We will analyze the roots z of [\(12\)](#page-3-0) for  $We = 1$  since the formula for the roots z for  $We \neq 1$  are obtained from those with  $We = 1$  by replacing  $\epsilon^2$  by  $\frac{\epsilon^2}{We}$ . Furthermore, since  $\lambda = \frac{z}{\epsilon}$ , proving Theorem [2](#page-2-0) is equivalent to showing that the roots of [\(12\)](#page-3-0) for  $We = 1$  satisfy the property

(13) 
$$
\Gamma(\epsilon) = \inf_{l \ge 2} \{-\text{Im } z\} = \frac{A_{\epsilon}}{\epsilon^2} \exp \left[-\frac{B}{\epsilon^2}\right]
$$

for  $0 < \epsilon \leq \epsilon_0$  sufficiently small.

In the following, we consider the roots in different regimes in  $l$  relative to  $\epsilon$ .

**Definition 3.**  $f = O(g)$  means that for any  $\epsilon$  sufficiently small, there exists a constant C independent of  $\epsilon$  so that  $|f| \leq C|g|$ . We write  $f = O_s(g)$ , i.e. f is strictly of order g, if  $(f = O(g)$  and  $g = O(f)$ . We write  $f \ll g$  if  $f = o(g)$ , that is  $f/g \to 0$ , as  $\epsilon \to 0$  and  $f \gg g$  if  $|f/g| \to \infty$  as  $\epsilon \to 0$ .

Outline of the proof: In section [3](#page-4-0) equation [\(12\)](#page-3-0) is rewritten as a coupled set of equations for the real and imaginary parts of  $z = x + iy$ . In section [4](#page-5-0) we show, via Lemma [4](#page-5-1) and Lemma [5,](#page-5-2) that the scattering resonance of minimal imaginary part does not occur for  $l = \mathcal{O}(1)$  or for  $l \gg \epsilon^{-2}$ . In section [5](#page-6-0) we show that the scattering

resonance of minimal imaginary part does not occur for  $1 \ll l \ll \epsilon^{-2}$ . Therefore, the scattering resonance of interest must satisfy  $l = \mathcal{O}_s(\epsilon^{-2})$ . The detailed analysis of this regime and the rigorous approximation of [\(5\)](#page-2-1) is in section [6.](#page-6-1) Section [7](#page-11-0) is an appendix containing many of the asymptotic forms of special functions.

# <span id="page-4-0"></span>3. DERIVATION OF A SYSTEM OF EQUATION FOR  $x, y$ , WHERE  $z = x + iy$

It turns out that for large l some of the roots  $z = x+iy$  of [\(12\)](#page-3-0) have exponentially small y (with also  $y/x$  is exponentially small) and therefore the asymptotic analysis is more delicate. We next re-express [\(12\)](#page-3-0) as an equivalent system of equations obtained by setting its real and imaginary parts equal to zero. We note that (14)

$$
h_l^{(1)}(z) = j_l(z) + iy_l(z), \text{ where } j_l(z) = \sqrt{\frac{\pi}{2z}} J_{l+1/2}(z) , \ y_l(z) = \sqrt{\frac{\pi}{2z}} Y_{l+1/2}(z),
$$

where  $J_{l+1/2}(z)$  and  $Y_{l+1/2}(z)$  are Bessel functions of order  $l + 1/2$ . Substituting into [\(12\)](#page-3-0) leads to

<span id="page-4-1"></span>(15) 
$$
(x+iy)j_l(x+iy) + i(x+iy)y_l(x+iy) + Qj'_l(x+iy) + iQj'_l(x+iy) = 0
$$

Defining the real valued functions

(16)  
\n
$$
A_1(x,y) = \frac{1}{2} [y_l(x+iy) + y_l(x-iy)] , A_2(x,y) = \frac{1}{2iy} [y_l(x+iy) - y_l(x-iy)]
$$
\n
$$
A_3(x,y) = \frac{1}{2} [y'_l(x+iy) + y'_l(x-iy)] , A_4(x,y) = \frac{1}{2iy} [y'_l(x+iy) - y'_l(x-iy)],
$$

$$
(17)
$$

$$
B_1(x,y) = \frac{1}{2} [j_l(x+iy) + j_l(x-iy)] , B_2(x,y) = \frac{1}{2iy} [j_l(x+iy) - j_l(x-iy)]
$$
  

$$
B_3(x,y) = \frac{1}{2} [j'_l(x+iy) + j'_l(x-iy)] , B_4(x,y) = \frac{1}{2iy} [j'_l(x+iy) - j'_l(x-iy)],
$$

and separating out the imaginary and real parts of [\(15\)](#page-4-1), we get

<span id="page-4-2"></span>(18) 
$$
xA_1 = -QA_3 - y(xB_2 + B_1 + QB_4 - yA_2)
$$

<span id="page-4-3"></span>(19) 
$$
y (A_1 + xA_2 + QA_4) = (xB_1 + QB_3 - y^2B_2)
$$

We rewrite [\(18\)](#page-4-2), [\(19\)](#page-4-3) in the form

<span id="page-4-4"></span>(20) 
$$
x = -\frac{QA_3}{A_1} - \frac{y(xB_2 + B_1 + QB_4 - yA_2)}{A_1}
$$

(21) 
$$
y = \frac{xB_1 + QB_3 - y^2B_2}{A_1 + xA_2 + QA_4}
$$

4. LOCATION OF THE ROOTS  $z$  OF  $(12)$ 

<span id="page-5-0"></span>As will be seen, the roots with smallest  $|\text{Im } z|$  occur for  $l = O(\epsilon^{-2}), z = O(l)$ ; for them, it turns out that  $\text{Im}z$  is exponentially small in  $\epsilon$ . First, we show that roots  $z$  in other regimes have larger  $|Im z|$ .

<span id="page-5-1"></span>**Lemma 4.** If  $l = O(1)$  as  $\epsilon \to 0$ , then any root of [\(12\)](#page-3-0) in the sector arg  $z \in (-\pi, \pi)$ satisfies  $z^2 \sim (l+1)Q$ , and  $y = \text{Im}z = O(\epsilon^{2l+2})$ .

See Theorem 6.2 of [\[5\]](#page-13-0) for detailed asymptotics.

*Proof.* Assume first that we had  $z \gg 1$ . Since in this regime  $\frac{h^{(1)'}(z)}{h_l(z)} \sim i$  (see [\(65\)](#page-12-0) in the Appendix), it follows that  $z \sim -iQ = O(\epsilon^2)$ , a contradiction. Similarly,  $z = O_s(1)$  implies  $z = O(\epsilon^2)$ , a contradiction again. Thus  $z \ll 1$ . Using the wellknown expansions of of  $h_l^{(1)}$  $l_1^{(1)}(z), h_l^{(1)}$  $(z)$  for small  $z$  ([\(57\)](#page-11-1) and [\(58\)](#page-11-2) in the Appendix) we get

<span id="page-5-4"></span>
$$
(22)
$$

$$
z \sim -Q \frac{h_l^{(1)'}(z)}{h_l^{(1)}(z)} = -Q \frac{y_l'(z)}{y_l(z)} \left(1 - \frac{ij_l'(z)}{y_l'(z)}\right) \left(1 - \frac{ij_l(z)}{y_l(z)}\right)^{-1}
$$
  
= 
$$
\frac{(l+1)Q}{z} \left[1 - \frac{z^2}{(2l-1)(l+1)} - \frac{2z^4}{(l+1)(2l-3)(2l-3)^2} + O(z^6/l^6)\right]
$$
  
(23) 
$$
\times \left[1 + O\left(\frac{z}{l}\right)^{2l+1}\right],
$$

implying by iteration

<span id="page-5-3"></span>
$$
(24) \ \ z^2 = (l+1)Q\left[1 - \frac{Q}{2l-1} + \frac{Q^2(l-4)}{(2l-3)(2l-1)^2} + O\left(Q^4l^{-4}, Q^{l+1/2}l^{-l-1/2}\right)\right],
$$

which is real and positive up to and including  $O(Q^l)$  Therefore, x can be computed within errors of  $O(Q^l) = O(\epsilon^{2l})$  from the series [\(24\)](#page-5-3). Using [\(7.1\)](#page-11-3) and the small x expansion of Bessel functions (see Appendix), the result follows from straightforward calculations.

<span id="page-5-5"></span>**Remark 2.** The expression [\(24\)](#page-5-3) holds also in the regime  $1 \ll l \ll \epsilon^{-2}$ 

<span id="page-5-2"></span>**Lemma 5.** If  $l \gg \epsilon^{-2} \gg 1$ , then any root of [\(12\)](#page-3-0) in the sector arg  $z \in (-\pi, \pi)$ satisfies  $z \sim -iQ$ , implying  $-\text{Im}z = O(l^2 \epsilon^2) \gg \epsilon^{-2}$ .

*Proof.* We first claim that there are no roots  $z \ll l$ . Indeed, otherwise the expansion of  $h_l^{(1)}$  $\binom{1}{l}(z)$  and  $h_l^{(1)}$ l  $(z)$  in this regime (see [\(57\)](#page-11-1), [\(58\)](#page-11-2) in the Appendix) would lead to a contradiction:  $z = O(\sqrt{lQ}) = O(l^{3/2} \epsilon) \gg l$ . On the other hand, the existence of roots  $z = O_s(l)$  implies, using the asymptotics of  $h_l^{(1)}$  $\binom{1}{l}$  and  $\binom{h^{(1)}}{l}$  $\binom{11}{l}$  (see §7.3- §7.5) in the Appendix for results in different ranges of  $\frac{z}{\sqrt{l(l+1)}}$  in [\(12\)](#page-3-0), a contradiction:  $\frac{z}{\sqrt{l(l+1)}} = O_s\left(\frac{Q}{l}\right) = O_s(l\epsilon^2) \gg 1.$  Thus,  $|z| \gg l$ . Using the asymptotic behavior of  $h_l^{(1)}$  $\ell_l^{(1)}(z)$  and its derivative in this regime leads (see section [7.2\)](#page-12-1) to  $z \sim -iQ$ .

# 5. ANALYSIS OF THE CASE  $1 \ll l \ll \epsilon^{-2}$

<span id="page-6-2"></span><span id="page-6-0"></span>**Lemma 6.** If  $1 \ll l \ll \epsilon^{-2}$ , then any root of [\(12\)](#page-3-0) is given by [\(24\)](#page-5-3), implying that  $z = O_s(l^{3/2} \epsilon) \ll l \ll \epsilon^{-2}.$ 

*Proof.* Using the asymptotics [\(65\)](#page-12-0) of  $h_l^{(1)}$  $\binom{1}{l}(z)$  and  $\binom{h^{(1)}}{l}$  $\binom{1}{l}(z)$ <sup>'</sup> in [\(12\)](#page-3-0), the assumption  $z \gg l$  leads to a contradiction:  $z \sim -iQ = O(l^2 \epsilon^2) \ll l$ . Assuming  $z = O_s(l)$  and using the asymptotics of  $h_l^{(1)}$  $\binom{1}{l}$  and  $\binom{h^{(1)}}{l}$  $\binom{1}{k}$  (Appendix, §7.3- §7.5) in [\(12\)](#page-3-0), also ends up in a contradiction:  $\frac{z}{\sqrt{l(l+1)}} = O_s\left(\frac{Q}{l}\right) = O_s(l\epsilon^2) \ll 1$ . In the regime  $z = o(l)$ , exploiting the asymptotics of  $h_l^{(1)}$  $\binom{1}{l}(z)$ , we obtain [\(22\)](#page-5-4) and therefore [\(24\)](#page-5-3).

By Lemma [6,](#page-6-2) any root of [\(12\)](#page-3-0) in this case satisfies  $z = o(l)$ . From [\(24\)](#page-5-3) (see Remark [2\)](#page-5-5), it follows that the roots are close to the real axis. In this section, we estimate  $y = \text{Im } z$  in this regime.

<span id="page-6-4"></span>**Lemma 7.** Any root  $z = x + iy$  of [\(12\)](#page-3-0) in the regime  $1 \ll l \ll \epsilon^{-2}$  satisfies

<span id="page-6-3"></span>(25) 
$$
y \sim -\frac{\sqrt{Q(l+1)}}{2e} \left( \frac{eQ^{1/2}(l+1)^{1/2}}{2l} \right)^{2l+1} \left[ 1 - \frac{Q}{(2l-1)} + O(Q^2/l^2) \right]^{l+1/2}
$$

*Proof.* Since  $z \ll l$ , we use [\(24\)](#page-5-3) and the expansions of Bessel functions in this regime (see Appendix) to conclude

(26) 
$$
\frac{j_l(z)}{y_l(z)} = -\frac{1}{2e} \left(\frac{ze}{2l}\right)^{2l+1} \left[1 + O\left(\frac{z^2}{l^2}\right)\right]
$$

$$
= -\frac{1}{2e} \left(\frac{eQ^{1/2}(l+1)^{1/2}}{2l}\right)^{2l+1} \left[1 - \frac{Q}{(2l-1)} + O(Q^2/l^2)\right]^{l+1/2}
$$

In the same way,

(27) 
$$
1 + \frac{QB_3}{xB_1} = 2 + O\left(\frac{1}{l}, \frac{Q}{l}\right), 1 + x\frac{A_2}{A_1} + \frac{QA_3}{A_1} = 2 + O(Q/l)
$$

Using  $(7.1)$ , one obtains  $(25)$ . П

**Remark 3.** Since  $Q^{1/2}/l^{1/2}$  scales like  $l^{1/2} \epsilon \ll 1$ , we have  $|y| \sim Cl (\epsilon l^{1/2})^{2l+2}$ , which is a decreasing function of l in the regime  $1 \ll l \ll \epsilon^{-2}$ . Therefore, |y| is minimal at the right end, and this minimal value turns out to be larger than the values obtained in the next section.

6. ANALYSIS OF THE CASE  $l = O_s(\epsilon^{-2})$ 

<span id="page-6-1"></span>From [\(11\)](#page-3-1), it follows that when  $l = O_s(\epsilon^{-2})$ ,  $Q = O_s(\epsilon^{-2})$ . If  $z = O_s(l)$ , but  $|z-l|\gg l^{1/3}$ , from the asymptotics [\(74\)](#page-12-2) and [\(75\)](#page-12-3), it follows that [\(12\)](#page-3-0) results in

(28) 
$$
z^2 \sim -iQ\sqrt{\frac{z^2}{l(l+1)}} - 1
$$

Hence  $|y| = |\text{Im}z| \gg 1$ . In the crossover regime,  $1 - \frac{z}{\sqrt{l(l+1)}} = O(l^{-2/3})$ , the asymptotics of Bessel functions involve the Airy functions Ai and Bi [\(76\)](#page-13-3) and [\(77\)](#page-13-4)) whose ratio is  $O<sub>s</sub>(1)$ , and thus y cannot be exponentially small.

So, we restrict to the sub-region  $\left| \frac{z}{\sqrt{l(l+1)}} \right|$  $\vert$  < 1 and  $1 - \frac{z}{\sqrt{l(l+1)}} \gg l^{-2/3}$ . In this regime, since  $\frac{j_l(z)}{y_l(z)}$  is exponentially small in l, by [\(7.1\)](#page-11-3), any root z must have y exponentially small. We introduce the scaled variables

<span id="page-7-0"></span>(29) 
$$
\sqrt{l(l+1)} = l_* \epsilon^{-2}, \ Q = l_* \epsilon^{-2} q, \ x = \frac{l_*}{\epsilon^2} \zeta, \ -\epsilon^2 \log \left[ -\epsilon^2 y \right] = \eta
$$

From the definition of Q, it follows that (for  $We = 1$ ),

$$
(30) \t\t q = l_* - \frac{2\epsilon^4}{l_*}
$$

From  $(20)$  and  $(7.1)$ , we can replace  $(7.1)$  by

<span id="page-7-1"></span>(31) 
$$
y = \frac{xB_1 + QB_3 - y^2B_2}{D}
$$
 where  
\n
$$
D = A_1 + xA_2 + QA_4 + \sqrt{\frac{l(l+1)}{x^2} - 1} [xA_1 + QA_3 + y (xB_2 + B_1 + QB_4 - yA_2)].
$$

Note: the square-bracketed quantity on the previous line vanishes by [\(7.1\)](#page-11-3). With the notation in  $(29)$ ,  $(20)$  and  $(31)$  imply

<span id="page-7-2"></span>(32) 
$$
0 = \zeta + \frac{qA_3}{A_1} - \frac{e^{-\eta/\epsilon^2}}{\epsilon^2 A_1} \left[ \frac{B_1}{l_* \epsilon^2} + \zeta B_2 + qB_4 + \frac{A_2}{\epsilon^4 l_*} e^{-\eta/\epsilon^2} \right] =: F(\zeta, \eta; \epsilon)
$$

(33) 
$$
0 = \eta + \epsilon^2 \log \left[ -\epsilon^2 \frac{(xB_1 + QB_3 - y^2B_2)}{D} \right] =: G(\zeta, \eta; \epsilon)
$$

<span id="page-7-6"></span>**Lemma 8.** For any fixed  $\delta \in (0,1)$ , if  $l_* \in \left(\frac{\delta^2}{\sqrt{1-\delta^2}}, \frac{(1-\delta)^2}{\sqrt{1-(1-\delta)^2}}\right)$  $\Big), \zeta \in (\delta, 1-\delta)$  and  $\eta > \eta_0 = 2l_* \int_{\frac{1}{\epsilon}}^1 \delta t^{-1} \sqrt{1-t^2} dt$ , then there exists  $\epsilon_0$  such that for  $|\epsilon| \leq \epsilon_0$ , F and G are smooth functions of  $\zeta$ ,  $\eta$ ,  $l_*$  and  $\epsilon$ , with

<span id="page-7-3"></span>(34) 
$$
F(\zeta, \eta; 0) = \zeta - l_* \frac{\sqrt{1 - \zeta^2}}{\zeta}, \ G(\zeta, \eta; 0) = \eta - 2l_* \int_{\zeta}^1 \frac{\sqrt{1 - t^2}}{t} dt
$$

and det  $\frac{\partial \{F,G\}}{\partial \{\zeta,\eta\}}$  $\begin{bmatrix} \epsilon_{0} \\ \epsilon_{1} \\ \end{bmatrix} \neq 0$  Furthermore, in these intervals, the system of equations  ${F(\zeta, \eta; \epsilon) = 0, G(\zeta, \eta; \epsilon) = 0}$  has a unique smooth solution  $(\zeta(l_*, \epsilon), \eta(l_*, \epsilon))$ , for sufficiently small  $\epsilon$ , with asymptotic behavior in  $\epsilon$  determined implicitly from

<span id="page-7-4"></span>(35) 
$$
l_* = \frac{\zeta^2}{\sqrt{1-\zeta^2}} - \frac{\epsilon^2(1-2\zeta^2)}{2(1-\zeta^2)^{3/2}} + O(\epsilon^4),
$$

<span id="page-7-5"></span>(36) 
$$
\eta = 2l_* \int_{\zeta}^{1} (t^{-2} - 1)^{1/2} dt - \epsilon^2 \log [l_* \zeta] + \epsilon^2 \log \left[ 2 + \frac{1}{2(1 - \zeta^2)} - \frac{l_*^2 (1 - 2\zeta^2)}{2\zeta^4} \right] + O(\epsilon^4)
$$

*Proof.* Using a standard integral representation of  $j_l(\sqrt{l(l+1)}\zeta)$  and  $y_l(\sqrt{l(l+1)}\zeta)$ , [\(54\)](#page-11-4) and [\(55\)](#page-11-5), and noting that  $\sqrt{l(l+1)} = l_* \epsilon^{-2}$ , smoothness of F and G in  $\epsilon$  follows provided  $A_1$  and  $D$  in [\(32\)](#page-7-2) and [\(7.2\)](#page-12-1) are nonzero, which is the case, as shall

be seen, for  $\epsilon = 0$ . Routine calculations, using the asymptotics [\(68\)](#page-12-4)-[\(73\)](#page-12-5) show that (37)

$$
\zeta + q \frac{A_3}{A_1} = \zeta + q \frac{y_l'(x)}{y_l(x)} + O(e^{-\eta_0 \epsilon^{-2}}) = \zeta - \frac{l_* \sqrt{1 - \zeta^2}}{\zeta} \left[ 1 + \epsilon^2 \frac{1 - 2\zeta^2}{2l_*(1 - \zeta^2)^{3/2}} + O(\epsilon^4) \right],
$$

Therefore,

<span id="page-8-0"></span>(38) 
$$
F(\zeta, \eta; \epsilon) = \zeta - l_{*} \frac{\sqrt{1 - \zeta^{2}}}{\zeta} - \frac{\epsilon^{2} (1 - 2\zeta^{2})}{2\zeta (1 - \zeta^{2})} + O(\epsilon^{4}),
$$

from which the formula for  $F(\zeta, \eta; 0)$  in [\(34\)](#page-7-3) follows. Now, setting  $F = 0$  in [\(38\)](#page-8-0) immediately leads to  $(35)$ . For the second part, from the asymptotics of  $j_l$ ,  $y_l$  in this regime,  $(68)-(73)$  $(68)-(73)$ , we get

(39) 
$$
xB_1 + QB_3 - y^2B_2 = xj_l(x)\left[1 + \frac{qj'_l(x)}{\zeta j_l(x)} + O(y^2)\right]
$$
  

$$
= xj_l(x)\left[1 + \frac{l_*\sqrt{1-\zeta^2}}{\zeta^2} - \frac{\epsilon^2(1-2\zeta^2)}{2\zeta^2(1-\zeta^2)} + O(\epsilon^4)\right]
$$

(40)

$$
D = A_1 + xA_2 + QA_4 + \sqrt{\frac{l(l+1)}{x^2} - 1} \left[ xA_1 + QA_3 + y \left( xB_2 + B_1 + QB_4 - yA_2 \right) \right]
$$
  
= 
$$
y_l \left\{ \frac{1}{2(1 - \zeta^2)} + \frac{2l_*}{\zeta^2} \sqrt{1 - \zeta^2} - \frac{l_*(1 - 2\zeta^2)}{2\zeta^2 \sqrt{1 - \zeta^2}} + O(\epsilon^2) \right\}
$$

Therefore, it follows that

(41) 
$$
G(\zeta, \eta; \epsilon) = \eta + \epsilon^2 \log \left\{ -\epsilon^2 \frac{xB_1 + QB_3 - y^2B_3}{D} \right\} = \eta + \epsilon^2 \log \left[ -\frac{l_* \zeta j_l(x)}{y_l(x)} \right] + \epsilon^2 \log \left[ 1 + \frac{l_* \sqrt{1 - \zeta^2}}{\zeta^2} \right] - \epsilon^2 \log \left[ \frac{1}{2(1 - \zeta^2)} + \frac{2l_*}{\zeta^2} \sqrt{1 - \zeta^2} - \frac{l_* (1 - 2\zeta^2)}{2\zeta^2 \sqrt{1 - \zeta^2}} \right] + O(\epsilon^4)
$$

From equation [\(73\)](#page-12-5) in the Appendix, we obtain

(42) 
$$
-2\frac{j_l(x)}{y_l(x)} = \exp\left[-2l_*\epsilon^{-2}\int_{\zeta}^1 (t^{-2}-1)^{1/2}dt\right],
$$

<span id="page-8-1"></span>(43) 
$$
G(\zeta, \eta; \epsilon) = \eta - 2l_* \int_{\zeta}^{1} (t^{-2} - 1)^{1/2} dt + \epsilon^2 \log \left( \frac{1}{2} l_* \zeta \left[ 1 + l_* \frac{\sqrt{1 - \zeta^2}}{\zeta^2} \right] \right) - \epsilon^2 \log \left[ \frac{1}{2(1 - \zeta^2)} + \frac{2l_*}{\zeta^2} \sqrt{1 - \zeta^2} - \frac{l_*^2 (1 - 2\zeta^2)}{2\zeta^4} \left( \frac{\zeta^2}{l_* \sqrt{1 - \zeta^2}} \right) \right] + O(\epsilon^4)
$$

from which the expression of  $G(\zeta, \eta; 0)$  in [\(34\)](#page-7-3) follows. The Jacobian of  $F(\zeta, \eta; 0)$ ,  $G(\zeta, \eta; 0)$  with respect to  $(\zeta, \eta)$  is clearly nonzero. The statement of existence and uniqueness of  $(\zeta(l_*, \epsilon), \eta(l_*, \epsilon))$  now follows from the implicit function theorem.<br>Furthermore, using (43) and (35) in  $F = 0$ ,  $G = 0$  immediately implies (36). Furthermore, using [\(43\)](#page-8-1) and [\(35\)](#page-7-4) in  $F = 0$ ,  $G = 0$  immediately implies [\(36\)](#page-7-5).

**Note 4.** We now comment on why the interval restrictions in Lemma [8](#page-7-6) for  $l_*$ ,  $\zeta$  and  $\eta$  do not matter for finding the smallest |y| as a function of  $l_*$ . First, if  $l_* < \frac{\delta^2}{\sqrt{1-\delta^2}}$ , then for sufficiently small  $\delta$ , the conclusions of Lemma [7](#page-6-4) hold and the corresponding |y| is not as small as implied by Lemma [8](#page-7-6) for  $l_*$  in the given interval. On the other hand if  $l_* > \frac{(1-\delta)^2}{\sqrt{1-(1-\delta)^2}}$  for sufficiently small  $\delta$ , then the conclusions of Lemma [5](#page-5-2) hold where  $y = \text{Im } z$  is no longer small. Thus, the restriction  $l_* \in \left(\frac{\delta^2}{\sqrt{1-\delta^2}} , \frac{(1-\delta)^2}{\sqrt{1-(1-\delta)^2}} \right)$  $\setminus$ is appropriate. Furthermore, for  $l_*$  in this interval, there is no need to consider the possibility  $\zeta \notin (\delta, 1-\delta)$  since we have already shown that  $l = O_s(\epsilon^{-2})$ , implies  $x = O_s\left(\epsilon^{-2}\right)$  and argued at the outset that we only need to consider  $1 - \zeta \gg l^{-2/3}$ . When  $(1 - \zeta)$  is small, the asymptotic behavior of  $\frac{j_l(x)}{y_l(x)}$  is given by a ratio of Airy functions, as is the case for  $1 - \zeta = O(l^{-2/3})$  (see [\(76\)](#page-13-3) and [\(77\)](#page-13-4)) and the resulting  $|y|$  is not as small as obtained in [\(53\)](#page-10-0) below, as a consequence of Lemma [8.](#page-7-6) Also, we need not worry about possible solutions for which  $\eta < \eta_0$ , since we are seeking to maximize  $\eta$  (minimizing |y|).

6.1. Maximizing  $\eta$  as a function of  $l_*$  and determination of  $\Gamma(\epsilon)$  of [\(5\)](#page-2-1). Since Im  $z =: y = -\epsilon^{-2} \exp\left[-\frac{\eta}{\epsilon^2}\right]$ , minimizing |y| as a function of  $l_*$  corresponds to maximizing  $\eta$ . Now, [\(35\)](#page-7-4) implies that  $l_*$  increases with  $\zeta$  in any compact subset of  $(\delta, 1 - \delta)$  for all sufficiently small  $\epsilon$ . For now, we consider  $l_*$  as a continuous variable for the purposes of finding the maximum value of  $\eta$ . We will show later that the maximal value of  $\eta$  to the order calculated is the same if  $l_*$  takes discrete values:  $\sqrt{l(l+1)}\epsilon^2$  for  $l \in \mathbb{N}$ .

We seek a critical value  $l_*$  for which  $\partial_{l_*}\eta = 0$  and  $\partial_{l_*l_*}^2 \eta < 0$  implying a maximum of  $\eta$ . We note that

<span id="page-9-1"></span>
$$
(44) \quad 0 = \partial_{l_*} \eta = 2 \int_{\zeta}^{1} (t^{-2} - 1)^{1/2} dt - \frac{2l_*}{\zeta} \zeta_{l_*} (1 - \zeta^2)^{1/2} - \frac{\epsilon^2}{l_*} - \frac{\epsilon^2 \zeta_{l_*}}{\zeta}
$$

$$
+ \epsilon^2 \zeta_{l_*} \partial_{\zeta} \log \left[ 2 + \frac{1}{2(1 - \zeta^2)} - \frac{l_*^2 (1 - 2\zeta^2)}{2\zeta^4} \right] + \epsilon^2 \partial_{l_*} \log \left[ 2 + \frac{1}{2(1 - \zeta^2)} - \frac{l_*^2 (1 - 2\zeta^2)}{2\zeta^4} \right] + O(\epsilon^4)
$$

We also note that [\(35\)](#page-7-4) implies

<span id="page-9-0"></span>(45) 
$$
\zeta_{l_*} = \frac{(1 - \zeta^2)^{3/2}}{\zeta(2 - \zeta^2)} - \frac{(1 + 2\zeta^2)\sqrt{1 - \zeta^2}}{2\zeta(2 - \zeta^2)^2} \epsilon^2 + O(\epsilon^4)
$$

Substituting  $(35)$  and  $(45)$  into  $(44)$ , we obtain (46)

$$
0 = \frac{1}{2}\eta_{l*} = \int_{\zeta}^{1} \sqrt{t^{-2} - 1} \, dt - \frac{(1 - \zeta^2)^{3/2}}{(2 - \zeta^2)} + \epsilon^2 \frac{\sqrt{1 - \zeta^2} (2\zeta^4 + 5\zeta^2 - 4)}{2\zeta^2 (2 - \zeta^2)^2} + O(\epsilon^4) =: g(\zeta; \epsilon)
$$

It is clear that the solution  $\zeta$  of  $g(\zeta; \epsilon) = 0$  has the behavior

(47) 
$$
\zeta = \zeta_{m,0} + \epsilon^2 \zeta_{m,2} + O(\epsilon^4)
$$

where

(48) 
$$
0 = \int_{\zeta_{m,0}}^1 \sqrt{t^{-2} - 1} dt - \frac{(1 - \zeta_{m,0}^2)^{3/2}}{(2 - \zeta_{m,0}^2)} = g(\zeta_{m,0}; 0) =: g_0(\zeta_{m,0}).
$$

Since

<span id="page-10-1"></span>(49) 
$$
g'_0(\zeta) = 2\sqrt{1-\zeta^2} \left( \frac{2-(2-\zeta^2)^2}{\zeta(2-\zeta^2)^2} \right),
$$

it follows that

<span id="page-10-2"></span>
$$
(50) \qquad \zeta_{m,2} = -\frac{\sqrt{1 - \zeta_{m,0}^2} \left(2\zeta_{m,0}^4 + 5\zeta_{m,0}^2 - 4\right)}{2g_0'(\zeta_{m,0})\zeta_{m,0}^2 \left(2 - \zeta_{m,0}^2\right)^2} = -\left\{\frac{\left(2\zeta_{m,0}^4 + 5\zeta_{m,0}^2 - 4\right)}{4\zeta_{m,0} \left(2 - \left(2 - \zeta_{m,0}^2\right)^2\right)}\right\}
$$

We now seek to find  $\zeta_{m,0}$  which corresponds to the maximal  $\eta$ . We note from [\(49\)](#page-10-1) that  $g'_0 < 0$  for  $\zeta \in (0, \sqrt{2-\sqrt{2}}) =: J$  and  $g'_0 > 0$  for  $\zeta \in (\sqrt{2-\sqrt{2}}, 1)$ . Since we are seeking a maximum for  $\eta$ , we must have  $g'_0 < 0$  at the corresponding  $\zeta$  (noting that  $\zeta_{l*} > 0$ ). Thus the roots of  $g_0(\zeta) = 0$  only need to be sought for  $\zeta \in J$ . At the right end of the interval J, explicit integration –in terms of elementary functions– gives  $g_0 < 0$  ( $\approx -0.0678$ ), while  $g_0 \to +\infty$  when  $\zeta \to 0$ . Since  $g_0$  is monotonic in J, there exists unique  $\zeta_{m,0}$  satisfying  $g_0(\zeta_{m,0}) = 0$ . The explicit calculation of  $g_0$ gives  $g_0(0.58) > 0$  and  $g_0(0.59) < 0$ , implying that the unique maximum  $\zeta_{m,0}$  is in  $(0.58, 0.59)$ –in fact  $\zeta_{m,0} = 0.58134...$  Using [\(50\)](#page-10-2) we get  $\zeta_{m,2} \approx -1.1743...$  Equation [\(35\)](#page-7-4) also implies that the critical  $l_* (= l_{*,m})$  that maximizes  $\eta$  is given by

$$
(51) \quad l_{*,m} = \frac{\zeta_{m,0}^2}{\sqrt{1 - \zeta m, 0^2}} + \epsilon^2 \left[ \frac{\zeta_{m,1} \zeta_{m,0} (2 - \zeta_{m,0}^2)}{(1 - \zeta_{m,0}^2)^{3/2}} - \frac{(1 - 2\zeta_{m,0}^2)}{2(1 - \zeta_{m,0}^2)^{3/2}} \right] + O(\epsilon^4)
$$

$$
=: l_{m,0} + \epsilon^2 l_{m,2} + O(\epsilon^4) \approx 0.41535 - 2.4071 \epsilon^2 + O(\epsilon^4)
$$

Therefore, it follows that the maximized  $\eta = \eta_m$  satisfies

<span id="page-10-3"></span>
$$
(52)
$$

$$
\eta_m = 2l_{m,0} \int_{\zeta_{m,0}}^1 (t^{-2} - 1)^{1/2} dt + \epsilon^2 \left\{ 2l_{m,2} \int_{\zeta_{m,0}}^1 (t^{-2} - 1)^{1/2} dt - 2l_{m,0} \zeta_{m,2} (\zeta_{m,0}^{-2} - 1)^{1/2} dt \right\}
$$

$$
- \log[l_{m,0} \zeta_{m,0}] + \log \left[ 2 + \frac{1}{2(1 - \zeta_{m,0}^2)} - \frac{l_{m,0}^2 (1 - 2\zeta_{m,0}^2)}{2\zeta_{m,0}^4} \right] \right\} =: \eta_{m,0} + \epsilon^2 \eta_{m,2} + O(\epsilon^4)
$$

$$
\approx 0.26924 + 2.1465 \epsilon^2 + O(\epsilon^4) .
$$

Therefore the minimum value of  $|y|$  is

<span id="page-10-0"></span>(53) 
$$
|y_m| = \epsilon^{-2} \exp\left[-\epsilon^{-2} \eta_{m,0} - \eta_{m,2}\right] \left[1 + O(\epsilon^2)\right]
$$

$$
\approx -\epsilon^{-2} \exp\left[-0.26924\epsilon^{-2} - 2.1465\right] \left[1 + O(\epsilon^2)\right]
$$

The proof of Theorem [2](#page-2-0) now follows from Remark [1.](#page-3-2)

**Remark 5.** The calculation [\(52\)](#page-10-3) assumed  $l_*$  to be a continuous variable rather than discrete,  $l_* = \sqrt{l(l+1)}\epsilon^2$  with  $l \in \mathbb{N}$ . This however makes no difference to the leading order result in [\(53\)](#page-10-0). Maximization of  $\eta$  over  $l \in \mathbb{N}$  results in an optimal  $l_*$  that is different from the computed  $l_{*,m}$  by  $O(ε^2)$ . Since  $η$  is a smooth function of  $l_*,$  for  $l_* - l_{*,m} = O(\epsilon^2), \eta(l_*) - \eta(l_{*,m}) = O(l_* - l_{*,m})^2 = O(\epsilon^4)$  and therefore discreteness does not affect the leading order asymptotic result [\(53\)](#page-10-0).

## 7. APPENDIX: ASYMPTOTICS OF  $j_l$  and  $y_l$  in different regimes

<span id="page-11-0"></span>The results quoted below are standard and either given in standard references such as [\[1\]](#page-13-5), [\[2\]](#page-13-6) or follow directly from them.

The spherical Bessel functions  $j_l(z) = \sqrt{\frac{\pi}{2z}} J_{l+1/2}(z)$  and  $y_l(z) = \sqrt{\frac{\pi}{2z}} Y_{l+1/2}(z)$ have the following integral representations which immediately follows from equa-tions (10.9.6), (10.9.7) in [\[1\]](#page-13-5) (see also [http://dlmf.nist.gov/10.9\)](http://dlmf.nist.gov/10.9) using  $\nu = l + 1/2$ 

<span id="page-11-4"></span>(54) 
$$
j_l(z) = \sqrt{\frac{1}{2\pi z}} \left\{ \int_0^{\pi} \cos\left[ (l + \frac{1}{2})\tau - z \sin \tau \right] d\tau + (-1)^{l+1} \int_0^{\infty} \exp\left[ -z \sinh t - (l + \frac{1}{2})t \right] dt \right\}
$$

<span id="page-11-5"></span>(55) 
$$
y_l(z) = \sqrt{\frac{1}{2\pi z}} \Big[ (-1)^{l+1} \int_0^{\pi} \cos\left[ (l+\frac{1}{2})\tau + z \sin \tau \right] d\tau - \int_0^{\infty} \exp\left[ -z \sinh t + (l+\frac{1}{2})t \right] dt \Big]
$$

which results in the asymptotic representations below by standard Laplace method for asymptotics of integrals [\[2\]](#page-13-6). The Bessel functions  $j_l(z)$ ,  $y_l(z)$  and the Hankel function  $h_l^{(1)}$  $l_1^{(1)}(z) = j_1(z) + iy_1(z)$  satisfy

(56) 
$$
u'' + \frac{2}{z}u' + \left(1 - \frac{l(l+1)}{z^2}\right)u = 0
$$

<span id="page-11-3"></span>7.1. The regime  $z \ll l$ : We only present the asymptotic regimes relevant to this analysis. For  $z \ll l$  we have (see equations (10.1.2) and (10.1.3), page 437 of [\[1\]](#page-13-5))

(57) 
$$
j_l(z) = \frac{z^l 2^l l!}{(2l+1)!} \left[ 1 + O\left(\frac{z^2}{(l+1)^2}\right) \right]
$$

$$
(58)
$$

<span id="page-11-2"></span><span id="page-11-1"></span>
$$
y_l(z) = -\frac{(2l-1)!}{2^{l-1}(l-1)!z^{l+1}} \left[ 1 + \frac{z^2}{2(2l-1)} + \frac{z^4}{8(2l-1)(2l-3)} + O\left(\frac{z^6}{(l+1)^3}\right) \right]
$$

(Three orders are indeed needed because of cancellations.) For  $l \gg z \gg 1$ , using  $\Gamma(x+1) \sim \sqrt{2\pi}e^{-x}x^{x+1/2}$  for large x we note the simplification:

(59) 
$$
j_l(z) \sim \frac{2^{-3/2}}{l} \left(\frac{ze}{2l}\right)^l \left[1 + O\left(\frac{z^2}{l^2}\right)\right]
$$
  
(60)

$$
y_l(z) \sim -\frac{e}{l\sqrt{2}} \left(\frac{ze}{2l}\right)^{-l-1} \left[1 + \frac{z^2}{2(2l-1)} + \frac{z^2}{2(2l-1)} + \frac{z^4}{8(2l-1)(2l-3)} + O\left(\frac{z^6}{l^3}\right)\right]
$$
  
Since the asymptotics is differentiable [6] this implies

Since the asymptotics is differentiable [\[6\]](#page-13-7), this implies

(61) 
$$
\frac{j'_l(z)}{j_l(z)} = \frac{l}{z} \left[ 1 + O\left(\frac{z^2}{l^2}\right) \right]
$$

(62) 
$$
\frac{zy_l'(z)}{y_l(z)} = -(l+1) + \frac{z^2}{2l-1} + \frac{z^4}{(2l-3)(2l-1)^2} + O\left(\frac{z^6}{l^5}\right)
$$

(63) 
$$
\frac{y_l''(z)}{y_l(z)} = (l+1)(l+2)z^{-2} - \frac{2l+1}{2l-1} - \frac{2z^2}{(2l-3)(2l-1)^2} + O\left(\frac{z^4}{l^4}\right)
$$

<span id="page-12-0"></span>LIFETIME OF SHAPE OSCILLATIONS 13

(64) 
$$
\frac{j_l(z)}{y_l(z)} = -\frac{1}{2e} \left(\frac{ze}{2l}\right)^{2l+1} \left[1 + O(\frac{z^2}{l^2})\right]
$$

<span id="page-12-1"></span>7.2. The regime  $z \gg l$ , arg  $z \in (-\pi, \pi)$ . In this case

(65) 
$$
h_l^{(1)}(z) \sim \frac{1}{z} \exp\left[i\left(z - \frac{(l+1)}{2}\pi\right)\right]
$$

(66) 
$$
j_l(z) \sim \frac{1}{z} \cos\left(z - \frac{(l+1)}{2}\pi\right)
$$

(67) 
$$
y_l(z) \sim \frac{1}{z} \sin\left(z - \frac{(l+1)}{2}\pi\right)
$$

7.3. The regime  $l \gg 1$ ,  $\xi = \frac{z}{\sqrt{l(l+1)}} = O_s(l)$ ,  $1 - |\xi| > l^{-2/3}$ ,  $|y| \ll l^{1/3}$ ,  $arg(1 - \xi) \in (-\pi/3, \pi/3)$  $arg(1 - \xi) \in (-\pi/3, \pi/3)$  $arg(1 - \xi) \in (-\pi/3, \pi/3)$ . From [\(54\)](#page-11-4) and [\(55\)](#page-11-5), by the saddle point method ([\[2\]](#page-13-6)), we obtain

<span id="page-12-4"></span>(68) 
$$
j_l(z) = \frac{1}{2\sqrt{l(l+1)}\xi} \left(\xi^{-2} - 1\right)^{-1/4}
$$

$$
\times \exp\left[-l^{1/2}(l+1)^{1/2}\int_{\xi}^{1} \left(t^{-2} - 1\right)^{1/2} dt\right] \left[1 + \frac{u(\xi)}{l} + O(l^{-2})\right]
$$
  
where

where

(69) 
$$
u(\xi) = -\frac{5}{24(1-\xi^2)^{3/2}} + \frac{1}{8(1-\xi^2)^{1/2}} + \frac{1}{16}\log\left(\frac{1-\sqrt{1-\xi^2}}{1+\sqrt{1-\xi^2}}\right)
$$

(70) 
$$
y_l(z) \sim -\frac{1}{\sqrt{l(l+1)}\xi} \left(\xi^{-2} - 1\right)^{-1/4}
$$
  
  $\times \exp\left[l^{1/2}(l+1)^{1/2} \int_{\xi}^{1} \left[t^{-2} - 1\right]^{1/2} dt\right] \left[1 - \frac{u(\xi)}{l} + O(l^{-2})\right]$   
(71)  $\frac{y_l'(z)}{l} = -\sqrt{\xi^{-2} - 1} - \frac{1 - 2\xi^2}{l} + O(l^{-2})$ 

(71) 
$$
\frac{y_l'(z)}{y_l(z)} = -\sqrt{\xi^{-2} - 1} - \frac{1 - 2\xi^2}{2l\xi(1 - \xi^2)} + O(l^{-2})
$$

(72) 
$$
\frac{j'_l(z)}{j_l(z)} = \sqrt{\xi^{-2} - 1} - \frac{1 - 2\xi^2}{2l\xi(1 - \xi^2)} + O(l^{-2})
$$

In this regime, we have

<span id="page-12-5"></span>(73) 
$$
-\frac{2j_l(z)}{y_l(z)} = \exp\left[-2\sqrt{l(l+1)}\int_{\xi}^{1} (t^{-2} - 1)^{1/2} dt\right] \left[1 + \frac{2}{l}u(\xi) + O(l^{-2})\right]
$$

7.4. The regime  $l \gg 1$ ,  $\xi = \frac{z}{\sqrt{l(l+1)}} = O_s(l)$ ,  $|\xi| - 1 \gg l^{-2/3}$ ,  $|y| \ll 1$ ,  $\arg(\xi - 1) \in$  $(-\pi/3, \pi/3)$ . In this case we have

<span id="page-12-3"></span><span id="page-12-2"></span>(74) 
$$
j_l(z) \sim \frac{1}{\sqrt{l(l+1)}\xi} (1-\xi^{-2})^{-1/4} \cos\left[\sqrt{l(l+1)} \int_1^{\xi} \sqrt{(1-t^{-2})} dt - \frac{(l+1)\pi}{2}\right]
$$
  
(75)  $y_l(z) \sim \frac{1}{\sqrt{l(l+1)}\xi} (1-\xi^{-2})^{-1/4} \sin\left[\sqrt{l(l+1)} \int_1^{\xi} \sqrt{(1-t^{-2})} dt - \frac{(l+1)\pi}{2}\right]$ 

7.5. The regime  $l \gg 1$ ,  $\xi = \sqrt{z} \sqrt{l(l+1)} = O_s(l)$ ,  $1 - \xi = O(l^{-2/3})$ . Here the asymptotic behavior is given by

(76) 
$$
j_l(z) \sim \frac{\sqrt{\pi}l^{1/6}}{2^{1/6}z} \text{Ai}\left(l^{2/3}2^{1/3}[1-\xi]\right)
$$

(77) 
$$
y_l(z) \sim -\frac{\sqrt{\pi}l^{1/6}}{2^{1/6}z} \text{Bi}\left(l^{2/3}2^{1/3}[1-\xi]\right)
$$

# <span id="page-13-4"></span><span id="page-13-3"></span>**REFERENCES**

- <span id="page-13-6"></span><span id="page-13-5"></span>[1] M. Abramowitz & I. Stegun, Handbook of Mathematical Functions, Dover, New York (1970).
- [2] F.W.J. Olver, Asymptotics and Special Functions, A.K. Peters, Wellesley, Massachusetts (1997).
- <span id="page-13-1"></span>[3] M.P. Brenner, D. Lohse and T.F. Dupont, Bubble Shape Oscillations and the Onset of Sonoluminescence, Phys. Rev. Lett. Volume 75, #5 954–957 (1995)
- <span id="page-13-2"></span>[4] T.G. Leighton, From seas to surgeries, from babbling brooks to baby scans: the acoustics of gas bubbles in liquids, International Journal of Modern Physics B Volume 18 #25 3267–3314 (2004)
- <span id="page-13-0"></span>[5] A.M. Shapiro and M.I. Weinstein, Radiative decay of bubble oscillations in a compressible fluid, SIAM J. Math. Analysis Volume 43, 828-876 (2011)
- <span id="page-13-7"></span>[6] W. Wasow, Asymptotic Expansions for Ordinary Differential Equations, Interscience, New York (1965).

Department of Mathematics, The Ohio State University, 231 W 18th Ave, Columbus, OH 43210

Department of Applied Physics and Applied Mathematics, Columbia University, New York, NY 10027