

Combinatorial Proof of the Inversion Formula on the Kazhdan-Lusztig R -Polynomials

William Y.C. Chen¹, Neil J.Y. Fan², Alan J.X. Guo³, Peter L. Guo⁴
Harry H.Y. Huang⁵, Michael X.X. Zhong⁶

^{1,3,4,5,6}Center for Combinatorics, LPMC-TJKLC
Nankai University, Tianjin 300071, P.R. China

²Department of Mathematics
Sichuan University, Chengdu, Sichuan 610064, P.R. China

¹chen@nankai.edu.cn, ²fan@scu.edu.cn, ³aalen@mail.nankai.edu.cn
⁴lguo@nankai.edu.cn, ⁵haoyanghuang@mail.nankai.edu.cn
⁶michaelzhong@mail.nankai.edu.cn

Abstract

Let W be a Coxeter group, and for $u, v \in W$, let $R_{u,v}(q)$ be the Kazhdan-Lusztig R -polynomial indexed by u and v . In this paper, we present a combinatorial proof of the inversion formula on R -polynomials due to Kazhdan and Lusztig. This problem was raised by Brenti. Based on Dyer's combinatorial interpretation of the R -polynomials in terms of increasing Bruhat paths, we reformulate the inversion formula in terms of V -paths. By a V -path from u to v with bottom w we mean a pair (Δ_1, Δ_2) of Bruhat paths such that Δ_1 is a decreasing path from u to w and Δ_2 is an increasing path from w to v . We find a reflection principle on V -paths, which leads to a combinatorial proof of the inversion formula. Moreover, we give two applications of the reflection principle. First, we restrict this involution to V -paths from u to v with maximal length. This provides a direct interpretation for the equi-distribution property that any nontrivial interval $[u, v]$ has as many elements of even length as elements of odd length. This property was obtained by Verma in his derivation of the Möbius function of the Bruhat order. Second, using the reflection principle for the symmetric group, we obtain a refinement of the inversion formula by restricting the summation to permutations ending with a given element.

Keywords: Kazhdan-Lusztig R -polynomial, inversion formula, Bruhat order, reflection ordering

AMS Classification: 05A19, 05E15, 20F55

1 Introduction

Let W be a Coxeter group. For $u, v \in W$, let $R_{u,v}(q)$ be the Kazhdan-Lusztig R -polynomial indexed by u and v . The following inversion formula was obtained by Kazhdan and Lusztig [8]:

$$\sum_{u \leq w \leq v} (-1)^{\ell(w) - \ell(u)} R_{u,w}(q) R_{w,v}(q) = \delta_{u,v}, \quad (1.1)$$

where \leq is the Bruhat order and ℓ is the length function. The aim of this paper is to present a combinatorial interpretation of this formula. This problem was raised by Brenti [3].

To give a combinatorial proof of the above inversion formula, we start with Dyer's combinatorial description of the R -polynomials in terms of increasing Bruhat paths [5]. Then we reformulate the inversion formula in terms of V -paths. For $u \leq w \leq v$, by a V -path from u to v with bottom w we mean a pair (Δ_1, Δ_2) of Bruhat paths such that Δ_1 is a decreasing path from u to w and Δ_2 is an increasing path from w to v . We construct a reflection principle on V -paths. This leads to a combinatorial proof of the inversion formula.

We give two applications of the reflection principle. First, we restrict the reflection principle to V -paths from u to v with maximal length. This induces an involution on the interval $[u, v]$ with $u < v$, which leads to a combinatorial proof of the equidistribution property that any nontrivial interval $[u, v]$ has as many elements of even length as elements of odd length. This property was first conjectured by Verma [11], and proved in [12] by induction. It has been used in Verma's derivation of the Möbius function of the Bruhat order [12]. Other proofs of the Möbius function formula for the Bruhat order can be found in [2, 4, 10]. Recently, Jones [6] found a combinatorial proof for the equidistribution property by constructing an involution on the intervals of a Coxeter group W . When W is finite, Jones [6] showed that this involution agrees with the construction of Rietsch and Williams [9] in their study of discrete Morse theory and totally nonnegative flag varieties. Our involution seems to be simpler as far as the equidistribution property is concerned.

As a second application, we find a refinement of the inversion formula when W is the symmetric group S_n . For a permutation $w \in S_n$, we write $w = w(1)w(2) \cdots w(n)$, where, for $1 \leq i \leq n$, $w(i)$ denotes the element in the i -th position. Let u and v be two permutations in S_n such that $u < v$ in the Bruhat order. For $1 \leq k \leq n$, let $[u, v]_k$ denote the set of permutations in the interval $[u, v]$ that end with k , that is,

$$[u, v]_k = \{w \in [u, v] : w(n) = k\}.$$

Applying a variation of the reflection principle, we show that the summation

$$\sum_{w \in [u, v]_k} (-1)^{\ell(w) - \ell(u)} R_{u,w}(q) R_{w,v}(q)$$

equals zero or a power of q up to a sign.

2 Preliminaries

In this section, we give an overview of the terminology and background on Coxeter groups and R -polynomials that will be used in this paper.

Let (W, S) be a Coxeter system, and

$$T = \{wsw^{-1} : s \in S, w \in W\}$$

its set of reflections. We use $\ell(w)$ to denote the length of $w \in W$. For two elements u and v in W , $u \leq v$ in the Bruhat order if and only if there exists a sequence of reflections t_1, t_2, \dots, t_r in T such that

- (i) $v = ut_1t_2 \cdots t_r$;
- (ii) $\ell(ut_1 \cdots t_i) > \ell(ut_1 \cdots t_{i-1})$ for $1 \leq i \leq r$.

The R -polynomials were introduced by Kazhdan and Lusztig [8] in order to describe the inverses of the basis elements in the Hecke algebra associated to the Coxeter system (W, S) . They obtained the following recurrence relation. For $w \in W$, let $D_R(w)$ be the right descent set of w , that is,

$$D_R(w) = \{s \in S : \ell(ws) < \ell(w)\}.$$

Theorem 2.1 *For any $u, v \in W$,*

- (i) $R_{u,v}(q) = 0$, if $u \not\leq v$;
- (ii) $R_{u,v}(q) = 1$, if $u = v$;
- (iii) If $u < v$ and $s \in D_R(v)$, then

$$R_{u,v}(q) = \begin{cases} R_{us,vs}(q), & \text{if } s \in D_R(u); \\ qR_{us,vs}(q) + (q-1)R_{u,vs}(q), & \text{if } s \notin D_R(u). \end{cases}$$

Notice that the R -polynomials may contain negative coefficients. A variant of the R -polynomials introduced by Dyer [5], which have been called the \tilde{R} -polynomials, have nonnegative coefficients. The following two theorems are due to [5], see also Björner and Brenti [1].

Theorem 2.2 *For any $u, v \in W$,*

- (i) $\tilde{R}_{u,v}(q) = 0$, if $u \not\leq v$;
- (ii) $\tilde{R}_{u,v}(q) = 1$, if $u = v$;

(iii) If $u < v$ and $s \in D_R(v)$, then

$$\tilde{R}_{u,v}(q) = \begin{cases} \tilde{R}_{us,vs}(q), & \text{if } s \in D_R(u); \\ \tilde{R}_{us,vs}(q) + q\tilde{R}_{u,vs}(q), & \text{if } s \notin D_R(u). \end{cases}$$

Theorem 2.3 For $u \leq v \in W$, we have

$$R_{u,v}(q) = q^{\frac{\ell(v)-\ell(u)}{2}} \tilde{R}_{u,v}(q^{\frac{1}{2}} - q^{-\frac{1}{2}}). \quad (2.1)$$

In this paper, we shall give a combinatorial interpretation of the inversion formula for \tilde{R} -polynomials. By Theorem 2.3, the inversion formula (1.1) can be restated as follows

$$\sum_{u \leq w \leq v} (-1)^{\ell(w)-\ell(u)} \tilde{R}_{u,w}(q) \tilde{R}_{w,v}(q) = \delta_{u,v}. \quad (2.2)$$

Dyer [5] has given a combinatorial interpretation of the \tilde{R} -polynomials in terms of increasing paths in the Bruhat graph of a Coxeter group. The Bruhat graph $BG(W)$ of a Coxeter group W is a directed graph whose nodes are the elements of W such that there is an arc from u to v if $v = ut$ for some reflection $t \in T$ and $\ell(u) < \ell(v)$.

The Bruhat graph of the symmetric group S_n can be described as follows. Let (i, j) denote a transposition in S_n , where $1 \leq i < j \leq n$. The reflection set T of S_n consists of the transpositions of S_n , that is,

$$T = \{(i, j) : 1 \leq i < j \leq n\}.$$

For two permutations u, v in S_n , there is an arc from u to v if $v = u(i, j)$ and $u(i) < u(j)$, see [1]. The Bruhat graph of S_3 is illustrated in Figure 2.1.




Figure 2.1: The Bruhat graph of S_3

An increasing path in the Bruhat graph is defined based on the reflection ordering on the positive roots of a Coxeter group. Let Φ be the root system of W , and Φ^+ be

the positive root system. A total ordering \prec on Φ^+ is called a reflection ordering on Φ^+ if for any $\alpha \prec \beta \in \Phi^+$ and two nonnegative real numbers λ, μ such that $\lambda\alpha + \mu\beta \in \Phi^+$, we have

$$\alpha \prec \lambda\alpha + \mu\beta \prec \beta.$$

Here is an example of a reflection ordering for the symmetric group S_n . Consider the following root system

$$\Phi = \{\pm(\varepsilon_i - \varepsilon_j) : 1 \leq i < j \leq n\},$$

where ε_i is the i -th coordinate vector, and consider the following positive root system

$$\Phi^+ = \{\varepsilon_i - \varepsilon_j : 1 \leq i < j \leq n\}.$$

It is easy to verify that the following lexicographic ordering on Φ^+ is a reflection ordering

$$\varepsilon_1 - \varepsilon_2 \prec \varepsilon_1 - \varepsilon_3 \prec \cdots \prec \varepsilon_1 - \varepsilon_n \prec \varepsilon_2 - \varepsilon_3 \prec \cdots \prec \varepsilon_{n-1} - \varepsilon_n. \quad (2.3)$$

Since positive roots in Φ^+ are in one-to-one correspondence with reflections, a reflection ordering induces a total ordering on the set T of reflections. For example, for the symmetric group S_n , a positive root $\varepsilon_i - \varepsilon_j$ corresponds to a transposition (i, j) . Thus, the total ordering on transpositions inheriting from the lexicographic ordering in (2.3) is

$$(1, 2) \prec (1, 3) \prec \cdots \prec (1, n) \prec (2, 3) \prec \cdots \prec (n-1, n). \quad (2.4)$$

We use $u \rightarrow v$ to denote an arc from u to v in the Bruhat graph. Let $\Delta = u_0 \rightarrow u_1 \rightarrow \cdots \rightarrow u_r$ be a path from u to v , where $u_0 = u$ and $u_r = v$. We say Δ is an increasing path with respect to a reflection ordering \prec if

$$t_1 \prec t_2 \prec \cdots \prec t_r,$$

where, for $1 \leq i \leq r$, t_i is the reflection determined by $u_i = u_{i-1}t_i$. For example, the following is an increasing path with respect to the lexicographic ordering in the Bruhat graph of S_4

$$2314 \xrightarrow{(12)} 3214 \xrightarrow{(14)} 4213 \xrightarrow{(24)} 4312.$$

Similarly, we say that $\Delta = u_0 \rightarrow u_1 \rightarrow \cdots \rightarrow u_r$ is a decreasing path with respect to a reflection ordering \prec if

$$t_1 \succ t_2 \succ \cdots \succ t_r,$$

where, for $1 \leq i \leq r$, t_i is the reflection determined by $u_i = u_{i-1}t_i$.

Let $\ell(\Delta)$ denote the length of Δ , namely, the number of arcs in Δ . The following combinatorial interpretation of the \tilde{R} -polynomials was obtained by Dyer [5].

Theorem 2.4 *Let W be a Coxeter group, and T its set of reflections. Then, for any fixed reflection ordering \prec on T and any $u, v \in W$ with $u \leq v$, we have*

$$\tilde{R}_{u,v}(q) = \sum_{\Delta} q^{\ell(\Delta)}, \quad (2.5)$$

where the sum ranges over increasing Bruhat paths from u to v with respect to \prec .

Since the reverse of a reflection ordering is also a reflection ordering, relation (2.5) can be restated as follows

$$\tilde{R}_{u,v}(q) = \sum_{\Delta'} q^{\ell(\Delta')}, \quad (2.6)$$

where the sum ranges over decreasing Bruhat paths from u to v with respect to any fixed reflection ordering. Both (2.5) and (2.6) will be used in our combinatorial interpretation of the inversion formula (2.2) in the next section.

3 A reflection principle on V -paths

In this section, we give a restatement of the inversion formula in terms of V -paths, and we construct a reflection principle on V -paths leading to a combinatorial proof of the inversion formula.

By a V -path from u to v with bottom w under a reflection ordering \prec , we mean a pair (Δ_1, Δ_2) of Bruhat paths such that Δ_1 is a decreasing path from u to w and Δ_2 is an increasing path from w to v . The sign of a V -path (Δ_1, Δ_2) is defined by

$$\text{sgn}(\Delta_1, \Delta_2) = (-1)^{\ell(\Delta_1)}.$$

It is known that the length of a Bruhat path from u to w has the same parity as $\ell(w) - \ell(u)$, see Björner and Brenti [1]. It follows that

$$\text{sgn}(\Delta_1, \Delta_2) = (-1)^{\ell(w) - \ell(u)}. \quad (3.1)$$

Using (2.5), (2.6) and (3.1), we are led to the following restatement of the inversion formula.

Theorem 3.1 *For a Coxeter group W and $u \leq v \in W$, we have*

$$\sum_{u \leq w \leq v} (-1)^{\ell(w) - \ell(u)} \tilde{R}_{u,w}(q) \tilde{R}_{w,v}(q) = \sum_{(\Delta_1, \Delta_2)} \text{sgn}(\Delta_1, \Delta_2) q^{\ell(\Delta_1) + \ell(\Delta_2)} = \delta_{u,v}, \quad (3.2)$$

where the sum ranges over V -paths from u to v under any fixed reflection ordering.

We shall construct a reflection principle I on V -paths from u to v , where $u < v$ in the Bruhat order, which turns out to be a length preserving and sign reversing involution. So we obtain a combinatorial interpretation of the inversion formula.

Reflection Principle on V -Paths: For $u < v$, let (Δ_1, Δ_2) be a V -path from u to v with bottom w . Write

$$\Delta_1 = u_0 \rightarrow u_1 \rightarrow \cdots \rightarrow u_i \quad \text{and} \quad \Delta_2 = v_0 \rightarrow v_1 \rightarrow \cdots \rightarrow v_j,$$

where $u_0 = u$, $u_i = v_0 = w$ and $v_j = v$. From (Δ_1, Δ_2) , we construct a V -path (Δ'_1, Δ'_2) by the following procedure:

- (1) $u = w$, that is, Δ_1 is an empty path. Set

$$\Delta'_1 = u \rightarrow v_1 \quad \text{and} \quad \Delta'_2 = v_1 \rightarrow v_2 \rightarrow \cdots \rightarrow v_j.$$

- (2) $v = w$, that is, Δ_2 is an empty path. Set

$$\Delta'_1 = u_0 \rightarrow u_1 \rightarrow \cdots \rightarrow u_{i-1} \quad \text{and} \quad \Delta'_2 = u_{i-1} \rightarrow v.$$

- (3) $u \neq w$ and $v \neq w$. In this case, we may assume that $u_r = u_{r-1} t_r$ for $1 \leq r \leq i$, and $v_s = v_{s-1} t'_s$ for $1 \leq s \leq j$. If $t_i \prec t'_1$, set

$$\Delta'_1 = u_0 \rightarrow u_1 \rightarrow \cdots \rightarrow u_{i-1} \quad \text{and} \quad \Delta'_2 = u_{i-1} \rightarrow v_0 \rightarrow v_1 \rightarrow \cdots \rightarrow v_j.$$

If $t_i \succ t'_1$, set

$$\Delta'_1 = u_0 \rightarrow u_1 \rightarrow \cdots \rightarrow u_i \rightarrow v_1 \quad \text{and} \quad \Delta'_2 = v_1 \rightarrow \cdots \rightarrow v_j.$$

The V -path (Δ'_1, Δ'_2) is defined to be $I(\Delta_1, \Delta_2)$.

An illustration of the reflection principle is given in Figure 3.2. Let us consider an




Figure 3.2: The reflection principle on V -paths.

example. Let W be the symmetric group S_4 . We choose the lexicographic order in

(2.4) as the reflection ordering. Let $u = 1234$ and $v = 4312$. The following is a V -path from u to v

$$1234 \xrightarrow{(23)} 1324 \xrightarrow{(13)} \underline{2314} \xrightarrow{(12)} 3214 \xrightarrow{(14)} 4213 \xrightarrow{(24)} 4312,$$

where we use underline to indicate the bottom node. Since $(1, 3) \succ (1, 2)$, we see that under the reflection principle I , this V -path is mapped to the V -path

$$1234 \xrightarrow{(23)} 1324 \xrightarrow{(13)} 2314 \xrightarrow{(12)} \underline{3214} \xrightarrow{(14)} 4213 \xrightarrow{(24)} 4312.$$

To conclude this section, we remark that the involution I also yields a combinatorial interpretation of the following result of Verma [12].

Theorem 3.2 *Let W be a Coxeter group and $u < v \in W$. Then the interval $[u, v]$ has the same number of elements of odd length as elements of even length.*

Combinatorial Proof. It is known that for $\sigma \leq \tau \in W$, there exists a unique maximal increasing (or, decreasing) Bruhat path from σ to τ [5]. Thus, for any $w \in [u, v]$ there is a unique maximal V -path from u to v whose bottom is w . So the maximal V -paths from u to v are in one-to-one correspondence with elements in the interval $[u, v]$.

By the construction of the involution I , it can be seen that the set of all maximal V -paths from u to v is invariant under I . Thus, the restriction of I to the maximal V -paths from u to v induces an involution on the elements in the interval $[u, v]$. Moreover, the induced involution on the interval $[u, v]$ changes parity of the length of each element. This completes the proof. \blacksquare

4 A refinement of the inversion formula for S_n

In this section, we use the reflection principle to give a refinement of the inversion formula for the symmetric group S_n . For $u < v \in S_n$ and $1 \leq k \leq n$, let $[u, v]_k$ denote the set of permutations in the interval $[u, v]$ that end with k . We show that the following summation

$$\sum_{w \in [u, v]_k} (-1)^{\ell(w) - \ell(u)} \widetilde{R}_{u,w}(q) \widetilde{R}_{w,v}(q), \quad (4.1)$$

equals zero or a power of q up to a sign. More precisely, let $D(u, v)$ denote the set of indices where u and v differ, that is,

$$D(u, v) = \{1 \leq i \leq n : u(i) \neq v(i)\}.$$

Suppose that $D(u, v) = \{i_1, i_2, \dots, i_j\}_{<}$, that is, $D(u, v) = \{i_1, i_2, \dots, i_j\}$ and $i_1 < i_2 < \dots < i_j$. Let $b_1 < b_2 < \dots < b_j$ be the values of $u(i_1), u(i_2), \dots, u(i_j)$ listed in increasing order. We say that $[u, v]$ is an S-interval if it satisfies the following conditions:

- (1) $i_j = n$ and $u(i_j) = b_j$;
- (2) the values in $\{b_1, b_2, \dots, b_j\}$ that are greater than $u(i_1)$ appear increasingly in u , whereas the values in $\{b_1, b_2, \dots, b_j\}$ that are less than $u(i_1)$ appear decreasingly in u . In other words, if $b_{j_0} = u(i_1)$ ($1 \leq j_0 < j$), then the subsequence $u(i_2) \cdots u(i_{j-1})$ of u is a shuffle of the decreasing sequence $b_{j_0-1} \cdots b_1$ and the increasing sequence $b_{j_0+1} \cdots b_{j-1}$.
- (3) v is obtained from u by rotating the values b_1, b_2, \dots, b_j . That is, for $m \notin D(u, v)$ we have $v(m) = u(m)$, and for $m \in D(u, v)$, if $m = i_j$ then $v(m) = b_1$, and if $m \neq i_j$ then we have $v(m) = b_{t+1}$, where t is the index such that $u(m) = b_t$.

For example, let $u = \mathbf{432596178}$ and $v = \mathbf{453792681}$. So we have $D(u, v) = \{2, 3, 4, 6, 7, 8, 9\}$ and $(b_1, \dots, b_7) = (1, 2, 3, 5, 6, 7, 8)$. The elements b_i are written in boldface. It is easily checked that u and v satisfy Conditions (1), (2) and (3). Notice that the subsequence $u(3)u(4)u(6)u(7)u(8) = 25617$ is a shuffle of 21 and 567. Thus $[u, v]$ is an S-interval.

Our main theorem of this section can be stated as follows.

Theorem 4.1 *Assume that $u < v \in S_n$. Let m be the smallest index such that $u(m) \neq v(m)$. If $[u, v]$ is an S-interval and $k = u(m)$ or $k = v(m)$, then we have*

$$\sum_{w \in [u, v]_k} (-1)^{\ell(w) - \ell(u)} \tilde{R}_{u, w}(q) \tilde{R}_{w, v}(q) = (-1)^r q^{s-1}, \quad (4.2)$$

where $s = |D(u, v)|$ and $r = |\{j \in D(u, v) : u(j) > k\}|$; Otherwise, we have

$$\sum_{w \in [u, v]_k} (-1)^{\ell(w) - \ell(u)} \tilde{R}_{u, w}(q) \tilde{R}_{w, v}(q) = 0. \quad (4.3)$$

We give an example before presenting a proof of Theorem 4.1. Let $n = 7$, $u = 2354167$ and $v = 3564271$. It is easy to check that $[u, v]$ is an S-interval. Consider the sum (4.1) over $[u, v]_2$, $[u, v]_3$ and $[u, v]_5$, where

$$\begin{aligned} [u, v]_2 &= \{3456172, 3465172, 3546172, 3564172\}, \\ [u, v]_3 &= \{2456173, 2465173, 2546173, 2564173\}, \\ [u, v]_5 &= \{2364175, 2463175, 3264175, 3462175\}. \end{aligned}$$

It can be checked that

$$\begin{aligned} \tilde{R}_{2354167, 3456172}(q) &= q^4, & \tilde{R}_{3456172, 3564271}(q) &= q^3, \\ \tilde{R}_{2354167, 3465172}(q) &= q^5, & \tilde{R}_{3465172, 3564271}(q) &= q^2, \\ \tilde{R}_{2354167, 3546172}(q) &= q^5, & \tilde{R}_{3546172, 3564271}(q) &= q^2, \\ \tilde{R}_{2354167, 3564172}(q) &= q^6 + q^4, & \tilde{R}_{3564172, 3564271}(q) &= q, \end{aligned}$$

and thus we have

$$\sum_{w \in [u, v]_2} (-1)^{\ell(w) - \ell(u)} \tilde{R}_{u, w}(q) \tilde{R}_{w, v}(q) = q^5.$$

Similarly, we have

$$\sum_{w \in [u, v]_3} (-1)^{\ell(w) - \ell(u)} \tilde{R}_{u, w}(q) \tilde{R}_{w, v}(q) = -q^5.$$

Since

$$\begin{aligned} \tilde{R}_{2354167, 2364175}(q) &= q^2, & \tilde{R}_{2364175, 3564271}(q) &= q^5 + q^3, \\ \tilde{R}_{2354167, 2463175}(q) &= q^3, & \tilde{R}_{2463175, 3564271}(q) &= q^4, \\ \tilde{R}_{2354167, 3264175}(q) &= q^3, & \tilde{R}_{3264175, 3564271}(q) &= q^4 + q^2, \\ \tilde{R}_{2354167, 3462175}(q) &= q^4, & \tilde{R}_{3462175, 3564271}(q) &= q^3, \end{aligned}$$

we find that

$$\sum_{w \in [u, v]_5} (-1)^{\ell(w) - \ell(u)} \tilde{R}_{u, w}(q) \tilde{R}_{w, v}(q) = 0.$$

To prove Theorem 4.1, let $P_k(u, v)$ denote the set of V -paths from u to v with bottom w contained in $[u, v]_k$. We shall give an involution \tilde{I} on $P_k(u, v)$. From now on, assume that \prec is the lexicographic ordering, namely, the transpositions are ordered by

$$(1, 2) \prec (1, 3) \prec \cdots \prec (1, n) \prec (2, 3) \prec \cdots \prec (n-1, n).$$

We also write a Bruhat path $\Delta = w_0 \rightarrow w_1 \rightarrow \cdots \rightarrow w_r$ as

$$\Delta = (t_1, t_2, \dots, t_r),$$

where $w_i = w_{i-1} t_i$ for $1 \leq i \leq r$. It is clear that $w_i = w_0 t_1 \cdots t_i$.

The Involution \tilde{I} : Let (Δ_1, Δ_2) be a V -path in $P_k(u, v)$ with bottom w . Write

$$\Delta_1 = u_0 \rightarrow u_1 \rightarrow \cdots \rightarrow u_i \quad \text{and} \quad \Delta_2 = v_0 \rightarrow v_1 \rightarrow \cdots \rightarrow v_j,$$

where $u_0 = u$, $u_i = v_0 = w$ and $v_j = v$. Assume that $u_r = u_{r-1} t_r$ for $1 \leq r \leq i$, and $v_s = v_{s-1} t'_s$ for $1 \leq s \leq j$. Since (Δ_1, Δ_2) is a V -path, we have

$$t_1 \succ t_2 \succ \cdots \succ t_i \quad \text{and} \quad t'_1 \prec t'_2 \prec \cdots \prec t'_j.$$

Let $t = \min\{t_i, t'_1\}$. To define \tilde{I} , here are three cases.

Case 1. t is an internal transposition, that is, $t = (a, b)$ and $1 \leq a < b < n$. Set

$$\tilde{I}(\Delta_1, \Delta_2) = I(\Delta_1, \Delta_2).$$

In this case, the bottom node of $\tilde{I}(\Delta_1, \Delta_2)$ is also contained in $[u, v]_k$. Thus $\tilde{I}(\Delta_1, \Delta_2)$ remains a V -path in $P_k(u, v)$.

Case 2. t is a boundary transposition, that is, $t = (a, n)$ for some $a < n$, and there is at least one internal transposition among the transpositions $t_1, \dots, t_i, t'_1, \dots, t'_j$. Let \tilde{t} be the smallest internal transposition among $t_1, \dots, t_i, t'_1, \dots, t'_j$. We consider the following two subcases.

Subcase 1. \tilde{t} belongs to $\{t_1, \dots, t_i\}$. Assume that $t_{i_0} = \tilde{t}$, where $1 \leq i_0 \leq i$. If the transpositions t'_1, \dots, t'_j are all smaller than \tilde{t} , then set

$$\Delta'_1 = (t_1, \dots, \widehat{t_{i_0}}, \dots, t_i) \quad \text{and} \quad \Delta'_2 = (t'_1, t'_2, \dots, t'_j, t_{i_0}),$$

where $\widehat{t_{i_0}}$ indicates that the element t_{i_0} is missing. Define $\tilde{I}(\Delta_1, \Delta_2) = (\Delta'_1, \Delta'_2)$.

We claim that (Δ'_1, Δ'_2) is contained in $P_k(u, v)$. By the choice of t_{i_0} , we see that the boundary transpositions t_{i_0+1}, \dots, t_i commute with t_{i_0} . Thus the bottom of (Δ'_1, Δ'_2) is given by

$$u t_1 \cdots t_{i_0-1} \widehat{t_{i_0}} t_{i_0-1} \cdots t_i = u t_1 \cdots t_{i_0-1} t_{i_0} t_{i_0+1} \cdots t_i t_{i_0} = w t_{i_0}.$$

Clearly, $w t_{i_0}$ is a permutation in $[u, v]_k$. Hence the claim is proved.

If there exists a transposition among t'_1, \dots, t'_j that is greater than \tilde{t} , let $1 \leq p \leq j$ be the smallest index such that $t'_p \succ \tilde{t}$. Set

$$\Delta'_1 = (t_1, \dots, \widehat{t_{i_0}}, \dots, t_i) \quad \text{and} \quad \Delta'_2 = (t'_1, \dots, t'_{p-1}, t_{i_0}, t'_p, \dots, t'_j).$$

Define $\tilde{I}(\Delta_1, \Delta_2) = (\Delta'_1, \Delta'_2)$.

Similarly, it can be checked that the bottom of (Δ'_1, Δ'_2) is $w t_{i_0}$. Hence (Δ'_1, Δ'_2) is a V -path in $P_k(u, v)$.

Subcase 2. \tilde{t} belongs to $\{t'_1, \dots, t'_j\}$. Assume that $t'_{j_0} = \tilde{t}$, where $1 \leq j_0 \leq j$.

If t_1, \dots, t_i are all smaller than \tilde{t} , then set

$$\Delta'_1 = (t'_1, t_1, \dots, t_i) \quad \text{and} \quad \Delta'_2 = (t'_1, \dots, \widehat{t'_{j_0}}, \dots, t'_j).$$

Define $\tilde{I}(\Delta_1, \Delta_2) = (\Delta'_1, \Delta'_2)$.

If there exists a transposition among t_1, \dots, t_i that is greater than \tilde{t} , let $1 \leq p \leq i$ be the largest index such that $t_p \succ \tilde{t}$. Set

$$\Delta'_1 = (t_1, \dots, t_p, t'_{j_0}, t_{p+1}, \dots, t_i) \quad \text{and} \quad \Delta'_2 = (t'_1, \dots, \widehat{t'_{j_0}}, \dots, t'_j).$$

Define $\tilde{I}(\Delta_1, \Delta_2) = (\Delta'_1, \Delta'_2)$. Using the argument in subcase 1, it can be seen that the bottom of (Δ'_1, Δ'_2) is $w t_{i_0}$. Thus (Δ'_1, Δ'_2) is a V -path in $P_k(u, v)$.

Case 3. The transpositions $t_1, \dots, t_i, t'_1, \dots, t'_j$ are all boundary transpositions. In this case, set (Δ_1, Δ_2) to be a fixed point of \tilde{I} , that is, $\tilde{I}(\Delta_1, \Delta_2) = (\Delta_1, \Delta_2)$.

The construction of \tilde{I} implies the following property.

Theorem 4.2 *The map \tilde{I} is a length preserving involution on $P_k(u, v)$. Moreover, \tilde{I} reverses the sign of (Δ_1, Δ_2) unless (Δ_1, Δ_2) is a fixed point.*

To finish the proof of Theorem 4.1, we need the following lemma.

Lemma 4.3 *The involution \tilde{I} has at most one fixed point in $P_k(u, v)$. Moreover, \tilde{I} has a fixed point in $P_k(u, v)$ if and only if $[u, v]$ is an S-interval and $k = u(m)$ or $k = v(m)$, where m is the smallest index such that $u(m) \neq v(m)$.*

Proof. We first show that \tilde{I} has at most one fixed point in $P_k(u, v)$. Suppose that $(\Delta_1, \Delta_2) \in P_k(u, v)$ is a V -path with bottom w that is fixed by \tilde{I} . Let

$$\Delta_1 = (t_1, t_2, \dots, t_i) \quad \text{and} \quad \Delta_2 = (t'_1, t'_2, \dots, t'_j).$$

By the construction of \tilde{I} , we see that t_1, \dots, t_i and t'_1, \dots, t'_j are all boundary transpositions. Assume that

$$t_1 = (p_1, n), \dots, t_i = (p_i, n) \quad \text{and} \quad t'_1 = (p'_1, n), \dots, t'_j = (p'_j, n).$$

Since Δ_1 and Δ_2 are Bruhat paths, we see that

$$u(n) > u(p_1) > \dots > u(p_i) = k = w(n) > w(p'_1) > \dots > w(p'_j). \quad (4.4)$$

Moreover, noting that $t_1 \succ t_2 \succ \dots \succ t_i$ and $t'_1 \prec t'_2 \prec \dots \prec t'_j$, we have

$$n > p_1 > \dots > p_i \quad \text{and} \quad p'_1 < \dots < p'_j < n.$$

By (4.4) and relation $w = u(p_1, n) \dots (p_i, n)$, it is easily seen that $\{p_1, \dots, p_i\} \cap \{p'_1, \dots, p'_j\} = \emptyset$. This implies that

$$w(p'_1) = u(p'_1), \dots, w(p'_j) = u(p'_j),$$

and so (4.4) becomes

$$u(n) > u(p_1) > \dots > u(p_i) = k = w(n) > u(p'_1) > \dots > u(p'_j). \quad (4.5)$$

Observing that $\{p_1, \dots, p_i\} \cup \{p'_1, \dots, p'_j\} \cup \{n\} = D(u, v)$, in view of (4.5) we see that once u, v and k are given, the values i, j as well as $p_1, \dots, p_i, p'_1, \dots, p'_j$ are uniquely determined. This leads to the conclusion that \tilde{I} has at most one fixed point in $P_k(u, v)$.

It remains to prove the assertion that $P_k(u, v)$ has a fixed point under \tilde{I} if and only if $[u, v]$ is an S-interval and $k = u(m)$ or $k = v(m)$. By the above argument, we find that if \tilde{I} has a fixed point in $P_k(u, v)$, then $[u, v]$ is an S-interval and $k = u(p_i) = v(p'_1)$. Since $m = \min\{p_i, p'_1\}$, we obtain that $k = u(m)$ if $p_i < p'_1$ and $k = v(m)$ if $p_i > p'_1$.

Conversely, if $[u, v]$ is an S-interval, it is easy to construct a V -path in $P_k(u, v)$ fixed by \tilde{I} , where $k = u(m)$ or $k = v(m)$. This completes the proof. ■

We give an example to illustrate the above lemma. Let $n = 9$, $u = 432596178$ and $v = 453792681$. It can be seen that $[u, v]$ is an S-interval. The fixed point in $P_3(u, v)$ is (Δ_1, Δ_2) with $\Delta_1 = ((8, 9), (6, 9), (4, 9), (2, 9))$ and $\Delta_2 = ((3, 9), (7, 9))$. The fixed point in $P_5(u, v)$ is (Δ_3, Δ_4) with $\Delta_3 = ((8, 9), (6, 9), (4, 9))$ and $\Delta_4 = ((2, 9)(3, 9), (7, 9))$.

We are now ready to complete the proof of Theorem 4.1.

Proof of Theorem 4.1. By Theorem 4.2 and Lemma 4.3, we only need to consider the case when $[u, v]$ is an S -interval and $k = u(m)$ or $k = v(m)$. In this case, we have

$$\sum_{w \in [u, v]_k} (-1)^{\ell(w) - \ell(u)} \tilde{R}_{u, w}(q) \tilde{R}_{w, v}(q) = (-1)^{\ell(\Delta_1)} q^{\ell(\Delta_1) + \ell(\Delta_2)},$$

where (Δ_1, Δ_2) is the unique V -path in $P_k(u, v)$ that is fixed by \tilde{I} . Evidently, we have

$$\ell(\Delta_1) + \ell(\Delta_2) = |D(u, v)| - 1.$$

Moreover, it is easy to check that

$$\ell(\Delta_1) = |\{j \in D(u, v) : u_j > k\}|.$$

This completes the proof. ■

Acknowledgments. This work was supported by the 973 Project, the PCSIRT Project of the Ministry of Education, and the National Science Foundation of China.

References

- [1] A. Björner and F. Brenti, *Combinatorics of Coxeter Groups*, Grad. Texts in Math., Vol. 231, Springer, New York, 2005.
- [2] A. Björner and M. Wachs, Bruhat order of Coxeter groups and shellability, *Adv. Math.* 43 (1982), 87-100.
- [3] F. Brenti, Kazhdan-Lusztig and R -polynomials from a combinatorial point of view, *Discrete Math.* 193 (1998), 93-116.
- [4] V.V. Deodhar, Some characterizations of Bruhat ordering on a Coxeter group and determination of the relative Möbius function, *Invent. Math.* 39 (1977), 187-198.
- [5] M. J. Dyer, Hecke algebras and shellings of Bruhat intervals, *Compos. Math.* 89 (1993), 91-115.
- [6] B.C. Jones, An explicit derivation of the Möbius function for Bruhat order, *Order* 26 (2009), 319-330.

- [7] J.E. Humphreys, Reflection groups and Coxeter groups, Cambridge Studies in Advanced Mathematics, No. 29, Cambridge Univ. Press, Cambridge, 1990.
- [8] D. Kazhdan and G. Lusztig, Representations of Coxeter groups and Hecke algebras, *Invent. Math.* 53 (1979), 165-184.
- [9] K. Rietsch and L. Williams, Discrete Morse theory for totally non-negative flag varieties, *Adv. Math.* 223 (2010), 1855-1884.
- [10] J. R. Stembridge, A short derivation of the Möbius function for the Bruhat order, *J. Algebraic Combin.* 25 (2007), 141-148.
- [11] D.-N. Verma, Structure of certain induced representations of complex semisimple Lie algebras, *Bull. Amer. Math. Soc.* 74 (1968), 160-166.
- [12] D.-N. Verma, Möbius inversion for the Bruhat ordering on a weyl group, *Ann. Scient. Ec. Norm. Sup.* (1971), 393-399.