

1ES 1927+654: a bare Seyfert 2

L. C. Gallo,¹ C. MacMackin,¹ R. Vasudevan,² E. M. Cackett,³ A. C. Fabian⁴ and F. Panessa⁵

¹ Department of Astronomy and Physics, Saint Mary's University, 923 Robie Street, Halifax, NS, B3H 3C3, Canada

² Department of Astronomy, University of Maryland, College Park, MD 20742-2421, USA

³ Department of Physics and Astronomy, Wayne State University, 666 W. Hancock St, Detroit, MI 48201, USA

⁴ Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA

⁵ Istituto di Astrofisica Spaziale e Fisica Cosmica (IASF-INAF), via del Fosso del Cavaliere 100, 00133 Roma, Italy

Accepted. Received.

ABSTRACT

1ES 1927+654 is an active galactic nucleus (AGN) that appears to defy the unification model. It exhibits a type-2 optical spectrum, but possesses little X-ray obscuration. *XMM-Newton* and *Suzaku* observations obtained in 2011 are used to study the X-ray properties of 1ES 1927+654. The spectral energy distribution derived from simultaneous optical-to-X-ray data obtained with *XMM-Newton* shows the AGN has a typical Eddington ratio ($L/L_{Edd} = 0.014 - 0.11$). The X-ray spectrum and rapid variability are consistent with originating from a corona surrounding a standard accretion disc. Partial covering models can describe the x-ray data; however, the narrow Fe $K\alpha$ emission line predicted from standard photoelectric absorption is not detected. Ionized partial covering also favours a high-velocity outflow ($v \approx 0.3c$), which requires the kinetic luminosity of the wind to be $\gtrsim 30$ per cent of the bolometric luminosity of the AGN. Such values are not unusual, but for 1ES 1927+654 it requires the wind is launched very close to the black hole ($\sim 10 r_g$). Blurred reflection models also work well at describing the spectral and timing properties of 1ES 1927+654 if the AGN is viewed nearly edge-on, implying that an inner accretion disc must be present. The high inclination is intriguing as it suggests 1ES 1927+654 could be orientated like a Seyfert 2, in agreement with its optical classification, but viewed through a tenuous torus.

Key words: galaxies: active – galaxies: nuclei – galaxies: individual: 1ES 1927+654 – X-ray: galaxies

1 INTRODUCTION

Previous *Chandra* and *ROSAT* observations of the active galactic nucleus (AGN) 1ES 1927+654 ($z = 0.017$) found that it did not fit within the standard AGN unification model (Boller et al. 2003, hereafter B03). Its optical spectrum is deficient of the broad emission lines associated with unabsorbed Seyfert 1 (Sy1) galaxies, but its X-ray spectrum lacks the absorption associated with Seyfert 2 (Sy2) galaxies. B03 provided a number of possible explanations including an underluminous broad line region (BLR); an X-ray absorber that is optically thick possibly with a higher than usual dust-to-gas ratio; partial covering absorption; or that 1ES 1927+654 was highly variable, and misclassified due to non-simultaneous X-ray and optical observations. Such “changing look” AGN have been previously observed (e.g. Matt et al. 2003; Bianchi et al. 2005), but simultaneous X-ray and optical observations by Panessa et al. (in prep) show this is not the case for 1ES 1927+654 and that the AGN is a *true* optical type 2 (e.g. Bianchi et al. 2008, 2012; Panessa et al. 2009). Several objects that exhibited this non-standard behaviour have been discovered (e.g. Panessa & Bassani 2002; Mateos et al. 2005; Balestra et al. 2005; Gallo et al. 2006) demonstrating that such objects may not be unusual. Panessa & Bassani (2002) specu-

late that as many as $\sim 10 - 30$ per cent of optically selected type 2 AGN could exhibit such unorthodox behaviour in X-rays (see also Trouille et al. 2009).

Nicastro (2000) considered the possibility of a class of objects that are truly void of a BLR (i.e. true Seyfert 2s). Assuming the BLR is a wind formed in a region of accretion disc instabilities where the disc changes from radiation-pressure dominated to gas-pressure dominated, the radius at which this occurs depends on the accretion rate and will decrease as the accretion rate falls. Once the accretion rate becomes sufficiently low this radius will not be conducive to stable orbits, the wind will cease, and the BLR will fade. This is compatible with an evolutionary scenario proposed by Wang & Zhang (2007), in which they postulate that non-HBLR (non-hidden broad line region) Sy2 galaxies, without absorption, are the end state of AGN development. These galaxies would have the most massive AGNs with the lowest accretion rates. It is worth mentioning that the three confirmed true type 2 AGN (NGC 3147, NGC 3660, Q 2131 – 427) all have low Eddington ratios (Bianchi et al. 2012). Tran et al. (2011) propose 1ES 1927+654 is such an object.

1ES 1927+654 may appear atypical even amongst this unusual

class. In the X-rays, Boller (2000) found the object possessed a steep *ROSAT* spectrum and large-amplitude variability similar to narrow-line Seyfert 1 galaxies (NLS1s), which are normally associated with high Eddington ratios. Even Nicastro et al. (2003) likened IES 1927+654 to a NLS1. Wang et al. (2012) proposed that IES 1927+654 could be a young AGN that has not yet had time to develop a BLR, and predict that such objects should be distinguished by high Eddington ratios.

In this work we examine the X-ray properties of IES 1927+654 making use of non-simultaneous *XMM-Newton* and *Suzaku* data obtained in 2011 in order to determine if the X-rays can elucidate the nature of IES 1927+654. While the AGN was discovered in the *Einstein* survey and observed with *ROSAT* and the *Chandra* LETG (B03), these most recent observations are the highest quality data obtained to date of IES 1927+654 above ~ 2 keV. In the next section we describe the observations and data processing. In Section 3 we fit the X-ray spectra of IES 1927+654 with traditional models in order to compare with other AGNs, and in Section 4 we model the UV-to-X-ray spectral energy distribution. We use this information in Section 5 to consider physical motivated spectral models for the X-ray emission. The X-ray variability over the past 20 years, as well as the shorter time scales during the *Suzaku* and *XMM-Newton* observations, are examined in Section 6. We discuss our results and conclusions in Section 7 and 8, respectively.

2 OBSERVATIONS AND DATA REDUCTION

IES 1927+654 was observed with *Suzaku* (Mitsuda et al. 2007) and *XMM-Newton* (Jansen et al. 2001) starting on 2011 April 16 and May 20, respectively. The duration of the *Suzaku* observation was 122 ks and that of the *XMM-Newton* observation was 29 ks. A summary of the observations is provided in Table 1.

The EPIC pn (Strüder et al. 2001) and MOS (MOS1 and MOS2; Turner et al. 2001) cameras were operated in small-window and full-window modes, respectively, and with the thin filter in place. The Reflection Grating Spectrometers (RGS1 and RGS2; den Herder et al. 2001) and the Optical Monitor (OM; Mason et al. 2001) also collected data during this time.

The *XMM-Newton* Observation Data Files (ODFs) from all observations were processed to produce calibrated event lists using the *XMM-Newton* Science Analysis System (SAS v12.0.0). EPIC response matrices were generated using the SAS tasks ARFGEN and RMFGEN. Light curves were extracted from these event lists to search for periods of high background flaring, which was deemed negligible. The total amount of good pn exposure is listed in Table 1. Source photons were extracted from a circular region $35''$ across and centered on the source. Pile up was examined for and determined to be unimportant. The background photons were extracted from an off-source region on the same CCD. Single and double events were selected for the pn detector, and single-quadruple events were selected for the MOS. The MOS and pn data at each epoch are compared for consistency and determined to be in agreement within known uncertainties (Guainazzi et al. 2010).

The RGS spectra were extracted using the SAS task RGSPROC and response matrices were generated using RGSRMFGEN. The OM operated in imaging mode and collected data in the *V*, *UVW1*, and *UVM2* filters.

During the *Suzaku* observation the two front-illuminated (FI) CCDs (XIS0 and XIS3), the back-illuminated (BI) CCD (XIS1),

and the HXD-PIN all functioned normally and collected data. The target was observed in the XIS-nominal position.

Cleaned event files from version 2 processed data were used in the analysis and data products were extracted using `xselect`. For each XIS chip, source counts were extracted from a $4'$ circular region centred on the target. Background counts were taken from surrounding regions on the chip. Response files (rmf and arf) were generated using `xisrmfgen` and `xissimarfgen`. After examining for consistency, the data from the XIS-FI were combined to create a single spectrum.

The PIN spectrum was extracted from the HXD data following standard procedures. A non-X-ray background (NXB) file corresponding to the observation was obtained to generate good-time interval (GTI) common to the data and NXB. The data were also corrected for detector deadtime. The resulting PIN exposure was 74330 s. The cosmic X-ray background (CXB) was modelled using the provided flat response files. The CXB and NXB background files were combined to create the PIN background spectrum. Examination of the PIN data yielded no detection of the AGN.

All parameters are reported in the rest frame of the source unless specified otherwise. The quoted errors on the model parameters correspond to a 90% confidence level for one interesting parameter (i.e. a $\Delta\chi^2 = 2.7$ criterion). A value for the Galactic column density toward IES 1927+654 of $6.87 \times 10^{20} \text{ cm}^{-2}$ (Kalberla et al. 2005) is adopted in all of the spectral fits and solar abundances from Anders & Grevesse (1989) are assumed unless stated otherwise (see Section 5.1). K-corrected luminosities are calculated using a Hubble constant of $H_0=70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and a standard flat cosmology with $\Omega_M = 0.3$ and $\Omega_\Lambda = 0.7$.

3 CHARACTERIZING THE X-RAY SPECTRA

In this section we attempt to characterize the X-ray spectra of IES 1927+654 with traditional phenomenological models. Initially, the spectra from all the *Suzaku* and *XMM-Newton* CCDs were fitted separately. It was found that *XMM-Newton* pn data agreed well with the MOS data. Similarly, it was found that the data from the *Suzaku* FI detectors agreed well with data from the BI CCD. Since all the data agreed within the calibration uncertainties we present only the *XMM-Newton* pn and *Suzaku* FI spectra for multi-epoch comparison and for ease of presentation. Due to uncertainties in the calibration the FI data are ignored below 0.7 keV and between 1.5 – 2.5 keV. In general, given the AGN spectrum is rather steep the data quickly become noisy at higher energies. This seems to be the main cause of the increased residuals above ~ 7 keV. The *Suzaku* data above 9 keV become completely background dominated and are consequently ignored. The pn data are used between 0.3 – 10 keV.

For comparison the pn and FI spectra are shown in Fig 1 corrected for instrumental differences. The spectra are comparable at both epochs except that the *Suzaku* spectrum appears slightly flatter and dimmer. Both spectra show a drop at lower energies commensurate with some level of absorption in addition to Galactic.

3.1 The 2.5 – 10 keV band

Fitting the 2.5 – 10 keV band at each epoch with a single power law modified by the Galactic column density resulted in a good fit ($\chi^2_\nu = 0.99$). The photon indices were $\Gamma \approx 2.41$ and $\Gamma \approx 2.27$ for the pn and FI, respectively. If the X-ray spectra were highly absorbed like in typical Seyfert 2s one would predict a strong narrow

Table 1. *Suzaku* and *XMM-Newton* observation logs for IES 1927+654. The start date of the observation is given in column (1). The telescope and instrument used is shown in column (2) and (3), respectively. Column (4) is the ID corresponding to the observations. Good exposure time and source counts (corrected for background) are given in column (5) and (6), respectively. For the EPIC instruments the counts correspond to the 0.3 – 10 keV band. For the RGS the counts correspond to the 0.3 – 2 keV band. The values for the *Suzaku* CCDs are taken in the 0.7 – 1.5 and 2.5 – 10 keV band.

(1)	(2)	(3)	(4)	(5)	(6)
Start Date (year.mm.dd)	Telescope	Instrument	Observation ID	Exposure (s)	Counts
2011.04.16	<i>Suzaku</i>	BI	706006010	71890	40611
		FI		71900	60132
		PIN		74330	NA
2011.05.20	<i>XMM-Newton</i>	PN	0671860201	19750	99132
		MOS1		28010	35096
		MOS2		28030	34728
		RGS1		28480	4814
		RGS2		28520	5356




Figure 1. The *XMM-Newton* pn (black) and *Suzaku* FI (red) spectra corrected for instrumental differences. Both spectra exhibit a steep slope and a drop at low energies, likely from absorption in addition to Galactic levels.

Fe $K\alpha$ feature at around 6.4 keV, however the residuals show no deviations around this band. Adding a narrow ($\sigma = 1$ eV) Gaussian profile at 6.4 keV to the model did not provide a significant improvement. The upper-limit on the flux of the narrow feature is $< 9.39 \times 10^{-15}$ erg cm $^{-2}$ s $^{-1}$, corresponding to an upper-limit on the equivalent width of $EW < 30$ eV and $EW < 28$ eV for pn and FI, respectively. Allowing the width and energy of the Gaussian profile to vary freely did not improve the fit over the simple power law.

3.2 The broadband X-ray spectra

When the 2.5 – 10 keV power law fit from Section 3.1 is extrapolated to lower energies, a clear soft excess is evident below ~ 2 keV (Fig 2 top panel).

Adding a blackbody component worked well to describe the




Figure 2. Top panel: The residuals remaining when fitting a power law absorbed by Galactic column density to the 2.5 – 10 keV band and extrapolating to 0.3 keV. A soft-excess below ~ 2 keV is revealed, which is then modified at energies below ~ 0.7 keV. Lower panel: The residuals after the addition of blackbody modified by additional absorption.

data. Since the pn data extend to lower energies than the FI data, the blackbody temperature was linked between the two epochs, but all other parameters were allowed to vary. The fit was statistically acceptable with $\chi^2_{\nu}/\text{dof} = 1.01/1226$. Replacing the blackbody with a second power law (i.e. a double power law) or refitting the spectra with a broken power law generated poorer fits ($\chi^2_{\nu}/\text{dof} = 2.20/1226$ and $1.20/1227$, for the double power law and broken power law, respectively).

Although reasonable fits were possible, in particular with the blackbody plus power law model, all three models showed a drop in the residuals at energies below about 0.5 keV. The addition of a neutral absorber intrinsic to the host galaxy (`ztbabs`) improved the residuals in all cases. The level of absorption was modest ($\sim 5 - 10 \times 10^{20}$ cm $^{-2}$) and it was in line with what was

reported with *ROSAT* observations (B03), which were sensitive to even lower energies than the pn. The parameters are reported for each model in Table 2. The residuals from the best-fitting blackbody plus power law model are shown in Fig 2 (lower panel). The observed 0.3 – 2 keV and 2 – 10 keV fluxes during the *XMM-Newton* observation are 6.8 and 3.7×10^{-12} erg cm $^{-2}$ s $^{-1}$, respectively. The intrinsic 2–10 keV luminosity, corrected for absorption in the Galaxy and host galaxy, is $L = 2.4 \times 10^{42}$ erg s $^{-1}$ during the *XMM-Newton* observation.

3.3 RGS data

The small level of cold absorption evident in the CCD spectra motivated investigation at higher spectral resolution with the *XMM-Newton* RGS data. The RGS data between 0.4–2.0 keV were examined 100 channel at a time and fitted with a simple power law (corrected for Galactic absorption). A Gaussian profile was used to examine for possible narrow emission and absorption features. A potential absorption feature was found around 0.85 keV, but was not constrained in the error analysis. No other features were detected within the available signal-to-noise.

4 OPTICAL-TO-X-RAY SPECTRAL ENERGY DISTRIBUTION

We construct a broadband SED using the *XMM-Newton* pn and OM data along with the *Suzaku* FI data. The OM points are corrected for Galactic dust extinction [$E(B - V)_{\text{Gal}} = 0.087 \pm 0.003$] (Schlegel, Finkbeiner & Davis 1998)¹ before fitting in XSPEC. We follow the procedure of Vasudevan et al. (2009) for constructing the SED, using the xspec model combination ZDUST(DISKPN)+ZTBABS(BKNPOWER), to allow for optical/UV reddening as well as X-ray absorption obscuring the intrinsic accretion disc and power-law emission, respectively. The power law falls in the UV where the disc dominates. We include an intrinsic (host galaxy) dust reddening component of $E(B - V) = 0.55$, corresponding to the upper limiting intrinsic extinction A_V of 1.71 identified in B03 from the H_α/H_β line ratio, using the standard relation $E(B - V) = A_V/R_V$ and assuming $R_V = 3.1$. We freeze the normalisation of the DISKPN model using the black hole mass estimate of $\log(M_{\text{BH}}/M_{\text{sun}}) = 7.34$ from Tran et al. (2011), and assume an accretion disc extending close to the innermost stable orbit, down to $6 R_g$. The photon index of the power law in the 2–10 keV regime is fixed at 2.3, as found from the analysis in Section 3.2 Under these assumptions, we obtain a bolometric (absorption-corrected) luminosity of 3.0×10^{44} erg s $^{-1}$ (integrated between 0.001 and 100 keV), and find that at least 53 per cent of this power is absorbed by dust. These assumptions yield an Eddington ratio of 0.11, larger than the value of 0.006 estimated from Tran et al. (2011) due to the extra disc contribution extrapolated here into the unobservable far-UV. If we consider an alternate disc geometry where the disc is truncated further out from the black hole (at, e.g. $60 R_g$), we find that the Eddington ratio reduces to 0.046 (with 80 per cent of the bolometric flux absorbed by dust reddening), but is still well above the Tran et al. estimate. If we simply extrapolate a power law from the X-rays down to the optical and UV, discarding any contribution from the disc and integrate over the same range,




Figure 3. Spectral energy distribution (SED) for 1ES 1927+654, from *XMM-Newton* PN and OM data and *Suzaku* FI data (black filled points). The dashed line shows an accretion disk and broken power law model fit with dust extinction in the optical–UV and photoelectric absorption in X-rays, and the solid line shows the unabsorbed fit (with dust reddening and absorption set to zero), with a prominent unobservable accretion disc component. The dot-dashed line shows a simple power law extrapolation from X-rays down to optical wavelengths.

we find a lower limit to the Eddington ratio of 0.014. It is also possible that the optical/UV flux is not from a canonical accretion disc, but we do not explore that possibility further here.

Overall, this implies that the accretion rate of the object (0.014 – 0.11) is in between the Wang et al. and Tran et al. estimates, regardless of the type of geometry we assume for the accretion disc. A substantial fraction of the intrinsic luminosity is probably absorbed by dust (between 50–80 per cent). We have not attempted a host-galaxy light correction to the XMM-OM fluxes, but this would serve to push the true bolometric luminosity down and reduce the Eddington accretion rate from the values seen here. The optical images reveal a crowded field, so a more refined analysis may be needed to more accurately estimate the bolometric luminosity and Eddington ratio.

5 MULTI-EPOCH SPECTRAL ANALYSIS OF 1ES 1927+654

Based on our phenomenological approach in Section 3, there are only slight differences between the *XMM-Newton* and *Suzaku* data sets (Table 2). In this section we attempt to model the two broadband X-ray data sets simultaneously and in a self-consistent manner while testing more physically motivated models. For example, while the blackbody plus power law model presented in Section 3 was statistically pleasing, the implied thermal origin for the soft excess is questionable. Given the relatively large black hole mass, even for the highest Eddington accretion rate measured in Section 4, a standard accretion disc extending down to $2 r_g$ would peak at a temperature of $kT \approx 45$ eV. Any blackbody disc component in 1ES 1927+654 should have a temperature that is less than one-

¹ <http://irsa.ipac.caltech.edu/applications/DUST/>

Table 2. Results from fitting *XMM-Newton* and *Suzaku* spectra with traditional models. The broadband X-ray model is stated in column (1). The model components and parameters are shown in column (2) and (3), respectively. The parameter values during the *XMM-Newton* and *Suzaku* epochs are reported in columns (4) and (5), respectively. Observed fluxes are reported for each component in units of $\times 10^{-12}$ erg cm $^{-2}$ s $^{-1}$ over the 0.3 – 10 keV band except in the case of the broken power law model where fluxes are reported above and below the break energy. Parameters that are linked between epochs are only reported in one column. The Galactic column density has been included in all models.

(1) Model	(2) Model component	(3) Model parameter	(4) <i>XMM-Newton</i>	(5) <i>Suzaku</i>	
Broken power law	Intrinsic Absorption	N_H (cm $^{-2}$)	$1.24 \pm 0.08 \times 10^{21}$		
		Power law	Γ_1	3.30 ± 0.06	
			Flux	25.3 ± 0.1	21.3 ± 0.3
			E_b (keV)	1.82 ± 0.07	
			Γ_2	2.47 ± 0.04	2.27 ± 0.04
			Flux	4.06 ± 0.02	3.99 ± 0.03
		Fit Quality	χ^2_ν/dof	1.11/1226	
Double power law	Intrinsic Absorption	N_H (cm $^{-2}$)	$1.22^{+0.11}_{-0.10} \times 10^{21}$		
		Power law 1	Γ	$3.40^{+0.12}_{-0.11}$	
			Flux	26.2 ± 0.2	20.6 ± 0.2
	Power law 2	Γ	$1.49^{+0.17}_{-0.20}$	$1.60^{+0.11}_{-0.13}$	
		Flux	3.13 ± 0.10	4.06 ± 0.10	
		Fit Quality	χ^2_ν/dof	1.23/1225	
Blackbody plus power law	Intrinsic Absorption	N_H (cm $^{-2}$)	$2.58 \pm 0.01 \times 10^{20}$		
		Power law	Γ	2.39 ± 0.04	2.27 ± 0.04
			Flux	3.35 ± 0.06	2.75 ± 0.10
	Blackbody	kT (keV)	0.170 ± 0.005		
		Flux	12.36 ± 0.09	10.7 ± 0.1	
		Fit Quality	χ^2_ν/dof	0.97/1225	

quarter of what we measure in the X-rays. In general, the lack of $T_\infty M^{-1/4}$ relation between the black hole mass and disc temperature observed in samples of unobscured AGN (e.g. Crummy et al. 2006) suggests a non-thermal origin for the soft excess. Both absorption (e.g. Gierliński & Done 2004) and blurred reflection models (e.g. Crummy et al. 2006) can be used to describe the shape of the soft excess via atomic processes, and we will attempt such models to describe the spectrum of IES 1927+654. In doing so, we will also be testing some of the scenarios that have been put forth to describe the characteristics of IES 1927+654.

5.1 Neutral partial covering

We consider the scenario in which the intrinsic power law continuum is partially obscured by a neutral absorber that lies along the line-of-sight. The observed X-ray emission is then the combination of the primary emitter and a highly obscured component. Such models have been proposed for type-1 Seyfert galaxies with relative success in fitting the spectrum (e.g. Gallo et al. 2004; Tanaka et al. 2005; Grupe et al. 2008).

Since the *XMM-Newton* and *Suzaku* observations were separated by only about one month, we assumed that the primary emitter (e.g. the power law component) remained constant in shape and normalisation and that only changes in the absorber (i.e. the covering fraction and column density) would be necessary to describe the differences at the two epochs. The simplest case of the single absorber resulted in a mediocre fit ($\chi^2_\nu/\text{dof} = 1.18/1226$; Fig 4 top panel). The addition of a second absorber (i.e. neutral double partial covering) was favorable, as an improvement could be achieved by changing only the covering fraction of the second absorber ($\Delta\chi^2 = 43$ for 1 additional free parameter). The model and

quality of fit are shown in Table 3 and Fig 4 (lower panel), respectively.

The two absorbers have considerably different densities, which could be indicative of a density gradient along the line-of-sight as opposed to two distinct absorbing regions (e.g. Tanaka et al. 2004). The slightly lower flux during the *Suzaku* observation can be attributed to slightly higher covering fraction while the primary power law emission remains unchanged between the epochs. Allowing the power law component to vary along with the absorbers does not produce a significant improvement.

While the fit is reasonable, there are clearly spectral regions where the model does not adequately fit the data, for example between 0.4 – 0.6 keV and 7 – 10 keV (Fig 4 top panel). The 0.4 – 0.6 keV residuals can be improved with the addition of an absorption edge with rest-frame energy $E = 0.39 \pm 0.01$ keV and optical depth $\tau = 0.66 \pm 0.12$. Given the energy the feature seems unlikely to originate from calibration issues around the oxygen edge. We considered the possibility that the residuals, could be improved by adjusting the elemental abundance. In other abundance tables, like those of Wilms, Allen & McCray (2000) or Grevesse & Sauval (1998), the oxygen abundance relative to hydrogen can be almost half of what is used in Anders & Grevesse (1989). However, while the adopted abundance table does influence some model parameters, the residuals between 0.4 – 0.6 keV do not change significantly.

5.2 Ionized partial covering

The negative residuals between 0.4 – 0.6 keV resulting from the neutral partial covering model (Fig 4) could be indicative of an absorber with some level of ionisation. A moderately ionised absorber




Figure 4. Top panel: The residuals remaining from fitting the *XMM-Newton* and *Suzaku* spectra with one neutral absorber partially covering the power law source. The changes are attributed to variations in the covering fraction and column density at each epoch. Lower panel: The residuals remaining from fitting the *XMM-Newton* and *Suzaku* spectra with two neutral absorbers partially covering the power law source. In this case the variations from one epoch to the next are attributed to changes in the covering fraction of only one absorber. See text and Table 3 for details.

will preferentially remove intermediate energy X-rays giving rise to a spectral break in the 0.3 – 10 keV band generating a soft-excess and it could account for some of the low-energy residuals.

As with single neutral partial covering, a single ionised absorber (`zxipcf` in `xspec`) was a modest fit to the data ($\chi^2_{\nu}/\text{dof} = 1.32/1225$; Fig 5 top panel). A considerably better fit could be achieved with the addition of a second ionised absorber. We tested various combinations of parameters to determine which were the most important to describe the spectra in a self-consistent manner. We determined that an excellent fit could be obtained ($\chi^2_{\nu}/\text{dof} = 1.01/1221$) when the covering fraction of each absorber was allowed to vary at each epoch (Table 3 and Fig 5 lower panel), but we also recognize the power law photon index is considerably steeper than is normally seen in AGN (likewise for the neutral partial covering model). In addition, we found the best fit was achieved when the absorbers were significantly blueshifted with respect to the source frame, consistent with an outflow of 90000 ± 3000 km/s (0.30 ± 0.01 c). The value of the blueshift was tightly constrained independent of the uncertainty in other covering parameters like the column density, covering fraction, and ionisation.

We note the ionisation parameter of the second absorber is rather low $\log(\xi) \approx -0.54$ and could perhaps be described with another neutral absorber instead (e.g. `ztbabs`). The fit was significantly poorer than the double ionised absorber, but acceptable ($\chi^2_{\nu}/\text{dof} = 1.05/1222$). We do not test various combinations of cold and warm absorbers any further. In all likelihood, the absorbers may not be distinct regions with different densities, but a single region with some density and/or ionisation gradient. Comparing the residuals from the ionized and neutral partial covering




Figure 5. Top panel: The residuals remaining from fitting the *XMM-Newton* and *Suzaku* spectra with one ionised absorber partially covering the power law source. The ionisation parameter, covering fraction and column density of the absorber are allowed to vary at each epoch. Lower panel: The improved residuals from the addition of a second ionised absorber are shown. The variations from one epoch to the next are attributed to changes in the covering fraction of each ionised absorber. See text and Table 3 for details.

models (Fig 4 and 5) shows the ionized partial covering is favoured. The deviations between 0.4 – 0.6 keV and 7 – 10 keV seen in the neutral absorber model are considerably improved with the ionized absorber.

If the partial covering model were correct it would only have a small impact on the Eddington ratio estimated in Section 4. The unabsorbed X-ray luminosity in this model is approximately 3.5 times greater than the value used in Section 4, thereby increasing the bolometric luminosity only slightly. The unabsorbed putative disc would still dominate and the upper limit on the Eddington ratio would only increase to about 0.13.

5.3 Compton-thick absorption

There has been speculation that 1ES 1927+654 could be a Compton-thick source. Indeed, Bianchi et al. (2012) found that some true Sy 2 candidates were in fact Compton-thick. With observations available in the 0.3 – 50 keV band (i.e. including the PIN null detection) we can test this model more robustly. For this purpose we used the model `MYTORUS` (Murphy & Yaqoob 2009), in which we view the central engine through some fraction of the torus. Reasonable fits were established, but only in cases with low, face-on, inclinations ($i \approx 0^\circ$), which would be indicative of no absorption from a torus. Attempts to fix the disc inclination more edge-on ($i \approx 85^\circ$) generated very poor fits (Fig 6).

5.4 Ionized Disc Reflection

Reflection from an ionized disc blurred for relativistic effects close to the black hole is often adopted to describe the origin of the soft




Figure 6. The residuals based on an edge-on dusty torus model fitted to the data.

excess (Ballantyne, Ross & Fabian 2001; Ross & Fabian 2005), and has been successfully fitted to the spectra of unabsorbed AGN (e.g. Fabian et al. 2004; Crummy et al. 2006; Ponti et al. 2010; Walton et al. 2013). We consider this scenario to describe the spectra of IES 1927+654 and the variations between the two epochs.

Given that the observations were obtained within about one-month of each other and that the AGN is in a similar flux state we do not expect significant variations. Initially, only the power law slope and normalisation; and the reflector ionisation and normalisation were permitted to vary. The blurring parameters were fixed to default values and linked between the two observations. The model produced a reasonable fit ($\chi^2_{\nu}/\text{dof} = 1.04/1224$).

We examined each individual parameter and then combinations of parameters, to determine which variables would improve the fit most significantly. We determined that the fit was substantially improved ($\Delta\chi^2 = 104$ for 2 additional free parameters) only when the inclination and iron abundance were free to vary (Fig 7 and see Blurred Reflection (A) in Table 3). The iron abundance was found to be about twice the solar value ($A_{Fe} = 2.78^{+1.03}_{-1.21}$) and the disc was significantly inclined (edge-on, $i = 86 \pm 1^\circ$). Permitting more than these two additional parameters to vary did not improve the quality of the fit. We further examined how the two parameters depend on each other and found that they influence each other rather modestly (Fig 8).

We note that the measured inclination is rather extreme while the emissivity index is fixed to $q = 3$ as expected from lamp-post illumination at a large distance. However, the quality of the fit does not change if the emissivity is fixed to values of $q > 3$. Similarly, while the disc inner edge was initially fixed to $R_{in} = 4.5 r_g$, the value in the current fit could be reduced to $R_{in} = 3.3 r_g$ and still achieve a perfectly acceptable fit ($\chi^2_{\nu} = 0.99$).

In fact, a second blurred reflection model that is more relativistic in nature fits the spectra equally well. In this case, we fixed the inner emissivity index to $q_{in} = 5$ and the disc inner edge to $R_{in} = 1.5 r_g$ (see Blurred Reflection (B) in Table 3). While the iron abundance and ionization parameter slightly decrease in this relativistic fit, the inclination still remains high ($i \approx 90^\circ$).

The two blurred reflection models demonstrate that compact geometries for the primary emitter cannot be ruled out, but the fit shows a preference for high inclinations. We also note there are no dominant Fe K α emission features in the spectrum, so the blurred reflection models are primarily fitting the soft excess.




Figure 7. Top panel: The blurred reflection model (A) for the *Suzaku* FI (red) and *XMM-Newton* (black) data. The source is power law dominated over the entire X-ray band at both epochs. Lower panel: The residuals from the model fit in the 0.3 – 10 keV band.




Figure 8. A contour plot of the blurred reflection model (A) fit with varying inclination and iron abundance. The contour lines are drawn around $\Delta\chi^2$ values of 3, 5, and 9.

Table 3. Results from fitting the *XMM-Newton* and *Suzaku* spectra with physically motivated models. The model is stated in column (1). The model components and parameters are shown in column (2) and (3), respectively. The parameter values during the *XMM-Newton* and *Suzaku* epochs are reported in columns (4) and (5), respectively. Parameters that are linked between epochs are only reported in one column. The Galactic column density has been included in all models.

(1) Model	(2) Component	(3) Parameter	(4) <i>XMM-Newton</i>	(5) <i>Suzaku</i>	
Neutral double partial covering	Intrinsic Absorption Absorber 1	N_H (cm $^{-2}$)	$(1.21 \pm 0.08) \times 10^{21}$		
		N_H (cm $^{-2}$)	$5.8^{+3.3}_{-2.0} \times 10^{23}$		
	Absorber 2	C_f	$0.52^{+0.11}_{-0.09}$		
		N_H (cm $^{-2}$)	$(7 \pm 1) \times 10^{22}$		
	Power Law	C_f	$0.57^{+0.03}_{-0.04}$		0.64 ± 0.03
		Γ	3.27 ± 0.06		
Fit Quality	χ^2_v/dof		$1.15/1225$		
Ionized double partial covering	Intrinsic Absorption Absorber 1	N_H (cm $^{-2}$)	$9.23^{+0.40}_{-1.16} \times 10^{20}$		
		N_H (cm $^{-2}$)	$6.6^{+1.6}_{-0.7} \times 10^{23}$		
	Absorber 2	$\log(\xi)$	$3.03^{+0.11}_{-0.16}$		0.34 ± 0.09
		C_f	0.49 ± 0.11		
		z^1	-0.30 ± 0.01		
		N_H (cm $^{-2}$)	$3.34^{+2.5}_{-1.0} \times 10^{22}$		
	Power Law	$\log(\xi)$	$-0.54^{+0.81}_{-0.70}$		$0.43^{+0.07}_{-0.11}$
		C_f	$0.28^{+0.11}_{-0.08}$		
		z^1	-0.30 ± 0.01		
	Fit Quality	χ^2_v/dof		$1.01/1221$	
Blurred reflection (A)	Intrinsic Absorption Power Law	N_H (cm $^{-2}$)	$5.00^{+0.39}_{-0.56} \times 10^{20}$		
		Γ	$2.48^{+0.05}_{-0.03}$	2.36 ± 0.04	
	Blurring	E_{cut}	300^f		
		q	3^f		
		R_{in} (r_g)	4.5^f		
		R_{out} (r_g)	400^f		
		i ($^\circ$)	86 ± 1		
	Reflection	A_{Fe} ($Fe/solar$)	$2.78^{+1.03}_{-1.21}$		310^{+197}_{-91}
		ξ	384^{+162}_{-131}		
	Fit Quality	χ^2_v/dof		$0.96/1222$	
Blurred reflection (B)	Intrinsic Absorption Power Law	N_H (cm $^{-2}$)	$(4.40 \pm 0.05) \times 10^{20}$		
		Γ	2.43 ± 0.05	2.31 ± 0.05	
	Blurring	E_{cut}	300^f		
		q_{in}	5^f		
		R_{in} (r_g)	1.5^f		
		R_{out} (r_g)	400^f		
		R_{break} (r_g)	5^f		
		q_{out}	3^f		
	Reflection	i ($^\circ$)	90^{+0}_{-1}		
		A_{Fe} ($Fe/solar$)	$0.76^{+0.37}_{-0.16}$		202^{+35}_{-120}
ξ		200^{+25}_{-88}			
Fit Quality	χ^2_v/dof		$0.96/1222$		

¹The redshifts of the partial covering components in this model are linked.

6 TIMING ANALYSIS

In Fig 9 the 0.5 – 2 keV light curve of IES 1927+654 over the past ~ 20 years is shown. The vertical bars mark the range between minimum and maximum flux that is observed at each epoch. The *XMM-Newton* and *Suzaku* fluxes are measured from the blackbody plus power law model described in Section 3.2. Fluxes from the earlier data are estimated from the count rates and models presented by B03 using *webPIMMS*.²

IES 1927+654 varies significantly during all observations. On average, the *XMM-Newton* and *Suzaku* observations catch IES 1927+654 at lower X-ray flux levels than have been previously observed.

In this section we examine the variability of the AGN during the *XMM-Newton* and *Suzaku* observations. The short, uninterrupted *XMM-Newton* observations will provide the opportunity to examine the rapid variability (i.e. over hundreds of seconds) with high count rates. The *Suzaku* observation is interrupted every ~ 5.7 ks due to Earth occultation, but the longer duration of the observation allows us to study the variability over ~ 1.5 -days.

² <http://heasarc.gsfc.nasa.gov/Tools/w3pimms.html>




Figure 9. The 0.5 – 2 keV light curve of IES 1927+654 over the past ~ 20 years. The mission from which the data are obtain is labeled beside each point. The vertical bars represent the range between minimum and maximum flux that is observed at that epoch.

6.1 Rapid and large amplitude variability

One of the distinguishing characteristics of IES 1927+654 from *ROSAT* observations was the rapid variability, comparable to what is seen in narrow-line Seyfert 1 galaxies (Boller 2000; B03). Even at this lower flux level, the AGN continues to exhibit impressive variability.

The broadband (0.2 – 12 keV) pn light curve is shown in Fig 10. The variability is persistent over the 30 ks. During the largest flares the count rate changes by more than a factor of 3 in about a thousand seconds. We calculated the radiative efficiency (η) assuming photon diffusion through a spherical mass of accreting material (Fabian 1979). The most rapid rise seen in the light curve corresponds to a luminosity change of $8.95 \times 10^{42} \text{ erg s}^{-1}$ in about 900 s (rest frame). However, as IES 1927+654 is a relatively low-luminosity AGN the lower limit on the radiative efficiency is $\eta \geq 0.5$ per cent. Anisotropic emission or a maximum spinning black hole is not required to describe the large amplitude variability in IES 1927+654.

Light curves were created in several energy bands between 0.2 – 12 keV to compare the variability at different energies. All of the energy bands examined exhibited persistent variability. Given the highly variable light curves from IES 1927+654 we decided to perform a search for lags in these data. To date, over a dozen AGN display reverberation lags (e.g., Fabian et al. 2009, 2012, Zoghbi et al. 2012, Emmanoulopoulos et al. 2011, De Marco et al. 2011, 2012). We initially search for lags between two bands, a soft band (0.4 – 1.5 keV) where the soft excess is strong, and a hard band (2 – 4 keV) dominated by the power-law. We extracted light curves in these bands using 20 s binning. From these light curves we calculate the cross spectrum, and determine the phase lag from the argument of the cross spectrum (see Nowak et al. 1999 for a detailed description). The time lag is simply the phase lag divided by $2\pi f$, where f is the Fourier frequency. We find no significant lag




Figure 10. The 0.2 – 12 keV *XMM-Newton* light curve binned in 200 s intervals.




Figure 11. The time lag as a function of Fourier frequency between the 0.4 – 1.5 keV and 2 – 4 keV bands. A negative lag would imply that the hard band leads the soft. No significant lag is detected in IES 1927+654.

between these two bands (see Fig 11) even when we combine the first two data points at the lowest frequencies.

Hardness ratios ($HR = H - S / H + S$; where H and S are the count rates in the hard and soft band, respectively) as a function of time were calculated between all the energy bands. A few examples are shown in Fig 12. The degree of variability depends on the energy bands being compared. The variations were most significant when the intermediate bands were compared to the softest bands (see the upper and middle panels of Fig 12).

The 0.5 – 10 keV FI CCD light curve is shown in Fig 13 with bins corresponding to orbital time scales (5760 s). The variability is very similar to that exhibited during the *XMM-Newton* observation.

The F_{var} in various energy bands is calculated following Edelson et al. (2002) and uncertainties are estimated following Ponti et




Figure 12. The hardness ratio plotted for various energy bands as a function of time. The light curves are binned in 1 ks intervals to improve statistics. The degree of variability differs depending on the energy bands being compared.




Figure 13. The 0.5 – 10 keV *Suzaku* light curve binned in orbital time intervals (5760 s).

al. (2004). The *XMM-Newton* light curves are in 500 s bins while the *Suzaku* light curves are in 5760 s bins. The F_{var} spectrum from the *XMM-Newton* and *Suzaku* observations are shown in Fig 14. The spectrum shows the amplitude of the variability peaks at intermediate energies resulting in the bell-shaped F_{var} that is commonly seen in unobscured Seyfert galaxies (e.g. Gallo et al. 2004; Ponti et al. 2010).




Figure 14. Upper panel: The fractional variability spectrum from the *XMM-Newton* observation calculated using 500 s binning of the light curves. The fractional variability expected from the the best-fitting spectral models assuming the fluctuations are dominated by the power law component are overplotted. Lower panel: The same as above, but for the *Suzaku* data using 5760 s binning of the light curves.

We considered if the blurred reflection, neutral partial covering, and ionized partial covering spectral models presented in Section 5 could describe the spectral variability seen in Fig 14. In all three cases we considered the possibility that the variability was caused by changing the normalization of the power law component alone. The models are overplotted on the F_{var} spectra in Fig 14. The ionized partial covering describes the variations very well. We note that in unabsorbed sources, the rapid variability is often attributed to variations in the power law component, and it is reasonable to expect that such variability is taking place even when line-of-sight absorption is present. However, variations in covering fraction and/or column densities have been observed, even on time scales of less than about 1 – 2 days (e.g. Turner et al. 2008; Risaliti et al. 2005). We return to this in Section 6.2 and consider variations in the covering fraction.

Varying the power law normalization in the blurred reflection model gives a reasonable approximation of the F_{var} spectrum. The model broadly describes the amplitude, peak, and general shape of the F_{var} spectrum, but there are clearly inaccuracies.

6.2 Flux-resolved spectroscopy

It stands to reason the variability model (i.e. power law varying in brightness) presented above is an oversimplification. For example, in the blurred reflection model, when varying the brightness of the primary emitter (whether by intrinsic fluctuations or via light bending, Miniutti & Fabian 2004), there should be variations in the dependent parameters of the reflector (e.g. ionisation and reflected flux).

To determine if more complicated (but more realistic) variability is required we created flux-resolved spectra of IES 1927+654 in low- ($< 6 \text{ count s}^{-1}$), medium- ($6 - 7 \text{ count s}^{-1}$), and high-flux ($> 7 \text{ count s}^{-1}$) states³. Fitting the three states with the average blurred reflection model, but allowing for changes in the power law normalization resulted in a good fit ($\chi^2_{\nu}/\text{dof} = 1.04/1421$). The fit was significantly improved ($\Delta\chi^2 = 83$ for 3 addition free parameters) when we allowed the ionization of the reflector to vary along with the power law. (Fig 15) The changes in the ionization are well correlated with the flux in the reflection component (Fig 16) as would be expected (e.g. Miniutti & Fabian 2004). The fits to the flux-resolved spectra demonstrate that in terms of the blurred reflection model the variability is more complicated than was assumed for Fig 14, and completely consistent with general AGN behaviour.

For the ionised partial covering model, we also considered if changes in the covering fraction alone (i.e. constant primary emitter) could describe the various flux states. The three spectra could be well fitted ($\chi^2_{\nu}/\text{dof} = 1.02/1419$; Fig 17) by allowing only the covering fraction of each absorber to vary. In this scenario, the covering fraction of the highly ionised absorber dropped from ~ 58 to ~ 42 per cent between the high and low state. The covering fraction of the colder absorber dropped significantly from ~ 40 to ~ 2 per cent between the high and low state.

7 DISCUSSION

These *XMM-Newton* and *Suzaku* data provide the first high signal-to-noise observations of IES 1927+654 above 2 keV. For the first time the existence of a soft-excess below about 1 keV has been confirmed, and an upper limit on a narrow Fe K α line and Compton hump have been established. The 0.3 – 10 keV spectra are well fitted with the standard traditional models, in particularly the blackbody plus power law model. The photon index ($\Gamma \approx 2.3$) and blackbody temperature ($kT \approx 170 \text{ eV}$) are both higher than the canonical values, but within observed ranges (e.g. Nandra & Pounds 1994; Crummy et al. 2005). There is a small level of cold absorption required ($5 - 10 \times 10^{20} \text{ cm}^{-2}$) for every model attempted, which could be associated with either the AGN or the host galaxy.

IES 1927+654 exhibits rapid and large amplitude flux and spectral variations. The bell-shape seen in the F_{var} spectrum resembles the typical curve seen in unabsorbed, type-1 Seyferts, in particularly those of narrow-line Seyfert 1 galaxies (e.g. Gallo et al. 2004; Ponti et al. 2010). The F_{var} spectra and flux-resolved spectra can be well described by either a blurred reflection model or by an ionized double partial covering model. In terms of blurred reflection the rapid variability could be described by changing brightness of the primary emitter and corresponding changes in the ionization

³ The average, observed 0.3 – 10 keV flux in each level from low-to-high is approximately 0.8, 1.1, and $1.5 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$




Figure 15. Top panel: The blurred reflection models used to describe the various flux states during the *XMM-Newton* observation. The differences can be described by changes in the ionisation parameter of the reflector as would arise from changes in the flux of the source illuminating the disc. Lower panel: The residuals remaining at each flux state based on the model described in the top panel.

of the reflector. Variability based on the ionized partial covering model can be described by changes in the covering fraction while the power law component remains constant. The variability seems to be in line with what is predicted by each model.

The partial covering models are pure absorption models in that they do not include Compton scattering (e.g. Miller & Turner 2013) or the fluorescent emission that accompanies photoelectric absorption (e.g. Reynolds et al. 2009). If the partial covering models could be described in this standard way, then a narrow Fe K α emission line is expected to accompany the absorber. The denser absorber ($N_H \approx 66 \times 10^{22} \text{ cm}^{-2}$) in the ionized partial covering model would remove about $1.8 \times 10^{-5} \text{ ph s}^{-1} \text{ cm}^{-2}$ ionizing photons (7.08 – 20 keV) from the *XMM-Newton* spectrum. Assuming the absorber is isotropically distributed around the source, then the strength of the iron line generated from photoelectric absorption is governed by the fluorescent yield. Adopting the fluorescent yield value for neutral iron (0.347; slightly higher for ionized iron) results in $6.4 \times 10^{-6} \text{ ph s}^{-1} \text{ cm}^{-2}$ line photons being created. Such a feature would have an $EW \approx 147 \text{ eV}$ in the *XMM-Newton* spectrum. A similar exercise for the *Suzaku* observation predicts a line with $EW \approx 101 \text{ eV}$. Such lines should be detected and are in-




Figure 16. The correlation between the ionisation parameter of the reflector and the 20–50 keV flux of the power law (top panel) and reflection (bottom panel) components derived from the blurred reflection model description for the *XMM-Newton* flux states (Fig 15). The correlation follows the expected behaviour that would occur as the flux illuminating the disc varies.

consistent with the upper-limit of 30 eV found for a narrow Fe $K\alpha$ feature in the data. We note that Yaqoob et al. (2010) argue that Compton-thick lines-of-sight could have much lower Fe $K\alpha$ fluxes (see also Miller et al. 2009), rendering our predicted fluxes overestimated.

The expected flux of the Compton bump in the *Suzaku* PIN band can also be estimated from the strength of the predicted iron line using `perxav`. The reflection fraction (R) is estimated from the equivalent width of the predicted iron line during the *Suzaku* observation: $R = EW/180 \text{ eV} = 0.56$. The 20 – 50 keV flux is estimated to be about $1.6 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$, more than double the flux in the pure absorption scenario. However, the source would still not be bright enough to be detected in the *Suzaku* PIN.

The ionized double partial covering model provides a good fit to the multi-epoch spectra and the rapid variability. The variability, both on short and long time scales, can be reasonably attributed to changes in covering fraction and a relatively constant X-ray source. However, the best-fit is achieved when the absorbers are outflowing at a velocity of $v \approx 0.3c$. We note the determination of this blueshift is not driven by any particular spectral feature, but rather by the broadband fit, specifically the soft excess. The ionised absorption model has been shown to replicate the soft excess in AGN quite well (e.g. Middleton, Done & Gierliński 2007). The shape of this soft excess in a sample of AGN is also well modelled with a black body with a temperature between $\sim 100 - 150 \text{ eV}$ (e.g. Crummy et al. 2006; Gierliński & Done 2004). The black-body temperature measured in IES 1927+654 was about 170 eV (see Table 2), which is consistent with an average blackbody spectrum seen in AGN (i.e. $kT \approx 130 \text{ eV}$) blueshifted by $\sim 0.3c$. The outflowing, ionised partial covering seems consistent with the high black body temperature measured in IES 1927+654.

If the wind is launched where v is the local escape velocity, then the wind must originate very close to the black hole ($r_{min} \sim$




Figure 17. Top panel: The ionised double partial covering model applied to the various flux states during the *XMM-Newton* observation. The differences can be attributed to changes in the covering fraction of a relatively constant primary emitter. Lower panel: The residuals remaining at each flux state based on the model described in the top panel.

$(c/v)^2 r_g \approx 10 r_g$). Following Fabian (2012), the kinetic luminosity of the wind, is $L_W = C_f/2(r_{min}/r_g)(v/c)N_H/N_T = 0.03L_{Edd}$, where $N_T = 1.5 \times 10^{24} \text{ cm}^{-2}$ and C_f and N_H are the covering fraction and column density of the denser absorber in Table 3. Based on the SED fitting in Section 4, the Eddington ratio is $L_{bol}/L_{Edd} \lesssim 0.1$, which would imply that the kinetic luminosity of the wind is about $\gtrsim 30$ per cent the bolometric luminosity of the AGN. Such values for L_W do not appear unreasonable for ultrafast outflows according to Tombesi et al. (2012). However, Tombesi et al. find the inner launching radius for such winds are on average > 10 -times farther from the black hole than r_{min} is in IES 1927+654.

The blurred reflection models fit the spectral data quite well. However we cannot distinguish between extremely relativistic models (e.g. rapidly spinning black hole and compact primary emitter) and less extreme ones. The Newtonian model (Blurred reflection (A) in Table 3) favours about a factor of two overabundance of iron, which is commonly seen in Seyfert X-ray spectra (e.g. Walton et al. 2013; Reynolds et al. 2012). The ionization parameter at both epochs is about $350 \text{ erg cm s}^{-1}$ and falls in the range where Auger effect is dominant over fluorescence rendering weak Fe $K\alpha$ features. In the relativistic model (Blurred reflection (B) in Table 3),

the iron abundance and ionisation parameters are both lower. However, the more intense blurring describes the absence of distinct features in the spectrum. If the soft excess is due to reflection then the inner disc must be present. The predicted 20 – 50 keV flux based on the reflection model is about $1.2 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ and would not be detectable in the *Suzaku* PIN. Unfortunately, the null PIN detection of IES 1927+654 alone does not allow us to eliminate any of the potential models.

Both reflection models predict a high inclination ($\sim 85^\circ$) consistent with an edge-on disc. As with the need for a high velocity outflow in the partial covering models, the preference for an edge-on disc is largely motivated by the “bluer” soft excess seen in IES 1927+654. Reflected emission will be more highly beamed in the disc plane thus the reflection component in the spectrum should appear shifted to higher energies. The high inclination would also be manifested in the blue wing of the Fe $K\alpha$ line, but since the fluorescent line is weak (either due to the Auger effect dominating or extreme blurring), the spectral feature is not significant.

This is intriguing given the Seyfert 2 nature of IES 1927+654 exhibited in the optical band (B03), and arguments that this is a true Sy2 based on optical and near-infrared observations (Panessa et al. in prep). At face-value this would indicate that IES 1927+654 is orientated like a Seyfert 2 in X-rays and optical, but void of the absorption that is associated with the torus. The small level of cold absorption ($\sim 10^{21} \text{ cm}^{-2}$) could be from a torus that is currently in some evolutionary state. This could be confirmed with mid-infrared data to measure the torus emission.

The works of Tran et al. (2011) and Wang et al. (2012) propose two different scenarios to explain the absence of the BLR in some AGN. Tran et al. predict such objects would be old systems and could be identified by very low L/L_{Edd} values, while Wang et al. suggests such objects are young and would exhibit high Eddington ratios. The SED measurements presented in this work show that IES 1927+654 may be a typical AGN with an Eddington ratio between 0.014 – 0.11. These values appear inconsistent with the predictions of both Tran et al. (2011) and Wang et al. (2012). Neglecting the effects of absorption momentarily, if the disc, and consequently the BLR, are seen edge-on as the blurred reflection models suggest, then the Doppler broadening of the BLR lines could be quite extreme. These very broad lines could be difficult to distinguish from the optical continuum. B03 measure the upper-limit on the flux of a broad $H\alpha$ component to be < 5 per cent that of the narrow component, but it is not clear what the width of this feature is.

8 CONCLUSION

IES 1927+654 is a perplexing object that challenges the standard AGN unification model exhibiting optical properties of absorbed systems and X-ray properties of unobscured systems. According to our SED measurements, IES 1927+654 exhibits a typical Eddington ratio and the absence of its BLR cannot be easily attributed to current models that call for very high or very low L/L_{Edd} . Based on X-ray spectral models we speculate that IES 1927+654 could be an edge-on system (i.e. like a Seyfert 2), with a standard accretion disc that extends to the inner regions, but viewed through a tenuous torus. Future optical and mid-infrared observations could test this hypothesis. Future observations with *NuSTAR* (Harrison et al. 2013) and *ASTRO-H* (Takahashi et al. 2012) would produce high-quality data above 10 keV that would distinguish between the two proposed x-ray models.

ACKNOWLEDGMENTS

The authors are grateful to the referee for valuable comments that improved the paper. LCG would like to thank Kirsten Bonson, Marcin Sawicki and Dave Thompson for discussions. The *XMM-Newton* project is an ESA Science Mission with instruments and contributions directly funded by ESA Member States and the USA (NASA). This research has made use of data obtained from the *Suzaku* satellite, a collaborative mission between the space agencies of Japan (JAXA) and the USA (NASA).

REFERENCES

Anders E., Grevesse N., 1989, *GeCoA*, 53, 197
 Arnaud K., 1996, in: *Astronomical Data Analysis Software and Systems*, Jacoby G., Barnes J., eds, ASP Conf. Series Vol. 101, p17
 Balestra I., Boller Th., Gallo L., Lutz D., Hess S., 2005, *A&A*, 442, 469
 Ballantyne D. R., Iwasawa K., Fabian A. C., 2001, *MNRAS*, 323, 506
 Bianchi S., Guainazzi M., Matt G., Chiaberge M., Iwasawa K., Fiore F., Maiolino R., 2005, *A&A*, 442, 185
 Bianchi S., Corral A., Panessa F., Barcons X., Matt G., Bassani L., Carrera F. J., Jiménez-Bailón, E., 2008, *MNRAS*, 385, 195
 Bianchi S. et al. 2012, *MNRAS*, 426, 3225
 Boller T., 2000, *New Astron. Rev.*, 44, 387
 Boller T., Voges W., Dennefeld M., Lehmann I., Predehl P., Burwitz V., Perlman E., Gallo L., Papadakis I.E., Anderson S., *A&A*, 397, 557 (B03)
 Crummy J., Fabian A., Gallo L., Ross R., 2006, *MNRAS*, 365, 1067
 De Marco B., Ponti G., Uttley P., Cappi M., Dadina M., Fabian A. C., Miniutti G., 2011, *MNRAS*, 417, 98
 De Marco B., Ponti G., Cappi M., Dadina M., Uttley P., Cackett E. M., Fabian A. C., Miniutti G., 2013, *MNRAS* (arXiv: 1201.0196)
 den Herder J. W. et al. 2001, *A&A*, 365, 7
 Edelson R., Turner T. J., Pounds K., Vaughan S., Markowitz A., Marshall H., Dobbie P., Warwick R., 2002, *ApJ*, 568, 610
 Emmanoulopoulos D., McHardy I. M., Papadakis I. E., 2011, *MNRAS*, 416, 94
 Fabian A. C., 1979, *RSPSA*, 366, 449
 Fabian A. C., Miniutti G., Gallo L., Boller Th., Tanaka Y., Vaughan S., Ross R. R., 2004, *MNRAS*, 353, 1071
 Fabian A. C. et al. , 2009, *Nature*, 459, 540
 Fabian A. C. et al. 2012, *MNRAS*, 424, 217
 Fabian A. C., 2012, *ARA&A*, 50, 455
 Gallo L. C., Tanaka Y., Boller Th., Fabian A. C., Vaughan S., Brandt W. N., 2004, *MNRAS*, 353, 1064
 Gallo L. C., Lehmann I., Pietsch W., Boller Th., Brinkmann W., Friedrich P., Grupe D., 2006, *MNRAS*, 365, 688
 Gierliński M. & Done C., 2004, *MNRAS*, 349, 7
 Grevesse N., Sauval A. J., 1998, *Space Science Reviews*, 85, 161
 Grupe D., et al. , 2008, *ApJ*, 681, 982
 Guainazzi M., 2010, *XMM-Newton Calibration Documents* (CAL-TN-0018)
 Harrison F. et al. 2013, submitted *ApJ* (arXiv: 1301.7307)
 Jansen F. et al. 2001, *A&A*, 365, L1
 Kalberla P. M. W., Burton W. B., Hartmann D., Arnal E. M., Bajaja E., Morras R., Pöppel W. G. L., 2005, *A&A*, 440, 775
 Mason K. O. et al. 2001, *A&A*, 365, 36
 Mateos S., Barcons X., Carrera F. J., Ceballos M. T., Hasinger G., Lehmann I., Fabian A. C., Streblyanska A., 2005, *A&A*, 444, 79
 Matt G., Guainazzi M., Maiolino R., 2003, *MNRAS*, 342, 422
 Middleton M., Done C., Gierliński M., 2007, *MNRAS*, 381, 1426
 Miller L., Turner T. J., Reeves J. N., 2009, *MNRAS*, 399, 69
 Miller L., Turner T. J., 2013, submitted to *ApJ* (arXiv: 1303.4309)
 Miniutti G., Fabian A. C., 2004, *MNRAS*, 349, 1435
 Mitsuda K. et al. 2007, *PASJ*, 59, 1
 Murphy K. D., Yaqoob T., 2009, *MNRAS*, 397, 1549

- Nandra K., Pounds K. A., 1994, MNRAS, 268, 405
Nicastro F., 2000, ApJL, 530, L65
Nicastro F., Martocchia A., Matt G., 2003, ApJL, 589, L13
Nowak M. A., Wilms J., Dove J. B., 1999, ApJ, 517, 355
Panessa F., Bassani L., 2002, A&A, 394, 435
Panessa F., Carrera F.J., Bianchi S., Corral A., Gastaldello F., Barcons X., Bassani L., Matt G., Monaco L., 2009, MNRAS, 398, 1951
Ponti G., Cappi M., Dadina M., Malaguti, G., 2004, A&A, 417, 451
Ponti G. et al. , 2010, MNRAS, 406, 2591
Reynolds C. S., Fabian A. C., Brenneman L. W., Miniutti G., Uttley P., Gallo L. C., 2009, MNRAS, 397, 21
Reynolds C. S., Brenneman L. W., Lohfink A. M., Trippe M. L., Miller J. M., Fabian A. C., Nowak M. A., 2012, ApJ, 755, 88
Risaliti G., Elvis M., Fabbiano G., Baldi A., Zezas A., 2005, ApJ, 623, 93
Ross R. R., Fabian A. C., 2005, MNRAS, 358, 211
Schlegel D., Finkbeiner D., Davis M., 1998, ApJ, 500, 525
Strüder L. et al. 2001, A&A, 365, L18
Takahashi T. et al. 2012, SPIE, 8443, 1
Tanaka Y., Boller T., Gallo L., Keil R., Ueda, Y., 2004, PASJ, 56, 9
Tanaka Y., Boller T., Gallo L., 2005, in Merloni A., Nayakshin S., Sunyaev R. A., eds, ESO Conf. Ser., Growing Black Holes: Accretion in a Cosmological Context. Springer-Verlag, Berlin, p. 290
Tombesi F., Cappi M., Reeves J. N., Braito V., 2012, MNRAS, 422, 1
Tran H.D., Lyke J.E., Mader J.A., 2011, ApJL, 726, L21
Trouille L., Barger A. J., Cowie L. L., Yang Y., Mushotzky R. F., 2009, ApJ, 703, 2160
Turner M. J. L. et al. 2001, A&A, 365, 27
Turner T. J., Reeves J., Kraemer S., Miller L., 2008, A&A, 483, 161
Walton D. J., Nardini E., Fabian A. C., Gallo L. C., Reis R. C., 2013, MNRAS, 428, 2901
Vasudevan R.V, Mushotzky R. F., Winter L. M., Fabian A. C., 2009, MNRAS, 399, 1553
Wang J.-M., Du P., Baldwin J.A., Ge J.-Q., Hu C., Ferland G.J., 2012, ApJ, 746, 137
Wilms J., Allen A., McCray R., 2000, ApJ, 542, 914
Wang J.-M., Zhang E.-P., 2007, ApJ, 660, 1072
Yaqoob T., Murphy K. D., Miller L., Turner T. J., 2010, MNRAS, 401, 411
Zoghbi A., Fabian A. C., Reynolds C. S., Cackett E. M., 2012, MNRAS, 422, 129

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.