DEFORMATION OF INVOLUTION AND MULTIPLICATION IN A $C^{*}\mbox{-}{\rm ALGEBRA}$

H. NAJAFI AND M. S. MOSLEHIAN

ABSTRACT. We investigate the deformation of involution and multiplication in a unital C^* -algebra when its norm is fixed. Our main result is to present all multiplications and involutions on a given C^* -algebra \mathcal{A} under which \mathcal{A} is still a C^* -algebra whereas we keep the norm unchanged. For each invertible element $a \in \mathcal{A}$ we also introduce an involution and a multiplication making \mathcal{A} into a C^* algebra in which a becomes a positive element. Further, we give a necessary and sufficient condition for that the center of a unital C^* -algebra \mathcal{A} is trivial.

1. INTRODUCTION

A C^* -algebra is a complex Banach *-algebra \mathcal{A} satisfying $||a^*a|| = ||a||^2$ $(a \in \mathcal{A})$. By the Gelfand-Naimark theorem, a C^* -algebra is a norm closed *-subalgebra of $\mathbb{B}(\mathcal{H})$ for some Hilbert space \mathcal{H} . A strongly closed *-subalgebra of $\mathbb{B}(\mathcal{H})$ containing the identity operator is called a von Neumann algebra. By the double commutant theorem a unital *-subalgebra \mathcal{A} of $\mathbb{B}(\mathcal{H})$ is a von Neumann algebra if and only if \mathcal{A} is equal to its double commutant \mathcal{A}^{cc} , where $\mathcal{A}^c = \{B \in \mathbb{B}(\mathcal{H}) : AB = BA$ for all $A \in \mathcal{A}\}$. By Sakai's characterization of von Neumann algebras, \mathcal{A} is a von Neumann algebra if and only if it is a W^* -algebra, i.e. it is a C^* -algebra being the norm dual of a Banach space \mathcal{A}_* . Throughout the paper \mathcal{A} denotes an arbitrary C^* -algebra and $\mathcal{Z}(\mathcal{A})$ stands for its center.

For a self adjoint element $a \in \mathcal{A}$, it holds that r(a) = ||a||, where r(a) denotes the spectral radius of \mathcal{A} . This implies that the norm of a C^* -algebra is unique when we fix the involution and the multiplication. Indeed, if \mathcal{A} is a C^* -algebra under two norms $||.||_1$ and $||.||_2$, then $||a||_1 = ||a^*a||_1^{\frac{1}{2}} = r(a^*a)^{\frac{1}{2}} = ||a^*a||_2^{\frac{1}{2}} = ||a||_2$ for all $a \in \mathcal{A}$. Bohnenblust and Karlin [BK] showed that there is at most one involution on a Banach algebra with the unit 1 making it into a C^* -algebra (see also [R]): Let * and # be two involutions on a unital Banach algebra \mathcal{A} making it into C^* -algebras.

²⁰¹⁰ Mathematics Subject Classification. Primary 46L05; Secondary 46L10.

Key words and phrases. C^* -algebra; von Neumann algebra; involution; multiplication; positive element, center, double commutant.

Let $x \in \mathcal{A}$. It follows from the fact "an element x of a unital C^* -algebra is selfadjoint if and only if $\tau(x)$ is real for every bounded linear functional τ on \mathcal{A} with $\| \tau \| = \tau(1) = 1$ ([KR 1, Proposition 4.3.3])" that x is self-adjoint with respect to * if and only if it is self-adjoint with respect to #. Now let $a \in \mathcal{A}$ be arbitrary and $a = a_1 + ia_2$ with self-adjoint parts a_1, a_2 with respect to *. Then $a_1^* = a_1^{\#}$ and $a_2^* = a_2^{\#}$ and $a^* = a_1 - ia_2 = a^{\#}$. There is another way to show the uniqueness of the involution. Indeed if * and # be two involutions on a unital Banach algebra \mathcal{A} making it into C^* -algebras, then the identity map from $(\mathcal{A}, *)$ onto $(\mathcal{A}, \#)$ is positive (see [P, Proposition 2.11]) and so $a^* = a^{\#}$ for all $a \in \mathcal{A}$.

There are several characterizations of C^* -algebras among involutive Banach algebras, see [DT] in which the authors start with a C^* -algebra and modify its structure. We however investigate a different problem in the same setting. In fact we investigate the deformation of involution and multiplication in a unital C^* -algebra when its norm is fixed. Our main result is to present all multiplications \circ and involutions \star on a given C^* -algebra \mathcal{A} under which \mathcal{A} is still a C^* -algebra whereas we keep the norm unchanged. As an application, for each invertible element $a \in \mathcal{A}$ we introduce an involution and a multiplication making \mathcal{A} into a C^* -algebra in which a becomes a positive element. Further, we give a necessary and sufficient condition for that the center of a unital C^* -algebra \mathcal{A} is trivial.

Recall that a Jordan *-homomorphism is a self-adjoint map preserving squares of self-adjoint operators. Jacobson and Rickart [JR] showed that for every Jordan *-homomorphism ρ of a C^* -algebra \mathcal{A} with the unit 1 into a von Neumann algebra \mathcal{B} there exist central projections $p_1, p_2 \in \mathcal{B}$ such that $\rho = \rho_1 + \rho_2$, $\rho(1) = p_1 + p_2$, $\rho_1(a) = \rho(a)p_1$ is a *-homomorphism and $\rho_2(a) = \rho(a)p_2$ is a *-antihomomorphism. Kadison [K] showed that an isometry of a unital C^* -algebra onto another C^* -algebra is a Jordan *-isomorphism.

2. Results

We start our work with the following lemma.

Lemma 2.1. Let \mathcal{A} be a unital C^* -algebra of operators acting on a Hilbert space \mathcal{H} . Let $p \in \mathcal{A}$ be a central projection and $u \in \mathcal{A}$ be a unitary. Let \circ be the multiplication and \star be the involution defined on \mathcal{A} by

$$a \circ b = paub + (1-p)bua$$
 and $a^* = u^*a^*u^*$ (2.1)

for $a, b \in A$, respectively. Then A equipped with the multiplication \circ and the involution \star is a unital C^* -algebra.

Proof. It is easy to check that \mathcal{A} is a complex Banach algebra under the multiplication \circ and u^* is the unit for this multiplication. By the decomposition

 $\mathcal{H} = p\mathcal{H} \oplus (1-p)\mathcal{H}, \text{ we can represent any element } a \in \mathcal{A} \text{ by the } 2 \times 2 \text{ matrix} \begin{pmatrix} pa & 0 \\ 0 & (1-p)a \end{pmatrix}. \text{ For } a, b \in \mathcal{A}, \text{ therefore } pa + (1-p)b \text{ can be identified by} \\ \begin{pmatrix} pa & 0 \\ 0 & (1-p)b \end{pmatrix}, \text{ whence } ||pa + (1-p)b|| = \max(||pa||, ||(1-p)b||). \text{ Hence}$

$$\begin{aligned} \|a^* \circ a\| &= \|p \ u^* a^* a + (1-p) \ a a^* u^* \| \\ &= \max(\|pa^*a\|, \|aa^*(1-p)\|) \\ &= \max(\|pa\|^2, \|(1-p)a\|^2) \\ &= \max(\|pa\|, \|(1-p)a\|)^2 = \|a\|^2 \end{aligned}$$

for all $a \in \mathcal{A}$.

The unital C^* -algebra \mathcal{A} equipped with the multiplication \circ and the involution \star is denoted by $\mathcal{A}(\circ, \star)$. Next we establish a converse of Lemma 2.1.

Theorem 2.2. Let \mathcal{A} be a unital C^* -algebra of operators acting on a Hilbert space \mathcal{H} and there exist a multiplication \circ and an involution \star on the normed space \mathcal{A} making it into a C^* -algebra. Then there exists a unitary element $u \in \mathcal{A}$ and a central projection p in the double commutant of \mathcal{A}^{cc} of \mathcal{A} such that both equalities (2.1) hold.

Proof. Since \mathcal{A} is unital, the closed unit ball of \mathcal{A} has an extreme point, hence the C^* algebra $\mathcal{A}(\circ, \star)$ is unital. Since $\iota(x) = x$ is an isometric linear map of \mathcal{A} onto $\mathcal{A}(\circ, \star)$, the unitary elements of $\mathcal{A}(\circ, \star)$ and those of \mathcal{A} coincide [KR 2, Exercise 7.6.17]. Thus if u^* is the unit of $\mathcal{A}(\circ, \star)$, then u is a unitary of \mathcal{A} . Define $\rho : \mathcal{A} \to \mathcal{A}(\circ, \star)$ by $\rho(a) = u^*a$. Clearly ρ is a unital isometric linear map of \mathcal{A} onto $\mathcal{A}(\circ, \star)$. Hence ρ is a positive map. This implies that $u^*a^* = \rho(a^*) = (u^*a)^*$ and so $a^* = u^*a^*u^*$. For determining the multiplication, define a multiplication \diamond on \mathcal{A}^{cc} (with respect to the original multiplication) by (2.1) as p = 1. Then \mathcal{A}^{cc} with the multiplication \diamond is a C^* -algebra. The space \mathcal{A}^{cc} as a Banach space is already the dual of a Banach space, so \mathcal{A}^{cc} with the new product and the new involution is a von Neumann algebra. Then the map $\rho(x) = x$ is a unital isometric linear map of $\mathcal{A}(\circ, \star)$ into the von Neumann algebra $\mathcal{A}^{cc}(\diamond, \star)$. By the result of Kadison [K] it is a Jordan *-isomorphism and by the Jacobson and Rickart theorem [JR] there exists a central projection p' in $\mathcal{A}^{cc}(\diamond, \star)$ such that $\rho_1(x) = p' \diamond \rho(x)$ is a *-homomorphism and

 $\rho_2(x) = (u^* - p') \diamond \rho(x)$ is a *-antihomomorphism. Therefore for each $a, b \in \mathcal{A}$ we

have

$$\begin{aligned} a \circ b &= \rho(a \circ b) \\ &= \rho_1(a \circ b) + \rho_2(a \circ b) \\ &= p' \diamond \rho_1(a) \diamond \rho_1(b) + (u^* - p') \diamond \rho_2(b) \diamond \rho_2(a) \\ &= p' \diamond a \diamond b + (u^* - p') \diamond b \diamond a. \\ &= p' uaub + (u^* - p')ubua \\ &= p' uaub + (1 - p'u)bua. \end{aligned}$$

Let p = p'u. Since $(p'u)^2 = p'up'u^2 = p' \diamond (p' \diamond 1) = p' \diamond 1 = p'u$ and $(p'u)^* = u^*p'^* = u^*p'^*u^*u = p'^*u = p'u$, so p is a projection in \mathcal{A}^{cc} . A similar argument shows that $\theta : \mathcal{A}^{cc}(\diamond, \star) \to \mathcal{A}^{cc}$ defined by $\theta(a) = au$ is a Jordan *-isomorphism. So, by [JR, Corollary 1], $\theta(\mathcal{Z}(\mathcal{A}^{cc}(\diamond, \star))) = \mathcal{Z}(\theta(\mathcal{A}^{cc}(\diamond, \star)))$. Therefore $pa = \theta(p')\theta(au^*) = \theta(au^*)\theta(p') = ap$ for each $a \in \mathcal{A}$. Hence p is a central projection in \mathcal{A}^{cc} .

Remark 2.3. Note that in general case, a C^* -algebra \mathcal{A} has many representations. However the proof of Theorem 2.2 shows that for any representation of \mathcal{A} , we can present all multiplications and involutions on \mathcal{A} which keep it still a C*-algebra with the same norm by a unitary and a central projection in the double commutant with respect to the same representation. Further, since p in Theorem 2.2 is in $\mathcal{A}^{cc} \subseteq \mathbb{B}(\mathcal{H})$, it depends on \mathcal{H} . If \mathcal{A} is a von Neumann algebra, then $p \in \mathcal{A}^{cc} = \mathcal{A}$.

Corollary 2.4. Let \mathcal{I} be an ideal of a von Neumann algebra \mathcal{A} . Then \mathcal{I} is also an ideal of the C^* -algebra $\mathcal{A}(\circ, \star)$ for any multiplication \circ and any involution \star .

Proof. It is sufficient to note that *paub* and (1-p)bua belong to \mathcal{I} when $a \in \mathcal{A}, b \in \mathcal{I}$ and so $a \circ b = paub + (1-p)bua \in \mathcal{I}$.

It is easy to see that $a \circ b = b \circ a$ if and only if aub = bua. We therefore have

Corollary 2.5. Suppose that \mathcal{A} is a unital C^* -algebra and the normed space \mathcal{A} is equipped with a multiplication \circ and an involution \star is a C^* -algebra with the unit u^* , where $u \in \mathcal{A}$ is a unitary. Then

(i) \mathcal{A} is commutative if and only if so is $\mathcal{A}(\circ, \star)$.

(ii) $\mathcal{Z}(\mathcal{A}) = \mathbb{C}1$ if and only if $\mathcal{Z}(\mathcal{A}(\circ, \star)) = \mathbb{C}u^*$.

Proof. (i) Let \mathcal{A} be commutative. By Theorem 2.2 there exist a unitary element $u \in \mathcal{A}$ and a central projection p in \mathcal{A}^{cc} such that

$$a \circ b = paub + (1-p)bua$$
 $(a, b \in \mathcal{A}).$

Hence

$$a \circ b = paub + (1-p)bua = pbua + (1-p)aub = b \circ a$$

Therefore $\mathcal{A}(\circ, \star)$ is commutative. Changing the role of \mathcal{A} by $\mathcal{A}(\circ, \star)$, we reach the reverse assertion.

(ii) Let $\mathcal{Z}(\mathcal{A}) = \mathbb{C}1$. If $a \in \mathcal{Z}(\mathcal{A}(\circ, \star))$, then for any $b \in \mathcal{A}$ we have $a \circ b = b \circ a$. As in the proof of Theorem 2.2 we observe that $\theta : \mathcal{A}^{cc}(\circ, \star) \to \mathcal{A}^{cc}$ defined by $\theta(a) = au$ is a Jordan *-isomorphism. Hence, by [JR, Corollary 1], $\theta(b)\theta(a) = \theta(a)\theta(b)$, so aubu = buau. Since each element of \mathcal{A} is of the form bu for some $b \in \mathcal{A}$, it follows that $au \in \mathcal{Z}(\mathcal{A})$. Hence $au = \lambda 1$ for some $\lambda \in \mathbb{C}$. Therefore $\mathcal{Z}(\mathcal{A}(\circ, \star)) = \mathbb{C}u^*$. Similarly we can deduce the converse.

Remark 2.6. The Arens product on $(c_0)^{**} = l^{\infty}$ coincide with the usual product in l^{∞} [D, Example 2.6.22]. It was extended to arbitrary C^* -algebras in [BD]. We reprove the fact in our own way: Let \mathcal{A} be a C^* -algebra and its second dual \mathcal{A}^{**} be also a C^* -algebra under a multiplication $(a, b) \mapsto a \cdot b$ whose restriction to $\mathcal{A} \times \mathcal{A}$ is the same multiplication of \mathcal{A} . We shall show that the Arens product (denoted by \diamond) on \mathcal{A}^{**} is the same as the multiplication \cdot on \mathcal{A}^{**} . It is known that \mathcal{A}^{**} is a von Neumann algebra under the Arens multiplication [D, Theorem 3.2.37]. By the Kaplansky density theorem, \mathcal{A} is dense in \mathcal{A}^{**} in the weak*-topology, so there exists a net u_{α} in \mathcal{A} such that $u_{\alpha} \to 1$ in the weak*-topology in which 1 denotes the unit of \mathcal{A}^{**} . So

$$b = w^* - \lim_{\alpha} u_{\alpha} b = w^* - \lim_{\alpha} u_{\alpha} \diamond b = 1 \diamond b$$

for each $b \in \mathcal{A}$. The Kaplansky density theorem implies that $1 \diamond x = x$ for each $x \in \mathcal{A}^{**}$. Therefore the units of both multiplications \cdot and \diamond are same. By Theorem 2.2 there exist a central projection $p \in \mathcal{A}$ such that

$$x \diamond y = pxy + (1-p)yx,$$

for each $x, y \in \mathcal{A}^{**}$. On the other hand for each $a, b \in \mathcal{A}$, we have $a \diamond b = ab$. So (1-p)ab = (1-p)ba. Since \mathcal{A} is dense in \mathcal{A}^{**} in the weak*-topology, we have (1-p)xy = (1-p)yx for each $x, y \in \mathcal{A}^{**}$. Therefore $x \diamond y = pxy + (1-p)yx = pxy + (1-p)xy = xy$ for each $x, y \in \mathcal{A}^{**}$. For instance, we deduce that the Arens product on $\mathbb{K}(\mathcal{H})^{**} = \mathbb{B}(\mathcal{H})$ is equal to the operator multiplication on $\mathbb{B}(\mathcal{H})$.

Theorem 2.7. Let \mathcal{A} be a unital C^* -algebra. Then the following assertions are equivalent:

(i) $\mathcal{Z}(\mathcal{A}) = \mathbb{C}1$

(ii) If for invertible operators $a, b \in \mathcal{A}$, ||axb|| = ||x|| holds for each $x \in \mathcal{A}$, then there exists $\lambda > 0$ such that both λa and $\frac{1}{\lambda}b$ are unitary. Proof. (i) \Rightarrow (ii) Note that if $||a^{-1}xa|| \leq ||x||$ for each $x \in \mathcal{A}$, then map $\varphi(x) = a^{-1}xa$ is a contractive unital linear map on \mathcal{A} . It follows from [P, Proposition 2.11] that φ is positive. Therefore $(a^{-1}xa)^* = a^{-1}x^*a$ and so $aa^*x^* = x^*aa^*$ for each $x \in \mathcal{A}$. Hence $aa^* \in \mathcal{Z}(\mathcal{A}) = \mathbb{C}1$. So $a^*a = \lambda 1$ for some $\lambda > 0$. Therefore $\frac{1}{\sqrt{\lambda}}a$ is unitary. First, assume that ||axb|| = ||x|| for positive invertible operators a, b and each $x \in \mathcal{A}$. Then $||b^{-1}a^{-1}|| = ||a^{-1}b^{-1}|| = ||aa^{-1}b^{-1}b|| = 1$, whence

$$||a^{-1}xa|| \le ||axb|| \ ||b^{-1}a^{-1}|| \le ||x||.$$

Therefore there exists $\lambda > 0$ such that $\frac{1}{\lambda}a$ is unitary. Since $\frac{1}{\lambda}a$ is positive and unitary we have $a = \lambda$. A similar argument shows that $b = \lambda'$. It follows from $1 = ||1|| = ||ab|| = \lambda'\lambda$ that $\lambda = \frac{1}{\lambda'}$. Second, assume that ||axb|| = ||x|| for invertible operators a, b and each $x \in \mathcal{A}$. Utilizing the polar decompositions of aand b^* , there exist unitary operators u, v such that a = u|a| and $b = |b^*|v$. Hence $|||a||x||b^*||| = ||u||a||x||b^*||v|| = ||axb|| = ||x||$ for each $x \in \mathcal{A}$. The above argument shows that $|a| = \lambda$ and $|b^*| = \frac{1}{\lambda}$ for some $\lambda > 0$, so $a = \lambda u$ and $b = \frac{1}{\lambda}v$.

(ii) \Rightarrow (i) Note that each central invertible element a of \mathcal{A} is a scalar multiple of a unitary element. In fact, we have $||a^{-1}xa|| = ||a^{-1}ax|| = ||x||$ for all $x \in \mathcal{A}$, so λa is unitary for some $\lambda > 0$. Let $a \in \mathcal{Z}(\mathcal{A})$ be a positive element and $\lambda_1, \lambda_2 \in \operatorname{sp}(a)$ are distinct. Then there exists an invertible continuous function f on $\operatorname{sp}(a)$ such that $f(\lambda_1) = \frac{1}{2}$ and $f(\lambda_2) = 1$. Hence f(a), which is a central invertible element should be a scalar multiple of a unitary. On the other hand, $\frac{1}{2}, 1 \in \operatorname{sp}(f(a))$, which is impossible. Hence the spectrum of a is singleton, so a = ||a||. Since $\mathcal{Z}(\mathcal{A})$ is a C^* -algebra, any one of its elements is a linear combination of four positive elements. Therefore $\mathcal{Z}(\mathcal{A}) = \mathbb{C}1$.

Let $\mathcal{A}(u, p)$ denote the C^* -algebra given via Lemma 2.1 corresponding to a unitary u and a central projection p in \mathcal{A} . The self-adjoint elements of $\mathcal{A}(u, p)$ are all elements a such that $au = u^*a^*$, a fact which is independent of the choice of p. Also a self-adjoint element a is positive in $\mathcal{A}(u, p)$ if and only if $a = b \circ b = pbub + (1-p)bub = bub$ for some self-adjoint element $b \in \mathcal{A}(u, p)$ and this occurs if and only if a is positive in $\mathcal{A}(u, 1)$.

Theorem 2.8. Let \mathcal{A} be a C^* -algebra and $a \in \mathcal{A}$ be invertible. Then there exists a unique unitary $u \in \mathcal{A}$ such that a is a positive element of the C^* -algebra $\mathcal{A}(u^*, p)$ for any central projection $p \in \mathcal{A}$.

Proof. Let a = u|a| be the polar decomposition of a. Then $u = a|a|^{-1} \in \mathcal{A}$. So $a = u|a|^{\frac{1}{2}}|a|^{\frac{1}{2}} = |a|^{\frac{1}{2}*} \circ |a|^{\frac{1}{2}}$, where \circ is defined in $\mathcal{A}(u^*, 1)$ by (2.1). So a is positive in $\mathcal{A}(u^*, p)$ for every central projection $p \in \mathcal{A}$. To see the uniqueness, note that if a is invertible and a, wa are positive for a unitary w, then $a = w^*(wa)$. By the

uniqueness of polar decomposition, we have w = 1. Now if a is positive in $\mathcal{A}(v^*, 1)$, then $a = b^* \circ b = vb^*b$. Hence $v^*u|a| = v^*a = b^*b$ is positive. Therefore $v^*u|a|$ and |a| are positive and so v = u according to what we just proved.

Remark 2.9. The invertibility condition in Proposition 2.8 is essential. For example let $\mathcal{A} = C[-1, 1]$ and f(t) = t. If f is positive in C[-1, 1](u, 1) for a unitary function u, then there exist $g \in C[-1, 1]$ such that $t = f(t) = u(t)|g(t)|^2$ for each $t \in [-1, 1]$. So u(t) = 1 for each $t \in (0, 1]$ and u(t) = -1 for each $t \in [-1, 0)$, which is impossible.

Acknowledgements. The authors would like to sincerely thank Professor Marcel de Jeu and Professor Jun Tomiyama for their valuable comments.

References

- [BK] H.F. Bohnenblust and S. Karlin, Geometrical properties of the unit sphere of Banach algebras, Ann. of Math. (2) 62, (1955), 217-229.
- [BD] F.F. Bonsall and J. Duncan, Complete Normed Algebras, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 80. Springer-Verlag, New York-Heidelberg, 1973.
- [D] H.G. Dales, Banach Algebras and Automatic Continuity, London Math. Soc. Monogr. Ser., vol. 24, Clarendon Press, Oxford, 2000.
- [DT] M. de Jeu and J. Tomiyama, A characterisation of C*-algebras through positivity of functionals, Ann. Funct. Anal. 4 (2013), no. 1, 61–63.
- [JR] N. Jacobson and C. Rickart, Homorphisms of Jordan rings, Trans. Amer. Math. Soc. 69 (1950), 479–502.
- [K] R.V. Kadison, Isometries of operator algebras, Ann. Math. 54 (1951), 325–338.
- [KR 1] R.V. Kadison and J.R. Ringrose, Fundamentals of the Theory of Operator Algebras, Vol. I. Elementary theory. Pure and Applied Mathematics, 100. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1983.
- [KR 2] R.V. Kadison and J.R. Ringrose, Fundamentals of the Theory of Operator Algebras, Volume II. Advanced theory. Pure and Applied Mathematics, 100. Academic Press, Inc., Orlando, FL, 1986.
- [P] V.I. Paulsen, Completely Bounded Maps and Operator Algebras, Cambridge Studies in Advanced Mathematics 78, Cambridge University Press, Cambridge, 2002.
- [R] C.E. Rickart, General Theory of Banach Algebras The University Series in Higher Mathematics D. van Nostrand Co., Inc., Princeton, N.J.-Toronto-London-New York, 1960.

DEPARTMENT OF PURE MATHEMATICS, CENTER OF EXCELLENCE IN ANALYSIS ON ALGE-BRAIC STRUCTURES (CEAAS), FERDOWSI UNIVERSITY OF MASHHAD, P. O. BOX 1159, MASH-HAD 91775, IRAN

E-mail address: hamednajafi20@gmail.com

E-mail address: moslehian@um.ac.ir and moslehian@member.ams.org