Sodium content as a predictor of the advanced evolution of globular cluster stars

Simon W. Campbell¹, Valentina D'Orazi^{2,1}, David Yong³, Thomas N. Constantino¹, John C. Lattanzio¹, Richard J. Stancliffe^{3,4}, George C. Angelou¹, Elizabeth C. Wylie-de Boer³, Frank Grundahl⁵

¹Monash Centre for Astrophysics, School of Mathematical Sciences, Monash University, Melbourne, VIC 3800, Australia.

²Department of Physics & Astronomy, Macquarie University, Balaclava Rd, North Ryde, Sydney, NSW 2109, Australia.

³Research School of Astronomy and Astrophysics, Australian National University, Weston, ACT 2611, Australia

⁴Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, 53121 Bonn, Germany. ⁵Stellar Astrophysics Centre, Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, DK-8000 Aarhus C, Denmark.

The asymptotic giant branch (AGB) phase is the final stage of nuclear burning for low mass stars. Although Milky Way globular clusters are now known to harbour (at least) two generations of stars^{1,2} they still provide relatively homogeneous samples of stars which are used to constrain stellar evolution theory^{3–5}. It is expected, and predicted by stellar models, that the majority of cluster stars with masses around the current turn-off mass will evolve through the AGB phase^{6,7}. However this has never been confirmed observationally. Here we show that all of the second generation stars in the globular cluster NGC 6752 – 70 % of the cluster population – fail to reach the AGB phase. Through spectroscopic abundance measurements we found that every AGB star in our sample has a low sodium abundance, with [Na/Fe] $\lesssim 0.18$ dex, indicating they are exclusively first generation stars. The extremely high AGB failure rate implies that many clusters cannot be reliably used for star counts to test stellar evolution timescales if the AGB population is involved. A thorough investigation is required to reconcile stellar theory with this observed phenomenon.

We obtained high resolution spectra ($R \sim 24,000$) for a sample of 20 AGB stars and 24 red giant branch stars in the Galactic globular cluster (GC) NGC 6752. The spectral coverage included the strong Na I doublet at 5680 Å. In Figure 1 we show the stellar sample. We include red giant branch stars as a control group, since it has previously been shown that this evolutionary population harbours the standard abundance distributions, including the well-known Na-O anticorrelation present in all GCs⁸.

Our sodium abundance results are shown in Figure 2. The red giant branch sample shows the usual spread in [Na/Fe] = $\log_{10}(N_{Na}/N_{Fe})_{star} - \log_{10}(N_{Na}/N_{Fe})_{Sun}$ of roughly 1 dex $(N_x$ is the number density of atoms of each elemental species). On the other hand the AGB result is very striking – every single one of the AGB stars in our sample lies at the low end of the red giant




Figure 1: Sample selection in the Strömgren uvby colour-magnitude diagram of NGC 6752. Small black dots show the whole photometric sample⁹. Our AGB and red giant branch (RGB) stellar samples are shown as blue squares and red triangles respectively. Part of the horizontal branch can be seen at bottom left, at y magnitudes ≥ 13.5 .




Figure 2: Sodium abundance results for NGC 6752. [Na/Fe] for our sample of red giant branch stars (red triangles, 24 stars) and AGB stars (blue squares, 20 stars) shown against stellar effective temperature T_{eff} (also see Supplementary Table S1). For comparison the red giant branch results of a previous study $(C07)^8$ are included (grey open circles). The horizontal dotted line at [Na/Fe] = 0.18 marks the upper envelope of AGB values, and divides Na-rich and Na-poor stars. The spectroscopic observations were carried out with the Very Large Telescope under European Southern Observatory programme 089.D-0038 (principal investigator SWC). The FLAMES/Giraffe HR11 grating was used, with a spectral coverage $\lambda = 5597 \rightarrow 5840$ Å. Na abundances (assuming local thermodynamic equilibrium) were obtained from the strong Na I doublet at 5680 Å, with the driver *abfind* in MOOG¹⁰ (2011 version) and the Kurucz model atmospheres with no overshooting¹¹. Stellar parameters were derived in the following way: T_{eff} values were calculated from a Strömgren colour (b-y) calibration¹²; gravities were then computed from stellar luminosities and the derived temperatures (with assumed stellar mass $M = 0.8 M_{\odot}$ and distance modulus of $(m - M)_V = 13.30$, while microturbulence values ξ were obtained from the literature¹³. A metallicity of $[Fe/H] = -1.54 \text{ dex}^8$ was adopted for all stars. Although the lines under scrutiny are known to be not largely affected by departures from local thermodynamic equilibrium (at most 0.15 dex), we applied non local thermodynamic equilibrium corrections¹⁴ to our Na abundances. Error bars show the random (internal) uncertainties (also see Supplementary Table S1). The uncertainties were estimated by adding, in quadrature, errors due to the equivalent width measurements and those related to stellar parameters. The latter were evaluated in the standard way, by varying one parameter at a time and inspecting the corresponding variation in the resulting abundances. We adopted errors of $\Delta T_{eff} = \pm 30$ K, $\Delta \log g = 0.1$, $\Delta \xi = 0.1$ km/s, and Δ [Fe/H] $= \pm 0.05 \, \text{dex}.$

branch distribution. The upper envelope of the AGB sodium abundances is located at about [Na/Fe] = 0.18 dex. Interestingly this corresponds very closely to a previous red giant branch study that defines the Na-poor, first generation population as having [Na/Fe] ≤ 0.2 dex (their 'Primordial' population)¹. We find the proportions of Na-poor to Na-rich red giant branch stars in our data to be 30:70. This also corresponds well to the roughly 30:70 proportions found previously¹. Thus, surprisingly, *all* of our AGB stars appear to be first generation stars, giving a first generation to second generation ratio change from 30:70 in the red giant branch population to 100:0 in the AGB population. The range in [Na/Fe] in our AGB sample is very small, with a mean of -0.07 dex and a standard deviation of 0.10 dex. This scatter is comparable to our internal uncertainties (Fig. 2 and Supplementary Table S1), which suggests that the AGB stars may have a uniform abundance of Na.

Our results indicate that the entire population of second generation stars, having elevated levels of Na, must fail to enter the AGB phase. This is a very significant effect since the second generation population contains the majority of the stars in NGC 6752 (70%). Although two studies have theorised that some AGB stars may not ascend the AGB in NGC 6752¹⁵, including the study by our group¹⁶, this has been based on low-resolution cyanogen band strength observations. It is known that these observations are affected by many uncertainties¹⁷, including the in situ variation of C and N along the red giant branch¹⁸. Measurement of elemental Na is more robust since Na is not affected by molecular band formation uncertainties and stars of these masses (~ 0.8 M_o) cannot alter their Na abundances in situ. In particular a reduction in surface Na would require very high temperatures at the base of the convective envelope. This is only achieved in much more massive stars ($\gtrsim 4 M_{o}$), via 'Hot Bottom Burning' nucleosynthesis¹⁹. Thus the result presented here is the first conclusive confirmation that stars of certain chemical composition do not ascend the AGB. Moreover, we can readily identify which stars avoid the AGB based on their Na content.

An obvious consequence of such a large proportion of stars avoiding the AGB is that there should currently be many fewer stars in the AGB phase than expected. A detailed study reporting star counts of GC populations finds a value of $R_2 = N_{AGB}/N_{HB} \sim 0.06$ for NGC 6752²⁰. This is one of the lowest R_2 values in their GC sample. The GCs with the two highest R_2 values in their sample (M 5 and M 55) could be assumed to provide an upper limit to R_2 since R_2 is fairly insensitive to metallicity, He abundance, and GC age²¹. Interestingly this upper value is roughly 0.18, i.e. a factor of 3 higher than that of NGC 6752. This is indeed consistent with our result of $\gtrsim 70\%$ of stars not ascending the AGB. Current model predictions for R_2 tend to be lower than 0.18, being around $0.12 \rightarrow 0.15^{21,22}$, however the models are known to suffer from significant uncertainties²². We note that the observed R_2 value for NGC 6752 is still at least a factor of 2 smaller than the model predictions.

Studies of the stage of evolution directly preceding the AGB, the horizontal branch phase, can shed light on the AGB avoidance phenomenon. One recent investigation into the Na abundances in a sample of horizontal branch stars in NGC 6752 showed that the redder end of the horizontal branch (NGC 6752 only has a blue horizontal branch) contains only Na-poor stars²³




Figure 3: Theoretical stellar model tracks overlain on the Strömgren colour-magnitude diagram of NGC 6752. The solid red line (with open circle symbols marking 5 Myr time intervals) is a model with an initial mass of 0.80 M_{\odot} and a helium content of Y=0.245. This Y value matches that reported for the redder end of the horizontal branch²³. This first generation model does indeed spend most of its horizontal branch evolution at the redder end of the horizontal branch. The solid black line (with open square symbols marking 5 Myr time intervals) is a model with an initial mass of 0.75 M_{\odot} and an enhanced helium content of Y = 0.285. This second generation model spends its horizontal branch evolution in a bluer part of the horizontal branch, but still ascends the AGB, contrary to the observational findings of the current study. The solid blue line (with open triangle symbols marking 5 Myr time intervals) shows the evolution of the Y = 0.285 model with an ad-hoc 20-fold increase in mass loss rate (M = dM/dt) initiated once the star settles on the horizontal branch. This model evolves downwards along the extreme blue end of the horizontal branch and fails to ascend the AGB. The arrow indicates the location of the Grundahl jump at y = 14.65 (see text for details). The stellar models were calculated using the Monash University stellar structure code MONSTAR²⁴. The code has been recently updated with low temperature opacity tables which follow variations in C, N and O²⁵. The usual Reimers mass loss rate²⁶ was used for the red giant branch and horizontal branch (with $\eta = 0.48$ for the models with normal mass-loss). Transforms from theoretical luminosity- T_{eff} plane to the colour-magnitude diagram have been made²⁷. 5

(see Supplementary Information section S2 for more discussion). This implies that it is the bluer (presumably Na-rich) horizontal branch stars that must avoid AGB ascent, leaving only the redder, Na-poor horizontal branch stars to populate the AGB of NGC 6752. Combining this information with our ratio of Na-poor to Na-rich stars (Figure 2) we can estimate the horizontal branch colour at which stars fail to ascend the AGB. Interestingly we find an 'ascension cut-off' colour that coincides exactly with the colour for the Grundahl Jump. The Grundahl Jump is a discontinuity in horizontal branch morphology which is seen in all GCs studied to date whose horizontal branch extends beyond $T_{eff} \approx 11,500 \text{ K}^{9,28}$. Thus it appears that all stars bluer than the Grundahl jump do not ascend the AGB, at least in NGC 6752. This may represent further evidence that there is some fundamental change in the stellar atmosphere structure and/or mass-loss physics occurring at the Grundahl jump temperature⁹.

We have calculated some stellar models to compare with the photometric observations. The model results are shown in Figure 3. Our first generation model populates the redder end of the horizontal branch, before the Grundahl jump, as expected. It then continues to the AGB. The He-rich second generation model (presumed to correspond to the Na-rich population) populates the bluer end of the horizontal branch (after the Grundahl jump), and also continues to the AGB. Thus our second generation model cannot account for the lack of ascension of the Na-rich blue horizontal branch stars in this part of the colour-magnitude diagram. We note that an increased mass loss rate during the red giant branch phase would result in a bluer zero-age horizontal branch star (see Supplementary Information section S3 for more detail), so this can not be a solution since the colour-magnitude diagram is clearly populated in this region. We speculate that one solution to this problem may be that the horizontal branch stars blueward of the Grundahl jump experience enhanced mass loss. To test this we artificially increased the mass loss rate in the second generation model during its horizontal branch phase by an ad-hoc factor of 20 (Fig. 33). Indeed this model can populate the blue end of the horizontal branch and also fail to become an AGB star. We note however that this test is purely hypothetical and, although this result may provide a starting point, a thorough investigation into the reasons behind the discordance between theory and observation is sorely needed. There is currently no clear explanation for such a high proportion of failed AGB stars.

Finally, since GCs are often used to test stellar evolution theory, the extremely high AGB failure rate reported here will affect any test which uses star counts of AGB stars. This is true of the R-Method used to check the lifetimes of various phases of evolution. In particular the R', R1 and R2 values^{3,4,20} all involve the number of AGB stars, so these values will be flawed (including the GC He values inferred from them). This is particularly important if the GCs in question have blue extensions to their horizontal branches, since it is the blue horizontal branch stars that appear not to ascend the AGB. Star number counts used to ascertain AGB lifetimes will also be misleading, unless the proportion of AGB ascenders is known somehow, for example via an ascension cut-off in Na abundance or horizontal branch colour.

References

- 1. Carretta, E., Bragaglia, A., Gratton, R. & Lucatello, S. Na-O anticorrelation and HB. VIII. Proton-capture elements and metallicities in 17 globular clusters from UVES spectra. *Astron. Astrophys.* **505**, 139–155 (2009).
- 2. Gratton, R. G., Carretta, E. & Bragaglia, A. Multiple populations in globular clusters. Lessons learned from the Milky Way globular clusters. *Astron. Astrophys. Rev.* **20**, 50 (2012).
- Iben, I. & Rood, R. T. Ratio of Horizontal Branch Stars to Red Giant Stars in Globular Clusters. *Nature* 224, 1006–1008 (1969).
- Buonanno, R., Corsi, C. E. & Fusi Pecci, F. The giant, asymptotic, and horizontal branches of globular clusters. II - Photographic photometry of the metal-poor clusters M15, M92, and NGC 5466. *Astron. Astrophys.* 145, 97–117 (1985).
- 5. Renzini, A. & Fusi Pecci, F. Tests of evolutionary sequences using color-magnitude diagrams of globular clusters. *Ann. Rev. Astron. Astrophys.* **26**, 199–244 (1988).
- 6. Kippenhahn, R. & Weigert, A. *Stellar Structure and Evolution* (Springer-Verlag, Berlin, 1990).
- Landsman, W. B. *et al.* Ultraviolet Imagery of NGC 6752: A Test of Extreme Horizontal Branch Models. *Astrophys. J.* 472, L93–L96 (1996).
- Carretta, E., Bragaglia, A., Gratton, R. G., Lucatello, S. & Momany, Y. Na-O anticorrelation and horizontal branches. II. The Na-O anticorrelation in the globular cluster NGC 6752. *Astron. Astrophys.* 464, 927–937 (2007).
- Grundahl, F., Catelan, M., Landsman, W. B., Stetson, P. B. & Andersen, M. I. Hot Horizontal-Branch Stars: The Ubiquitous Nature of the "Jump" in Strömgren u, Low Gravities, and the Role of Radiative Levitation of Metals. *Astrophys. J.* 524, 242–261 (1999).
- 10. Sneden, C. A. *Carbon and Nitrogen Abundances in Metal-Poor Stars*. Ph.D. thesis, The University of Texas at Austin. (1973).
- 11. Kurucz, R. ATLAS9 Stellar Atmosphere Programs and 2 km/s grid. *CD-ROM No. 13., Smithsonian Astrophysical Observatory, Cambridge, Mass.* **13** (1993).
- Alonso, A., Arribas, S. & Martínez-Roger, C. Erratum: The effective temperature scale of giant stars (F0-K5). II. Empirical calibration of T_{eff} versus colours and [Fe/H]. *Astron. Astrophys.* **376**, 1039–1039 (2001).
- 13. Gratton, R. G., Carretta, E. & Castelli, F. Abundances of light elements in metal-poor stars. I. Atmospheric parameters and a new T_{eff} scale. *Astron. Astrophys.* **314**, 191–203 (1996).
- Gratton, R. G., Carretta, E., Eriksson, K. & Gustafsson, B. Abundances of light elements in metal-poor stars. II. Non-LTE abundance corrections. *Astron. Astrophys.* 350, 955–969 (1999).

- 15. Norris, J., Cottrell, P. L., Freeman, K. C. & Da Costa, G. S. The abundance spread in the giants of NGC 6752. *Astrophys. J.* **244**, 205–220 (1981).
- 16. Campbell, S. W. *et al.* The Case of the Disappearing CN-strong AGB stars in Galactic Globular Clusters Preliminary Results. *Mem. Soc. Astron. Ital.* **81**, 1004 (2010).
- 17. Yong, D., Grundahl, F., Johnson, J. A. & Asplund, M. Nitrogen Abundances in Giant Stars of the Globular Cluster NGC 6752. *Astrophys. J.* **684**, 1159–1169 (2008).
- 18. Smith, G. H. & Tout, C. A. The production of surface carbon depletions among globular cluster giants by interior mixing. *Mon. Not. R. Astron. Soc.* **256**, 449–456 (1992).
- 19. Boothroyd, A. I., Sackmann, I.-J. & Ahern, S. C. Prevention of High-Luminosity Carbon Stars by Hot Bottom Burning. *Astrophys. J.* **416**, 762 (1993).
- 20. Sandquist, E. L. & Bolte, M. Exploring the Upper Red Giant and Asymptotic Giant Branches: The Globular Cluster M5. *Astrophys. J.* **611**, 323–337 (2004).
- Cassisi, S., Salaris, M. & Irwin, A. W. The Initial Helium Content of Galactic Globular Cluster Stars from the R-Parameter: Comparison with the Cosmic Microwave Background Constraint. *Astrophys. J.* 588, 862–870 (2003).
- 22. Cassisi, S., Castellani, V., Degl'Innocenti, S., Piotto, G. & Salaris, M. Asymptotic Giant Branch predictions: Theoretical uncertainties. *Astron. Astrophys.* **366**, 578–584 (2001).
- 23. Villanova, S., Piotto, G. & Gratton, R. G. The helium content of globular clusters: light element abundance correlations and HB morphology. I. NGC 6752. *Astron. Astrophys.* **499**, 755–763 (2009).
- 24. Campbell, S. W. & Lattanzio, J. C. Evolution and nucleosynthesis of extremely metal-poor and metal-free low- and intermediate-mass stars. I. Stellar yield tables and the CEMPs. *Astron. Astrophys.* **490**, 769–776 (2008).
- 25. Marigo, P. & Aringer, B. Low-temperature gas opacity. ÆSOPUS: a versatile and quick computational tool. *Astron. Astrophys.* **508**, 1539–1569 (2009).
- 26. Reimers, D. Circumstellar absorption lines and mass loss from red giants. *Memoires of the Societe Royale des Sciences de Liege* **8**, 369–382 (1975).
- 27. Clem, J. L., VandenBerg, D. A., Grundahl, F. & Bell, R. A. Empirically Constrained Color-Temperature Relations. II. uvby. *Astron. J.* **127**, 1227–1256 (2004).
- 28. Momany, Y. *et al.* A New Feature along the Extended Blue Horizontal Branch of NGC 6752. *Astrophys. J.* **576**, L65–L68 (2002).
- 29. Marino, A. F. *et al.* Sodium-Oxygen Anticorrelation Among Horizontal Branch Stars in the Globular Cluster M4. *Astrophys. J.* **730**, L16–L21 (2011).

- D'Antona, F., Caloi, V., Montalbán, J., Ventura, P. & Gratton, R. Helium variation due to selfpollution among Globular Cluster stars. Consequences on the horizontal branch morphology. *Astron. Astrophys.* 395, 69–75 (2002).
- 31. Greggio, L. & Renzini, A. Clues on the hot star content and the ultraviolet output of elliptical galaxies. *Astrophys. J.* **364**, 35–64 (1990).
- 32. Dalessandro, E. *et al.* Ultraviolet Properties of Galactic Globular Clusters with Galex. II. Integrated Colors. *Astron. J.* **144**, 126 (2012).
- 33. D'Cruz, N. L., Dorman, B., Rood, R. T. & O'Connell, R. W. The Origin of Extreme Horizontal Branch Stars. *Astrophys. J.* **466**, 359–371 (1996).
- 34. Milone, A. P. *et al.* A WFC3/HST view of the three stellar populations in the Globular Cluster NGC6752. *ArXiv e-prints* (2013). 1301.7044.

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Acknowledgements We thank Dr. Yazan Momany of the European Southern Observatory (ESO, Chile) for providing his *UBV* photometric dataset which is mentioned in Supplementary Information section S2. S.W.C. acknowledges support from the Australian Research Council's Discovery Projects funding scheme (project DP1095368). RJS is the recipient of a Sofja Kovalevskaja Award from the Alexander von Humboldt Foundation. F.G. acknowledges funding for the Stellar Astrophysics Centre provided by The Danish National Research Foundation. The research is supported by the ASTERISK project funded by the European Research Council (Grant agreement no.: 267864). This work was based on observations made with ESO telescopes at the La Silla Paranal Observatory under programme ID 089.D-0038 and made extensive use of the SIMBAD, Vizier, 2MASS, and NASA ADS databases.

Author Contributions S.W.C. designed and prepared the ESO Very Large Telescope (VLT) observing proposal, collected the spectroscopic data, and prepared the paper. V.D. reduced and analysed the spectroscopic data, and prepared the paper. D.Y. designed and prepared the ESO/VLT observing proposal and assisted in the paper preparation. T.N.C. calculated the stellar models and prepared figures for the paper. J.C.L. assisted in the preparation of the observing proposal and with the paper preparation. R.J.S., G.C.A. and E.C.W. assisted in the paper preparation and made preliminary observations with the Anglo-Australian Telescope. F.G. provided the *uvby* photometric data for the AGB and red giant branch sample and assisted in the paper preparation.

Author Information Reprints and permissions information is available at www.nature.com/reprints. Correspondence and requests for materials should be addressed to simon.campbell@monash.edu.

Supplementary Information

Table S1: The stellar sample including atmospheric parameters and Na abundances.

Туре	ID	T_{eff}	log (g)	ξ	$\log(N_{Na})$	[Na/Fe]	Error
AGB	FGJ000022	4607	1.414	1.765	4.85	0.06	0.09
AGB	FGJ000025	4371	1.146	1.851	4.71	-0.08	0.09
AGB	FGJ000031	4460	1.285	1.806	4.73	-0.06	0.09
AGB	FGJ000044	4629	1.537	1.725	4.63	-0.16	0.07
AGB	FGJ000052	4787	1.740	1.660	4.74	-0.05	0.09
AGB	FGJ000053	4688	1.636	1.693	4.73	-0.06	0.10
AGB	FGJ000059	4772	1.719	1.666	4.82	0.03	0.06
AGB	FGJ000060	4685	1.654	1.687	4.58	-0.21	0.10
AGB	FGJ000061	4714	1.689	1.676	4.69	-0.10	0.10
AGB	FGJ000065	4677	1.540	1.724	4.91	0.12	0.14
AGB	FGJ000075	4763	1.764	1.652	4.55	-0.24	0.07
AGB	FGJ000076	4881	1.850	1.624	4.74	-0.05	0.05
AGB	FGJ000078	4868	1.855	1.623	4.82	0.03	0.07
AGB	FGJ000080	4829	1.843	1.627	4.67	-0.12	0.08
AGB	FGJ000083	4825	1.849	1.625	4.66	-0.13	0.08
AGB	FGJ000089	4861	1.868	1.618	4.80	0.01	0.04
AGB	FGJ000094	4925	1.937	1.596	4.75	-0.04	0.10
AGB	FGJ000097	4946	1.978	1.583	4.74	-0.05	0.05
AGB	FGJ000104	4874	1.907	1.606	4.56	-0.23	0.08
AGB	FGJ201620	4864	1.938	1.596	4.74	-0.05	0.11
RGB	FGJ000012	4270	1.062	1.878	5.06	0.27	0.12
RGB	FGJ000023	4360	1.181	1.840	5.18	0.39	0.12
RGB	FGJ000027	4425	1.290	1.805	4.75	-0.04	0.11
RGB	FGJ000029	4298	1.102	1.865	4.67	-0.12	0.12
RGB	FGJ000030	4294	1.070	1.876	5.13	0.34	0.10
RGB	FGJ000035	4439	1.353	1.784	5.41	0.62	0.10
RGB	FGJ000043	4443	1.359	1.782	5.49	0.70	0.10
RGB	FGJ000050	4404	1.267	1.812	5.02	0.23	0.13
RGB	FGJ000054	4496	1.487	1.741	4.92	0.13	0.12
RGB	FGJ000064	4436	1.353	1.784	5.42	0.63	0.12
RGB	FGJ000069	4583	1.587	1.709	5.34	0.55	0.12
RGB	FGJ000091	4665	1.776	1.648	5.12	0.33	0.11
RGB	FGJ000092	4612	1.711	1.669	4.73	-0.06	0.10
RGB	FGJ000107	4662	1.822	1.633	5.01	0.22	0.11
RGB	FGJ000129	4717	1.939	1.596	5.03	0.24	0.09
RGB	FGJ000155	4726	1.992	1.579	4.62	-0.17	0.07
RGB	FGJ000161	4775	2.052	1.559	5.17	0.38	0.05
RGB	FGJ000170	4794	2.083	1.549	5.33	0.54	0.12
RGB	FGJ000186	4800	2.117	1.538	4.70	-0.09	0.10
RGB	FGJ000193	4806	2.134	1.533	4.55	-0.24	0.04
RGB	FGJ000217	4813	2.161	1.524	5.22	0.43	0.05
RGB	FGJ000262	4855	2.252	1.495	5.13	0.34	0.09
RGB	FGJ000276	4858	2.260	1.492	5.15	0.36	0.11
RGB	FGJ200619	4760	1.940	1.595	5.43	0.64	0.11

The evolutionary status of each star is indicated in column 1. ID codes are designations of the current study. T_{eff} , log (g), and ξ are the surface temperature, gravity, and microturbulence values used in the abundance determinations. log (N_{*na*}) and [Na/Fe] are the final Na abundances. The final column shows the internal errors in [Na/Fe].

Discussion

Relationship between horizontal branch morphology and stellar composition

It has long been speculated that the composition differences between the first generation and second generation populations could have an effect on the colour-magnitude diagram structure of GCs⁹. Recent work has begun to provide some evidence to support this. For example, a new study on M4, which has both a red horizontal branch and a blue horizontal branch, has shown that all red horizontal branch stars in their sample are Na-poor, whilst all their blue horizontal branch stars are Na-rich²⁹. They infer that the He content must be different between the two Na populations since it is not expected that Na (or N) could affect the position of stars in the horizontal branch, while He can³⁰ (see Supplementary Discussion S3 below). As mentioned in the main text, a sample of horizontal branch stars from the redder end of the horizontal branch of NGC 6752 (it only has a blue horizontal branch) was shown to exclusively contain Na-poor stars¹⁷ (that study also reports one star with elevated Na abundance, but, as noted by the authors, the star has evolved off the horizontal branch and probably started from a much bluer position). The same stars have a uniform He abundance that is consistent with Big Bang theory predictions (Y = 0.245), as expected for a first generation population. Thus it appears that the bluer (presumably Na-rich) horizontal branch stars must avoid AGB ascent – leaving only the redder, Na-poor horizontal branch stars to populate the AGB of NGC 6752.

As also mentioned in the main text, if we combine this information with our estimate of the proportion of stars that do not ascend the AGB, we can estimate what horizontal branch colour delineates the border between the two groups. Since our *uvby* colour-magnitude diagram dataset is not complete at the bluest end of the horizontal branch, we obtained a very high quality UBVphotometric dataset¹⁹ for this purpose, courtesy of Dr. Yazan Momany at the European Southern Observatory (Chile). We counted horizontal branch stars in the U, U - V plane, starting at the red edge of the horizontal branch at U - V = 0.25. The total number of horizontal branch stars was found to be 320. Thus we expect the reddest 96 stars (30%) to eventually ascend the AGB. We find that this number of stars corresponds to an 'ascension cut-off' in U-V of -0.30. Interestingly this is exactly the colour for the Grundahl jump, a discontinuity in horizontal branch morphology which is seen in all GCs studied to date whose horizontal branch extends beyond $T_{eff} \approx 11,500 \text{ K}^{18,19}$. Explanations of this discontinuity include radiative levitation of elements heavier than carbon and nitrogen in the high-temperature atmospheres of these stars¹⁸, or the combination of post-zero-age horizontal branch evolution and diffusion effects¹⁹. At face value, it appears that all stars bluer than the Grundahl jump do not ascend the AGB, at least in NGC 6752. This may represent further evidence that there is some fundamental change in the stellar atmosphere structure and/or massloss physics occurring at the Grundahl jump temperature¹⁸. We note that it is these extremely blue horizontal branch stars that are considered to be the source of excess UV flux in the spectra of elliptical galaxies³¹ as well as Galactic and extra-galactic GCs³².

Stellar model experiments

Theoretical models show that for a given metallicity and core mass at the tip of the red giant branch the position of a star along the horizontal branch is determined by the mass of the hydrogen-rich envelope. The lower the envelope mass, the bluer the star will be. If the envelope mass is extremely low ($\leq 10^{-2} M_{\odot}$)³³, a horizontal branch star will not ascend the AGB. These stars instead evolve directly to the WD cooling track. A horizontal branch star can have such a low envelope mass if it suffered extra mass-loss during the preceding red giant branch phase, or if it formed with an elevated helium abundance. In the latter case the higher He affects the evolution of the star such that it arrives on the horizontal branch with a lower total mass (for a given GC age). In the former case the mechanisms that might affect the mass loss rates are unknown, although rotation is a possibility. In both cases the stars populate the blue end of the horizontal branch³⁰.

Our stellar model experiments (see Figure 3 in main text) are based on standard assumptions for mass-loss rates, cluster age, and a possible helium variation between the two generations of $\delta Y = 0.04$ (although we note a very recent study has just reported stronger constraints on Y)³⁴. Different assumptions of mass-loss rate or cluster age could produce bluer (or redder) horizontal branch morphologies. Our simulations show that standard models cannot reproduce the observations and thus non-standard/improved models are needed. We are currently working on improved horizontal branch models and expect that other groups will also investigate the very high AGB failure rate phenomenon reported in this Letter. Finally we note that our rough test of enhanced mass-loss on the horizontal branch (Figure 3 in main text) resulted in a quite different evolution in the CMD – the star with a 20-fold increase in mass-loss evolved 'downwards' (towards lower luminosities), practically along the zero-age horizontal branch line, then spent a significant amount of time at higher luminosities after leaving the zero-age line. This different evolution should be taken into account in future theoretical and observational investigations. We thank a thoughtful referee for inspiring this final paragraph of discussion.